Search for neutrinoless τ decays involving the K_S^0 meson

K.K. Gan
a $\,^*$

^aThe Ohio State University, Columbus, OH 43210, U.S.A.

We have searched for lepton flavor violating decays of the τ lepton with at least one K_S^0 meson in the final state. The data used in the search were collected with the CLEO II and II.V detectors at the Cornell Electron Storage Ring (CESR) and correspond to an integrated luminosity of 13.9 fb^{-1} at the $\Upsilon(4S)$ resonance. No evidence for signals were found, therefore we have set 90% confidence level (C.L.) upper limits on the branching fractions $\mathcal{B}(\tau^- \to e^- K_S^0) < 9.5 \times 10^{-7}$, $\mathcal{B}(\tau^- \to \mu^- K_S^0) < 9.5 \times 10^{-7}$, $\mathcal{B}(\tau^- \to \mu^- K_S^0) < 9.5 \times 10^{-7}$, $\mathcal{B}(\tau^- \to e^- K_S^0 K_S^0) < 2.4 \times 10^{-6}$, and $\mathcal{B}(\tau^- \to \mu^- K_S^0 K_S^0) < 3.5 \times 10^{-6}$. These represent significantly improved upper limits on the two-body decays and first upper limits on the three-body decays.

1. INTRODUCTION

In physics, all fundamental conservation laws are expected to have associated symmetries. Lepton flavor conservation, however, is an experimentally observed phenomena with no associated symmetry in the Standard Model. Lepton flavor violation (LFV) is expected in many extensions of the Standard Model such as lepto-quark, supersymmetry, superstring, left-right symmetric models, and models that include heavy neutral leptons [1,2]. Experimentally, both Super Kamiokande [3] and SNO [4] observe neutrino oscillation, which may imply LFV in the neutrino sector, therefore LFV is expected to occur in charged lepton decay at some branching fraction, albeit very small. The τ lepton provides a clean laboratory for such searches. Ilakovac [5] has calculated upper limits on branching fractions for many neutrinoless LFV modes within a model involving heavy Dirac neutrinos. The branching fractions depend on the heavy neutrino masses and mixings. For the decays $\tau^- \to \ell^- K_S^0$ [6], the branching fractions are of $\mathcal{O}(10^{-16})$, where ℓ can be e or μ . For the decays $\tau^- \to \ell^- K^0_S K^0_S$ or $\tau^- \to \ell^- K^+ K^-$, the branching fractions are of $\mathcal{O}(10^{-7})$. The decays with two kaons in the final state are therefore of particular experimental interest. Previous published upper limits on the branching fractions for the decays

 $\tau^- \to \ell^- K_S^0$ are of $\mathcal{O}(10^{-4})$ [7]. There are no previous results for the decays $\tau^- \to \ell^- K_S^0 K_S^0$. In this paper, we present the result of a search for the decays into one lepton and one or two K_S^0 mesons, with the K_S^0 decaying into two charged pions.

The data used in this analysis were collected using the CLEO detector [8,9] from e^+e^- collisions at the Cornell Electron Storage Ring (CESR) at a center-of-mass energy $\sqrt{s} \sim 10.6$ GeV. The total integrated luminosity of the data sample is 13.9 fb^{-1} corresponding to the production of $N_{\tau\tau} = 1.27 \times 10^7 \tau^+\tau^-$ events. The CLEO detector is a general purpose spectrometer with excellent charged particle and shower energy detection.

2. EVENT SELECTION CRITERIA

The $\tau^+\tau^-$ candidate events must contain four or six charged tracks with zero net charge. To reject beam-gas events, the distance of closest approach of each non- K_S^0 track to the IP must be within 0.5 cm transverse to the beam and 5 cm along the beam direction. The polar angle θ of each track with respect to the beam must satisfy $|\cos \theta| < 0.90$. Photons are defined as energy clusters in the calorimeter with at least 60 MeV in the barrel ($|\cos \theta| < 0.80$) or 100 MeV in the endcap ($0.80 < |\cos \theta| < 0.95$). We further require every photon to be separated from

^{*}Representing the CLEO Collaboration.

the projection of any charged track by at least 30 cm unless its energy is greater than 300 MeV. In order to diminish QED background such as radiative Bhabha and μ -pair events with photon conversion, we require each event to have total energy less than 95% of the center-of-mass energy. This requirement rejects most of the QED background, while incurring a small loss in detection efficiency.

We divide each event into two hemispheres (signal and tag), one containing one charged track and the other containing three or five charged tracks, using the plane perpendicular to the thrust axis [10]. The thrust axis is calculated from both charged tracks and photons. The invariant mass of the tag hemisphere must be less than the τ mass, $M_{\tau} = 1.777 \text{ GeV}/c^2$ [11]. The signal hemisphere must contain an electron or a muon and one or two K_S^0 mesons. The electron candidate must have shower energy to momentum ratio in the range, 0.85 < E/p < 1.10, and when available, the specific ionization lost must be consistent with an electron. The muon candidate must penetrate at least three absorbtion lengths of iron. The K_S^0 candidate is reconstructed in the π^+ π^- final state with a detached vertex, and the invariant mass must be within approximately three standard deviations of the nominal mass, $485 < m_{\pi\pi} < 510 \text{ MeV}/c^2$. In order to diminish radiative Bhabha and μ -pair events further, neither pion should be consistent with identification as an electron.

Since there is no neutrino in the signal hemisphere while there is at least one neutrino undetected in the tag hemisphere, the missing momentum of the event must point toward the tag hemisphere, $0 < \cos \theta_{tag-missing} < 1.0$. In order to suppress the background from radiative Bhabha and μ -pair events, the direction of the missing momentum of the event is required to satisfy $|\cos \theta_{missing}| < 0.90$. For the decay $\tau^- \to e^- K_S^0$, $\cos \theta_{tag-missing}$ is further restricted to be less than 0.99 to reduce the radiative backgrounds. This corresponds to the minimum ratio of background to detection efficiency. The background is estimated from the sidebands in the invariant mass vs. total energy distribution of the decay candidates.

To search for decay candidates, we select τ candidates with invariant mass and total energy consistent with the expectations. The following kinematic variables are used to select the candidate events:

$$\Delta E = E - E_{beam}$$
$$\Delta M = M - M_{\tau}$$

where E_{beam} is the beam energy, and E and M are the reconstructed τ energy and mass. The ΔE vs. ΔM distributions of the decay candidates in the data and signal Monte Carlo samples are shown in Figs. 1 and 2.

In the Monte Carlo simulation, one τ lepton decays according to two- or three-body phase space for the mode of interest, and the other τ lepton decays generically according to the $KORALB/TAUOLA \ \tau$ event generator [12]. The phase space model is appropriate for an unpolarized tau. If the Lorentz structure of the neutrinoless decay is V-A (Standard Model-like), correlations between the spins of the two τ 's in the event will lead to slightly higher detection efficiency than phase space, while V+A decays will lead to lower detection efficiency. The detector response is simulated using the *GEANT* program [13]. The estimated detection efficiencies (ϵ) [14] are summarized in Table 1.

Table 1

Summary of detection efficiency (with statistical error), 90% C.L. upper limits on the branching fraction with and without including systematic errors.

Mode	$\epsilon~(\%)$	$\mathcal{B}(10^{-7})$	$\mathcal{B}(10^{-7})$
- 720	100104	(stat.)	0 5
$e^{-}K_{S}^{0}$	18.8 ± 0.4	8.8	9.5
$\mu \kappa_{\tilde{S}}$ $e^{-K^0 K^0}$	19.0 ± 0.4 12.0 ± 0.1	0.7 91	9.0 94
$\mu^- K^0_G K^0_G$	12.0 ± 0.1 8.0 ± 0.1	$\frac{21}{31}$	$\frac{24}{35}$





Figure 1. ΔE vs. ΔM distribution of the (a) data and (b) signal Monte Carlo sample for the decay $\tau^- \rightarrow e^- K_S^0$; (c) and (d) show the corresponding distributions for $\tau^- \rightarrow \mu^- K_S^0$. The normalization of the signal Monte Carlo sample is arbitrary. The ellipses indicate the signal region (see text).

Figure 2. ΔE vs. ΔM distribution of the (a) data and (b) signal Monte Carlo sample for the decay $\tau^- \rightarrow e^- K_S^0 K_S^0$; (c) and (d) show the corresponding distributions for $\tau^- \rightarrow \mu^- K_S^0 K_S^0$. The normalization of the signal Monte Carlo sample is arbitrary. The ellipses indicate the signal region (see text).

3. RESULTS

The upper limit on the branching fraction is related to the upper limit λ on the number of signal events by

$$\mathcal{B} = \frac{\lambda}{2\epsilon N_{\tau\tau} \mathcal{B}_1 (\mathcal{B}_{K_S^0 \to \pi^+ \pi^-})^n}$$

where $\mathcal{B}_1 = (84.71 \pm 0.13)\%$ is the inclusive 1-prong branching fraction [11], $\mathcal{B}_{K_S^0 \to \pi^+\pi^-} = (68.61 \pm 0.28)\%$ is the branching fraction for K_S^0 to decay to two charged pions, and n is the number of K_S^0 mesons in the final state. No candidate decays are observed, so we take λ as 2.44 events in each mode at 90% confidence level, according to the frequentist method [15]. The upper limits on the branching fractions with and without systematic errors are shown in Table 1.

The systematic errors include the uncertainties in the $\tau^+\tau^-$ cross section (1%), luminosity (1%), track reconstruction efficiency (1% per charged track), K_S^0 detection efficiency (2% per K_S^0), lepton identification (1.5% for electron and 4% for muon), and the statistical uncertainties in the detection efficiencies due to limited Monte Carlo samples (1-2%).

Black *et al.* [16] have analyzed the constraints on the new physics scale for dimensionsix fermionic effective operators involving tau-mu mixing, motivated by the observed $\nu_{\mu} - \nu_{\tau}$ oscillation. The most stringent lower limits from exotic heavy quarks and tau decays on the physics scale of the operators involving quarks are ~10 TeV. The new upper limit on $\mathcal{B}(\tau^- \to \mu^- K_S^0)$ presented in this paper yields a lower limit of 17.3 and 18.2 TeV for the axial vector and pseudoscalar operators, respectively.

In conclusion, we have searched for τ decays involving K_S^0 mesons that violate lepton flavor, but find no evidence for a signal. This results in improved upper limits for the decays $\tau^- \to \ell^- K_S^0$ and first upper-limits for the decays $\tau^- \to \ell^- K_S^0 K_S^0$. The upper limits for the $\tau^- \to \ell^- K_S^0 K_S^0$ final states are more stringent than those found previously for $\ell^- K^+ K^-$ [17].

This work was supported in part by the U.S. Department of Energy.

REFERENCES

- M.C. Gonzalez-Garcia and J.W.F. Valle, Mod. Phys. Lett. A 7, 477 (1992); A 9, 2569 (E) (1994).
- 2. J.E. Kim et al., Phys. Rev. D 56, 100 (1997).
- Super Kamiokande Collaboration, Y. Fukuda et al., Phys. Rev. Lett. 81, 1562 (1998).
- SNO Collaboration, Q.R. Ahmad *et al.*, Phys. Rev. Lett. 87, 071301 (2001).
- A. Ilakovac *et al.*, Phys. Rev. D **52**, 3993 (1995); Phys. Rev. D **62**, 036010 (2000).
- 6. Throughout this paper, the charge conjugate state is implied.
- MARK II Collaboration, K.G. Hayes *et al.*, Phys. Rev. D 25, 2869 (1982).
- Y. Kubota *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **320**, 66 (1992).
- T. Hill, Nucl. Instrum. Methods Phys. Res., Sect. A 418, 32 (1998).
- 10. E. Farhi, Phys. Rev. Lett. **39**, 1587 (1977).
- Particle Data Group, J. Bartels *et al.*, Eur. Phys. J. C **15**, 1 (2000).
- S. Jadach and Z. Was, Comput. Phys. Commun. **36**, 191 (1985); **64**, 267 (1991);
 S. Jadach, J.H. Kuhn, and Z. Was, *ibid.* **64**, 275 (1991).
- R. Brun *et al.*, CERN Report No. CERN-DD/EE/84-1, 1987 (unpublished).
- 14. The detection efficiencies for $\tau^- \rightarrow \mu^- K_S^0 K_S^0$ depends on the mass of the K_S^0 -pair produced. The detection efficiency is approximately constant up to $M_{K_S^0 K_S^0} \sim 1.25 \text{ GeV}/c^2$, falling to zero near the kinematic limit, $M_{K_S^0 K_S^0} = M_{\tau}$. The kinematic limit corresponds to the muon being produced at rest in the center-ofmass frame of the τ -lepton. In the laboratory frame, the muon has low momentum, hence would not be able to penetrate enough material to be classified as a muon.
- G.J. Feldman and R.D. Cousins, Phys. Rev. D 57, 3873 (1998).
- D. Black, T. Han, H.J. He and M. Sher, Phys. Rev. D 66, 053002 (2002).
- CLEO Collaboration, D.W. Bliss *et al.*, Phys. Rev. D 57, 5903 (1998).