

## RELEVANT RESULTS FROM THE NA48 EXPERIMENT

E. Imbergamo

*Department of Physics, University of Perugia, Italy*

M. Piccini

*Department of Physics, University of Perugia, Italy*

M.C. Petrucci

*INFN, section of Perugia, Italy*

### ABSTRACT

We report relevant results from NA48 experiment at CERN SPS. NA48 was proposed in 1990 [1] to study direct CP violation in  $K^0 \rightarrow \pi\pi$  to a level of accuracy sufficient to resolve the inconclusive status left by the previous measurements performed by NA31 [2] and E731 [3]. In 2002 NA48 published the final result [4]. Small modification to the experimental setup have allowed NA48 to go forward with an extensive investigation of  $K^0$  rare decays and hyperon decays. Some results are already available and reported here together with the final CP violation measurement.

### 1 The measurement of $Re(\epsilon'/\epsilon)$

It is well known that the decay of  $K^0/\overline{K^0}$  into two pions violates CP. Such a violation can have two contributions: one, indirect, associated to  $K^0/\overline{K^0}$  mixing and another, direct, coming from the decay amplitude. The amount of direct CP violation in this decay is parametrized by the parameter  $Re(\epsilon'/\epsilon)$ , which can be computed in the framework of the standard electro-weak-model, albeit with large theoretical uncertainties. Typical theoretical predictions of  $Re(\epsilon'/\epsilon)$  varies from few  $10^{-4}$  to about  $2 \times 10^{-3}$ , even though with large exceptions.

$Re(\epsilon'/\epsilon)$  is connected to the double ratio of decay rates according to the following formula:

$$R = \frac{\Gamma(K_L \rightarrow \pi^0\pi^0)}{\Gamma(K_S \rightarrow \pi^0\pi^0)} / \frac{\Gamma(K_L \rightarrow \pi^+\pi^-)}{\Gamma(K_S \rightarrow \pi^+\pi^-)} \simeq 1 - 6 \cdot Re(\epsilon'/\epsilon) \quad (1)$$




Figure 1: *World measurements of  $Re(\epsilon'/\epsilon)$*

In order to exploit the cancellation in the double ratio of systematic uncertainties, the experimental apparatus consists of two concurrent and almost co-linear beams, one providing the experiment with  $K_L$  decays and the other one with  $K_S$  decays and the  $K^0$  to  $\pi\pi$  decays are reconstructed in the same decay region. A tagging station is devoted to identify the  $K^0$  decay as a  $K_S$  or a  $K_L$  decay. In 2002 NA48 has published [4] the final result of the measurement:

$$Re(\epsilon'/\epsilon) = (15.3 \pm 2.6) \times 10^{-4} \quad (2)$$

It is the most precise measurement ever done as shown in Fig.1. The world average value is also reported together with the uncertainty (yellow band).

## 2 Selected items about rare decays

The radiative kaon decays are ideally suited to test the validity of the Chiral Perturbation Theory ( $\chi PhT$ ). The processes may be described in a perturbative expansion of momenta:  $\mathcal{O}(p^2)$ ,  $\mathcal{O}(p^4)$ . Examples of this type are  $K_S \rightarrow \gamma\gamma$  and  $K_L \rightarrow \pi^0\gamma\gamma$ . In both cases there is no contribution from the  $\mathcal{O}(p^2)$  term, while the  $\mathcal{O}(p^4)$  contribution is predicted to better than 5% by  $\chi PhT$ . NA48 has measured [5],[6]:

$$BR(K_S \rightarrow \gamma\gamma) = (2.78 \pm 0.06_{stat} \pm 0.02_{MCstat} \pm 0.04_{syst}) \times 10^{-6} \quad (3)$$

$$BR(K_L \rightarrow \pi^0 \gamma \gamma) = (1.36 \pm 0.03_{stat} \pm 0.03_{syst} \pm 0.03_{norm}) \times 10^{-6} \quad (4)$$

The value of  $BR(K_S \rightarrow \gamma \gamma)$  deviates from  $\mathcal{O}(p^4)$  prediction and indicates a large  $\mathcal{O}(p^6)$  contribution. The  $\mathcal{O}(p^4)$  contribution to  $K_L \rightarrow \pi^0 \gamma \gamma$  turns out to be also an underestimation of the decay rate. Anyway at  $\mathcal{O}(p^6)$  the rate may be reproduced by adding a contribution from the VDM mechanism, via the coupling constant  $a_v$  that NA48 has measured to be:

$$a_v = -0.46 \pm 0.03_{stat} \pm 0.04_{syst} \quad (5)$$

### 3 Hyperon decays in NA48

The target used for the production of  $K^0$ s is also a huge source of hyperons. By using the small fraction of hyperons that passed the standard triggers in the previous years, the NA48 collaboration has already published results on hyperon physics [7]:

$$m(\Xi^0) = [1314.82 \pm 0.06(stat.) \pm 0.20(syst.)] MeV/c^2 \quad (6)$$

$$BR(\Xi^0 \rightarrow \Lambda \gamma) = [1.90 \pm 0.34(stat.) \pm 0.19(syst.)] \times 10^{-3} \quad (7)$$

$$BR(\Xi^0 \rightarrow \Sigma^0 \gamma) = [3.14 \pm 0.76(stat.) \pm 0.32(syst.)] \times 10^{-3} \quad (8)$$

In the 2002 special triggers have been dedicated to hyperon decays and NA48 claims main achievements to:

1. study form factors and flavor symmetry violations in the  $\Xi^0$  decays;
2. give an alternative measurement of  $V_{us}$  (CKM parameter) using  $\Xi^0$  (and  $\Lambda$ ) beta decay instead of kaon beta decays.

### References

1. NA48 Collaboration (A. Lay *et al.*) CERN-SPSC-90-22.
2. NA48 Collaboration (G.D. Barr *et al.*) Phys. Lett. B **317**, 233-242 (1993).
3. NA48 Collaboration (L.K. Gibbons *et al.*) Phys. Rev. D **55**, 6625-6715 (1997).
4. NA48 Collaboration (J.R. Batley *et al.*) Phys. Lett. B **544**, 97-112 (2002).
5. NA48 Collaboration (A. Lay *et al.*) Phys. Lett. B **551**, 7-15 (2003).
6. NA48 Collaboration (A. Lay *et al.*) CERN-EP/2002-030, hep-ex/0205010.
7. NA48 Collaboration (A. Lay *et al.*) Eur. Phys. J. C **12**, 69-76 (2000).