

A Measure of the Goodness of Fit in Unbinned Likelihood Fits; End of Bayesianism?

Rajendran Raja
 Fermilab, Batavia, IL 60510, USA

Maximum likelihood fits to data can be done using binned data (histograms) and unbinned data. With binned data, one gets not only the fitted parameters but also a measure of the goodness of fit. With unbinned data, currently, the fitted parameters are obtained but no measure of goodness of fit is available. This remains, to date, an unsolved problem in statistics. Using Bayes' theorem and likelihood ratios, we provide a method by which both the fitted quantities and a measure of the goodness of fit are obtained for unbinned likelihood fits, as well as errors in the fitted quantities. The quantity, conventionally interpreted as a Bayesian prior, is seen in this scheme to be a number not a distribution, that is determined from data.

1. INTRODUCTION

As of the Durham conference [1], the problem of obtaining a goodness of fit in unbinned likelihood fits was an unsolved one. In what follows, we will denote by the vector s , the theoretical parameters (s for "signal") and the vector c , the experimentally measured quantities or "configurations". For simplicity, we will illustrate the method where both s and c are one dimensional, though either or both can be multi-dimensional in practice. We thus define the theoretical model by the conditional probability density $P(c|s)$. Then an unbinned maximum likelihood fit to data is obtained by maximizing the likelihood [2],

$$\mathcal{L} = \prod_{i=1}^{i=n} P(c_i|s) \quad (1)$$

where the likelihood is evaluated at the n observed data points $c_i, i = 1, n$. Such a fit will determine the maximum likelihood value s^* of the theoretical parameters, but will not tell us how good the fit is. The value of the likelihood \mathcal{L} at the maximum likelihood point does not furnish a goodness of fit, since the likelihood is not invariant under change of variable. This can be seen by observing that one can transform the variable set c to a variable set c' such that $P(c'|s^*)$ is uniformly distributed between 0 and 1. Such a transformation is known as a hypercube transformation, in multi-dimensions. Other datasets will yield different values of likelihood in the variable space c when the likelihood is computed with the original function $P(c|s^*)$. However, in the original hypercube space, the value of the likelihood is unity regardless of the dataset $c'_i, i = 1, n$, thus the likelihood \mathcal{L} cannot furnish a goodness of fit by itself, since neither the likelihood, nor ratios of likelihoods computed using the same distribution $P(c|s^*)$ is invariant under variable transformations. The fundamental reason for this non-invariance is that only a single distribution, namely, $P(c|s^*)$ is being used to compute the goodness of fit.

2. LIKELIHOOD RATIOS

In binned likelihood cases, where one is comparing a theoretical distribution $P(c|s)$ with a binned histogram, there are two distributions involved, the theoretical distribution and the data distribution. The *pdf* of the data is approximated by the bin contents of the histogram normalized to unity. If the data consists of n events, the *pdf* of the data $P^{data}(c)$ is defined in the frequentist sense as the normalized density distribution in c space of n events as $n \rightarrow \infty$. In the binned case, we can bin in finer and finer bins as $n \rightarrow \infty$ and obtain a smooth function, which we define as the *pdf* of the data $P^{data}(c)$. In practice, one is always limited by statistics and the binned function will be an approximation to the true *pdf*. We can now define a likelihood ratio $\mathcal{L}_{\mathcal{R}}$ such that

$$\mathcal{L}_{\mathcal{R}} = \frac{\prod_{i=1}^{i=n} P(c_i|s)}{\prod_{i=1}^{i=n} P^{data}(c_i)} \equiv \frac{P(\mathbf{c}_n|s)}{P^{data}(\mathbf{c}_n)} \quad (2)$$

where we have used the notation \mathbf{c}_n to denote the event set $c_i, i = 1, n$. Let us now note that $\mathcal{L}_{\mathcal{R}}$ is invariant under the variable transformation $c \rightarrow c'$, since

$$P(c'|s) = \left| \frac{dc}{dc'} \right| P(c|s) \quad (3)$$

$$P^{data}(c') = \left| \frac{dc}{dc'} \right| P^{data}(c) \quad (4)$$

$$\mathcal{L}'_{\mathcal{R}} = \mathcal{L}_{\mathcal{R}} \quad (5)$$

and the Jacobian of the transformation $\left| \frac{dc}{dc'} \right|$ cancels in the numerator and denominator in the ratio. This is an extremely important property of the likelihood ratio $\mathcal{L}_{\mathcal{R}}$ that qualifies it to be a goodness of fit variable. Since the denominator $P^{data}(\mathbf{c}_n)$ is independent of the theoretical parameters s , both the likelihood ratio and the likelihood maximize at the same point s^* . One can also show [3] that the maximum value of the likelihood ratio occurs when the theoretical likelihood $P(c_i|s)$ and the data likelihood $P^{data}(c_i)$ are equal for all c_i .

3. BINNED GOODNESS OF FIT

In the case where the *pdf* $P^{data}(c)$ is estimated by binned histograms and the statistics are Gaussian, it is readily shown [3] that the commonly used goodness of fit variable $\chi^2 = -2\log\mathcal{L}_{\mathcal{R}}$. It is worth emphasizing that the likelihood ratio as defined above is needed and not just the negative log of theoretical likelihood $P(\mathbf{c}_n|s)$ to derive this result. The popular conception that χ^2 is $-2\log P(\mathbf{c}_n|s)$ is simply incorrect!. It can also be shown that the likelihood ratio defined above can describe the binned cases where the statistics are Poissonian [4]. In order to solve our problem of goodness of fit in unbinned likelihood cases, one needs to arrive at a method of estimating the data *pdf* $P^{data}(c)$ without the use of bins.

4. UNBINNED GOODNESS OF FIT

One of the better known methods of estimating the probability density of a distribution in an unbinned case is by the use of Probability Density Estimators (*PDE's*), also known as Kernel Density Estimators [*KDE's*]. The *pdf* $P^{data}(c)$ is approximated by

$$P^{data}(c) \approx PDE(c) = \frac{1}{n} \sum_{i=1}^{i=n} \mathcal{G}(c - c_i) \quad (6)$$

where a Kernel function $\mathcal{G}(c - c_i)$ is centered around each data point c_i , is so defined that it normalizes to unity and for large n approaches a Dirac delta function [3]. The choice of the Kernel function can vary depending on the problem. A popular kernel is the Gaussian defined in the multi-dimensional case as

$$\mathcal{G}(c) = \frac{1}{(\sqrt{2\pi}h)^d \sqrt{\det(E)}} \exp\left(\frac{-H^{\alpha\beta} c^\alpha c^\beta}{2h^2}\right) \quad (7)$$

where E is the error matrix of the data defined as

$$E^{\alpha,\beta} = \langle c^\alpha c^\beta \rangle - \langle c^\alpha \rangle \langle c^\beta \rangle \quad (8)$$

and the $\langle \rangle$ implies average over the n events, and d is the number of dimensions. The Hessian matrix H is defined as the inverse of E and the repeated indices imply summing over. The parameter h is a ‘‘smoothing parameter’’, which has[6] a suggested optimal value $h \propto n^{-1/(d+4)}$, that satisfies the asymptotic condition

$$\mathcal{G}_\infty(c - c_i) \equiv \lim_{n \rightarrow \infty} \mathcal{G}(c - c_i) = \delta(c - c_i) \quad (9)$$

The parameter h will depend on the local number density and will have to be adjusted as a function of the local density to obtain good representation of the data by the *PDE*. Our proposal for the goodness of fit in unbinned likelihood fits is thus the likelihood ratio

$$\mathcal{L}_{\mathcal{R}} = \frac{P(\mathbf{c}_n|s)}{P^{data}(\mathbf{c}_n)} \approx \frac{P(\mathbf{c}_n|s)}{P^{PDE}(\mathbf{c}_n)} \quad (10)$$

evaluated at the maximum likelihood point s^* .

5. AN ILLUSTRATIVE EXAMPLE

We consider a simple one-dimensional case where the data is an exponential distribution, say decay times of a radioactive isotope. The theoretical prediction is given by

$$P(c|s) = \frac{1}{s} \exp\left(-\frac{c}{s}\right) \quad (11)$$

We have chosen an exponential with $s = 1.0$ for this example. The Gaussian Kernel for the *PDE* would be given by

$$\mathcal{G}(c) = \frac{1}{(\sqrt{2\pi}\sigma h)} \exp\left(-\frac{c^2}{2\sigma^2 h^2}\right) \quad (12)$$

where the variance σ of the exponential is numerically equal to s . To begin with, we chose a constant value for the smoothing parameter, which for 1000 events generated is calculated to be 0.125. Figure 1 shows the generated events, the theoretical curve $P(c|s)$ and the *PDE* curve $P(c)$ normalized to the number of events. The *PDE* fails to reproduce the data near the origin due to the boundary effect, whereby the Gaussian probabilities for events close to the origin spill over to negative values of c . This lost probability would be compensated by events on the exponential distribution with negative c if they existed. In our case, this presents a drawback for the *PDE* method, which we will remedy later in the paper using *PDE* definitions on the hypercube and periodic boundary conditions. For the time being, we will confine our example to values of $c > 1.0$ to avoid the boundary effect.

In order to test the goodness of fit capabilities of the likelihood ratio $\mathcal{L}_{\mathcal{R}}$, we superimpose a Gaussian on the exponential and try and fit the data by a simple exponential. Figure 2 shows the ‘‘data’’ with 1000 events generated as an exponential in the fiducial range $1.0 < c < 5.0$. Superimposed on it is a Gaussian of 500 events. More events in the exponential are generated in the interval $0.0 < c < 1.0$ to avoid the boundary effect at the fiducial boundary at $c=1.0$. Since the number density varies significantly, we have had to introduce a method of iteratively determining the smoothing factor as a function of c as described in [3]. With this modification in the *PDE*, one gets a good description of the behavior of the data by the *PDE* as shown in Figure 2. We now vary the number of events in the Gaussian and obtain the value of the negative log likelihood ratio \mathcal{NLLR} as a function of the strength of the Gaussian. Table I summarizes the results. The number of standard deviations the unbinned likelihood fit is from what is expected is determined empirically by plotting the value of \mathcal{NLLR} for a large number of fits where no Gaussian is superimposed (i.e. the null hypothesis) and determining the mean and *RMS* of this distribution and using these




Figure 1: Figure shows the histogram (with errors) of generated events. Superimposed is the theoretical curve $P(c|s)$ and the PDE estimator (solid) histogram with no errors.




Figure 2: Figure shows the histogram (with errors) of 1000 events in the fiducial interval $1.0 < c < 5.0$ generated as an exponential with decay constant $s=1.0$ with a superimposed Gaussian of 500 events centered at $c=2.0$ and width=0.2. The PDE estimator is the (solid) histogram with no errors.

to estimate the number of σ 's the observed \mathcal{NLLR} is from the null case. Table I also gives the results of a binned fit on the same “data”. It can be seen that the unbinned fit gives a 3σ discrimination when the number of Gaussian events is 85, where as the binned fit gives a χ^2/ndf of 42/39 for the same case. We intend to make these tests more sophisticated in future work.

Figure 3 shows the variation of $-\log P(\mathbf{c}_n|s)$ and $-\log P^{PDE}(\mathbf{c}_n)$ for an ensemble of 500 experiments each with the number of events $n = 1000$ in the exponential and no events in the Gaussian (null hypothesis).

Table I Goodness of fit results from unbinned likelihood and binned likelihood fits for various data samples. The negative values for the number of standard deviations in some of the examples is due to statistical fluctuation.

Number of Gaussian events	Unbinned fit \mathcal{NLLR}	Unbinned fit $N\sigma$	Binned fit χ^2 39 d.o.f.
500	189.	103	304
250	58.6	31	125
100	11.6	4.9	48
85	8.2	3.0	42
75	6.3	1.9	38
50	2.55	-0.14	30
0	0.44	-1.33	24

It can be seen that $-\log P(\mathbf{c}_n|s)$ and $-\log P^{PDE}(\mathbf{c}_n)$ are correlated with each other and the difference between the two ($-\log \mathcal{NLLR}$) is a much narrower distribution than either and provides the goodness of fit discrimination.




Figure 3: (a) shows the distribution of the negative log-likelihood $-\log_e(P(\mathbf{c}_n|s))$ for an ensemble of experiments where data and experiment are expected to fit. (b) Shows the negative log PDE likelihood $-\log_e(P(\mathbf{c}_n))$ for the same data (c) Shows the correlation between the two and (d) Shows the negative log-likelihood ratio \mathcal{NLLR} that is obtained by subtracting (b) from (a) on an event by event basis.

5.1. Improving the PDE

The PDE technique we have used so far suffers from two drawbacks; firstly, the smoothing parameter has to be iteratively adjusted significantly over the full range of the variable c , since the distribution $P(c|s)$ changes significantly over that range; and secondly, there are boundary effects at $c=0$ as shown in figure 1. Both these flaws are remedied if we define the

PDE in hypercube space. After we find the maximum likelihood point s^* , for which the *PDE* is not needed, we transform the variable $c \rightarrow c'$, such that the distribution $P(c'|s^*)$ is flat and $0 < c' < 1$. The hypercube transformation can be made even if c is multi-dimensional by initially going to a set of variables that are uncorrelated and then making the hypercube transformation. The transformation can be such that any interval in c space maps on to the interval $(0, 1)$ in hypercube space. We solve the boundary problem by imposing periodicity in the hypercube. In the one dimensional case, we imagine three “hypercubes”, each identical to the other on the real axis in the intervals $(-1, 0)$, $(0, 1)$ and $(1, 2)$. The hypercube of interest is the one in the interval $(0, 1)$. When the probability from an event kernel leaks outside the boundary $(0, 1)$, we continue the kernel to the next hypercube. Since the hypercubes are identical, this implies the kernel re-appearing in the middle hypercube but from the opposite boundary. Put mathematically, the kernel is defined such that

$$\mathcal{G}(c' - c'_i) = \mathcal{G}(c' - c'_i - 1); c' > 1 \quad (13)$$

$$\mathcal{G}(c' - c'_i) = \mathcal{G}(c' - c'_i + 1); c' < 0 \quad (14)$$

Although a Gaussian Kernel will work on the hypercube, the natural kernel to use considering the shape of the hypercube would be the function $\mathcal{G}(c')$

$$\mathcal{G}(c') = \frac{1}{h}; |c'| < \frac{h}{2} \quad (15)$$

$$\mathcal{G}(c') = 0; |c'| > \frac{h}{2} \quad (16)$$

This kernel would be subject to the periodic boundary conditions given above, which further ensure that every event in hypercube space is treated exactly as every other event irrespective of their co-ordinates. The parameter h is a smoothing parameter which needs to be chosen with some care. However, since the theory distribution is flat in hypercube space, the smoothing parameter may not need to be iteratively determined over hypercube space to the extent that data distribution is similar to the theory distribution. Even if iteration is used, the variation in h in hypercube space is likely to be much smaller.

Figure 4 shows the distribution of the \mathcal{NLLR} for the null hypothesis for an ensemble of 500 experiments each with 1000 events as a function of the smoothing factor h . It can be seen that the distribution narrows considerably as the smoothing factor increases. We choose an operating value of 0.2 for h and study the dependence of the \mathcal{NLLR} as a function of the number of events ranging from 100 to 1000 events, as shown in figure 5. The dependence on the number of events is seen to be weak, indicating good behavior. The *PDE* thus arrived computed with $h=0.2$ can be transformed from the hypercube space to c space and will reproduce data smoothly and with no edge effects. We note




Figure 4: The distribution of the negative log likelihood ratio \mathcal{NLLR} for the null hypothesis for an ensemble of 500 experiments each with 1000 events, as a function of the smoothing factor $h=0.1, 0.2$ and 0.3

that it is also easier to arrive at an analytic theory of \mathcal{NLLR} with the choice of this simple kernel.




Figure 5: The distribution of the negative log likelihood ratio \mathcal{NLLR} for the null hypothesis for an ensemble of 500 experiments each with the smoothing factor $h=0.2$, as a function of the number of events

6. END OF BAYESIANISM?

By Bayesianism, we mean the practice of “guessing” a prior distribution and introducing it into the calculations. In what follows we will show that what is conventionally thought of as a Bayesian prior distribution is in reality a number that can be calculated from the data. We are able to do this since we use two *pdf*'s, one for theory and one for data. In what

follows, we will interpret the probability distribution of the parameter s in a strictly frequentist sense. The *pdf* of s is the distribution of the best estimator of the true value s_T of s from an ensemble of an infinite number of identical experiments with the same statistical power n .

6.1. Calculation of fitted errors

After the fitting is done and the goodness of fit is evaluated, one needs to work out the errors on the fitted quantities. One needs to calculate the posterior density $P(s|\mathbf{c}_n)$, which carries information not only about the maximum likelihood point s^* , from a single experiment, but how such a measurement is likely to fluctuate if we repeat the experiment. The joint probability density $P(s, \mathbf{c}_n)$ of observing the parameter s and the data \mathbf{c}_n is given by

$$P^{data}(s, \mathbf{c}_n) = P(s|\mathbf{c}_n)P^{data}(\mathbf{c}_n) \quad (17)$$

where we use the superscript *data* to distinguish the joint probability $P^{data}(s, \mathbf{c}_n)$ as having come from using the data *pdf*. If we now integrate the above equation over all possible datasets \mathbf{c}_n , we get the expression for the *pdf* of s .

$$\mathcal{P}_n(s) = \int P^{data}(s, \mathbf{c}_n)d\mathbf{c}_n = \int P(s|\mathbf{c}_n)P^{data}(\mathbf{c}_n)d\mathbf{c}_n \quad (18)$$

where we have used the symbol \mathcal{P} to distinguish the fact that it is the true *pdf* of s obtained from an infinite ensemble. We use the subscript n in $\mathcal{P}_n(s)$ to denote that the *pdf* is obtained from an ensemble of experiments with n events each. Later on we will show that $\mathcal{P}_n(s)$ is indeed dependent on n . Equation 18 states that in order to obtain the *pdf* of the parameter s , one needs to add together the conditional probabilities $P(s|\mathbf{c}_n)$ over an ensemble of events, each such distribution weighted by the “data likelihood” $P^{data}(\mathbf{c}_n)$. At this stage of the discussion, the functions $P^{data}(s|\mathbf{c}_n)$ are unknown functions. We have however worked out $\mathcal{L}_R(s)$ as a function of s and have evaluated the maximum likelihood value s^* of s . We can choose an arbitrary value of s and evaluate the goodness of fit at that value using the likelihood ratio. When we choose an arbitrary value of s , we are in fact hypothesizing that the true value s_T is at this value of s . $\mathcal{L}_R(s)$ then gives us a way of evaluating the relative goodness of fit of the hypothesis as we change s . Let us now take an arbitrary value of s and hypothesize that that is the true value. Then the joint probability of observing \mathbf{c}_n and s_T being at this value of s is given from the data end by equation 17.

Similarly, from the theoretical end, one can calculate the joint probability of observing the dataset \mathbf{c}_n , with the true value being at s . The true value s_T is

taken to be the maximum likelihood point of the *pdf* $\mathcal{P}_n(s)$. It may coincide with the mean value of the *pdf* $\mathcal{P}_n(s)$. These statements are assertions of the unbiased nature of the data from the experiment. At this point, there is no information available on where the true value s_T lies. One can make the hypothesis that a particular value of s is the true value and the probability of obtaining a best estimator s^* from experiments of the type being performed in the interval s_T and $s_T + ds_T$ is $\mathcal{P}_n(s_T)ds_T$. The actual value of this number is a function of the experimental resolution and the statistics n of the experiment. The joint probability $P^{theory}(s, \mathbf{c}_n)$ from the theoretical end is given by the product of the probability density of the *pdf* of s at the true value of s , namely $\mathcal{P}_n(s_T)$, and the theoretical likelihood $P(\mathbf{c}_n|s)$ evaluated at the true value, which by our hypothesis is s .

$$P^{theory}(s, \mathbf{c}_n) = P^{theory}(\mathbf{c}_n|s)\mathcal{P}_n(s_T) \quad (19)$$

The joint probability $P(s, \mathbf{c}_n)$ is a joint distribution of the theoretical parameter s and data \mathbf{c}_n . The two ways of evaluating this (from the theoretical end and the data end) must yield the same result, for consistency. This is equivalent to equating $P^{data}(s, \mathbf{c}_n)$ and $P^{theory}(s, \mathbf{c}_n)$. This gives the equation

$$P(s|\mathbf{c}_n)P^{data}(\mathbf{c}_n) = P^{theory}(\mathbf{c}_n|s)\mathcal{P}_n(s_T) \quad (20)$$

which is a form of Bayes’ theorem, but with two *pdf*’s (theory and data). Let us note that the above equation can be immediately re-written as a likelihood ratio

$$\mathcal{L}_R = \frac{P(s|\mathbf{c}_n)}{\mathcal{P}_n(s_T)} = \frac{P^{theory}(\mathbf{c}_n|s)}{P^{data}(\mathbf{c}_n)} \quad (21)$$

which is what is used to obtain the goodness of fit. In order to get the fitted errors, we need to evaluate $P(s|\mathbf{c}_n)$ which necessitates a better understanding of what $\mathcal{P}_n(s_T)$ is in equation 20. Rearranging equation 20, one gets

$$P(s|\mathbf{c}_n) = \mathcal{L}_R(s)\mathcal{P}_n(s_T) = \frac{P^{theory}(\mathbf{c}_n|s)}{P^{data}(\mathbf{c}_n)}\mathcal{P}_n(s_T) \quad (22)$$

6.1.1. To show that $\mathcal{P}_n(s_T)$ depends on n

In practice, in both the binned and unbinned cases, one only has an approximation to $P^{data}(\mathbf{c}_n)$. As $n \rightarrow \infty$, in the absence of experimental bias, one expects to determine the parameter set s to infinite accuracy; and $P(s|\mathbf{c}_n) \rightarrow \delta(s - s_T)$, where s_T is the true value of s . However, for the null hypothesis, as $n \rightarrow \infty$, the statistical error introduced by our use of *PDE* in the unbinned case or by binning in the binned case becomes negligible with the result that the theory *pdf* describes the data for all c at the true value s_T . i.e.

$$\frac{P^{theory}(c|s_T)}{P^{data}(c)} \rightarrow 1 \text{ as } n \rightarrow \infty \quad (23)$$

When one evaluates the likelihood ratio $\mathcal{L}_{\mathcal{R}}$ over n events, with $n \rightarrow \infty$, the likelihood ratio does not necessarily remain unity. This is due to fluctuations in the data which grow as $\sqrt{(n)}$. For the binned likelihood case with n_b bins, one can show that as $n \rightarrow \infty$,

$$\mathcal{L}_{\mathcal{R}} \rightarrow e^{-\sum_{i=1}^{i=n_b} \chi_i^2/2} \rightarrow e^{-n_b/2} \quad (24)$$

This is just an example of the likelihood ratio theorem. If one uses a binned χ^2 fit, which can also be thought of as maximizing a likelihood ratio, one gets the same limit as when using binned likelihood fits. The point is that $\mathcal{L}_{\mathcal{R}}$ is finite as $n \rightarrow \infty$. In the unbinned case, we have currently no analytic theory available. However, one can argue that the binned case with the number of bins $n_b \rightarrow \infty$ and $n_b \ll n$ should approach the unbinned limit. In this case, the unbinned $\mathcal{L}_{\mathcal{R}}$ also is finite for infinite statistics. This implies that $\mathcal{P}_n(s_T) \rightarrow \infty$ as $n \rightarrow \infty$. i.e $\mathcal{P}_n(s_T)$ depends on n . This puts an end to the notion of a monolithic Bayesian prior interpretation for $\mathcal{P}_n(s)$.

6.1.2. To show that $\mathcal{P}_n(s_T)$ is constant with respect to s

When one varies the likelihood ratio in equation 22 as a function of s , for each value of s , one is making a hypothesis that $s = s_T$. As one changes s , a new hypothesis is being tested that is mutually exclusive from the previous one, since the true value can only be at one location. So as one changes s , one is free to move the *distribution* $\mathcal{P}_n(s)$ so that s_T is at the value of s being tested. This implies that $\mathcal{P}_n(s_T)$ does not change as one changes s and is a constant *wrt* s , which we can now write as α_n . Figure 6 illustrates these points graphically. Thus $\mathcal{P}_n(s_T)$ in our equations is a number, not a function. The distribution $\mathcal{P}_n(s)$ should not be thought of as a ‘‘prior’’ but as an ‘‘unknown concomitant’’, which depends on the statistics and the measurement capabilities of the apparatus. For a given apparatus, there are a denumerable infinity of such distributions, one for each n . These distributions become narrower as n increases and $\mathcal{P}_n(s_T) \rightarrow \infty$ as $n \rightarrow \infty$.

6.2. New form of equations

Equation 22 can now be re-written

$$P(s|\mathbf{c}_n) = \frac{P(\mathbf{c}_n|s)\alpha_n}{P^{data}(\mathbf{c}_n)} \quad (25)$$

Since $P(s|\mathbf{c}_n)$ must normalize to unity, one gets for α_n ,

$$\alpha_n = \frac{P^{data}(\mathbf{c}_n)}{\int P(\mathbf{c}_n|s)ds} = \frac{1}{\int \mathcal{L}_{\mathcal{R}}(s) ds} \quad (26)$$

We have thus determined α_n , the value of the ‘‘unknown concomitant’’ at the true value s_T using our




Figure 6: Comparison of the usage of Bayesian priors with the new method. In the upper figure, illustrating the Bayesian method, an unknown distribution is guessed at by the user based on ‘‘degrees of belief’’ and the value of the Bayesian prior changes as the variable s changes. In the lower figure, an ‘‘unknown concomitant’’ distribution is used whose shape depends on the statistics. In the case of no bias, this distribution peaks at the true value of s . As we change s , we change our hypothesis as to where the true value of s lies, and the distribution shifts with s as explained in the text. The value of the distribution at the true value is thus independent of s .

data set c_n . This is our *measurement* of α_n and different datasets will give different values of α_n , in other words α_n will have a sampling distribution with an expected value and standard deviation. As $n \rightarrow \infty$, the likelihood ratio $\mathcal{L}_{\mathcal{R}}$ will tend to a finite value at the true value and zero for all other values, and $\alpha_n \rightarrow \infty$ as a result.

Note that it is only possible to write down an expression for α_n dimensionally when a likelihood ratio $\mathcal{L}_{\mathcal{R}}$ is available. This leads to

$$P(s|\mathbf{c}_n) = \frac{\mathcal{L}_{\mathcal{R}}}{\int \mathcal{L}_{\mathcal{R}} ds} = \frac{P(\mathbf{c}_n|s)}{\int P(\mathbf{c}_n|s)ds} \quad (27)$$

The last equality in equation 27 is the same expression that ‘‘frequentists’’ use for calculating their errors after fitting, namely the likelihood curve normalized to unity gives the parameter errors. If the likelihood curve is Gaussian shaped, then this justifies a change of negative log-likelihood of $\frac{1}{2}$ from the optimum point to get the 1σ errors. Even if it is not Gaussian, as we show in section (8), we may use the expression for $P(s|\mathbf{c}_n)$ as a *pdf* of the parameter s to evaluate the errors.

The normalization condition

$$P(\mathbf{c}_n) = \int P^{theory}(s, \mathbf{c}_n)ds = \int P(c_n|s)\mathcal{P}_n(s_T)ds \quad (28)$$

is obeyed by our solution, since

$$\int P(\mathbf{c}_n|s)\mathcal{P}_n(s_T) ds = \int \alpha_n P(\mathbf{c}_n|s) ds \equiv P^{data}(\mathbf{c}_n) \quad (29)$$

The expression $\int \alpha_n P(\mathbf{c}_n|s) ds$ in the above equation may be thought of as being due to an “unknown concomitant” whose peak value is distributed uniformly in s space. The likelihoods of the theoretical prediction $P(\mathbf{c}_n|s)$ contribute with equal probability each with a weight α_n , to sum up to form the data likelihood $P^{data}(\mathbf{c}_n)$. i.e. the data, due to its statistical inaccuracy will entertain a range of theoretical parameters. However, equation 29 does not give us any further information, since it is obeyed identically. Fitting for the maximum likelihood value s^* of s is attained by maximizing the likelihood ratio $\mathcal{L}_R = \frac{P(\mathbf{c}_n|s)}{P^{data}(\mathbf{c}_n)}$. The goodness of fit is obtained using the value of \mathcal{L}_R at the maximum likelihood point. The best theoretical prediction is $P(c|s^*)$, and this prediction is used to compare to the data *pdf* $P^{data}(c)$. Note that the maximum likelihood value s is also the same point at which the posterior density $P(s|c)$ peaks. This is true only in our method. When an arbitrary Bayesian prior is used, the maximum likelihood value is not the same point at which the posterior density will peak. Note also that the normalization equation $\int \mathcal{P}_n(s) ds=1$ is still valid. The integral

$$\int \alpha_n ds \neq 1 \quad (30)$$

since α_n is our measurement of the value of $\mathcal{P}_n(s)$ at the true value. It is a measure of the statistical accuracy of the experiment. The larger the value of α_n , the narrower the distribution $\mathcal{P}_n(s)$ and the more accurate the experiment.

7. COMBINING RESULTS OF EXPERIMENTS

Each experiment should publish a likelihood curve for its fit as well as a number for the data likelihood $P^{data}(\mathbf{c}_n)$. Combining the results of two experiments with m and n experiments each, involves multiplying the likelihood ratios.

$$\mathcal{L}_{R_{m+n}}(s) = \mathcal{L}_{R_m}(s) \times \mathcal{L}_{R_n}(s) = \frac{P(\mathbf{c}_m|s)}{P^{data}(\mathbf{c}_m)} \times \frac{P(\mathbf{c}_n|s)}{P^{data}(\mathbf{c}_n)} \quad (31)$$

Posterior densities and goodness of fit can be deduced from the combined likelihood ratio.

8. INTERPRETING THE RESULTS OF ONE EXPERIMENT

After performing a single experiment with n events, we now can calculate $P(s|\mathbf{c}_n)$, using equation 27.

Equation 18 gives the prescription for arriving at $\mathcal{P}_n(s)$, given an ensemble of such experiments, the contribution from each experiment being weighted by the “data likelihood” $P^{data}(\mathbf{c}_n)$ for that experiment. The “data likelihoods” integrate to unity, i.e. $\int P^{data}(\mathbf{c}_n) d\mathbf{c}_n = 1$. In the case of only a single experiment, with the observed \mathbf{c}_n being denoted by \mathbf{c}_n^{obs} ,

$$P^{data}(\mathbf{c}_n) = \delta(\mathbf{c}_n - \mathbf{c}_n^{obs}) \quad (32)$$

Equation 18, for a single experiment, then reduces to

$$\mathcal{P}_n(s) = \int P(s|\mathbf{c}_n) P^{data}(\mathbf{c}_n) d\mathbf{c}_n = P(s|\mathbf{c}_n^{obs}) \quad (33)$$

i.e. given a single experiment, the best estimator for $\mathcal{P}_n(s)$, the *pdf* of s , is $P(s|\mathbf{c}_n^{obs})$ and thus the best estimator for the true value s_T is s^{*obs} deduced from the experiment. We can thus use $P(s|\mathbf{c}_n^{obs})$ as though it is the *pdf* of s and deduce limits and errors from it. The proviso is of course that these limits and errors as well as s^{*obs} come from a single experiment of finite statistics and as such are subject to statistical fluctuations.

9. COMPARISON WITH THE BAYESIAN APPROACH

In the Bayesian approach, an unknown Bayesian prior $P(s)$ is assumed for the distribution of the parameter s in the absence of any data. The shape of the prior is guessed at, based on subjective criteria or using other objective pieces of information. However, such a shape is not invariant under transformation of variables. For example, if we assume that the prior $P(s)$ is flat in s , then if we analyze the problem in s^2 , we cannot assume it is flat in s^2 . This feature of the Bayesian approach has caused controversy. Also, the notion of a *pdf* of the data does not exist and $P(c)$ is taken to be a normalization constant. As such, no goodness of fit criteria exist. In the method outlined here, we have used Bayes’ theorem to calculate posterior densities of the fitted parameters while being able to compute the goodness of fit. The formalism developed here shows that what is conventionally thought of as a Bayesian prior distribution is in fact a normalization constant and what Bayesians think of as a normalization constant is in fact the *pdf* of the data. Table II outlines the major differences between the Bayesian approach and the new one.

Table II The key points of difference between the Bayesian method and the new method.

Item	Bayesian Method	New Method
Goodness of fit	Absent	Now available in both binned and unbinned fits
Data	Used in evaluating theory <i>pdf</i> at data points	Used in evaluating theory <i>pdf</i> at data points as well as evaluating data <i>pdf</i> at data points
Prior	Is a distribution that is guessed based on “degrees of belief” Independent of data, monolithic	No prior needed. One calculates a constant from data $\alpha_n = \frac{P^{data}(\mathbf{c}_n)}{\int P(\mathbf{c}_n s)ds}$ $\rightarrow \infty$ as $n \rightarrow \infty$
Posterior density $P(s \mathbf{c}_n)$	Depends on Prior. $\frac{P(\mathbf{c}_n s)P(s)}{\int P(\mathbf{c}_n s)P(s) ds}$	Independent of prior. same as frequentists use $\frac{P(\mathbf{c}_n s)}{\int P(\mathbf{c}_n s) ds}$

10. FURTHER WORK TO BE DONE

Equation 18 can be used to show that the expectation value of $E(s)$ of the parameter s is given by

$$E(s) = \int sP_n(s)ds = \int d\mathbf{c}_n P(\mathbf{c}_n) \int sP(s|\mathbf{c}_n)ds \quad (34)$$

$$= \int \bar{s}(\mathbf{c}_n)P(\mathbf{c}_n)d\mathbf{c}_n \quad (35)$$

where $\bar{s}(\mathbf{c}_n)$ is the average of s for individual experiments. Equation 35 states $E(s)$ is the weighted average of $\bar{s}(\mathbf{c}_n)$ obtained from individual measurements, the weight for each experiment being the “data likelihood” $P(\mathbf{c}_n)$ for that experiment. In the absence of experimental bias, $E(s)$ would be identical to the true value s_T . It remains to be shown that the weighted average of maximum likelihood values s^* from individual experiments also converge to the maximum likelihood point of $P_n(s)$.

Also one needs to develop an analytic theory of the goodness of fit for unbinned likelihood fits. Finally, one needs to investigate a bit more closely the transformation properties of $P_n(s)$ under change of variable.

11. CONCLUSIONS

To conclude, we have proposed a scheme for obtaining the goodness of fit in unbinned likelihood fits.

This scheme involves the usage of two *pdf*'s, namely data and theory. In the process of computing the fitted errors, we have demonstrated that the quantity in the joint probability equations that has been interpreted as the “Bayesian prior” is in reality a number and not a distribution. This number is the value of the *pdf* of the parameter, which we call the “unknown concomitant” at the true value of the parameter. This number is calculated from a combination of data and theory and is seen to be an irrelevant parameter. If this viewpoint is accepted, the controversial practice of guessing distributions for the “Bayesian Prior” can now be abandoned, as can be the terms “Bayesian” and “frequentist”. We show how to use the posterior density to rigorously calculate fitted errors.

Acknowledgments

This work is supported by Department of Energy. The author wishes to thank Jim Linnemann and Igor Volobouev for useful comments.

Editor’s Note: The reviewer of this article found the discussion in Sections 6-11 unconvincing, and considered that some statements about frequentism were inconsistent with accepted Frequentist statistical practice.

Author’s Response: Sections 1-6 of the paper motivate the need for two pdf’s for obtaining the goodness of fit. Sections 6-11 deal with the mathematical consequences of the two pdf’s in calculating posterior densities of fitted parameters. The conclusions of the paper are based on mathematical equations (Equation 18 is a key equation). To rebut the conclusions of the paper, it is necessary to point to errors in the mathematics.

References

- [1] K. Kinoshita, “Evaluating Quality of Fit in Unbinned Maximum Likelihood fitting”, Proceedings of the Conference on Advanced Statistical Techniques in Particle Physics, Durham, March 2002 IPPP/02/39, DCPT/02/78.
B. Yabsley, “Statistical Practice at the BELLE Experiment, and some questions”, *ibid*.
R. D. Cousins, “Conference Summary”, *ibid*.
- [2] R. A. Fisher, “On the mathematical foundations of theoretical statistics”, *Philos. Trans. R. Soc. London Ser. A* **222**, 309-368(1922);
R. A. Fisher, “Theory of statistical estimation”, *Proc. Cambridge Philos. Soc.* **22**, 700-725 (1925).
- [3] “A measure of the goodness of fit in unbinned likelihood fits”, R. Raja, long write-up, <http://www.slac.stanford.edu/econf/C030908/>

- [4] “End of Bayesianism?”, R. Raja,
<http://www.slac.stanford.edu/econf/C030908/>
- [5] E. Parzen, “On estimation of a probability density function and mode” *Ann.Math.Statis.* **32**, 1065-1072 (1962).
- [6] D. Scott. *Multivariate Density Estimation*. John Wiley & Sons, 1992.
- M. Wand and M. Jones, *Kernel Smoothing*. Chapman & Hall, 1995.