

Weak B Decays into Orbitally Excited Charmed Mesons

C. Albertus, **J. Segovia**, D.R. Entem, F. Fernández, E. Hernández and
M.A. Pérez-García

The XIV International Conference on Hadron Spectroscopy (Hadron 2011)
München, 13-17th of June 2011



Universidad de Salamanca
Spain

Contents

- 1 Introduction
- 2 Theoretical framework
 - Constituent quark model
 - Weak decays
 - Strong decays
- 3 Results
 - Semileptonic $B \rightarrow D_2^* l \nu_l$ decay
 - Semileptonic $B \rightarrow D_1 l \nu_l$ decay
 - Comparison with experiment
- 4 Summary and conclusions

1.- Introduction

1.1.- Overview

- Accuracy on the knowledge of $|V_{cb}|$ and $|V_{ub}|$ demands detailed measurements of b -hadron decays
- A substantial contribution to the semileptonic decay width of b -hadrons is provided by decays including orbitally excited charmed mesons in their final state
- Additionally, the analysis of signals and backgrounds of inclusive and exclusive measurements of b -hadron decays calls for an improved understanding of these processes
- In this scenario, data reported by Belle and BaBar offer new theoretical possibilities to test meson models as far as they include both weak and strong decays

1.- Introduction

1.2.- Belle and BaBar measurements

	Belle [1] ($\times 10^{-3}$)	BaBar [2] ($\times 10^{-3}$)
$D_2^*(2460)$		
$\mathcal{B}(B^+ \rightarrow D_2^{*0} l^+ \nu_l) \mathcal{B}(D_2^{*0} \rightarrow D^+ \pi^-)$	$2.2 \pm 0.3 \pm 0.4$	$1.4 \pm 0.2 \pm 0.2^{(*)}$
$\mathcal{B}(B^+ \rightarrow D_2^{*0} l^+ \nu_l) \mathcal{B}(D_2^{*0} \rightarrow D^{*+} \pi^-)$	$1.8 \pm 0.6 \pm 0.3$	$0.9 \pm 0.2 \pm 0.2^{(*)}$
$\mathcal{B}(B^0 \rightarrow D_2^{*-} l^+ \nu_l) \mathcal{B}(D_2^{*-} \rightarrow D^0 \pi^-)$	$2.0 \pm 0.7 \pm 0.5$	$1.1 \pm 0.2 \pm 0.1^{(*)}$
$\mathcal{B}(B^0 \rightarrow D_2^{*-} l^+ \nu_l) \mathcal{B}(D_2^{*-} \rightarrow D^{*0} \pi^-)$	< 3	$0.7 \pm 0.2 \pm 0.1^{(*)}$
$\mathcal{B}_{D/D^{(*)}}$	0.55 ± 0.9	0.62 ± 0.03
$D_1(2420)$		
$\mathcal{B}(B^+ \rightarrow D_1^0 l^+ \nu_l) \mathcal{B}(D_1^0 \rightarrow D^{*+} \pi^-)$	$4.2 \pm 0.7 \pm 0.7$	$2.97 \pm 0.17 \pm 0.17$
$\mathcal{B}(B^0 \rightarrow D_1^- l^+ \nu_l) \mathcal{B}(D_1^- \rightarrow D^{*0} \pi^-)$	$5.4 \pm 1.9 \pm 0.9$	$2.78 \pm 0.24 \pm 0.25$

1 D. Liventsev et al. (Belle Collaboration), *Phys. Rev. D* **77**, 091503 (2008)

2 B. Aubert et al. (BaBar Collaboration), *Phys. Rev. Lett.* **103**, 051803 (2009)

2.- Theoretical framework

2.1.- Constituent quark model. Main features

- Spontaneous chiral symmetry breaking (Goldstone-Boson exchange):

$$L = \bar{\psi} (i\gamma^\mu \partial_\mu - MU\gamma^5) \psi \rightarrow U\gamma^5 = 1 + \frac{i}{f_\pi} \gamma^5 \lambda^a \pi^a - \frac{1}{2f_\pi^2} \pi^a \pi^a + \dots$$

$$M(q^2) = m_q F(q^2) = m_q \left[\frac{\Lambda^2}{\Lambda^2 + q^2} \right]^{1/2}$$

- QCD perturbative effects (One-Gluon Exchange):

$$L = i\sqrt{4\pi\alpha_s} \bar{\psi} \gamma_\mu G^\mu \lambda^c \psi$$

- Confinement (screened potential):

$$V_{CON}^C(\vec{r}_{ij}) = [-a_c(1 - e^{-\mu_c r_{ij}}) + \Delta] (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c)$$

$$\begin{cases} V_{CON}^C(\vec{r}_{ij}) = (-a_c \mu_c r_{ij} + \Delta) (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c) & r_{ij} \rightarrow 0 \\ V_{CON}^C(\vec{r}_{ij}) = (-a_c + \Delta) (\vec{\lambda}_i^c \cdot \vec{\lambda}_j^c) & r_{ij} \rightarrow \infty \end{cases}$$

2.- Theoretical framework

2.1.- Constituent quark model. Some applications

• N-N interaction

- D.R. Entem, F. Fernández and A. Valcarce, Phys. Rev. C **62**, 034002 (2000)
- B. Julia-Diaz, J. Haidenbauer, A. Valcarce and F. Fernández, Phys. Rev. C **65**, 034001 (2002)

• Baryon spectrum

- H. Garcilazo, A. Valcarce and F. Fernández, Phys. Rev. C **63**, 035207 (2001)
- H. Garcilazo, A. Valcarce and F. Fernández, Phys. Rev. C **64**, 058201 (2001)

• Meson spectrum

- J. Vijande, A. Valcarce and F. Fernández, J. Phys. G **31**, 481 (2005)
- J. Segovia, D.R. Entem and F. Fernández, Phys. Rev. D **78** 114033 (2008)
- J. Segovia, D.R. Entem and F. Fernández, accepted by Phys. Rev. D

• Molecular states

- P. G. Ortega, J. Segovia, D. R. Entem and F. Fernández, Phys. Rev. D **81**, 054023 (2010)


2.- Theoretical framework

2.1.- Constituent quark model. Some applications (Continuation)



Deuteron

	CQM	NijmII	Bonn B	Exp.
ϵ_d (MeV)	-2.2242	-2.2246	-2.2246	-2.224575
P_D (%)	4.85	5.64	4.99	-
Q_d (fm ²)	0.276	0.271	0.278	0.2859 ± 0.0003
A_S (fm ^{-1/2})	0.891	0.8845	0.8860	0.8846 ± 0.0009
A_D/A_S	0.0257	0.0252	0.0264	0.0256 ± 0.0004



Light mesons



Charmonium reactions



X(3872)



2.- Theoretical framework

2.1.- Constituent quark model. Mass predictions involved in the reactions

Quark masses	m_n (MeV)	313
	m_c (MeV)	1763
	m_b (MeV)	5110
Confinement	a_c (MeV)	507.4
	μ_c (fm ⁻¹)	0.576
	Δ (MeV)	184.432
	a_s	0.81
One-gluon exchange	α_0	2.118
	Λ_0 (fm ⁻¹)	0.113
	μ_0 (MeV)	36.976
	\hat{r}_0 (fm)	0.181
	\hat{r}_g (fm)	0.259
GBE	taken from Ref. [1]	

Meson	J^{PC}	CQM (MeV)	EXP (MeV)
B	0^-	5275	5279.3 ± 0.64
D	0^-	1896	1867.20 ± 0.44
D^*	1^-	2017	2008.60 ± 0.21
$D_1(2420)$	1^+	2466	2422.85 ± 3.36
$D_2^*(2460)$	2^+	2513	2460.5 ± 5.59
π	0^{-+}	138	137.27339 ± 0.00069


[1] *J. Vijande, F. Fernández and A. Valcarce J. Phys. G 31 481 (2005)*

2.- Theoretical framework

2.2.- Weak decays. Total decay width

Study of the weak process based on

- E. Hernández, J. Nieves and J.M. Verde-Velasco, *Phys. Rev. D* **74**, 074008 (2006)
- M.A. Ivanov, J.G. Körner and P. Santorelli, *Phys. Rev. D* **73**, 054024 (2006)



2.- Theoretical framework

2.2.- Weak decays. Case $0^- \rightarrow 2^+$

$$\langle D(2^+), \lambda \vec{P}_D \left| J_\mu^{cb}(0) \right| B(0^-) \vec{P}_B \rangle = \epsilon_{\mu\nu\alpha\beta} \epsilon_{(\lambda)}^{\nu\delta*}(\vec{P}_D) P_\delta P^\alpha q^\beta T_4(q^2) \\ - i \left\{ \epsilon_{(\lambda)\mu\delta}^*(\vec{P}_D) P^\delta T_1(q^2) + P^\nu P^\delta \epsilon_{(\lambda)\nu\delta}^*(\vec{P}_D) \left[P_\mu T_2(q^2) + q_\mu T_3(q^2) \right] \right\}$$

$$T_1(q^2) = -i \frac{2m_D}{m_B |\vec{q}|} A_{T\lambda=+1}^1(|\vec{q}|),$$

$$T_2(q^2) = i \frac{1}{2m_B^3} \left\{ -\sqrt{\frac{3}{2}} \frac{m_D^2}{|\vec{q}|^2} A_{T\lambda=0}^0(|\vec{q}|) - \sqrt{\frac{3}{2}} \frac{m_D^2}{|\vec{q}|^3} (E_D(-\vec{q}) - m_B) A_{T\lambda=0}^3(|\vec{q}|) \right. \\ \left. + \frac{2m_D}{|\vec{q}|} \left(1 - \frac{E_D(-\vec{q})(E_D(-\vec{q}) - m_B)}{|\vec{q}|^2} \right) A_{T\lambda=+1}^1(|\vec{q}|) \right\}$$

$$T_3(q^2) = i \frac{1}{2m_B^3} \left\{ -\sqrt{\frac{3}{2}} \frac{m_D^2}{|\vec{q}|^2} A_{T\lambda=0}^0(|\vec{q}|) - \sqrt{\frac{3}{2}} \frac{m_D^2}{|\vec{q}|^3} (E_D(-\vec{q}) + m_B) A_{T\lambda=0}^3(|\vec{q}|) \right. \\ \left. + \frac{2m_D}{|\vec{q}|} \left(1 - \frac{E_D(-\vec{q})(E_D(-\vec{q}) + m_B)}{|\vec{q}|^2} \right) A_{T\lambda=+1}^1(|\vec{q}|) \right\}$$

$$T_4(q^2) = i \frac{m_D}{m_B^2 |\vec{q}|^2} V_{T\lambda=+1}^1(|\vec{q}|)$$

2.- Theoretical framework

2.2.- Weak decays. Case $0^- \rightarrow 1^+$

$$\langle D(1^+), \lambda \vec{P}_D | J_\mu^{cb}(0) | B(0^-), \vec{P}_B \rangle = \frac{-1}{m_B + m_D} \epsilon_{\mu\nu\alpha\beta} \epsilon_{(\lambda)}^*(\vec{P}_D) P^\alpha q^\beta A(q^2) - i \left\{ (m_B - m_D) \epsilon_{(\lambda)\mu}^*(\vec{P}_D) V_0(q^2) - \frac{P \cdot \epsilon_{(\lambda)}^*(\vec{P}_D)}{m_B + m_D} [P_\mu V_+(q^2) + q_\mu V_-(q^2)] \right\}$$

$$A(q^2) = - \frac{i}{\sqrt{2}} \frac{m_B + m_D}{m_B |\vec{q}|} A_{\lambda=-1}^1(|\vec{q}|)$$

$$V_+(q^2) = + i \frac{m_B + m_D}{2m_B} \frac{m_D}{|\vec{q}| m_B} \left\{ V_{\lambda=0}^0(|\vec{q}|) - \frac{m_B - E_D(-\vec{q})}{|\vec{q}|} V_{\lambda=0}^3(|\vec{q}|) + \sqrt{2} \frac{m_B E_D(-\vec{q}) - m_D^2}{|\vec{q}| m_D} V_{\lambda=-1}^1(|\vec{q}|) \right\}$$


$$V_-(q^2) = - i \frac{m_B + m_D}{2m_B} \frac{m_D}{|\vec{q}| m_B} \left\{ - V_{\lambda=0}^0(|\vec{q}|) - \frac{m_B + E_D(-\vec{q})}{|\vec{q}|} V_{\lambda=0}^3(|\vec{q}|) + \sqrt{2} \frac{m_B E_D(-\vec{q}) + m_D^2}{|\vec{q}| m_D} V_{\lambda=-1}^1(|\vec{q}|) \right\}$$

$$V_0(q^2) = + i \sqrt{2} \frac{1}{m_B - m_D} V_{\lambda=-1}^1(|\vec{q}|)$$

2.- Theoretical framework


2.3.- Strong decays. 3P_0 and microscopic models

- 3P_0 decay model



$$H_I = g \int d^3x \bar{\psi}(\vec{x}) \psi(\vec{x})$$

- Microscopic decay model



$$H_I = \frac{1}{2} \int d^3x d^3y J^a(\vec{x}) K(|\vec{x} - \vec{y}|) J^a(\vec{y})$$

2.- Theoretical framework

2.3.- Strong decays. 3P_0 and microscopic models (Continuation)

- 3P_0 decay model

- L. Micu, *Nucl. Phys. B* **10**, 521 (1969)
- A. Le Yaouanc, L. Olivier, O. Pène, and J.C. Raynal, *Phys. Rev. D* **8**, 2223 (1973)
- R. Bonnaz, and B. Silvestre-Brac, *Few-Body Syst.* **27**, 163 (1999)

- Microscopic decay model

- E. Eichten et al. *Phys. Rev. D* **17** 3090 (1978); **21** 203 (1980)
 → update: *Phys. Rev. D* **73** 014014 (2006)
- E.S. Ackleh et al. *Phys. Rev. D* **54**, 6811 (1996)
- Yu.A. Simonov *arXiv:1103.4028v1 [hep-ph]* 21 Mar 2011
- Bao-Fei Li et al. *arXiv:1105.1620v1 [hep-ph]* 9 May 2011

$$\Gamma_{A \rightarrow BC} = 2\pi \frac{E_B E_C}{M_A k_0} \sum_{J_{BC}, l} |\mathcal{M}_{A \rightarrow BC}(k_0; J_{BC}, l)|^2$$

$$\mathcal{M}_{A \rightarrow BC} = M_{A \rightarrow BC} + (-1)^{l_B + l_C - l_A + J_B + J_C - J_{BC} + l} M_{A \rightarrow CB}$$

$$M_{A \rightarrow BC} = \mathcal{I}_{color} \mathcal{I}_{flavor} (\mathcal{I}_{signature} \mathcal{I}_{spin-space})$$

2.- Theoretical framework

2.3.- Strong decays. Factors for the 3P_0 model

- Color term \Rightarrow

$$\mathcal{I}_{color} = \frac{1}{\sqrt{3}}$$

- Flavor term \Rightarrow

$$\mathcal{I}_{flavor} = (-1)^{t_\alpha + t_\beta + I_A} \delta_{f_\alpha f_\delta} \delta_{f_\beta f_\rho} \delta_{f_\mu f_\lambda} \delta_{f_\nu f_\epsilon} \sqrt{(2I_B + 1)(2I_C + 1)(2t_\mu + 1)} \begin{Bmatrix} t_\beta & I_C & t_\mu \\ I_B & t_\alpha & I_A \end{Bmatrix}$$

- Spin-space term \Rightarrow

$$\begin{aligned} \mathcal{I}_{spin-space} &= \frac{1}{\sqrt{1 + \delta_{BC}}} \int d^3 K_B d^3 K_C d^3 p_\alpha d^3 p_\beta d^3 p_\mu d^3 p_\nu \delta^{(3)}(\vec{K} - \vec{K}_0) \\ &\delta^{(3)}(\vec{K}_B - \vec{P}_B) \delta^{(3)}(\vec{K}_C - \vec{P}_C) \delta^{(3)}(\vec{p}_\mu + \vec{p}_\nu) \delta^{(3)}(\vec{P}_A) \frac{\delta(k - k_0)}{k} \\ &\langle \{ [\phi_B(\vec{p}_B)(s_\alpha s_\nu) S_B] J_B [\phi_C(\vec{p}_C)(s_\mu s_\beta) S_C] J_C]_{BC} Y_l(\hat{k}) \} J_A | \\ &\{ [\phi_A(\vec{p}_A)(s_\alpha s_\beta) S_A] J_A [\gamma_{\mu,(1)} \left(\frac{\vec{p}_\mu - \vec{p}_\nu}{2} \right) (s_\mu s_\nu) 1] 0 \} J_A \rangle \end{aligned}$$

2.- Theoretical framework

2.3.- Strong decays. Factors for the microscopic model

- Color term \Rightarrow

$$\mathcal{I}_{color} = \frac{2^2}{3^{\frac{3}{2}}}$$

- Flavor term \Rightarrow



$$\mathcal{I}_{flavor} = (-1)^{t_\alpha + t_\beta + I_A} \delta_{f_\alpha f_\delta} \delta_{f_\beta f_\rho} \delta_{f_\mu f_\lambda} \delta_{f_\nu f_\epsilon} \sqrt{(2I_B + 1)(2I_C + 1)(2t_\mu + 1)} \begin{Bmatrix} t_\beta & I_C & t_\mu \\ I_B & t_\alpha & I_A \end{Bmatrix}$$

- Spin-space term \Rightarrow

$$\begin{aligned} \mathcal{I}_{spin-space} &= \frac{-2}{\sqrt{1 + \delta_{BC}}} \int d^3K_B d^3K_C \sum_{m, M_{BC}} \langle J_{BC} M_{BC} | J_A M_A \rangle \delta^{(3)}(\vec{K} - \vec{K}_0) \delta(k - k_0) \\ &\frac{Y_{lm}(\hat{k})}{k} \sum_{M_B, M_C} \langle J_B M_B J_C M_C | J_{BC} M_{BC} \rangle \int d^3p_\delta d^3p_\epsilon d^3p_\lambda d^3p_\rho \delta^{(3)}(\vec{K}_B - \vec{P}_B) \\ &\delta^{(3)}(\vec{K}_C - \vec{P}_C) \phi_B(\vec{p}_B) \phi_C(\vec{p}_C) \delta_{\rho\beta} \delta^{(3)}(\vec{p}_\rho - \vec{p}_\beta) \delta^{(3)}(\vec{p}_\lambda + \vec{p}_\epsilon + \vec{p}_\delta - \vec{p}_\alpha) \\ &K(|\vec{p}_\lambda + \vec{p}_\epsilon|) \lim_{v/c \rightarrow 0} [\bar{u}_\lambda(\vec{p}_\lambda) \Gamma v_\epsilon(\vec{p}_\epsilon)] \lim_{v/c \rightarrow 0} [\bar{u}_\delta(\vec{p}_\delta) \Gamma u_\alpha(\vec{p}_\alpha)] \\ &\int d^3p_\alpha d^3p_\beta \delta^{(3)}(\vec{P}_A) \phi_A(\vec{p}_A) \end{aligned}$$

3.- Results

3.1.- Semileptonic $B \rightarrow D_2^* l \nu_l$ decay



- Semileptonic decay widths

$$\Gamma(B^+ \rightarrow D_2^{*0} l^+ \nu_l) = 1.3426 \times 10^{-15} \text{ GeV}$$



$$\Gamma(B^0 \rightarrow D_2^{*-} l^+ \nu_l) = 1.3454 \times 10^{-15} \text{ GeV}$$

- Some results about strong decays

B	Exp.	3P_0	Microscopic
$\frac{\Gamma(D_2^{*+} \rightarrow D^0 \pi^+)}{\Gamma(D_2^{*+} \rightarrow D^{*0} \pi^+)}$	$1.9 \pm 1.1 \pm 0.3$	1.80	1.96
$\frac{\Gamma(D_2^{*0} \rightarrow D^+ \pi^-)}{\Gamma(D_2^{*0} \rightarrow D^{*+} \pi^-)}$	2.4 ± 0.5	1.87	1.97
$\frac{\Gamma(D_2^{*0} \rightarrow D^+ \pi^-)}{\Gamma(D_2^{*0} \rightarrow D^{(*)+} \pi^-)}$	$0.62 \pm 0.03 \pm 0.02$	0.64	0.66

3.- Results

3.2.- Semileptonic $B \rightarrow D_1 l \nu_l$ decay



- Semileptonic decay widths

$$\Gamma(B^+ \rightarrow D_1^0 l^+ \nu_l) = 1.0226 \times 10^{-15} \text{ GeV}$$

$$\Gamma(B^0 \rightarrow D_1^- l^+ \nu_l) = 1.0217 \times 10^{-15} \text{ GeV}$$

Only one open-charm decay $\Rightarrow \mathcal{B}(D_1^0 \rightarrow D^{*+} \pi^-) = \mathcal{B}(D_1^- \rightarrow D^{*0} \pi^-) = 1$

3.- Results

3.3.- Comparison with experiment

	Belle ($\times 10^{-3}$)	BaBar ($\times 10^{-3}$)	3P_0 ($\times 10^{-3}$)	Mic. ($\times 10^{-3}$)
$D_2^*(2460)$				
$\mathcal{B}(B^+ \rightarrow D_2^{*0} l^+ \nu_l) \mathcal{B}(D_2^{*0} \rightarrow D^+ \pi^-)$	2.2 ± 0.5	1.4 ± 0.3	2.2	2.2
$\mathcal{B}(B^+ \rightarrow D_2^{*0} l^+ \nu_l) \mathcal{B}(D_2^{*0} \rightarrow D^{*+} \pi^-)$	1.8 ± 0.7	0.9 ± 0.3	1.2	1.1
$\mathcal{B}(B^+ \rightarrow D_2^{*0} l^+ \nu_l) \mathcal{B}(D_2^{*0} \rightarrow D^{(*)+} \pi^-)$	4.0 ± 0.9	2.3 ± 0.4	3.4	3.3
$\mathcal{B}(B^0 \rightarrow D_2^{*-} l^+ \nu_l) \mathcal{B}(D_2^{*-} \rightarrow D^0 \pi^-)$	2.0 ± 0.9	1.1 ± 0.2	2.0	2.1
$\mathcal{B}(B^0 \rightarrow D_2^{*-} l^+ \nu_l) \mathcal{B}(D_2^{*-} \rightarrow D^{*0} \pi^-)$	< 3	0.7 ± 0.2	1.1	1.0
$\mathcal{B}(B^0 \rightarrow D_2^{*-} l^+ \nu_l) \mathcal{B}(D_2^{*-} \rightarrow D^{(*)0} \pi^-)$	< 5	1.8 ± 0.3	3.1	3.1
\mathcal{B}_{D/D^*}	0.55 ± 0.9	0.62 ± 0.03	0.64	0.66
$D_1(2420)$				
$\mathcal{B}(B^+ \rightarrow D_1^0 l^+ \nu_l) \mathcal{B}(D_1^0 \rightarrow D^{*+} \pi^-)$	4.2 ± 1.0	2.97 ± 0.24	2.5	2.5
$\mathcal{B}(B^0 \rightarrow D_1^- l^+ \nu_l) \mathcal{B}(D_1^- \rightarrow D^{*0} \pi^-)$	5.4 ± 2.1	2.78 ± 0.35	2.4	2.4

4.- Summary and conclusions

- We have studied semileptonic B decays into orbitally excited charmed mesons
- These data offer new theoretical possibilities to test meson models as far as they include weak and strong processes
- Weak decays: Studied within spectator approximation and in the helicity formalism.
- Strong decays: We study these processes within the context of the 3P_0 and microscopic models
- Predictions for $B \rightarrow D_2^* l \nu_l$: good agreement with Belle results and slightly higher than those reported by BaBar
- However, predictions for $B \rightarrow D_1 l \nu_l$: better agreement with BaBar data