Low-Energy Pion-Photon Reactions and Chiral Symmetry

N. Kaiser

International Conference HADRON 2011: Munich, 13. June 2011

- Tests of chiral perturbation theory via low-energy $\pi^-\gamma$ reactions
- COMPASS@CERN: Primakoff effect to extract $\pi^-\gamma$ cross sections
- Pion Compton scattering in ChPT: electric/magnetic polarizabilities
- Radiative corrections to $\pi^-\gamma \rightarrow \pi^-\gamma$, isospin-breaking correction
- Neutral and charged pion-pair production: $\pi^- \gamma \rightarrow \pi^- \pi^0 \pi^0 \& \pi^+ \pi^- \pi^-$
- $\bullet\,$ Total cross sections and $2\pi\,$ invariant mass spectra at one-loop order
- Radiative corrections to $\pi^- \gamma \rightarrow \pi^- \pi^0 \pi^0$ (simpler case)

ヘロト ヘ戸ト ヘヨト ヘヨト

Introduction: some ChPT highlights

- Pions $\pi^{\pm 0}$: Goldstone bosons of spontaneous chiral symmetry breaking
- Their low-energy dynamics: systematically (and accurately) calculable in Chiral Perturbation Theory (= loop-expansion with effective Lagrangian)
- 2-loop prediction for *I* = 0 ππ-scattering length: a₀m_π = 0.220 ± 0.005 confirmed by NA48/2@CERN: K⁺ → π⁺π⁻e⁺ν_e (π⁺π⁻ mass distribut.)
- Implications: quark condensate $\langle 0|\bar{q}q|0\rangle$ is large, linear term dominates quark mass expansion of m_{π}^2 : $m_{\pi}^2 f_{\pi}^2 = -\langle 0|\bar{q}q|0\rangle m_q + \mathcal{O}(m_q^2 \ln m_q)$
- DIRAC@CERN: Pionium lifetime $\tau_{pred} = (2.9 \pm 0.1) \cdot 10^{-15}$ sec

$$\Gamma((\pi^{+}\pi^{-})_{atom} \to \pi^{0}\pi^{0}) = \frac{2}{9}\alpha^{3}\rho_{cm}m_{\pi}^{2}(a_{0} - a_{2})^{2} + \dots$$

- Cusp effect in $2\pi^0$ mass spectrum of $K^+ \to \pi^+ \pi^0 \pi^0$ at $\pi^+ \pi^-$ threshold: $(a_0 - a_2)m_{\pi} = 0.257 \pm 0.006$, ChPT: $(a_0 - a_2)m_{\pi} = (0.265 \pm 0.005)$
- Electromagnetic processes with pions allow for further tests of ChPT
- Pion polarizability difference (2-loops): $\alpha_{\pi} \beta_{\pi} = (5.7 \pm 1.0) \cdot 10^{-4} \text{ fm}^3$, experimental determinations from Serpukhov and Mainz in conflict with it

Introduction: Primakoff effect





 Scattering of high energy pions in nuclear Coulomb field (high Z) allows to extract cross sections for π⁻γ reactions (equivalent-photon method)

$$\frac{d\sigma}{ds\,dQ^2} = \frac{Z^2\alpha}{\pi(s-m_\pi^2)}\frac{Q^2-Q_{\min}^2}{Q^4}\,\sigma_{\pi^-\gamma}(s)\,,\qquad Q_{\min} = \frac{s-m_\pi^2}{2E_{beam}}$$

- $s = (\pi^- \gamma \text{ invariant mass})^2$, $Q \rightarrow 0$ momentum transfer by virtual photon
- isolate Coulomb peak from strong interaction background
- COMPASS@CERN: (E18@TUM, S. Paul, J. Friedrich,...)
- π -Compton scattering $\pi^-\gamma \rightarrow \pi^-\gamma$: electric and magnetic polarizabilities
- π^0 -production $\pi^- \gamma \to \pi^- \pi^0$: test QCD chiral anomaly, $F_{\gamma 3\pi} = e/(4\pi^2 t_\pi^3)$
- pion-pair product. $\pi^-\gamma \rightarrow 3\pi$: $\sqrt{s} > 1$ GeV meson spectroscopy, exotics, high statistics allows to continue event rates even down to threshold



・ロト ・四ト ・ヨト・ヨト

Pion Compton-scattering in ChPT

• Pion Compton-scattering: $\pi^{-}(p_1) + \gamma(k_1, \epsilon_1) \rightarrow \pi^{-}(p_2) + \gamma(k_2, \epsilon_2)$ T-matrix in center-of-mass frame in Coulomb gauge $\epsilon_{1,2}^0 = 0$:

$$T_{\pi\gamma} = 8\pi\alpha \Big\{ -\vec{\epsilon_1} \cdot \vec{\epsilon_2} A(s,t) + \vec{\epsilon_1} \cdot \vec{k_2} \vec{\epsilon_2} \cdot \vec{k_1} \frac{2}{t} \Big[A(s,t) + B(s,t) \Big] \Big\}$$

Mandelstam variables: $s = (p_1 + k_1)^2$, $t = (k_1 - k_2)^2$ Differential cross section:

Differential cross section:

$$\frac{d\sigma}{d\Omega_{cm}} = \frac{\alpha^2}{2s} \Big\{ \big| A(s,t) \big|^2 + \big| A(s,t) + (1+z)B(s,t) \big|^2 \Big\}$$

 $t = (s - m_{\pi}^2)^2 (z - 1)/2s$ with $z = \cos \theta_{cm}$, scattering angle • Tree diagrams:

$$egin{aligned} egin{aligned} egi$$



Pion Compton-scattering in ChPT



• Pion-loop diagrams (photon scattering off the pion's "pion cloud"):

$${\cal A}({
m s},t)^{(loop)} = {1\over (4\pi f_\pi)^2} igg\{ -{t\over 2} - 2m_\pi^2 \ln^2 {\sqrt{4m_\pi^2 - t} + \sqrt{-t}\over 2m_\pi} igg\} \sim t^2 > 0$$

with $f_{\pi}=$ 92.4 MeV, expression corresponds to isospin limit: $m_{\pi^0}=m_{\pi}$

• Electric/magnetic polarizabilities = low-energy const. with $\alpha_{\pi} + \beta_{\pi} = 0$

$$A(s,t)^{(\text{pola})} = -\frac{\beta_{\pi}m_{\pi}t}{2\alpha} < 0, \qquad \alpha_{\pi} - \beta_{\pi} = \frac{\alpha}{24\pi^{2}f_{\pi}^{2}m_{\pi}}(\bar{\ell}_{6} - \bar{\ell}_{5})$$

Pion Compton-scattering in ChPT

- Combination $\bar{\ell}_6 \bar{\ell}_5 = 3.0 \pm 0.3$ determined via radiative pion decay $\pi^+ \rightarrow e^+ \nu_e \gamma$, PIBETA@PSI: axial-to-vector coupl. ratio $F_A/F_V \simeq 0.44$
- Current-algebra relation: $\langle 0|A^{\mu}V^{\nu}|\pi\rangle \simeq f_{\pi}\langle \pi|V^{\mu}V^{\nu}|\pi\rangle$ plus corrections
- One-loop "prediction": $\alpha_{\pi} = -\beta_{\pi} \simeq 3.0 \cdot 10^{-4} \, {\rm fm}^3$
- $\sigma_{tot}(s)$ insensitive to pion's low-energy structure
- Small effect on <u>backward</u> angular distributions of $d\sigma/d\Omega_{cm}$



- Pion-loop compensates partly reduction of $d\sigma/d\Omega_{cm}$ by polarizabilities
- Effect of pion polarizabilities on π -Compton cross section: less than 20%
- 2-loop corrections to $d\sigma/d\Omega_{cm}$ are very small (Gasser, Ivanov)

Pion polarizabilities in ChPT

- Gasser et al., NPB745, 84 (2006): Pion polarizabilities to 2 loops
- Analytical expression in terms of low-energy constants $\bar{\ell}_j$:

$$\begin{aligned} \alpha_{\pi} - \beta_{\pi} &= \frac{\alpha(\bar{\ell}_{6} - \bar{\ell}_{5})}{24\pi^{2}f_{\pi}^{2}m_{\pi}} + \frac{\alpha m_{\pi}}{(4\pi f_{\pi})^{4}} \Big\{ c^{r} + \frac{8}{3} \Big(\bar{\ell}_{2} - \bar{\ell}_{1} + \bar{\ell}_{5} - \bar{\ell}_{6} + \frac{65}{12} \Big) \ln \frac{m_{\pi}}{m_{\rho}} \\ &+ \frac{4}{9} (\bar{\ell}_{1} + \bar{\ell}_{2}) - \frac{\bar{\ell}_{3}}{3} + \frac{4\bar{\ell}_{4}}{3} (\bar{\ell}_{6} - \bar{\ell}_{5}) - \frac{187}{81} + \Big(\frac{53\pi^{2}}{48} - \frac{41}{324} \Big) \Big\} \end{aligned}$$

- Improved values of $\bar{\ell}_j$ from $\pi\pi$ data, $c^r \simeq 0$ via resonance saturation
- 2-loop prediction including realistic estimate of theoretical errors:

$$\alpha_{\pi} - \beta_{\pi} = (5.7 \pm 1.0) \cdot 10^{-4} \, \text{fm}^3, \qquad \alpha_{\pi} + \beta_{\pi} = (0.16 \pm 0.1) \cdot 10^{-4} \, \text{fm}^3$$

- Good reasons to believe that chiral prediction is stable against higher order corrections: ChPT at 2-loop order works very well for γγ → π⁰π⁰
- Existing expt. determinations α_π − β_π = (15.6 ± 7.8) · 10⁻⁴ fm³ from Serpukhov (via Primakoff) and α_π − β_π = (11.6 ± 3.4) · 10⁻⁴ fm³ from Mainz (via γp → γπ⁺n) violate chiral low-energy theorem by a factor 2!

• $\alpha_{\pi} + \beta_{\pi} = \frac{1}{2\pi^2} \int \frac{d\omega}{\omega^2} \sigma_{abs}^{\pi\gamma}(\omega)$ agrees with results from dispersion sum rules



Radiative corrections to pion Compton scattering

- Pion-structure effects small: necessary to include <u>radiative corr.</u> of $\mathcal{O}(\alpha)$
- Start with structureless pion: extensive calculation in 1-loop scalar QED
- Advantage of Coulomb gauge: all s-channel pole diagrams vanish



 Dimensional regularization to treat both ultraviolet divergencies (d < 4) and infrared divergencies (d > 4):

$$\xi = \frac{1}{d-4} + \frac{1}{2}(\gamma_E - \ln 4\pi) + \ln \frac{m_\pi}{\mu}$$

Alternative: introduce regulator photon mass m_{γ} , $\xi_{IR} = \ln(m_{\pi}/m_{\gamma})$

Radiative corrections to pion Compton scattering

• Infrared-finite after inclusion of soft photon bremsstrahlung: $d\sigma/d\Omega_{\rm cm} \cdot \delta_{\rm soft}$

$$\delta_{\text{soft}} = \alpha \, \mu^{4-d} \int_{|\vec{h}_{0,1}| < \lambda} \frac{d^{d-1}l}{(2\pi)^{d-2} \, l_0} \left\{ \frac{2m_{\pi}^2 - t}{p_1 \cdot l \, p_2 \cdot l} - \frac{m_{\pi}^2}{(p_1 \cdot l)^2} - \frac{m_{\pi}^2}{(p_2 \cdot l)^2} \right\}$$

 Evaluated in dim. regularization: ξ_{IR} from photon loops gets canceled, radiative correction depends on a small energy resolution scale λ:

$$\delta_{\text{real}} = \frac{\alpha}{\pi} \Biggl\{ \Biggl[2 + \frac{4\hat{t} - 8}{\sqrt{\hat{t}^2 - 4\hat{t}}} \ln \frac{\sqrt{4 - \hat{t}} + \sqrt{-\hat{t}}}{2} \Biggr] \ln \frac{m_{\pi}}{2\lambda} + \frac{\hat{s} + 1}{\hat{s} - 1} \ln \hat{s} + \int_{0}^{1/2} dx \frac{(\hat{s} + 1)(\hat{t} - 2)}{\sqrt{W} [1 - \hat{t}x(1 - x)]} \ln \frac{\hat{s} + 1 + \sqrt{W}}{\hat{s} + 1 - \sqrt{W}} \Biggr\}$$

where $\hat{s} = s/m_{\pi}^2$, $\hat{t} = t/m_{\pi}^2$ and $W = (\hat{s} - 1)^2 + 4\hat{s}\hat{t}x(1 - x)$

- Terms beyond $\ln(m_{\pi}/2\lambda)$ specific for evaluation in center-of-mass frame
- Idealized experiment with undetected soft photons filling in momentum space a small sphere of radius λ in the center-of-mass frame
- Further experiment-specific soft/hard γ -radiation can be accounted for

・ロット (雪) ・ (目) ・ (日)

Radiative corrections to pion Compton scattering

- QED radiative corrections are maximal in backward directions $z\simeq -1$
- Same kinematical signature as pion polarizability difference $\alpha_{\pi} \beta_{\pi}$
- $\bullet~$ Suppressed by a factor of $\lesssim 10$
- In long wavelength limit $k_1, k_2 \rightarrow 0$: all strong and radiative corrections vanish, pure Thomson amplitude $T_{\pi^-\gamma}^{(0)} = -8\pi\alpha \,\vec{\epsilon_1} \cdot \vec{\epsilon_2}$ survives

A (10) F (10) F (10)

Radiative corrections including pion structure

- Include leading pion-structure in form of polarizability difference $\alpha_{\pi} \beta_{\pi}$
- Reinterpret $\gamma\gamma$ contact vertex as representing the pion polarizabilities:

$$\sim F_{\mu\nu}F^{\mu\nu}, \qquad 8\pi i\beta_{\pi}m_{\pi}\Big(k_{1}\cdot k_{2}\epsilon_{1}\cdot\epsilon_{2}-\epsilon_{1}\cdot k_{2}\epsilon_{2}\cdot k_{1}\Big)$$

• Reference cross section: point-like $d\sigma^{(pt)}/d\Omega_{cm}$ + polarizability improved

$$\frac{d\sigma^{(\text{pola})}}{d\Omega_{\text{cm}}} = \frac{\alpha\beta_{\pi}m_{\pi}^{3}(s-m_{\pi}^{2})^{2}(1-z)^{2}}{2s^{2}[s(1+z)+m_{\pi}^{2}(1-z)]}$$

Isospin-breaking in pion Compton scattering

• Isospin-breaking induced by charged/neutral pion mass difference (elm)

$$A(s,t)^{(isobr)} = \frac{m_{\pi}^2 - m_{\pi^0}^2}{(2\pi f_{\pi})^2} \bigg\{ -\frac{1}{2} - \frac{2m_{\pi}^2}{t} \ln^2 \frac{\sqrt{4m_{\pi}^2 - t} + \sqrt{-t}}{2m_{\pi}} \bigg\} \sim t$$

entirely from dependence of chiral $\pi\pi$ interaction on $m_{\pi^0}^2$,

• Small contribution to pion polarizability difference

$$\delta(lpha_{\pi} - eta_{\pi}) = rac{lpha(m_{\pi}^2 - m_{\pi^0}^2)}{24\pi^2 f_{\pi}^2 m_{\pi}^3} \simeq 1.3 \cdot 10^{-5} \, {
m fm}^3$$

- Affects backward cross section at level of few permille at most
- One order of magnitude smaller than "genuine" radiative corrections

Tree level cross sections for $\pi^-\gamma \rightarrow 3\pi$

- Coulomb gauge $\epsilon \cdot p_1 = \epsilon \cdot k = 0$, photon does not couple to incoming π^-
- No $\gamma 4\pi$ vertex at leading order

(s - w)/f²_π factor: chiral ππ-interaction, rest from 3-body phase space
 How large are next-to-leading order corrections from chiral loops + cts?

• 3-body process: $\pi^{-}(p_1) + \gamma(k, \epsilon) \to \pi^{-}(p_2) + \pi^{0}(q_1) + \pi^{0}(q_2)$

• general form of T-matrix (in Coulomb gauge)

$$T_{3\pi} = \frac{2e}{t_{\pi}^2} \Big[\vec{\epsilon} \cdot \vec{q}_1 A_1 + \vec{\epsilon} \cdot \vec{q}_2 A_2 \Big], \qquad A_2 = A_1 \Big| (\mathbf{s}_1 \leftrightarrow \mathbf{s}_2, t_1 \leftrightarrow t_2)$$

• amplitudes A₁ and A₂ depend on five (independ.) Mandelstam variables:

$$s = (p_1 + k)^2$$
, $s_1 = (p_2 + q_1)^2$, $s_2 = (p_2 + q_2)^2$, $t_1 = (q_1 - k)^2$, $t_2 = (q_2 - k)^2$

• convenient for permutation of identical neutral pions ($s_1 \leftrightarrow s_2, t_1 \leftrightarrow t_2$)

tree-level amplitudes:

$$A_1^{(\text{tree})} = A_2^{(\text{tree})} = rac{2m_\pi^2 + s - s_1 - s_2}{3m_\pi^2 - s - t_1 - t_2}$$

Pion-loop corrections (example I)

$$A_{1}^{(I)} = \frac{1}{(4\pi f_{\pi})^{2}} \frac{2m_{\pi}^{2} + s - s_{1} - s_{2}}{3m_{\pi}^{2} - s - t_{1} - t_{2}} \left\{ \left(\xi + \ln \frac{m_{\pi}}{\mu}\right) (s_{1} + s_{2} + t_{1} + t_{2} - 11m_{\pi}^{2}) + (s_{1} + s_{2} + t_{1} + t_{2} - 7m_{\pi}^{2}) \left[J(3m_{\pi}^{2} + s - s_{1} - s_{2}) - \frac{1}{2}\right] \right\}$$

• Loop function (from loop with two pion-propagators)

$$J(s) = \sqrt{\frac{s - 4m_{\pi}^2}{s}} \left[\ln \frac{\sqrt{|s - 4m_{\pi}^2|} + \sqrt{|s|}}{2m_{\pi}} - \frac{i\pi}{2}\theta(s - 4m_{\pi}^2) \right], \ s < 0 \text{ or } s > 4m_{\pi}^2$$

イロト イポト イヨト イヨト

1

Loop function (from loop with three pion-propagators)

$$G(s) = \left[\ln \frac{\sqrt{|s - 4m_{\pi}^2|} + \sqrt{|s|}}{2m_{\pi}} - \frac{i\pi}{2}\theta(s - 4m_{\pi}^2) \right]^2, \quad s < 0 \text{ or } s > 4m_{\pi}^2$$

 $+2m_{\pi}^{2}\left[G(m_{\pi}^{2}+s-s_{1}-s_{2}+t_{1}+t_{2})-G(3m_{\pi}^{2}+s-s_{1}-s_{2})\right]\right\}$

• Chiral loop and counterterm corrections (completed)

- Chiral 6π -vertex: challenging combinatorics involved, 6! = 720
- Pion wavefunction renormalization factor, chiral counterterms $\sim \ell_1, \ell_2, \ell_4$
- First crucial check: ultraviolet divergence *ξ* drops out in total sum for *A*_{1,2}

• Introduce low-energy constants that subsume chiral logarithm $\ln(m_{\pi}/\mu)$

$$\ell_j^r = \frac{\gamma_j}{32\pi^2} \left(\bar{\ell}_j + 2\ln\frac{m_{\pi}}{\mu} \right), \quad \gamma_1 = \frac{1}{3}, \ \gamma_2 = \frac{2}{3}, \ \gamma_3 = -\frac{1}{2}, \ \gamma_4 = 2$$

Complete counterterm contribution:

$$\begin{aligned} \mathcal{A}_{1}^{(\mathrm{ct})} &= \frac{1}{(4\pi f_{\pi})^{2}} \frac{1}{3m_{\pi}^{2} - s - t_{1} - t_{2}} \left\{ \frac{\bar{\ell}_{1}}{3} (s_{1} + s_{2} - s - m_{\pi}^{2})^{2} + \frac{\bar{\ell}_{2}}{3} \left[s^{2} + s_{1}^{2} + s_{2}^{2} \right. \\ &+ t_{2}^{2} - 2ss_{1} + (s - 2s_{1} + 2s_{2} - t_{1})t_{2} + m_{\pi}^{2} (s - 6s_{2} + t_{1} - 2t_{2} + 6m_{\pi}^{2}) \right] \\ &- \frac{\bar{\ell}_{3}}{2} m_{\pi}^{4} + 2\bar{\ell}_{4} m_{\pi}^{2} (s + 2m_{\pi}^{2} - s_{1} - s_{2}) \right\} \end{aligned}$$

- Finite loop corrections with $\xi + \ln(m_{\pi}/\mu)$ terms deleted altogether
- Values of low-energy constants: $\bar{\ell}_1 = -0.4 \pm 0.6$, $\bar{\ell}_2 = 4.3 \pm 0.1$, $\bar{\ell}_3 = 2.9 \pm 2.4$, $\bar{\ell}_4 = 4.4 \pm 0.2$, determined with improved empirical input

- enhancement of $\sigma_{tot}(s)$ by factor 1.5 1.8 through chiral corrections
- suggestive explanation: $\pi^+\pi^- \to \pi^0\pi^0$ final state interaction $(1 + 0.20)^2$

$$\frac{1}{3}(a_0 - a_2) = \frac{3m_\pi}{32\pi f_\pi^2} \left[1 + \frac{m_\pi^2}{36\pi^2 f_\pi^2} \left(\bar{\ell}_1 + 2\bar{\ell}_2 - \frac{3\bar{\ell}_3}{8} + \frac{9\bar{\ell}_4}{2} + \frac{33}{8} \right) \right]$$

- Uncertainty induced by errorbars of $\bar{\ell}_i$: about $\pm 5\%$ for $\sigma_{tot}(s)$, mainly $\bar{\ell}_1$
- More exclusive observables: two-pion mass spectra
- $\pi^0 \pi^0$ invariant mass²: $\mu^2 = s s_1 s_2 + 3m_{\pi}^2$, $\pi^0 \pi^-$ invariant mass²: s_1 , range of invariant masses: $2m_{\pi} < \mu, \sqrt{s_1} < \sqrt{s} m_{\pi}$

- Mass spectra reproduce enhancement by chiral correct. seen in $\sigma_{tot}(s)$
- No further specific dynamical details visible in two-pion mass spectra

Charged pion-pair production

- 3-body process: $\pi^-(p_1) + \gamma(k, \epsilon) \rightarrow \pi^+(p_2) + \pi^-(q_1) + \pi^-(q_2)$
- $\bullet\,$ Photon couples to all charged pions: $\rightarrow\,$ many more diagrams

3

Charged pion-pair production

• $\sigma_{tot}(s)$ for $\sqrt{s} < 6m_{\pi}$ almost unchanged in comparison to tree approx.

• suggestive explanation: $\pi^-\pi^- \rightarrow \pi^-\pi^-$ final state interaction $(1 - 0.02)^2$

$$\mathbf{a}_{2} = -\frac{m_{\pi}}{16\pi f_{\pi}^{2}} \left[1 - \frac{m_{\pi}^{2}}{12\pi^{2}f_{\pi}^{2}} \left(\bar{\ell}_{1} + 2\bar{\ell}_{2} - \frac{3\bar{\ell}_{3}}{8} - \frac{3\bar{\ell}_{4}}{2} + \frac{3}{8} \right) \right]$$

 Analysis of COMPASS data agrees with tree approximation of ChPT (see next talk by Sebastian Neubert)

More exclusive observables: two-pion mass spectra

• Dip in $\pi^+\pi^-$ mass spectr. at intermediate $\sqrt{s_1}$ produced by chiral correc.

- Squared T-matrix $|\hat{k} \times (\vec{q}_1 A_1 + \vec{q}_2 A_2)|^2$ with its full dependence on pion energies and angles includes still more dynamical information
- It is expected that high statistics COMPASS data can reveal such details
- Role of ρ (770) resonance ($\Gamma_{\rho} = 150 \text{ MeV}$) needs to be investigated
- For π⁻γ → π⁺π⁻π⁻ inclusion of ρ⁰ resonance (consistent with chiral symmetry) does not affect total cross section σ_{tot}(s) below √s = 5m_π

ヘロト ヘ戸ト ヘヨト ヘヨト

Radiative corrections to neutral pion-pair production

• Radiative corr. to total cross section vary between about +2% and -2%

• Radiative corrections to $\pi^-\gamma \rightarrow \pi^+\pi^-\pi^-$ could be much more sizeable, Coulomb singularity from γ -exchange between charged pions: $\alpha \pi / v_{rel}$

nan