Precision Measurement of the Neutron Spin Structure Function.

A Dissertation Submitted to
the Temple University Graduate Board

in Partial Fulfillment of the Requirements for the Degree
DOCTOR OF PHILOSOPHY

by
Piotr A. Zyla
January, 1998

ABSTRACT

The SLAC E154 experiment measured the neutron spin structure function $g_1^n(x, Q^2)$. Scattering the 48.3 GeV longitudinally polarized electrons off the longitudinally polarized ^{3}He target allowed us to extend (in comparison with the previous SLAC E142 measurement) the kinematic range to $0.014 \leq x \leq 0.7$ and $1\,\mathrm{GeV}^2 \leq \mathrm{Q}^2 \leq 17\,\mathrm{GeV}^2$. We report the integral over the measured region $\int_{0.014}^{0.7} dx\; g_1^n(x) = -0.036 \pm 0.004(\mathrm{stat.}) \pm 0.005(\mathrm{syst.})$ at the average $Q^2 = 5\,\mathrm{GeV}^2$ of the experiment. We found large and strongly x dependent values of g_1^n at low x that call into question the traditional Regge theory method for extrapolating to $x = 0$ needed for the evaluation of QCD sum rules. Within perturbative QCD we performed the next to leading order global analysis of all polarized DIS structure function measurements. It explains the perturbative origin of the observed low x behavior. Using this next to leading order parametrization we evolve the neutron g_1^n and proton g_1^p experimental data to a common value of $Q^2 = 5\,\mathrm{GeV}^2$, evaluate their integrals and confirm the Bjorken sum rule.

TABLE OF CONTENTS

LIST OF TABLES

LIST OF FIGURES

CHAPTER 1

INTRODUCTION TO SPIN-DEPENDENT DEEP INELASTIC SCATTERING

1.1 Historical background

Since the beginnings of ancient Greek philosophy, natural sciences have been in search for the atoms of Democritus, the indivisible building blocks from which more complicated objects are built. Physics considers the question of the elementary objects that constitute matter. It has only been in the sixties that, based on hadron spectroscopy, Gell-Mann [1] and Zweig [2] suggested that quarks are the fundamental building blocks of hadrons. The quark model, based on $SU(3)_{\mathrm{flavor}}$, was formulated, and the search for its dynamics began. The simplest experimental method of probing the hadrons (protons) seemed to be with a beam of structureless leptons (electrons). The first experiments, called Deep Inelastic Scattering (DIS), were performed at

SLAC's End Station A in the late 1960's [3]. They revealed the approximate Bjorken scaling [4] of the structure functions in the large momentum transfer Q^2. One of the easiest ways of understanding this feature was to assume that the electrons scattered off almost free pointlike partons, as proposed by Bjorken and Paschos [5] and Feynman [6]. Thus the underlying dynamics should exhibit weaker coupling at shorter distances (larger momenta). At that time none of the proposed quantum field theories for quarks possessed that property. The long waited Yang-Mills [7] non-Abelian gauge theory was examined, and soon t'Hooft [8], Gross and Wilczek [9], and Politzer [10] reported its property of asymptotic freedom. In the meantime Han and Nambu [11] had suggested the color quantum number for quarks to resolve several difficulties of the quark model. Fritzsch and Gell-Mann [12] proposed the $\mathrm{SU}(3)_{\mathrm{color}}$ group for the symmetry of the non-Abelian gauge theory. Quantum Chromodynamics (QCD) was born, and the quarks were identified with partons. We were reminded of the importance of the early DIS experiments at SLAC when experimenters Jerome I. Friedman, Henry W. Kendall and Richard E. Taylor received the 1990 Nobel Prize in Physics [13].

The pioneering DIS experiments probed the momentum distribution of partons within the nucleon. The polarized experiments that dealt with the partonic spin distribution were performed by Yale-SLAC group [14, 15] soon after them. This work received renewed attention when EMC [16] extended the SLAC measurements on the proton to smaller x values and concluded that the quarks carry very little of the nucleon spin, in contradiction with the expectation from naive quark models. Since then much more data have been obtained by SLAC [17, 18, 19], CERN [20, 21] and DESY [22].

Our experiment E154 took place at SLAC's End Station A in October-November 1995. By scattering longitudinally polarized electrons from a longitudinally and transversely polarized ^{3}He targets, we measured the spin structure functions of the neutron $g_1^n(x, Q^2)$ and $g_2^n(x, Q^2)$. The high statistical precision and the broad kinematic range of the data gives us a better insight into the dynamics of quarks and gluons within the nucleon.

1.2 Structure functions

To explore the internal structure of the nucleon it is necessary to use a probe which has a wavelength smaller than the size of the nucleon. Since the wavelength of a particle is proportional to the inverse of its momentum, the probe must be very energetic. It is also convenient if the probe itself is structureless, say a lepton. In addition, if we want to look at the spin distribution within the nucleon both, as we shall see later, the object and the probe need to have a definite polarization. Let's consider a typical fixed target DIS experiment in which the polarized electrons scatter on a polarized proton target. In the lowest order in the electromagnetic coupling one virtual photon is exchanged in that process. This is shown in figure 1.1. The incident electron has an initial four-momentum $k^\mu = (E, \vec{k})$ and a polarization four-vector[1] s^μ. The electron scatters with respect to $\vec{k}$ through the lab angle

[1]In the rest frame the polarization four-vector s^μ has the form $s^\mu = (0, \vec{s})$ where $\vec{s}$ is a polarization. It is orthogonal to the momentum, $s \cdot p = 0$, and normalized, $s^2 = -1$.



Figure 1.1: Tree level diagram for DIS lepton-hadron scattering.

θ and has final momentum $k'^{\mu} = (E', \vec{k}')$. The exchanged virtual photon carries momentum $q^{\mu} = (k - k')^{\mu}$. The proton has momentum $P^{\mu} = (M, \vec{0})$ in the lab frame and polarization $S^{\mu} = (0, \vec{S})$. The final hadronic states X with the same invariant mass $W^2 = (P + q)^2$ are not detected. The process is described by two independent kinematic invariants, $Q^2 = -q^2$ and $\nu = P \cdot q/M$ ($= E - E'$ in the lab) or equivalently by Q^2 and $x = \frac{Q^2}{2M\nu}$. From the experimental point of view it is convenient to use the lab frame and work with the measurable quantities, E' and θ. The kinematic variables are summarized in table 1.1. The differential cross section for the considered process, calculated from the leading order diagram in fig. 1.1, can be written as [23]

$$\frac{d^2\sigma}{d\Omega\, dE'} = \frac{4\alpha^2}{Q^4}\frac{E'}{E} L_{\mu\nu} W^{\mu\nu}, \tag{1.1}$$

Table 1.1: Kinematic Variables for Inclusive Electron-Proton Scattering.

Symbol	Definition	Lab frame	Description
E			energy of incident e
$\vec{k}$			momentum of incident e
k^μ	$(E,\vec{k})$		4-momentum of incident e
s^μ			polariz. 4-vector of incident e
E'			energy of scattered e
$\vec{k'}$			momentum of scattered e
k'^μ	$(E',\vec{k'})$		4-momentum of scattered e
m			mass of e
θ	$\arccos\left(\frac{\vec{k}\cdot\vec{k'}}{\lvert\vec{k}\rvert\lvert\vec{k'}\rvert}\right)$		scattering angle of e
M			mass of proton
P^μ		$(M,0)$	4-momentum of proton
S^μ		$(0,\vec{S})$	polariz. 4-vector of proton
q^μ	$(k-k')^\mu$	$(\nu,\vec{q})$	4-momentum transfer
Q^2	$-q^2$	$4EE'\sin^2(\theta/2)$	4-mom. transfer squared
ν	$P\cdot q/M$	$E-E'$	energy transfer
x	$\frac{Q^2}{2M\nu}$	$\frac{4EE'\sin^2(\theta/2)}{2M(E-E')}$	Bjorken x
			of the final hadronic system

where α is the fine structure constant and $L_{\mu\nu}$, and $W^{\mu\nu}$ are leptonic and hadronic tensors, respectively. The leptonic part describes matrix elements of currents of elementary particles and has the form

$$\begin{aligned} L_{\mu\nu} &= 2\left[k_\mu k'_\nu + k_\nu k'_\mu - g_{\mu\nu}(k\cdot k' - m^2)\right] + 2\,i\,m\epsilon_{\mu\nu\rho\sigma}q^\rho s^\sigma \\ &\equiv L^S_{\mu\nu} + i\,L^A_{\mu\nu}. \end{aligned} \tag{1.2}$$

The part of $L_{\mu\nu}$ symmetric with respect to the interchange $\mu \leftrightarrow \nu$ is denoted by $L^S_{\mu\nu}$ and the antisymmetric part by $L^A_{\mu\nu}$. Note that only the antisymmetric part contains the electron polarization s^μ. The tensor $W_{\mu\nu}$ is the Fourier transform of the correlator of unknown hadronic currents,

$$W_{\mu\nu} = \frac{1}{2\pi M}\int d^4x\; e^{iq\cdot x}\langle P,S|\,[J_\mu(x),J_\nu(0)]\,|P,S\rangle. \tag{1.3}$$

We can use Lorentz invariance and impose the symmetry requirements of time reversal invariance, parity and current conservation, and hermiticity to parametrize it in terms of the structure functions F_1, F_2, g_1 and g_2. Decomposing $W_{\mu\nu}$ into symmetric and antisymmetric tensors,

$$W_{\mu\nu} = W^S_{\mu\nu} + i\,W^A_{\mu\nu}, \tag{1.4}$$

we obtain

$$W^S_{\mu\nu} = (\frac{q_\mu q_\nu}{q^2} - g_{\mu\nu})\frac{F_1(x,Q^2)}{M} + \frac{1}{M^2}(P_\mu - \frac{P\cdot q}{q^2}q_\mu)(P_\nu - \frac{P\cdot q}{q^2}q_\nu)\frac{F_2(x,Q^2)}{\nu}, \tag{1.5}$$

and

$$W^A_{\mu\nu} = M\epsilon_{\mu\nu\rho\sigma}q^\rho S^\sigma\frac{g_1(x,Q^2)}{M^2\nu} + \frac{1}{M}\epsilon_{\mu\nu\rho\sigma}q^\rho(P\cdot qS^\sigma - S\cdot qP^\sigma)\frac{g_2(x,Q^2)}{M\nu^2}. \tag{1.6}$$

We note again that only the antisymmetric part contains the proton polarization S^μ. Since

$$L_{\mu\nu}W^{\mu\nu} = L^S_{\mu\nu}W^{S\mu\nu} - L^A_{\mu\nu}W^{A\mu\nu}, \tag{1.7}$$

we need to polarize both the electron beam and the nucleon target to measure the spin structure functions g_1 and g_2.

Let us now consider longitudinally polarized electrons along ($\uparrow$) or opposite ($\downarrow$) to the direction of the beam, with the nucleon target polarized longitudinally ($\Uparrow$, $\Downarrow$) or transversely ($\Rightarrow$) to the beam direction. The differential cross sections have the form

$$\frac{d^2\sigma^{\downarrow\Uparrow}}{d\Omega dE'} + \frac{d^2\sigma^{\uparrow\Uparrow}}{d\Omega dE'} = \frac{8\alpha^2}{Q^4}E'^2\left[2\sin^2(\frac{\theta}{2})\frac{F_1(x,Q^2)}{M} + \cos^2(\frac{\theta}{2})\frac{F_2(x,Q^2)}{\nu}\right] \quad (1.8)$$

and

$$\frac{d^2\sigma^{\downarrow\Uparrow}}{d\Omega dE'} - \frac{d^2\sigma^{\uparrow\Uparrow}}{d\Omega dE'} = \frac{4\alpha^2}{Q^2}\frac{E'}{E\nu}\left[\frac{(E + E'\cos\theta)}{M}\; g_1(x,Q^2) - 2x\; g_2(x,Q^2)\right] \quad (1.9)$$

for the longitudinally polarized electrons and target and

$$\frac{d^2\sigma^{\downarrow\Rightarrow}}{d\Omega dE'} - \frac{d^2\sigma^{\uparrow\Rightarrow}}{d\Omega dE'} = \frac{4\alpha^2}{Q^2}\frac{E'^2}{E}\frac{\sin\theta}{M\nu}\left[\; g_1(x,Q^2) + 2\frac{E}{\nu}\; g_2(x,Q^2)\right] \quad (1.10)$$

for the longitudinally polarized electrons and transversely polarized target. The longitudinal asymmetry, see eq. (1.9), is dominated by the structure function g_1, as g_2 is suppressed by a factor $\sim \frac{M}{E} \approx 0.02$ for a $E = 50$ GeV beam. On the other hand the transverse asymmetry, see eq. (1.10), is primarily sensitive to g_2.

In polarized experiments one usually measures asymmetries rather than cross sections. The raw asymmetries are computed from the detector rates according to

$$A_{\parallel}^{raw} = \frac{(N/Q)^{\downarrow\Uparrow} - (N/Q)^{\uparrow\Uparrow}}{(N/Q)^{\downarrow\Uparrow} + (N/Q)^{\uparrow\Uparrow}} \quad (1.11)$$

and

$$A_{\perp}^{raw} = \frac{(N/Q)^{\downarrow\Rightarrow} - (N/Q)^{\uparrow\Rightarrow}}{(N/Q)^{\downarrow\Rightarrow} + (N/Q)^{\uparrow\Rightarrow},} \quad (1.12)$$

where N is the detected number of electrons normalized by the total charge of the beam Q in a given electron-target spin configuration. They are simply related to the cross section asymmetries

$$A_{\parallel} = \frac{\left(\frac{d^2\sigma}{d\Omega dE'}\right)^{\downarrow\Uparrow} - \left(\frac{d^2\sigma}{d\Omega dE'}\right)^{\uparrow\Uparrow}}{\left(\frac{d^2\sigma}{d\Omega dE'}\right)^{\downarrow\Uparrow} + \left(\frac{d^2\sigma}{d\Omega dE'}\right)^{\uparrow\Uparrow}}, \tag{1.13}$$

and

$$A_{\perp} = \frac{\left(\frac{d^2\sigma}{d\Omega dE'}\right)^{\downarrow\Rightarrow} - \left(\frac{d^2\sigma}{d\Omega dE'}\right)^{\uparrow\Rightarrow}}{\left(\frac{d^2\sigma}{d\Omega dE'}\right)^{\downarrow\Rightarrow} + \left(\frac{d^2\sigma}{d\Omega dE'}\right)^{\uparrow\Rightarrow}}, \tag{1.14}$$

by correcting for the beam polarization P_b target polarization P_t and the dilution factor f, which is the fraction of events generated by polarizable target material under consideration (^{3}He in our experiment),

$$A_{\parallel} = \frac{A_{\parallel}^{raw}}{P_b P_t f}, \tag{1.15}$$

$$A_{\perp} = \frac{A_{\perp}^{raw}}{P_b P_t f}. \tag{1.16}$$

The structure functions g_1 and g_2, eqs. (1.8)-(1.10), can now be written in terms of parallel and transverse asymmetries

$$g_1(x,Q^2) = \frac{F_1(x,Q^2)}{D'}\left[A_{\parallel}(x,Q^2) + \tan\left(\frac{\theta}{2}\right) A_{\perp}(x,Q^2)\right], \tag{1.17}$$

$$g_2(x,Q^2) = \frac{F_1(x,Q^2)y}{2D'\sin\theta}\left[-\sin\theta A_{\parallel}(x,Q^2) + \frac{E+E'\cos\theta}{E'}A_{\perp}(x,Q^2)\right]. \tag{1.18}$$

The various kinematic factors are defined below:

$$D' = \frac{(1-\epsilon)(2-y)}{y(1+\epsilon R)}, \tag{1.19}$$

$$\epsilon = \frac{1}{1+2(1+(\nu^2/Q^2))\tan^2(\theta/2)}, \tag{1.20}$$

$$y = \frac{E-E'}{E}, \tag{1.21}$$

$$R(x,Q^2) = \frac{1+\gamma^2}{2xF_1(x,Q^2)} F_2(x,Q^2) - 1, \tag{1.22}$$

$$\gamma = \sqrt{\frac{Q^2}{\nu^2}}. \tag{1.23}$$

Note that the above expressions are given in terms of measurable quantities, assuming that $F_2(x,Q^2)$ [124] and $R(x,Q^2)$ [24, 25] are known from other experiments.

1.3 Virtual photon-nucleon asymmetries

The hadronic tensor $W_{\mu\nu}$ given by eq. (1.3) is also known as the virtual forward Compton amplitude, since its contraction with the virtual photon polarization four-vector ϵ^μ gives the forward amplitude $\mathcal{M}$ for virtual photon-proton scattering:

$$\mathcal{M}(\gamma^* p \to \gamma^* p) = \epsilon^*_\mu \; \epsilon_\nu \; W^{\mu\nu}. \tag{1.24}$$

For the virtual photons, which have three polarization states, and nucleons, which have two polarizations there are four independent virtual Compton amplitudes [23, 26] (just as there are four independent structure functions), $\mathcal{M}_{ab\to cd}$, where a, b represent the incident helicities of the virtual photon and nucleon, and c, d the scattered helicities, respectively. We have chosen the following independent amplitudes:

$$\mathcal{M}_{1-\frac{1}{2}\to 1-\frac{1}{2}}, \quad \mathcal{M}_{1\frac{1}{2}\to 1\frac{1}{2}}, \quad \mathcal{M}_{0\frac{1}{2}\to 0\frac{1}{2}}, \quad \mathcal{M}_{1-\frac{1}{2}\to 0\frac{1}{2}}. \tag{1.25}$$

Using those amplitudes ,we can obtain, via the optical theorem, the total absorption cross sections of the virtual photon by the nucleon target,

$$
\begin{aligned}
\sigma_{1/2} &= \frac{4\pi^2\alpha}{K}\mathcal{M}_{1-\frac{1}{2}\to 1-\frac{1}{2}} = \frac{4\pi^2\alpha}{MK}\left[F_1 + g_1 - \frac{2Mx}{\nu}g_2\right], && (1.26)\\
\sigma_{3/2} &= \frac{4\pi^2\alpha}{K}\mathcal{M}_{1\frac{1}{2}\to 1\frac{1}{2}} = \frac{4\pi^2\alpha}{MK}\left[F_1 - g_1 + \frac{2Mx}{\nu}g_2\right], && (1.27)\\
\sigma_L &= \frac{4\pi^2\alpha}{K}\mathcal{M}_{0\frac{1}{2}\to 0\frac{1}{2}} = \frac{4\pi^2\alpha}{K}\left[\frac{F_2}{\nu}(1 + \frac{\nu^2}{Q^2}) - \frac{F_1}{M}\right], && (1.28)\\
\sigma_{TL} &= \frac{4\pi^2\alpha}{K}\mathcal{M}_{1-\frac{1}{2}\to 0\frac{1}{2}} = \frac{4\pi^2\alpha}{K}\frac{\sqrt{Q^2}}{M\nu}\left[g_1 + g_2\right], && (1.29)
\end{aligned}
$$

where K represents the incoming photon flux. Using Hand's convention for K we have,

$$K \equiv \frac{W^2 - M^2}{2M} = \nu - \frac{Q^2}{2M}. \tag{1.30}$$

The quantities $\sigma_{1/2}$ and $\sigma_{3/2}$ are the virtual photo-absorption cross sections with the projection of the spin of the photon-nucleon system along the virtual photon axis equal to 1/2 and 3/2, respectively. The total transverse cross section σ_T is defined as

$$\sigma_T \equiv \frac{1}{2}\left(\sigma_{1/2} + \sigma_{3/2}\right) = \frac{4\pi^2\alpha}{MK}\ F_1(x, Q^2). \tag{1.31}$$

The cross section σ_L corresponds to the longitudinally polarized photon, while σ_{TL} comes from the interference between transverse and longitudinal amplitudes.

The virtual photon-nucleon asymmetries, A_1 and A_2, can now be defined in terms of the cross sections and related to the structure functions,

$$A_1(x, Q^2) \equiv \frac{\sigma_{1/2} - \sigma_{3/2}}{\sigma_{1/2} + \sigma_{3/2}} = \frac{g_1(x, Q^2) - \gamma^2 g_2(x, Q^2)}{F_1(x, Q^2)}, \tag{1.32}$$

and

$$A_2(x,Q^2) \equiv \frac{\sigma_{TL}}{\sigma_T} = \frac{\gamma(g_1(x,Q^2)+g_2(x,Q^2))}{F_1(x,Q^2)}, \tag{1.33}$$

where γ has been defined previously, eq. (1.23). Combining the above equations with formulas (1.17), (1.18) for g_1 and g_2, we can also write directly in terms of measurable longitudinal and transverse asymmetries,

$$\begin{aligned} A_1(x,Q^2) &= \frac{1}{D(1+\zeta\eta)}A_{||}(x,Q^2) - \frac{\eta}{d(1+\zeta\eta)}A_{\perp}(x,Q^2), & (1.34)\\ A_2(x,Q^2) &= \frac{\zeta}{D(1+\zeta\eta)}A_{||}(x,Q^2) + \frac{1}{d(1+\zeta\eta)}A_{\perp}(x,Q^2), & (1.35)\end{aligned}$$

where in addition to the previously defined kinematic factors, eq. (1.19-1.23),

$$\begin{aligned} D &= \frac{1-E'\epsilon/E}{1+\epsilon R}, & (1.36)\\ d &= D\sqrt{\frac{2\epsilon}{1+\epsilon}}, & (1.37)\\ \eta &= \frac{\epsilon\sqrt{Q^2}}{E-E'\epsilon}, & (1.38)\\ \zeta &= \eta\frac{1+\epsilon}{2\epsilon}. & (1.39)\end{aligned}$$

We can also interprate $R(x,Q^2)$ as the ratio of the longitudinal and transverse cross sections,

$$R(x,Q^2) \equiv \frac{\sigma_L}{\sigma_T} = \frac{1+\gamma^2}{2xF_1(x,Q^2)}F_2(x,Q^2) - 1, \tag{1.40}$$

that relates two unpolarized structure functions,

$$F_1(x,Q^2) = \frac{1+\gamma^2}{2x(1+R(x,Q^2))}\ F_2(x,Q^2). \tag{1.41}$$

Note that definition (1.32) for A_1 gives the positivity (of cross sections) constraint,

$$|A_1(x,Q^2)| \ \leq\ 1, \tag{1.42}$$

and the definition (1.33) of A_2 combined with the Schwartz inequality

$$\|v \cdot w\|^2 \;\leq\; \|v\|^2 \, \|w\|^2$$

in the form of $\sigma_{TL}^2 \leq \sigma_T \, \sigma_L$ and the definition (1.40) of R gives the upper bound on A_2,

$$|A_2(x, Q^2)| \;\leq\; \sqrt{R(x, Q^2)}. \tag{1.43}$$

1.4 Models of the He^3 nucleus

In our experiment we measure the asymmetries of the polarized ^{3}He. In the simplest model the nucleons in the polarized ^{3}He are in a spatially symmetric S state. Due to the Pauli principle the total wave-function must be antisymmetric, and thus the two protons have opposite spins. In this picture the spin of the ^{3}He nucleus is carried exclusively by the neutron and the neutron asymmetries are directly proportional to those of ^{3}He.

A more realistic approach [27] includes other components of the wave-function. The dominant componenets are S' and the average of the D state components. The S' component originates in the small difference between the tensor T=0 and T=1 forces distorting the primary S wave configuration. The P wave contributions are suppressed since they have opposite parity.

In the S wave approximation the probability P_n^+ that the neutron spin is aligned with the spin of ^{3}He equals 1, while the probability P_n^- for the anti-aligned spin is 0. For the proton $P_p^+ = P_p^- = 1/2$. Including other waves contributions we write

$$P_n^+ \;=\; 1 - \Delta,$$

$$
\begin{aligned}
P_n^- &= \Delta, && (1.44)\\
P_p^+ &= \frac{1}{2} - \delta, &&\\
P_n^- &= \frac{1}{2} + \delta, && (1.45)
\end{aligned}
$$

where $\Delta = [P(S') + 2P(D)]/3$ and $\delta = [P(D) - P(S')]/6$ and $P(S')$, $P(D)$ are the probabilities of each partial wave component. The polarization of the neutron is $p_n = 1 - 2\,\Delta$ and of the proton $p_p = -2\,\delta$. The best fit to nucleon polarizations were extracted from many models [27] yielding $\Delta = 0.07 \pm 0.01$ and $\delta = 0.014 \pm 0.002$, and thus polarizations $p_n = 0.86 \pm 0.02$ and $p_p = -0.027 \pm 0.004$. This leads to the relation between asymmetries [28]

$$
A_{\|(\perp)}^{^3He} = f_n\, p_n\, A_{\|(\perp)}^n + 2\, f_p\, p_p\, A_{\|(\perp)}^p, \tag{1.46}
$$

where $f_{n(p)} = F_2^{n(p)}/[(F_2^n + 2F_2^p)\, f_{EMC}]$ with the EMC effect factor f_{EMC} [29], and between structure functions

$$
g_1^{^3He}(x) = p_n\, g_1^n(x) + 2\, p_p\, g_1^p(x). \tag{1.47}
$$

The above results neglect nucleon binding effects, Fermi motion and shadowing. The more complete calculations [28] suggest that these corrections are only important at $x > 0.9$ or very low $x \sim 0$.

1.5 Partonic interpretation

In the parton model of Bjorken and Paschos [5], and Feynman [6] a nucleon consists of pointlike constituents, partons. Each parton carries a fraction ξ of the total nucleon momentum P. When the nucleon is viewed from the infinite momentum frame [30], one argues, ignoring the transverse motion of constituents, that the interaction between partons is time dilated. If the probing momentum (and energy) of the virtual photon is large, i.e. in the Bjorken limit, the virtual photon-parton interaction is short in comparison with parton-parton interactions. The inclusive scattering may be considered as incoherent elastic scattering off effectively free pointlike constituents. For the scattered parton of mass m_q we write

$$(\xi P + q)^2 = m_q^2. \tag{1.48}$$

In the limit of large momentum transfer compared to the parton and nucleon masses,

$$\xi = \frac{Q^2}{2\,M\,\nu} \equiv x. \tag{1.49}$$

The virtual photon with x defined by kinematics only scatters off the parton with momentum fraction $\xi = x$. Or, equivalently, the Bjorken variable x is the fraction of the nucleon momentum carried by the struck quark.

The remarkable feature of this model is Bjorken scaling. We treat the parton as a spin 1/2 Dirac particle, the quark, and calculate structure functions for scattering off such a particle. We then sum incoherently over contributions from all single quarks weighted by the probabilities $q^{\uparrow(\downarrow)}(x)$ ($\bar{q}^{\uparrow(\downarrow)}(x)$) of finding a quark (antiquark) with a spin direction the same ($\uparrow$) or opposite ($\downarrow$) to the spin of the nucleon, carrying the fraction x of the nucleon momentum. Taking the Bjorken limit

$$Q^2 \to \infty,\ \nu \to \infty \quad \text{with} \quad x = \frac{Q^2}{2M\nu} = \text{const}, \tag{1.50}$$

of the structure functions, one finds the Bjorken scaling:

$$f(x,Q^2) \quad \xrightarrow{Bj} \quad f(x), \tag{1.51}$$

where f stands for F_1, F_2, g_1 or g_2. The structure functions only depend on one variable x. This x dependence is defined by the quark probability distributions,

$$q(x) \;\; \equiv \;\; \left[q^{\uparrow}(x)+\bar{q}^{\uparrow}(x)\right] \;\; + \;\; \left[q^{\downarrow}(x)+\bar{q}^{\downarrow}(x)\right], \tag{1.52}$$

$$\Delta q(x) \;\; \equiv \;\; \left[q^{\uparrow}(x)+\bar{q}^{\uparrow}(x)\right] \;\; - \;\; \left[q^{\downarrow}(x)+\bar{q}^{\downarrow}(x)\right]. \tag{1.53}$$

Namely,

$$F_1(x) = \frac{1}{2}\sum_f e_f^2 \; q_f(x), \tag{1.54}$$

and

$$g_1(x) = \frac{1}{2}\sum_f e_f^2 \; \Delta q_f(x), \tag{1.55}$$

where the sum is over all quark flavors f, and e_f denotes the quark charge.

Furthermore, in the Bjorken limit, the Callan-Gross [31] relation

$$F_2(x) \;\; = \;\; 2\,x\,F_1(x), \tag{1.56}$$

holds. When substituted into the definition of $R(x,Q^2)$ in eq. (1.40) taken in the Bjorken limit, ($\gamma \xrightarrow{Bj} 0$, eq. (1.23)), it gives

$$R(x,Q^2) \;\; = \;\; 0, \tag{1.57}$$

that is

$$\sigma_L(x,Q^2) \;\; = \;\; 0, \tag{1.58}$$

as seen from eq. (1.28). Also, see eq. (1.29),

$$\sigma_{TL}(x, Q^2) \; = \; 0. \tag{1.59}$$

The last equation implies that from the definition (1.33)

$$A_2(x, Q^2) \; = \; 0. \tag{1.60}$$

It is instructive to look at the Bjorken limit of the virtual photon asymmetry A_1. From the definition (1.32) and the partonic expressions on g_1 and F_1 we get

$$A_1(x, Q^2) = \frac{g_1(x, Q^2)}{F_1(x, Q^2)} = \frac{\sum_f e_f^2 \; \Delta q_f(x)}{\sum_f e_f^2 \; q_f(x)}. \tag{1.61}$$

The last equality can also be inferred from the following considerations [26]: Let the nucleon $P^\uparrow$ with the projection $J_z = +\frac{1}{2}$ collide with a transversely polarized photon $\gamma^{\uparrow(\downarrow)}$ with $J_z = +1\,(\uparrow)$ or $-1\,(\downarrow)$. The total spin angular momentum is $J_z = \frac{3}{2}$ or $\frac{1}{2}$:

$$\gamma^\uparrow + P^\uparrow \quad \to \quad \sigma_{3/2}, \tag{1.62}$$

$$\gamma^\downarrow + P^\uparrow \quad \to \quad \sigma_{1/2}. \tag{1.63}$$

The photon-quark interaction is collinear (neglecting the transverse motion of parton) so that the projection of orbital angular momentum is zero. Hence, from angular momentum conservation, the transverse photon will flip the spin of the quark with which it interacts,

$$\gamma^\uparrow + q^\downarrow \quad \to \quad q^\uparrow, \tag{1.64}$$

$$\gamma^\downarrow + q^\uparrow \quad \to \quad q^\downarrow, \tag{1.65}$$

but $\gamma^\uparrow q^\uparrow$ and $\gamma^\downarrow q^\downarrow$ are suppressed. So we see that

$$\sigma_{3/2} \sim \gamma^\uparrow P^\uparrow \sim \sum_f e_f^2 \, q_f^\downarrow, \tag{1.66}$$

$$\sigma_{1/2} \sim \gamma^\downarrow P^\uparrow \sim \sum_f e_f^2 \, q_f^\uparrow, \tag{1.67}$$

leading via definition (1.32) of $A_1 = \frac{\sigma_{1/2}-\sigma_{3/2}}{\sigma_{1/2}+\sigma_{3/2}}$ to equation (1.61).

As an example [32] of $A_1(x)$ asymmetry let us consider the SU(6) symmetric constituent quark model. In this model the proton consists of three valence quarks. The spin-flavor part of its wave-function can be written as

$$P^\uparrow = \frac{1}{\sqrt{6}} (2 \ \uparrow\uparrow\downarrow - \uparrow\downarrow\uparrow - \downarrow\uparrow\uparrow) \ \mathrm{uud}, \tag{1.68}$$

in the uds basis. The momentum part of the wave-function is totally symmetric, implying that the x dependence of partonic distributions, denoted by $p(x)$, is the same. The probability distribution for finding a down quark with the spin parallel to the proton spin can be obtained from the spin-flavor composition of the proton in eq. (1.68) as $d^\uparrow(x) = \frac{1}{6}[(-1)^2 + (-1)^2] \, p(x) = \frac{1}{3} \, p(x)$, where $p(x)$ is normalized to $1 = \int_0^1 dx \, p(x)$, and the probability of finding a quark with any flavor and spin is normalized to the number of quarks (to 3 in our case). Similarly, we get

$$u^\uparrow(x) = \frac{5}{3} \, p(x), \tag{1.69}$$

$$u^\downarrow(x) = \frac{1}{3} \, p(x), \tag{1.70}$$

$$d^\uparrow(x) = \frac{1}{3} \, p(x), \tag{1.71}$$

$$d^\downarrow(x) = \frac{2}{3} \, p(x). \tag{1.72}$$

Substituting these distributions into (1.61) we find

$$A_1^p(x) = \frac{\frac{4}{9}(u^{\uparrow}(x) - u^{\downarrow}(x)) + \frac{1}{9}(d^{\uparrow}(x) - d^{\downarrow}(x))}{\frac{4}{9}(u^{\uparrow}(x) + u^{\downarrow}(x)) + \frac{1}{9}(d^{\uparrow}(x) + d^{\downarrow}(x))} = \frac{5}{9} \tag{1.73}$$

for the proton. The neutron asymmetry is related to the proton asymmetry via isospin exchange $u \leftrightarrow d$, so

$$A_1^n(x) = 0. \tag{1.74}$$

In a more realistic model with SU(6) breaking admixtures [33] one obtains a nontrivial x dependence of the nucleon asymmetries [34].

The total spin $\Delta\Sigma$ carried by the quarks can be defined as the sum of the net spin of all contributing flavors:

$$\Delta\Sigma \;=\; \sum_f \int_0^1 dx \; \Delta q_f(x) \;\equiv\; \sum_f \Delta q_f. \tag{1.75}$$

In the three flavor model

$$\Delta\Sigma = \Delta u + \Delta d + \Delta s. \tag{1.76}$$

For the nucleon, with the values of $\Delta q_f(x)$ from the SU(6) model given in eqs. (1.69 - 1.72) we obtain

$$\Delta\Sigma = 1. \tag{1.77}$$

The spin of the nucleon is entirely carried by constituent quarks.

1.6 QCD improved parton model

The quark-parton model considered in the previous section can be formulated more rigorously on the basis of QCD. The main modifications come from the radiative corrections. Quarks within a nucleon can emit gluons, which in turn can produce $q\bar{q}$ pairs. These processes lead to scaling violations logarithmic in Q^2. The structure functions and the partonic distributions are no longer functions of just the x variable. They acquire additional Q^2 dependence.

The g_1 structure function, for example, can still (in leading order) be written in the partonic form of eq. (1.55)

$$g_1(x, Q^2) = \frac{1}{2} \sum_f e_f^2 \; \Delta q_f(x, Q^2). \tag{1.78}$$

That the structure function, directly related to the cross section, can be written in terms of partonic distributions and some coefficients is a manifestation of a more general property, the factorization of DIS cross sections (see section 4.1, page 177).

The Q^2 dependence of the partonic distributions is described by the Altarelli-Parisi equations (cf. section 4.2, page 179),

$$\begin{aligned} Q^2 \frac{d}{dQ^2} \; \Delta q_f(x, Q^2) &= \frac{\alpha_s(Q^2)}{2\pi} \left[P_{qq} \otimes \Delta q_f + P_{qG} \otimes \Delta G \right], && (1.79) \\ Q^2 \frac{d}{dQ^2} \; \Delta G(x, Q^2) &= \frac{\alpha_s(Q^2)}{2\pi} \left[P_{Gq} \otimes \sum_f^{2N_f} \Delta q_f + P_{GG} \otimes \Delta G \right]. && (1.80) \end{aligned}$$

The so called splitting functions $P_{ij}(x)$ are the hard scattering cross sections of the radiative processes. The convolution $\otimes$ is defined as

$$P \otimes \Delta q = \int_x^1 \frac{dy}{y} P\left(\frac{x}{y}\right) \Delta q(y, Q^2). \tag{1.81}$$

The total spin $\Delta q = \int_0^1 dx \Delta q(x)$ carried by the quark can be defined in the operator language as the matrix element of the axial current,

$$\Delta q S^\mu = \langle PS|\bar{q}\gamma^\mu\gamma_5 q|PS\rangle = \langle PS|\bar{q}_R\gamma^\mu q_R - \bar{q}_L\gamma^\mu q_L|PS\rangle, \tag{1.82}$$

where $q_{R,L} \equiv \frac{1}{2}(1\pm\gamma_5)\,q$. With this definition we can decompose the $\mathrm{SU}(3)_{\mathrm{flavor}}$ axial-vector currents j_a^μ, defined as

$$j_a^\mu = \bar{\psi}\gamma^\mu\gamma_5\tau_a\psi. \tag{1.83}$$

Here ψ is the flavor triplet,

$$\psi = \begin{pmatrix} u \\ d \\ s \end{pmatrix}, \tag{1.84}$$

and τ_a are the SU(3) generators. With the flavor indices shown explicitly the currents read,

$$j_a^\mu = \sum_{i,j=1}^{3} \bar{\psi}_i\gamma^\mu\gamma_5(\tau_a)_{ij}\psi_j. \tag{1.85}$$

Similarly to eq. (1.82) let us introduce axial charges Δq_a as appropriate matrix elements of the j_a^μ currents:

$$\Delta q_a S^\mu = \langle PS|\; j_a^\mu \;|PS\rangle. \tag{1.86}$$

Of special interest to DIS are the $a=0$, $\mathrm{SU}(3)_{\mathrm{flavor}}$ singlet and the $a=3,8$ non-singlet axial-vector currents from the octet. With our choice of the normalization of the τ_a generators,

$$\tau_0 = \mathrm{Id}(3), \qquad \tau_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \qquad \tau_8 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix}, \tag{1.87}$$

and with the help of eqs. (1.82-1.86) we can rewrite the corresponding charges in a more familiar form,

$$\Delta q_0 = \Delta u + \Delta d + \Delta s \equiv \Delta\Sigma, \tag{1.88}$$

$$\Delta q_3 = \Delta u - \Delta d, \tag{1.89}$$

$$\Delta q_8 = \Delta u + \Delta d - 2\,\Delta s. \tag{1.90}$$

The octet currents j_3^μ and j_8^μ are conserved. Thus the associated charges Δq_3 and Δq_8 are Q^2 independent. The singlet axial current is conserved in the leading order only. In general it acquires an anomalous contribution. The classical QCD lagrangian for massless quarks is invariant under the global U(1) axial (chiral) transformation

$$\psi \to \psi' = e^{-i\theta\gamma_5}\,\psi. \tag{1.91}$$

The singlet axial current is a Noether current arising from this symmetry. Thus on the classical level it is conserved and

$$\partial^\mu j_\mu^0 = 0. \tag{1.92}$$

However, in the full quantum theory the axial U(1) transformation is not a symmetry. The current divergence has an anomaly [35] , which arises from triangular diagrams of quantum corrections,

$$\partial^\mu j_\mu^0 = \frac{N_f \alpha_s}{8\pi} F_{\mu\nu}^a \widetilde{F}^{a\mu\nu}, \tag{1.93}$$

where $\widetilde{F}^a$ is a dual to F^a, i.e. $\widetilde{F}^{a\mu\nu} = \epsilon^{\mu\nu\rho\sigma} F_{\rho\sigma}^a$, and N_f is the number of flavors. In the path integral formulation of the quantum theory one sees that the path integral measure is not invariant under the U(1) transformation although the lagrangian is [36].

One can still construct a divergence-free axial current by noting that the anomalous contribution on the right hand side of eq. (1.93) can be written as the divergence of an axial gluon current, K^μ [37],

$$\partial_\mu K^\mu = \frac{1}{4} F^a_{\mu\nu} \widetilde{F}^{a\mu\nu}. \tag{1.94}$$

We can now redefine j^0_μ,

$$\widetilde{j}^0_\mu = j^0_\mu - N_f \frac{\alpha_s}{2\pi} K_\mu, \tag{1.95}$$

so that the new current is conserved,

$$\partial^\mu \widetilde{j}^0_\mu = 0. \tag{1.96}$$

Note that the axial gluon current is not gauge invariant. In the lightcone gauge $A^+ = 0$, its expectation value in the polarized nucleon is identified with the gluon spin contribution [38],

$$\Delta G\ S^\mu = -\ \langle PS|K^\mu|PS\rangle. \tag{1.97}$$

The conserved charge, $\Delta\widetilde{q}_0$, differs from the old charge by the anomalous gluon contribution,

$$\Delta\widetilde{q}_0 = \Delta q_0 + N_f \frac{\alpha_s}{2\pi} \Delta G. \tag{1.98}$$

One can argue that $\Delta\widetilde{q}_0$ should be identified with the total quark helicity,

$$\Delta\widetilde{\Sigma} = \Delta\Sigma + N_f \frac{\alpha_s}{2\pi} \Delta G. \tag{1.99}$$

This ambiguity is a manifestation of factorization scheme dependence, which will be treated in some detail in section 4.6 (page 197). We note that $\alpha_s(Q^2)\ \Delta G(Q^2) \approx constant$, so the ambiguity persists, even in the large Q^2 limit.

There is one more source of the scaling violation of structure functions in addition to the Q^2 evolution. It is referred to as higher twist corrections and gives contributions which are suppressed by powers of Q^2 with respect to the leading twist. They could be interpreted [39] as coming from the diagrams involving interaction between different quarks or from quark-gluon correlations. These corrections are expected to be important at low values of W^2 for high x and low values of Q^2 for low x. For example, for the F_2 structure function we can write

$$F_2(x, Q^2) = F_2^{lt}(x, Q^2)\ \left[1 + \sum_{m=1} \frac{D_m}{(Q^2(1-x))^m}\right], \tag{1.100}$$

where F^{lt} denotes the leading twist part. The unpolarized data from SLAC [24, 25] show sizeble higher twist contributions at high x. Thus we conclude that higher twist contributions to the polarized structure functions are important at low W^2.

1.7 Sum rules

1.7.1 Helicity sum rule

Since the nucleon is a spin 1/2 particle, $|PS\rangle$ is an eigenstate of the Pauli-Lubanski vector operator $W_\mu = \epsilon_{\mu\nu\rho\sigma} J^{\nu\rho} P^\sigma/(2M)$ projected onto the polarization S^μ,

$$S^\mu W_\mu \; |PS\rangle = \frac{1}{2} \; |PS\rangle, \tag{1.101}$$

or equivalently,

$$\frac{1}{2} = \langle PS| \; S^\mu W_\mu \; |PS\rangle. \tag{1.102}$$

For $\vec{S}$ along the direction of $\vec{P}$ (chosen to be the z axis) the nucleon is in the 1/2 helicity eigenstate, and

$$S^\mu W_\mu \;= J^z, \tag{1.103}$$

so one can construct a helicity sum rule,

$$\frac{1}{2} = \langle PS| \; J^z \; |PS\rangle. \tag{1.104}$$

To be more specific we need to consider the transformation properties of the QCD lagrangian under spatial rotations and construct the generators of rotations, i.e. the angular momentum operator [40]. In a shorthand notation the angular momentum operator can be written as

$$\vec{J} = \int d^3x \left[\frac{1}{2}\bar{\psi}\vec{\gamma}\gamma_5\psi + \psi^\dagger(\vec{x} \times (-i\vec{\nabla}))\psi + \vec{E} \times \vec{A} + E_i(\vec{x} \times \vec{\nabla})A_i\right]. \tag{1.105}$$

The physical interpretation of the above equation is obvious. The first term is the quark spin contribution to the nucleon helicity, the second is the quark orbital angular momentum, the third is the gluon spin and the last term is the gluon orbital momentum. Taking the expectation value of J^z as in eq. (1.104) and choosing the light cone gauge ($A^+ = 0$) in the infinite momentum frame, we obtain the sum rule

$$\frac{1}{2} = \frac{1}{2}\Delta\Sigma(Q^2) + L_q(Q^2) + \Delta G(Q^2) + L_G(Q^2) \tag{1.106}$$

for the nucleon spin. It is evident from the decomposition (1.105) that L_q, ΔG and L_G are gauge dependent. They are also scale dependent. One can write the evolution equations for those quantities [41, 42] and find the asymptotic, $Q^2 \to \infty$, behavior

$$\begin{aligned}
\Delta\Sigma &\to \text{const.} \\
L_q &\to \text{const.} \\
\Delta G &\to \quad \lambda \log(Q^2) + \text{const.} \\
L_G &\to -\lambda \log(Q^2) + \text{const.}
\end{aligned} \tag{1.107}$$

The asymptotic increase of gluon helicity is canceled by the gluon orbital contribution. Recently, [43], an explicitly gauge invariant decomposition of the angular momentum operator

$$\vec{J} = \int d^3x \left[\frac{1}{2}\bar{\psi}\vec{\gamma}\gamma_5\psi + \psi^\dagger(\vec{x}\times(-i\vec{D}))\psi + \vec{x}\times(\vec{E}\times\vec{B}) \right], \tag{1.108}$$

has been proposed, leading to a gauge invariant spin sum rule,

$$\frac{1}{2} = (\frac{1}{2}\Delta\Sigma(Q^2) + L'_q(Q^2)) + J_G(Q^2), \tag{1.109}$$

or

$$\frac{1}{2} = J_q(Q^2) + J_G(Q^2). \tag{1.110}$$

There are two additional interesting features of the above sum rule [42, 44]. Study of the Q^2 dependence of J_q and J_G reveals that the anomalous dimension matrix for these two operators has a zero eigenvalue. This implies that in the large Q^2 limit, where all other eigenvectors of the anomalous dimension matrix have evolved to zero, the ratio of quark and gluon contributions goes to a definite limit, independent of hadronic target. There is an analogous result for the momentum sum rule [45] which gives the same fractions,

$$\begin{aligned} J_q(Q^2 \to \infty) &\to \frac{1}{2}\,\frac{3N_f}{16+3N_f}, \\ J_G(Q^2 \to \infty) &\to \frac{1}{2}\,\frac{16}{16+3N_f}. \end{aligned} \tag{1.111}$$

Study of the J_q in a factorization scheme in which $\Delta\Sigma$ has an explicit triangle anomaly added to it, $\Delta\Sigma + N_f \frac{\alpha}{2\pi}\Delta G = \Delta\widetilde{\Sigma}$, shows that at the same time the quark orbital angular momentum has a compensating gluonic contamination,

$$\begin{aligned} \widetilde{J}_q &= \frac{1}{2}\Delta\widetilde{\Sigma} + (L_q - N_f \frac{\alpha}{4\pi}\Delta G) \\ &= \frac{1}{2}\Delta\Sigma + L_q = J_q. \end{aligned} \tag{1.112}$$

Thus the net effect of the factorization scheme dependence is to shift a contribution between the quark spin and orbital angular momentum, leaving J_q unchanged.

1.7.2 Bjorken sum rule

The Bjorken sum rule [46] follows from isospin symmetry and the short distance analysis of QCD. It tests QCD at a very fundamental level. Originally derived in the context of current algebra before the advent of QCD, it connects the low Q^2 physics of the neutron beta decay with the high Q^2 physics of deep inelastic scattering. In the scaling limit it has the form,

$$\int_0^1 dx\ \left(g_1^p(x) - g_1^n(x)\right) = \frac{1}{6}g_A, \tag{1.113}$$

where g_A is the axial coupling constant of the neutron beta decay. Let us briefly sketch the idea underlying the derivation [47]. The neutron beta decay is governed by the matrix element of the isovector axial current,

$$\langle p|\bar{\psi}T^{+}\gamma^{\mu}\gamma_5\psi|n\rangle = S^{\mu}\, g_A, \tag{1.114}$$

where T^{+} is the raising operator of the SU(2) isospin algebra. Substituting $|n\rangle = T^{-}|p\rangle$ and using $T^{+}|p\rangle = 0$ and $[T^{+}, T^{-}] = T^{3}$, we write,

$$S^{\mu}\, g_A = \langle p|\bar{\psi}T^{3}\gamma^{\mu}\gamma_5\psi|p\rangle. \tag{1.115}$$

Now, recalling

$$T^3 \begin{pmatrix} u \\ d \\ s \end{pmatrix} = \begin{pmatrix} u \\ -d \\ 0 \end{pmatrix}, \tag{1.116}$$

we connect the last equation with the helicities of the u and d quarks,

$$S^{\mu}\, g_A = \langle p|\bar{\psi}T^{3}\gamma^{\mu}\gamma_5\psi|p\rangle = \langle p|\bar{u}\gamma^{\mu}\gamma_5 u - \bar{d}\gamma^{\mu}\gamma_5 d|p\rangle = S^{\mu}\,(\Delta u - \Delta d), \tag{1.117}$$

that is

$$\Delta u - \Delta d \equiv \Delta q_3 = g_A = 1.2601 \pm 0.0025. \tag{1.118}$$

To proceed let us express the integrals of $g_1^p(x)$ and $g_1^n(x)$, see eq. (1.55), in terms of the axial charges (1.88-1.90),

$$\Gamma^p \equiv \int_0^1 dx\ g_1^p(x) = \frac{1}{12}\Delta q_3 + \frac{1}{36}\Delta q_8 + \frac{1}{9}\Delta q_0, \tag{1.119}$$

$$\Gamma^n \equiv \int_0^1 dx\ g_1^n(x) = -\frac{1}{12}\Delta q_3 + \frac{1}{36}\Delta q_8 + \frac{1}{9}\Delta q_0, \tag{1.120}$$

and arrive at the final result,

$$\int_0^1 dx\ \left(g_1^p(x) - g_1^n(x)\right) \equiv \frac{1}{6}\Delta q_3 = \frac{1}{6}g_A. \tag{1.121}$$

The right hand side of the last equation has been established from the neutron beta decay in the limit of $Q^2 \to 0$. Fortunately, the left hand side is proportional to Δq_3, which is conserved, and thus can be established at any value of Q^2, e.g. on the scale of DIS experiments. As mentioned earlier, the above formulation of the Bjorken sum rule is valid in the scaling limit. The coefficient $\frac{1}{6}$, called the Wilson coefficient, comes from the factorization (see section 4.1, page 177) of the scattering cross section (and thus structure functions) into soft partonic distributions and hard scattering cross sections (Wilson coefficients). At finite Q^2 the perturbatively calculated coefficient is expressed as a series in the strong coupling constant [48]. This gives the Q^2 dependence to the Bjorken sum rule,

$$\Gamma^p - \Gamma^n = \frac{1}{6} g_A \left[1 - \frac{\alpha_s(Q^2)}{\pi} + C_2 \left(\frac{\alpha_s(Q^2)}{\pi} \right)^2 + C_3 \left(\frac{\alpha_s(Q^2)}{\pi} \right)^3 \right], \quad (1.122)$$

where $C_2 = -3.5833$ and $C_3 = -20.2153$ in the three flavor theory. The corresponding expansion for Γ^p reads [49] up to $O(\alpha_s^2)$,

$$\Gamma^p(Q^2) = \frac{1}{6} \left[(\frac{1}{2}\Delta q_3 + \frac{1}{6}\Delta q_8)(1 - \frac{\alpha_s}{\pi}) + \frac{2}{3}\Delta q_0 (1 - \frac{\alpha_s}{\pi} \frac{33 - 8N_f}{33 - 2N_f}) \right]. \quad (1.123)$$

1.7.3 Ellis-Jaffe sum rule

In addition to the value of Δq_3 and with the assumption of the $\mathrm{SU}(3)_{\mathrm{flavor}}$ symmetry for the octet baryon currents, one obtains Δq_8 from the hyperon beta decays [50],

$$\Delta q_8 = 0.579 \pm 0.025. \quad (1.124)$$

In a simple model the nucleon consists of u and d valence quarks, and perturbative QCD generates quark-antiquark pairs, perhaps not highly correlated with the spin of the nucleon. Furthermore, the greater mass of the strange quark suggests that the strange sea is suppressed in comparison with the light $q\bar{q}$ pairs. Thus it has been conjectured [51] that Δs contribution can be neglected, which implies

$$\Delta s = 0 \;\Rightarrow\; \Delta q_0 = \Delta q_8, \tag{1.125}$$

so that Γ^p and Γ^n can be computed. At $Q^2 = 5$ GeV^2 the values are

$$\Gamma^p \;\equiv\; \int_0^1 dx\; g_1^p(x) = \quad 0.172 \pm 0.003, \tag{1.126}$$

$$\Gamma^n \;\equiv\; \int_0^1 dx\; g_1^n(x) = -0.020 \pm 0.003. \tag{1.127}$$

The results for the integrals of $g_1^p(x)$ and $g_1^n(x)$ with the assumption of eq. (1.125) are known as Ellis-Jaffe sum rules.

We note that from the values of Δq_3 and Δq_8 from beta decays and measuring Γ^p or Γ^n one can extract $\Delta\Sigma$ $(= \Delta q_0)$ or, equivalently, the total spin carried by different flavors, Δu, Δd and Δs.

1.7.4 Sum rules for g_2

The transverse spin structure function g_2 is a higher twist structure function that has been recently measured [53, 54]. The operator product expansion (OPE) analysis reveals that g_2 consists of two pieces: one entirely twist two part [55] g_2^{ww} and the true twist three part $\bar{g}_2$, with

$$g_2(x, Q^2) = g_2^{ww}(x, Q^2) + \bar{g}_2(x, Q^2). \tag{1.128}$$

The function g_2^{ww} is related to thestructure function g_1 by

$$g_2^{ww}(x,Q^2) = -\,g_1(x,Q^2) + \int_x^1 \frac{dy}{y}\, g_1(y,Q^2). \tag{1.129}$$

The part $\bar{g}_2$ measures quark-gluon correlations in the nucleon. Its moments are related to the local operator products such as

$$\int_0^1 dx\, x^2 \bar{g}_2(x,Q^2) \;\propto\; \langle PS|\; \bar{\psi}\widetilde{G}_{\sigma(\mu_1}\gamma_{\mu_2)}\psi\; |PS\rangle. \tag{1.130}$$

The interesting Burkhardt-Cottingham [56] sum rule for g_2 is a consequence of the rotational invariance and some assumptions about the Regge behavior of the Compton amplitude,

$$\int_0^1 dx\;\, g_2(x,Q^2) = 0. \tag{1.131}$$

Its test could provide some insight into the applicability of Regge theory to DIS. Note that Wandzura-Wilczek piece of g_2 obeys the Burkhardt-Cottingham sum rule,

$$\int_0^1 dx\;\, g_2^{ww}(x,Q^2) = 0, \tag{1.132}$$

provided g_1 is sufficiently well behaved when approaching x=0 to allow the exchange of the x and y integrations.

The OPE also provides us with the moment sum rules for g_1 and g_2

$$\begin{aligned}
\int_0^1 dx\, x^n\, g_1(x,Q^2) &= \frac{1}{4}\, a_n & n = 0,2,4,\ldots \\
\int_0^1 dx\, x^n\, g_1(x,Q^2) &= \frac{1}{4}\frac{n}{n+1}(d_n - a_n) & n = 2,4,\ldots
\end{aligned} \tag{1.133}$$

There is no information about the n=0 moment of g_2 i.e. the OPE does not say anything about the Burkhardt-Cottingham sum rule. The knowledge of all the $\{a_n,\, n = 0, 2, 4, \ldots\}$ and all the $\{d_n,\, n = 2, 4, \ldots\}$ does not completely determine $g_2(x, Q^2)$ because the lowest moment is unknown.

1.8 Traditional evaluation of sum rules from measurements

The measurements of the structure functions only cover a limited range in x and, due to spectrometer kinematics, each point x_i is obtained at a different (averaged over the x bin) scale Q_i^2. To evaluate any of the sum rules, one needs to evolve the data to a single value of Q^2 and then extrapolate the result to x=0 and x=1.

1.8.1 Evolution in Q^2

The value of Q^2 to which all the measurements are evolved is arbitrary. However, each experiment has a characteristic scale, namely, the statistically weighted average $\langle Q^2 \rangle$. The evolved values are quoted at this average $\langle Q^2 \rangle$ of the experiment.

The traditional evolution technique is based on the assumption that the ratio $g_1(x, Q^2)/F_1(x, Q^2)$, which is approximately equal to $A_1(x, Q^2)$, scales, i.e. is constant in Q^2. Here the function $F_1(x, Q^2)$ is considered to be known. For the present kinematic range in Q^2 and the precision of the data, this assumption is compatible with experimental evidence. The common wisdom for A_1 scaling is based on the following 'hand waving' argument. We know from perturbative QCD (pQCD) that both structure functions g_1 and F_1

vary with Q^2. If these variations are 'somewhat similar' (if they factor, one should say) then when taking the ratio g_1/F_1 the variations cancel out at least partially, so the g_1/F_1 varies less strongly with Q^2 than g_1 and F_1. The 'somewhat similar evolution' phrase is sometimes misleadingly quantified in terms of the identity of the splitting functions P_{qq} for polarized and unpolarized evolutions in the leading order. In the x region where the valence quarks dominate, it is argued, the variation of g_1 and F_1 is the same (meaning only that the splitting functions are the same). However, because the Altarelli-Parisi equations are integro-differential equations, see eq. (1.79), (1.80), the x dependence of the distributions matters, and 'the same splitting functions' argument is valid only if the x shapes of both structure functions are the same, i.e. $F_1(x, Q_i^2) = const. * g_1(x, Q_i^2)$ at the initial Q_i^2.

For the evolution of the precise data one has to refer directly to the pQCD formalism, as in Chapter 4 (page 176).

1.8.2 The low x extrapolation

The low x extrapolation has traditionally been performed with the assumption of Regge behavior of $g_1(x, Q^2)$. At a constant Q^2 and $x \approx 0$, the $g_1(x, Q^2)$ converges as

$$g_1(x, Q^2) \underset{x \to 0}{\sim} x^{-\alpha}, \tag{1.134}$$

where the leading intercept α, estimated to lay between -0.5 and 0.0, is believed to come from the a_1 meson trajectory [37, 57]. The experiments have assumed that at their Q^2 the few lowest x measurements are in a kinematic region of Regge behavior and have extrapolated data to $x = 0$ with $\alpha = 0.0$. The variation in α has been used to estimate the extrapolation error. There is no theoretical guidance as tp the values of x and Q^2 at which the Regge behavior sets in. Furthermore there are also other possibilities for the low x behavior, even within the Regge theory. We come back to that subject in more detail in section 4.3.2 (page 187).

1.8.3 The high x extrapolation

The high x behavior used for extrapolations seems to be much less controversial, perhaps mainly due to the fact that the structure functions decrease rapidly with x and the extrapolation contribution to the integral is typically small. From QCD arguments (see section 4.3.1, page 182 for details) formulated in terms of the counting rules [58], we expect the fall off

$$g_1(x) \underset{x \to 1}{\sim} (1-x)^3, \tag{1.135}$$

or stronger, due to the QCD evolution. The same considerations lead to the conclusion that the quark that carry almost all of the nucleon momentum also carry its spin, the opposite helicity being suppressed by another $(1-x)^2$. Thus at $x \sim 1$ F_1 is dominated by the quark distribution $q(x) = q^{\uparrow}(x)$, which is also dominant in g_1, $\Delta q(x) = q^{\uparrow}(x)$, and

$$A_1 \sim \frac{\Delta q(x)}{q(x)} \underset{x \to 1}{\sim} \frac{q^{\uparrow}(x)}{q^{\uparrow}(x)} = 1. \tag{1.136}$$

Both extrapolations, eq. (1.135) and (1.136), have been used and have given consistent results.

1.9 Summary of polarized DIS experiments

As mentioned in section 1.1, the first polarized DIS experiment, SLAC E80 [14] was initiated by the Yale-SLAC group and performed in 1976. A beam of polarized electrons was scattered off a polarized proton target, and the first measurement of the proton asymmetry A_1^p was made. An atomic beam source was used for the production of polarized electrons. A beam of ^{6}Li atoms was polarized by filtering through a Stern-Gerlach apparatus. An intense argon flash lamp light with the wavelength 170-230 nm was focused onto the polarized ^{6}Li atoms, giving polarized ionization photoelectrons that were injected into the accelerator. The accelerated beam energy ranged from 9 to 12 GeV. The average beam polarization, measured in the experimental area by the single arm Møller polarimeter, was 50%, but the beam intensity was only around $5 \cdot 10^8$ electrons per pulse. The proton target was polarized, 50% on average, by dynamic nuclear orientation [59] using a butanol sample with a paramagnetic dopant. The scattered electrons were detected in a fixed 9° angle spectrometer, and their momentum was determined. The experiment was repeated (as E130 [15]) by the same group in 1980 with higher beam energies of 16.2 and 22.7 GeV. The observed large A_1^p asymmetries were in agreement with the expectations of the quark-parton model and its sum rules.

After completion of the early SLAC series of experiments, a new polarized program was developed at CERN. In a 1984-85 run the EMC collaboration extended the SLAC results, performing polarized muon-proton scattering. The μ^+ beam with intensity $4 \cdot 10^7$ particles per spill and with average energy 190 GeV is naturally polarized. It originates from π^+ decay, and its polarization was calculated to be 80%. The target consisted of two cells filled with solid NH_3 and polarized in opposite directions via dynamic nuclear polarization to 75% on average. The higher beam energy allowed measurements to obtain a much lower x than the SLAC experiments. In the overlap region $x > 0.1$ the results of both experiments agree well. However, The low $0.01 < x < 0.1$ data fall well below the theoretical prediction. As a consequence of this, the measured integral with the Regge type $g_1^p(x) \sim x^{0.12}$ extrapolation at low x, at $\langle Q^2 \rangle = 10\,\mathrm{GeV}^2$,

$$\Gamma_1^p = 0.126 \pm 0.010(\mathrm{stat.}) \pm 0.015(\mathrm{syst.}) \tag{1.137}$$

is in disagreement with the Ellis-Jaffe prediction by three standard deviations of the experimental error. With the assumption of isospin invariance and $\mathrm{SU}(3)_{flavor}$ symmetry in the baryon octet decays, i.e. with the value of axial charges Δq_3 and Δq_8 given in eqs. (1.118), (1.124), the Γ_1^p result gives

$$\Delta\Sigma = 0.12 \pm 0.17. \tag{1.138}$$

The spin carried by the quarks was found to be surprisingly small, and this result became known as 'spin crisis'. It triggered a lot of theoretical activity and new experimental programs at CERN, SLAC and DESY.

The measurements at CERN continued, and the SMC collaboration took deuteron data [21] in 1992, 1994 and 1995, and proton data [20] in 1993 and 1996. The first year of running was devoted to the deuteron to have the first measurement of the neutron g_1^n needed to test the Bjorken sum rule. The high energy muon beam extended the coverage down to $x = 3 \times 10^{-3}$ at the high average $\langle Q^2 \rangle$ of 10 GeV^2. However, the low intensity of the beam limited the statistical precision of the data. The results, in terms of integrals of structure functions, are quoted in table 1.2.

Table 1.2: Integrals of $g_1^{n,p,d}$ From CERN, SLAC and DESY Experiments.

Experiment	Target	Target Material	$\langle Q^2 \rangle$ (GeV^2)	$\Gamma_1 \pm$ stat. $\pm$ syst.
CERN EMC	p	ammonia	10	0.126 $\pm$ 0.010 $\pm$ 0.015
CERN SMC	p	butanol ammonia	10	[2]0.149 $\pm$ 0.006 $\pm$ 0.010
CERN SMC	d	butanol	10	0.041 $\pm$ 0.006 $\pm$ 0.005
SLAC E142	n	^{3}He	2	-0.031 $\pm$ 0.006 $\pm$ 0.009
SLAC E143	p	ammonia	3	0.129 $\pm$ 0.004 $\pm$ 0.009
SLAC E143	d	ammonia	3	0.042 $\pm$ 0.003 $\pm$ 0.004
DESY HERMES	n	^{3}He	2.5	-0.037 $\pm$ 0.013 $\pm$ 0.008

Another series of SLAC experiments started with E142 [17] in the fall of 1992. A beam of high intensity was produced, typically $2 \cdot 10^{11}$ electrons per pulse, with three different energies 19, 23 and 26 GeV. The polarized electrons were obtained at the source by illuminating the AlGaAs photocathode with circularly polarized light. The polarization measured by the single arm Møller detector was about 36%. The new method of obtaining the polarized beam of electrons allowed for a pseudo random choice of the beam helicity from pulse to pulse. This feature, unique to SLAC, resulted in strong suppression of the

[2]1993 and preliminary 1996 data combined [52].

systematic effects related to the beam asymmetry. A high density ^{3}He target was developed and polarized, to 33% on average, via spin-exchange optical pumping. About 300 million deep inelastic events were observed in two magnetic spectrometers centered at 4.5° and 7°. The neutron asymmetries were measured directly for the first time, in the kinematic range of $0.03 \leq x \leq 0.6$, $\langle Q^2 \rangle = 2$ GeV^2, giving the most precise measurement of the neutron spin structure function $g_1^n(x)$ at that time. We note that for the first time the $A_\perp$ asymmetry was measured to minimize the systematic uncertainties.

Next came the winter of 1993/94 and SLAC experiment E143 [18, 19]. With the advent of the new strained GaAs photocathodes the beam polarization was 80%. Also the beam energy was raised to 29 GeV. The spectrometer setup of E142 was used with new polarized at 65% ammonia (for proton) and polarized at 25% deuterated ammonia (for deuteron) targets. The data were also taken at the beam energies of 9 and 16 GeV to study the Q^2 dependence [60]. The longitudinally and transversely [53] polarized proton and deuteron structure functions were measured in the kinematic range of $0.03 \leq x \leq 0.7$ and the average $\langle Q^2 \rangle$ of 3 GeV^2.

A new spin structure program HERMES [22] was started recently at the DESY collider. It utilizes a 28 GeV beam of positrons in the HERA storage ring polarized to an average 55% by the Sokolov-Ternov mechanism [61]. The polarized ^{3}He atoms, 46% on average, were injected directly into the windowless cell located inside the positron ring allowing for almost no dilution at the target. The g_1^n structure function was measured in the kinematic range



Figure 1.2: Proton data for $x\, g_1^p(x)$.

$0.02 \leq x \leq 0.6$, $\langle Q^2 \rangle = 2.5$ GeV^2. We note that the HERMES possibility of tagging the leading hadron (so called semi inclusive scattering) allows one to probe the valence and sea distributions directly. This program will continue beyond the year 2000.

The results of these experiments for the xg_1^p and xg_1^n structure functions are presented in figs. 1.2 and 1.3. We see that the proton data are positive, relatively large and rather well established in the common x region of experiments. At $x \lesssim 0.01$ the data is sparse, and we have to rely entirely on models for extrapolation to $x = 0$. Note that the lowest x point for xg_1^p



Figure 1.3: Neutron data for $x\, g_1^n(x)$.

suggests a rise, which would mean that $g_1^p(x)$ is more divergent than $1/x$ and thus nonintegrable. This measurement comes from the 1993 run of SMC. The preliminary analysis of the 1996 run [52] lowers that point to a value consistent with zero. The neutron values of xg_1^n are negative and relatively small. The experimental data are consistent in the region of overlapping x, but the errors are still significant in comparison with the values. And, as in the proton case, behavior of xg_1^n for $x \lesssim 0.01$ remains unconstrained.

The traditional analysis of all data existing at the time was performed by Ellis and Karliner [62] in 1995. The of integrals of g_1's given by the different experiments were evolved to the same Q^2 of 3 GeV^2 and then combined to find the Bjorken sum rule

$$\Gamma_1^p - \Gamma_1^n = 0.164 \pm 0.011 \qquad \text{at } 3 \text{ GeV}^2, \tag{1.139}$$

in agreement with the prediction. The authors go further and use the Bjorken sum rule to extract $\alpha_s(3\text{GeV}^2) = 0.328^{+0.026}_{-0.037}$, see eq. (1.122), or equivalently,

$$\alpha_s(M_Z^2) = 0.119^{+0.003}_{-0.005}. \tag{1.140}$$

They also make a global fit and obtain the flavor decomposition of the nucleon spin,

$$\begin{aligned} \Delta u &= 0.82 \pm 0.03 \\ \Delta d &= -0.44 \pm 0.03 \\ \Delta s &= -0.11 \pm 0.03 \end{aligned} \tag{1.141}$$

and

$$\Delta\Sigma = 0.27 \pm 0.04. \tag{1.142}$$

One should keep in mind the Q^2 evolution assumption, the uncertainty of the low x extrapolation assumptions, and the neglect of higher twist contributions and higher order corrections in the analysis.

CHAPTER 2

EXPERIMENTAL APPARATUS

The SLAC E154 experiment collected data for two months during October-November 1995.

The polarized electrons were produced at the source, accelerated in 3.2 km of linac to 48.3 GeV and delivered, longitudinally polarized, to the experimental area of the End Station A. The Møller detector determined that the beam polarization was 82%. The electrons scattered off the optically polarized, 38% on average, ^{3}He target and were detected in two independent magnetic spectrometers positioned at the central angles 2.75° and 5.5° with respect to the beam line. Each spectrometer consisted of Cherenkov detectors, hodoscopes and an electromagnetic calorimeter. The gas threshold Cherenkov detectors were used for particle identification. Several planes of scintillator hodoscopes served for tracking particles, allowing the determination of the direction and the value of the electron momentum. The lead glass calorimeter measured the deposited energy of a scattered particle. The kinematic range of the results on $g_1^n(x)$ and $g_2^n(x)$ determined from these measurements was $0.014 \leq x \leq 0.7$ and $1\,\mathrm{GeV}^2 \leq \mathrm{Q}^2 \leq 17\,\mathrm{GeV}^2$ (average $\langle Q^2 \rangle = 5\,\mathrm{GeV}^2$).

The experiment was an effort of 81 physicists representing 23 institutions.

2.1 SLAC polarized source

Since 1992 both the SLAC accelerator and SLC have been operating solely with polarized electrons. The electrons are produced at the source [63, 64, 65] with 120 Hz frequency. For the fixed target experiments the first 119 pulses, which are 10 μsec long, are produced by a flashlamp pumped titanium-sapphire (Ti:S) dye laser operating at 850 nm wavelength. The last witness pulse, used for monitoring purposes, only lasts 2 ns and is generated by another YAG-pumped Ti:S laser. The schematic view of the source is shown in fig. 2.1. The linearly polarized light goes through the slicer and shaper (laser pulse chopper) where it is cut to 240 ns and shaped slightly non-uniformly to compensate for beam loading effects of the linac. Next it is circularly polarized via a system of two Pockels cells with axes rotated 45° with respect to each other. The first cell operates at its quarterwave voltage, and the second one at zero, with small corrections generated by the feedback loop to maintain the charge asymmetry below 10^{-4}. The positive quarterwave HV produces one helicity, while the negative HV pulse produces the opposite helicity. The sign of the applied HV, and thus the helicity,



Figure 2.1: SLAC polarized source setup for SLC runs. For E154 operation one of the lasers was replaced by a flashlamp pumped Ti:S laser.

follows a pseudo-random pattern from pulse to pulse, a unique feature of the SLAC accelerator which reduces the false asymmetries correlated with the beam helicity. The photons with a well defined helicity hit the cathode of the electron gun, photoemitting polarized electrons.

Several types of gallium arsenide related cathodes have been tried. The E142 experiment used $Al_{0.12}Ga_{0.88}As$ material. The filled top valence band of AlGaAs has $P_{3/2}$ symmetry, while the empty conduction band has $S_{1/2}$ symmetry. The addition of Al changes the energy gap to 1.63 eV and makes it easier to tune the frequency of lasers to the gap. The light of +1 helicity excites electrons from $|P_{3/2}(m_j = -3/2)\rangle$ to $|S_{1/2}(m_j = -1/2)\rangle$ with probability 3 times larger than the competing transition from $|P_{3/2}(m_j = -1/2)\rangle$ to $|S_{1/2}(m_j = +1/2)\rangle$, giving the theoretical 50% polarization limit of the emitted electrons. In practice polarizations on the level of 40% were achieved. To get the electrons from the conduction band to the vacuum one needs to pass the work function barrier, approximately 4 eV for pure GaAs. By deposition of cesium and an oxidizer (O_2 or NF_3) the potential of the vacuum can be lowered below the conduction band potential. A material with such characteristic is said to have a negative affinity. Now the conducting electrons can tunnel through the potential barrier at the surface out of the material guided by 60 kV applied to the cathode. During the experiment the cesium layer gets slowly deactivated, probably due to the residual gas in the vacuum system, making the penetration of the surface barrier more difficult. There is a drop in the quantum efficiency of the cathode, since only electrons closer to the surface can escape. These electrons are less subject to latice interactions on the way out. That produces the observed increase in the polarization.

For our experiment a strained GaAs material for the cathode was used. This is achieved by growing a thin (e.g. 0.25μm) layer of GaAs on a thick layer of $GaAs_{1-x}P_x$ (e.g. 25μm). The x fraction of replaced As atoms was varied smoothly from 0 to 28%. The lattice constant of the phosphorus doped GaAs is 1% smaller than pure GaAs. That produces a compression in GaAs lattice and breaks the degeneracy between the $|P_{3/2}(m_j = \pm 3/2)\rangle$ and

$|P_{3/2}(m_j = \pm 1/2)\rangle$ valence band levels producing 50 mV energy difference. In figure 2.2 we show the energy structure of strained GaAs. The theoretical



Figure 2.2: The energy level structure of strained GaAs.

limit for the polarization of photo-emitted electrons is 100%. In practice polarizations above 80% were achieved. The level splitting is not large enough to allow tuning of the laser frequency completely away from the energy of the competing transition. Furthermore, even 99.5% circularly polarized light has 10% linearly polarized component. There is also some strain relaxation, and thermally excited electrons in the conduction band contribute to the lower than theoretical limit polarization.

The $0.5-2\times10^{11}$ photoemitted electrons were bunched together and sent down the accelerator.

2.2 Beam acceleration and transport

The 3.2 km of the linear accelerator consists of 30 sectors. Each sector contains steering magnets: dipoles and a quadrupole, some elements for monitoring beam position and current, and, above all, eight klystrons, each accompanied by a system of RF resonant cavities, the so called SLAC Energy Doubler (SLED). Before our experiment, when the SLED technique was not implemented, a beam energy of 29 GeV was achieved.

Every klystron feeds microwave radiation at 2856 MHz into copper cavities. The fields in the cavities build up, and a wave of increasing amplitude is emitted from the coupling apertures of each cavity. At 0.8 μsec before the end of a 3.5 μsec long RF pulse, the 180° phase shifter reverses the phase of the output wave from the klystron [66]. The emitted and reversed waves add in phase at the accelerator. Immediately after the phase reversal the amplitude of the superposition is twice the amplitude of each single wave. That allows for the maximum measured unloaded energy of 55 GeV and maximum practical beam energy up to 52 GeV. This setup can only accelerate about a 150 ns beam pulse within an energy spread of 0.5% with 10^{11} electrons. By applying two additional 180° phase inversions for 20% of klystrons [67], the classical SLED pulse is flattened to achieve an energy spread of 0.5% over 240 ns.

The electrons are transported from the linac to End Station A through the A-Line, which had to be upgraded to handle beam with energies up to 50 GeV. The E154 experiment was the first after the modification. The

48.75 GeV electrons from the linac are first bent over 0.5° by a set of five pulsed and two DC magnets that offer the option of switching the beam between four beam lines on a pulse-by-pulse basis [68]. This area is called the Beam Switchyard. The next 24° bend, for the total of 24.5°, is provided by twelve identical dipole magnets. Nine quadrupoles are used to control the beam divergence and the spot size at the target. Two massive high-power collimators limit the energy spread of the beam.

2.3 Beam parameters and monitoring

2.3.1 Energy

The beam energy in the experimental area of the End Station A (ESA) was established in two independent ways.

The first uses the optics of the A-Line [68]. Twelve dipole magnets were connected in series with an identical, off line dipole equipped with a flip-coil. The current induced in the flip-coil measures the magnetic field of the dipoles which, with the knowledge of the bending angle, determines the momentum of the electrons to about 0.1%. Some corrections to the flip-coil measurement are needed: for the synchrotron radiation loss, stray magnetic field beyond the coil, a bias in the coil readout electronics and the steel support plate present in all the beam line magnets except in the reference magnet. Including the corrections, the energy in the ESA was determined as 48.325 GeV with an accuracy of 40 MeV.

The second measurement was done by the energy scan of the longitudinal beam polarization in the Møller detector. The details are given in the next section 2.4.6 (page 60). This determination gave the energy in the ESA equal to 48.30 GeV, in excellent agreement with the first.

The value E of the energy was monitored throughout the experiment via the flip-coil signal. The energy spread of the beam $\Delta E/E$ was restricted to 1% by the A-line collimators.

2.3.2 Current

The beam current was measured for each pulse with two independent ferromagnetic toroids located more than 30 m apart, upstream from the target. The beam passing through the ferrite core of the toroid induced a signal in the loops of wire wrapped around the core, which in turn was sent to a resonant LC circuit. The induced current was then amplified and integrated. The total charge is proportional to the beam current. Several times a day a precisely charged capacitor was discharged through each of the toroids, giving the calibration coefficients for the beam measurements. In this way the total beam charge was established with about 0.5% accuracy.

2.3.3 Position and steering

The quick coarse centering and focusing of the beam was made with two mylar roller screens, coated with fluorescent ZnS. The screens, one in front of the target and the second 25 m upstream, were moved into the beam, observed with television monitors, and then moved back before collecting the data. They were placed periodically in the beam for monitoring purposes. The main information about the beam position and width was obtained from a wire array located 10.5 m downstream of the target. There were 24 wires in the x direction and 24 in the y direction, made of 0.127 mm diameter CuBe wire and spaced 1.1 mm apart. Each pulse of current induced in a single wire by the beam was integrated by ADC's, read out by our Data Acquisition System (DAQ) and logged on tape. The information was also available to the linac operators at the Main Control Center (MCC) for minor adjustments and monitoring.

Additional beam quality monitoring was performed with the use of two scintillation counters. The first was positioned upstream from the target in the ESA alcove, close to the beamline. It was sensitive to the beam scraping and thus dubbed 'bad spill monitor'. The other counter, the 'good spill monitor', was downstream from the target, several meters away from the beampipe. It was sensitive to particles created at the target. Because its signal (and also the signal from the first counter) was displayed on the oscilloscope (both in the Counting House and MCC) it was a sensitive indicator of the time structure of the beam. The ADC integrated signals were also collected by DAQ and stored on tape for the analysis.

Other monitors of the beam included two traveling wave beam position monitors (TWBPM) and a secondary emission monitor (SEM) for target protection. Traveling wave BPM's were located in front of the target. One microwave cavity produced an RF signal proportional to the horizontal deviation of the beam from the cavity center. The second was sensitive to vertical displacements.

The SEM consisted of a thin aluminum foil with a 2 cm circular aperture for the beam. The opening matched the diameter of the target. Thus, large signals from the monitor indicated a potentially destructive displacement of the beam and triggered an automatic beam shutdown. This prevented the beam from hitting the thick side walls of the target.

2.3.4 Spin precession

The helicity of electrons entering the linac is the same as at the source, and thus the same as the helicity of the photons hitting the photocathode. It remains unchanged until the entrance to the A-line. In the A-line the beam is bent by θ_{bend}=24.5°, and the spins of electrons undergo a precession [69]

$$\theta_{prec} = \gamma \left(\frac{g-2}{2} \right) \theta_{bend}, \tag{2.1}$$

relative to the beam direction. Here $\gamma = E/m$, and $g = 2.00231\ldots$ [70] is the gyromagnetic ratio of the electron. For the full A-line bend the above formula can be interpreted as follows: The electron helicity is flipped every 3.2374 GeV of beam energy. The E154 experiment was set to run at 48.56 GeV linac energy. That corresponds to $\theta_{prec} = 15\pi$. The helicity of the electrons in the End Station A was opposite to that at the source.

The A-line energy is modified due to synchrotron radiation, which, being proportional to E^4, becomes important at high energies. The synchrotron energy loss ΔE is given by [71]

$$\Delta E = \frac{2}{3}\frac{e^2}{\rho}\beta^3 \left(\frac{E}{m}\right)^4 \theta_{bend}, \tag{2.2}$$

where $\rho = 85.927$ m [72] is the bending radius in meters, θ_{bend} is expressed in radians and $\beta \approx 1$ for our energy. Taking into account this loss the End Station A beam energy $E = 48.362$ GeV corresponds to 15π spin rotations. The measured energy was $E = 48.325 \pm 0.040$ GeV, as explained in section 2.3.1.

The synchrotron radiation also increases the beam emittance, resulting in a larger spot size at the target. To reduce this effect and bring the beam diameter back to the required ~1 mm at the target, a quadrupole magnet (Q41) was installed at the entrance to the ESA with a focal point at the target.

2.3.5 Helicity reversal

The reversal of the beam polarization on a pulse-to-pulse basis with the Pockels cell was very important for reducing systematic errors. Possible false asymmetries arising from slow changes in spectrometer acceptance were averaged out by the rapid beam helicity reversal. Also, by changing the target polarization direction, we could further average asymmetries due to possible helicity related differences in beam properties. The absolute helicity of the beam in the ESA was determined by the sign of the asymmetry measured by the Møller polarimeter.

The beam helicity was chosen on a pulse-to-pulse basis using a pseudo-random number generator. The seed began with the choice of a 32 bit random number. The last bit, the 32nd, determined the helicity of the beam. For the next pulse, the 19th and 32nd bits were subject to a logical XOR operation. After the remaining bits had been shifted up, the result was placed as the 1-st bit. The new 32nd bit determined the helicity of this pulse. Then the cycle was repeated. The generator ran continuously with the source on or off. Thus, knowledge of the polarization bits within the 33 consecutive pulses was needed to determine the random sequence seed. Then the prediction for the polarization of each spill could be made based on the spill number and could be used to test the integrity of the system.

The polarization state measurements were sent to the Counting House via four physically distinct paths and then recorded by the Data Acquisition System. They were referred to as the PMON Line, Mach Line, HV Line (short for the Pockels Cell High Voltage Line) and Veto Line (short for the Klystron Veto Module Line). Each line delivered two bits. The combination '01' referred to the positive helicity photons incident on the photocathode, while '10' to the negative helicity. The '00' result meant that the unpolarized beam and '11' indicated an error. Typically all four measurements agreed with the prediction [73] and if they did not, the spill was discarded. If the rate of disagreement exceeded $5 \cdot 10^{-4}$ for a run, the entire run was excluded from the analysis.

2.4 Møller polarimeter

A single arm Møller polarimeter was used throughout the experiment to measure the beam polarization. It consisted of polarized ferromagnetic target foils, a momentum-analyzing dipole magnet and five silicon detectors. It was also used to optimize the wavelength of the source laser with respect to the beam polarization and to calibrate the flip coil energy (see also section 2.3.1 on page 47).

2.4.1 Møller asymmetry

The cross section for the Møller scattering of polarized e^- on polarized e^- has the form

$$\frac{d\sigma}{d\Omega} = \frac{\alpha^2}{s}\,\frac{(3+\cos^2\theta)^2}{\sin^4\theta}\,\left[1 - P_z^B P_z^T A_{zz}(\theta)\right] \tag{2.3}$$

in the CM frame [74], where s is the total CM momentum squared. The z-axis is along the beam direction, and θ is the CM scattering angle. Here the longitudinal beam and target polarizations are denoted by P_z^B and P_z^T, respectively, and the asymmetry $A_{zz}(\theta)$ is given by

$$A_{zz}(\theta) = \frac{7+\cos^2\theta}{(3+\cos^2\theta)^2}\,\sin^2\theta. \tag{2.4}$$

By counting the rates for the electron beam and target spins aligned , $\sigma^{\uparrow\uparrow}$, and antialigned, $\sigma^{\uparrow\downarrow}$, we can measure the asymmetry

$$A(\theta) = \frac{\sigma^{\uparrow\downarrow} - \sigma^{\uparrow\uparrow}}{\sigma^{\uparrow\downarrow} + \sigma^{\uparrow\uparrow}} = P_z^B P_z^T A_{zz}(\theta). \tag{2.5}$$

If the target polarization P_z^T is known independently, the above formula can be used to determine the beam polarization P_z^B. The asymmetry has a maximum at $\theta = 90°$, where the unpolarized laboratory cross section is 0.179 b/sr and $A_{zz} = 7/9$. With a typical target foil polarization of 0.08 and the beam polarization of 0.80, the expected measured asymmetry $A(90°)$ is roughly 0.05 [75].

2.4.2 Polarimeter design

The design was an evolution of previous End Station A Møller polarimeters [76] and utilized many ideas and components of previous designs.

The E154 Polarimeter consisted of polarized target foils, a mask to define the azimuthal and vertical acceptance, a magnet to momentum analyze the scattered electrons, and detectors to measure the scattering rate. It was located upstream from the ^{3}He target. Schematic top and side views are shown in fig. 2.3. Six target foils were installed at a 20.7° angle to the beamline [75]. The foils were 20(2 of them), 30, 40(2) and 154 μm thick and made of ferromagnetic Vacoflux (49% Fe, 49% Co, 2% Va by weight). They were magnetized to near saturation by Helmholtz coils providing nearly 100 G in the target center. The mask was made from 25 radiation lengths of tungsten. It had a central hole for the unscattered beam and two wedge-shaped holes to select vertically scattered electrons. That gave the ϕ acceptance of 0.20 rad (top hole) and 0.22 rad (bottom) over the range of allowed scattering angles, $3.59 < \theta_{\mathrm{lab}} < 8.96$ mrad in the lab frame. Additional Pb shielding and a collimator prevented the single-scattered target photons from reaching

Figure 2.3: Top (top picture) and side (bottom picture) views of the E-154 Møller Polarimeter.

the detectors. The momentum selection of the scattered electrons was accomplished with a dipole magnet producing $\int Bdl$ 33 kG-m at 900 A current.

The electrons were observed by five silicon detectors mounted 28.9 m from the target. The top detector had 48 channels with a fine segmentation of 2.18 mm in the vertical (θ) direction. It was mounted on a remotely controlled stage and could be positioned anywhere within the Møller acceptance. Typically the Møller peak corresponded to $\theta = 94°$ in the CM. Each of the four bottom detectors had 12 channels with a coarser separation of 8.69 mm. They covered the θ CM scattering angle range of $93° - 104°$. The silicon channels were connected to charge sensitive preamplifiers and then to ADC's read by DAQ. The unscattered electrons continued to travel the beam pipe that went through the magnet. The beam pipe was enclosed in the iron septum to eliminate the magnetic field.

2.4.3 Foil polarization measurement

The targets were made of Vacoflux, as mentioned above, composed of 49% Fe, 49% Co and 2% Va. The foils were typically 3 cm wide by 35 cm long by 20-154 μm thick. They were polarized to saturation by a 100 G magnetic field. Only two electrons in Fe and Va contribute to the magnetization, and on average the electrons were $\sim$ 8% polarized. This polarization was determined measuring the magnetization M [77]. An integrating voltmeter was connected to a pick-up coil made of $N = 500$ turns wrapped around the foil. The bipolar Helmholtz coil power supply ramped the H field from -100 to +100 Gauss, and the voltmeter recorded the flux change

$$\Phi_{\text{foil in}} \equiv \int V(t)\, dt = 2N\ [A_f B + (A_c - A_f)\ H], \tag{2.6}$$

where A_c and A_f are the cross-sectional areas of the pick-up coil and the foil respectively. The induction B at the coil has contributions from the magnetic field H and the magnetization M in accordance with

$$B = H + 4\pi M, \tag{2.7}$$

so that the flux $\Phi_{\text{foil in}}$ can be re-written as

$$\Phi_{\text{foil in}} = 2N\ (A_c H + 4\pi A_f M). \tag{2.8}$$

Because the foils were very thin, the H field was almost unaffected by their presence. Repeating the procedure without the foils in place,

$$\Phi_{\text{foil out}} = 2N A_c H, \tag{2.9}$$

was measured. The magnetization M is given by

$$M = \frac{\Phi_{\text{foil in}} - \Phi_{\text{foil out}}}{8\pi N A_f}. \tag{2.10}$$

The magnetization M has to be corrected for the orbital angular momentum contribution to the electron polarization. Let us recall the definition of g', the magneto-mechanical ratio [78], $M = g'\mu_B(L + S)$. The spin contributes $M_S = g\,\mu_B\,S$ to M, and the orbital angular momentum contributes $M_L = \mu_B\,L$. Obviously, $M = M_S + M_L$. The equations can be solved for M_S in terms of M, i.e. for the fraction of the magnetization due to the electron polarization, and the electron polarization $P^T = M_S/(n_e\mu_B)$, where n_e is the electron density, can be computed using

$$P^T = \left(\frac{g'-1}{g'}\right)\ \left(\frac{g}{g-1}\right)\ \frac{M}{n_e\ \mu_B}. \tag{2.11}$$

The gyromagnetic ratio $g = 2.00231\ldots$ [70]. The g' ratio was measured for the alloy consisting of 50% Fe and 50% Co to be $g' = 1.916 \pm 0.002$ [79]. We assume that it could only be slightly affected by the presence of 2% Va and increase the error to ± 0.02. The polarization of the foils was measured twice, before and after the experiment, giving the same results within 0.1% on average. The measured average polarization of the foils was 8.2% with relative error 1.7% [75].

2.4.4 Beam polarization results

The Møller data were taken during about 140 special, dedicated runs in the E154 experiment. The runs were typically ten minutes long. The preamplified outputs of the silicon detector channels were integrated by ADC's and recorded by the Data Acquisition System. In the analysis stage the averages for each ADC channel were made separately for pulses with left (L) and right (R) beam helicities. Then the sums (R+L) and differences (R-L) were formed. The background under the unpolarized (R+L) Møller scatters was estimated by fitting the (R+L) line-shape to an arbitrary quadratic background plus the line-shape expected from the unpolarized Møller scattering. The technique for estimating the unpolarized line-shape was based on the measured polarized (R-L) line shape with corrections for the atomic motion of the target electrons. It is in excellent agreement with Monte Carlo results [80]. The atomic motion correction comes from the atomic momentum of unpolarized electrons inside Fe and Co atoms. Although this momentum p is only on the order of 100 KeV, it considerably smears ($\sim 10\%$) the lab frame scattering angle by a maximum amount [81]

$$\tan(\theta_{\text{smeared}}) \approx \tan(\theta)\ \sqrt{1 - \frac{p}{m}}, \tag{2.12}$$

because it is to be compared with the electron mass m and not with the beam energy. The elastic peak from the unpolarized inner shell electrons is significantly wider than the peak from the polarized electrons. Experimentally the consequences of this effect depend on how much of the Møller peak is detected. As the fraction of the detected peak gets smaller, the measured asymmetry gets larger since the relative contribution of events from the unpolarized electrons decreases.

The measured asymmetry was calculated from the ADC averages as

$$A = \frac{\sum_i (R - L)_i}{\sum_i (R + L)_i - \sum_i (background)_i}, \tag{2.13}$$

where the sum is over channels including the Møller peak. The sum range was chosen large enough to avoid the necessity of additional corrections for the target motion effect. The background subtraction increased the measured asymmetry by 17%-24% relative.

The beam polarization history is given in table 2.1 and is shown in figure 2.4. The statistical errors for each run were typically $0.003 - 0.006$. The average beam polarization (weighted by statistics of $A_{||}$) for the experiment was $81.8\% \pm 2.3\%$.

2.4.5 Systematic uncertainties

The small statistical relative errors of 0.4%-0.7% per run made possible precise studies of the systematic errors. The overall systematic error has

Table 2.1: Measured Longitudinal Beam Polarization For Different Running Periods. The Error is Statistical Only.

Run range	Polarization
1329–1411	0.759 ± 0.004
1456–1684	0.775 ± 0.005
1691–2311	0.814 ± 0.002
2316–3371	0.824 ± 0.001
3377–3788	0.826 ± 0.002

contributions from the foil polarization due to uncertainty of the g' factor, uncertainties in the expected Møller asymmetry for each detector, and uncertainties in the background subtraction. The sizes of various contributions are given in table 2.2. The helicity related correlations were cancelled out

Table 2.2: Relative Systematic Error Contributions to the Beam Polarization Measurement.

Systematic error contribution	Value
Foil magnetization	1.9%
Kinematic acceptance	0.3%
Background correction	2.0%
Fit range	0.3%
TOTAL	2.8%

by pulse-by-pulse changes of the beam helicity and by almost always taking runs in pairs with opposite target helicities.

2.4.6 Beam energy scan and beam helicity determination

The Møller polarimeter permitted an independent measurement of the beam energy [75]. The electron spin precesses at the A-line bend by an



Figure 2.4: Measured longitudinal beam polarization versus E154 run number. The errors are statistical only. Changes in the measured average value of polarization indicate various adjustments made to the source during the run.

amount given by (g-2) and the beam energy E (see section 2.3.1 on page 47). The expected sinusoidal dependence is complicated by the E^4 synchrotron radiation loss. The longitudinal polarization of a beam with polarization $P_\circ$ at the source is expected to vary in the End Station A as

$$P_z = P_\circ \cos(\pi(E + 0.198(E/48.362)^4)/3.2374). \tag{2.14}$$

Data taken at different A-line energies were fitted with the expected E (from the flip coil) dependence with the polarization magnitude $P_\circ$ and the energy offset Δ as fit parameters, as shown in figure 2.5. The flip coil reading was



Figure 2.5: Measured beam polarization versus the nominal A-line beam energy as read out by a flip coil.

determined to be 74 MeV below the true energy of 48.30 GeV in the End Station A.

The polarimeter also established the absolute sign of the beam polarization in the End Station A. The Møller cross section is larger if the electron spin of the target and beam are anti-parallel. Knowing the sign of the asymmetry (negative) and the target polarization direction (the electron spin is opposite to the direction of electron magnetic moment, which is established by the direction of the current in the Helmholtz coils) the beam helicity orientation is extracted. The '01' polarization state was assigned to negative helicity in the End Station A and '10' to the positive one, in agreement with the conventions from section 2.3.5 (page 51).

2.5 Polarized ^{3}He target

The high density polarized ^{3}He target was a source of polarized neutrons. It was developed from a similar target used in E142 experiment [82]. The target was a two-chambered 30 cm long and 2 cm in diameter glass tube holding 2.6×10^{20} atoms/cm^3 of ^{3}He. The ^{3}He was polarized via spin exchange with optically pumped rubidium vapors and was kept aligned in 10-30 Gauss holding field. The average polarization was 38% for the experiment with a maximum of nearly 50% [83].

2.5.1 Optical pumping and spin exchange

The targets with a high density of ^{3}He are commonly polarized using the technique of optical pumping [84]. In this two step process the rubidium vapor is first optically polarized. Then the polarization of the rubidium electrons is transferred to the ^{3}He nucleus by the spin-exchange hyperfine interaction.

The circularly polarized laser light at 794.7 nm with positive, for example, helicity excites the $5S_{1/2}(m = -1/2)$ ground state to the $5P_{1/2}(m = +1/2)$

state in rubidium vapor (the D1 line). A simplified diagram of the process is shown in fig. 2.6. Radiative decays to the ground state favor the



Figure 2.6: Optical pumping of Rubidium vapor by circularly polarized light.

$5S_{1/2}(m = -1/2)$ state over $5S_{1/2}(m = +1/2)$ state by a factor of two. Every three photons absorbed would increase the population of $5S_{1/2}(m = +1/2)$ by one. By adding 65 torr of N_2 the $5P_{1/2}(m = \pm 1/2)$ are collision mixed at a rate exceeding the radiative decay. The relaxation into the $5S_{1/2}$ ground state now occurs equally into the two magnetic substates. Only two photons are required to increase the population of $5S_{1/2}(m = +1/2)$ by one. The polarization of the valence electrons of Rb competes with several depolarizing mechanisms. The most important are spin destruction from Rb–^{3}He and Rb – Rb collisions (both dominate and contribute approximately equal amounts of relaxation), and interactions of Rb with walls of the target cell. The optical pumping rate of the optically thick Rb vapor (10^{14} atoms/cm^3) with sufficient laser power is $1/\Gamma_{\rm pump} \approx 10^{-6}$ s and it dominates over the competing $1/\Gamma_{\rm relax} \approx 10^{-3}$ s spin relaxation rate. A Rb polarization of almost 100% could be achieved.

The spin exchange between the electrons of Rb and the nucleus of ^{3}He is mainly governed by the Fermi contact term in the hyperfine hamiltonian [85] which corresponds to the overlap of the Rb electron with the ^{3}He nucleus. The rate of this exchange, Γ_{SE}, is proportional to the relative velocity v between the ^{3}He and Rb, and to the number density of Rb, n_{Rb}:

$$\Gamma_{SE} = \langle v\, \sigma_{SE} \rangle\, n_{Rb}, \tag{2.15}$$

where the velocity averaged cross section is 1.2×10^{-19} cm^3/s [86]. With $n_{Rb} \approx 10^{14}$ cm^{-3} polarization times for ^{3}He are on the order of $1/\Gamma_{SE} \approx 25 - 40$ hours. The nuclear polarization $P_{(^3He)}$ of ^{3}He increases in time according to the following equation:

$$P_{(^3He)}(t) \;=\; P_{(Rb)}\, \frac{\Gamma_{SE}}{\Gamma_{SE} + \Gamma_R}\; \left[\, 1 - e^{-(\Gamma_{SE}+\Gamma_R)t}\, \right], \tag{2.16}$$

where $P_{(Rb)}$ is the equilibrium polarization of Rb ($\approx$ 100%) and Γ_R is the total spin relaxation rate of ^{3}He. The polarization curve of the best cell, Picard, is shown in fig. 2.7.

The biggest source of the ^{3}He depolarization were the collisions between ^{3}He atoms causing the exchange of nuclear spin and orbital angular momentum. For the cell Picard they resulted in $1/\Gamma_D = 84$ hours. Other, smaller



Figure 2.7: Polarization of the target cell Picard during the run.

contributions were caused by paramagnetic gaseous impurities embeded in the cell walls, gradients of the magnetic field perpendicular to the alignment field, and from the beam ionization of ^{3}He. In our experiment the other sources resulted in $1/\Gamma_O \equiv \sum_i (1/\Gamma_i) > 400$ hours. For the Picard cell the total relaxation was measured to be [83]

$$1/\Gamma_R = 1/\Gamma_D + 1/\Gamma_O = 84 \pm 5 \text{ hrs}, \tag{2.17}$$

and a maximum polarization of 47.3% was achieved (see fig. 2.7).

2.5.2 Target cells

The high polarization of the target that translates into long relaxation time is one of two goals of the target construction. The other is a large dilution factor f. Let us recall that f is the ratio of the number of electrons scattered by the ^{3}He to the total number of electrons scattered by the target (see sec. 3.10, page 154). Both goals are equally important because the running time for a given statistical error is inversely proportional to the square of the product of the polarization and the dilution. The dilution factor can be increased by increasing the density of ^{3}He or by decreasing the thickness of the cell windows traversed by the beam.

The target consisted of two chambers. The upper pumping cell had a 68-84 cm^3 volume and contained ^{3}He and a few tens of mg of Rb. It was connected with the target cell by a 10 mm diameter transfer tube. The cells

Table 2.3: Chemical Composition of Corning 1720 Glass by Weight.

Compound	SiO_2	Na_2O	CaO	MgO	B_2O_3	Al_2O_3	K_2O	As_2O_3
	60.7	1.0	8.6	7.4	5.0	17.3	0.2	0.5
	57	1.0	5.5	12	4.0	20.5		
	62	1	8	7	5	17		
Ave:	59.9	1.0	7.4	8.8	4.7	18.2		

were built from 1720 Corning glass, primarily because of its low permeability to ^{3}He. The chemical composition of the glass coming from three different sources is given in table 2.3 [87]. The new concave windows for the target cells were designed for our experiment. The windows had half of the thickness of the E142 convex design. At the same time they were able to withstand higher ^{3}He pressures, since glass is stronger under compression than under tension. However, they could not sustain the high beam intensity. Six target cells and one reference cell broke after less than a week of operation in the beam. The exact cause of the breaking remains unknown. The targets and some of their parameters are listed in table 2.4.

The thickness of the target cell windows was measured by three different methods: mechanically with a micrometer, by X-ray absorption with a ^{55}Fe and by using the interference between laser light reflections from the two surfaces of the window. The optical methods were the most precise, and their uncertainty of 3% was limited by the non-uniformity of the windows. The mechanical method, abandoned because it scratched or broke some windows, was accurate to 5% (the windows of cell Riker were measured mechanically only, for all others one or both optical methods were used).

The ^{3}He density was measured by two methods. The first relied on measurement of the amount of ^{3}He used for filling the cell. Then, after the cell was sealed, its density was determined by the broadening and shift of the Rb absorption lines by ^{3}He. This way all the surviving cells could be re-measured after the run. The uncertainties of the ^{3}He density measurements were 1.5%. The N_2 densities were measured during the filling process and were known with 5% uncertainty.

Table 2.4: E154 Target Cells.

Cell Name	Runs	Windows [μm]	Length [mm]	^{3}He density [10^{20} cm^{-3}]	N_2 density [10^{18} cm^{-3}]	Survived?
Dave	1201 – 1388	45.0, 53.5	291	2.70	2.3	no
Riker	1412 – 1757	66.5, 60.0	300	2.68	1.4	no
Bob	1777 – 2043	70.1, 69.7	299	2.68	2.3	no
SMC	2050 – 2311	68.8, 64.2	302	2.68	2.4	yes
Generals	2316 – 2594	52.2, 56.7	297	2.63	2.4	no
Hermes	2597 – 2902	46.9, 56.6	295	2.64	2.4	no
Prelims	2903 – 3100	45.1, 42.2	300	2.68	2.3	no
Chance	3101 – 3371	83.0, 83.0	299	2.68	2.3	yes
Picard	3377 – 3788	69.3, 61.6	299	2.64	2.3	yes
Ref. Cell #1	733 – 1757	58.4, 46.5	304			no
Ref. Cell #2	1777 – 2047	73.5, 46.3	297			yes
Ref. Cell #3	2050 – 2596	50.9, 48.5	304			yes
Ref. Cell #4	2597 – 3788	70.8, 64.0	302			yes

2.5.3 Target setup

The pumping chamber of the target was enclosed in a plastic oven and heated to 170-200 °C. That vaporized the Rb to a density of about 10^{14} atoms/cm^3. The Rb was polarized by the lasers located in a specially constructed laser hut near the target. The light from each laser passed through a set of optical components, including a quarter-wave plate, to produce circularly polarized light, before reaching the pumping cell. There were four Argon Ion pumped Ti:Saphire lasers with 20 W total power, and three fiber-coupled diode arrays, each with about 15-17 W of power, continuously polarizing Rb. The ^{3}He, polarized via spin exchange with the Rb in the upper cell, diffused through a 60 mm long and 10 mm in diameter transfer cell to the lower target cell with a time constant of about 10 minutes. The schematic of the target apparatus is shown in fig. 2.8. The endcaps of the cell were cooled by ^{4}He jets to relieve the heating caused by the beam. The temperatures of the target were monitored by seven Resistive Thermal Devices (RTDs) mounted in various places, five on the target and two on the pumping cells. The temperature of the target remained at 60-80 °C. The residual Rb density was on the order of 10^{11} atoms/cm^3, negligible in comparison to the primary target components: 2.6×10^{20} atoms/cm^3 of ^{3}He and 2.4×10^{18} molecules/cm^3 of N_2. The target was mounted in a movable holder and could be placed in the beam with a sub-millimeter accuracy. The holder had another position with a reference cell placed below the target. This cell could be filled remotely with a variable pressure of ^{3}He and was used for dilution factor studies. The



Figure 2.8: Schematic of the E154 target.

whole target assembly was enclosed in the scattering chamber kept under a few mtorr of vacuum. Outside of the scattering chamber were two 1.4 m diameter Helmholtz coils. They produced a 10-30 G holding field to align the nuclear spins of ^{3}He along the beam direction. A similar set of coils was used to produce a holding field perpendicular to the beam direction used for the runs with transverse target polarization for the measurement of $A_\perp$. There were also other sets of coils for polarization measurements.

2.5.4 Polarimetry and polarization results

The target polarization was measured by two independent methods [88]. One was the NMR technique using Adiabatic Fast Passage (AFP). The other used the shift of the Rb Zeeman resonance (EPR) frequency due to the ^{3}He polarization.

The AFP-NMR was used regularly during the run. A set of 42.75 cm diameter Helmholtz drive coils above and below the target provided a 72 mG RF field at 92.0 kHz, while the main holding field was swept from 18 to 36 G, through the Larmor resonance at 28.4 G. The sweep rate of 1.2 G/s was slow enough for the nuclear spins to follow the changing field, but fast enough to avoid the de-phasing of the spins while passing through the resonance. The resulting nuclear spin flip induced a signal in a set of 200 turn copper pick-up coils, wound on a 76.2 mm by 25.4 mm Teflon form and centered around the target cell. The coils connected with a 100 pF capacitor formed an LC resonant circuit. The signal was pre-amplified and the 92 kHz modulation was subtracted by the lock-in amplifier. The resulting NMR signal, see fig. 2.9, was proportional to the ^{3}He polarization. The proportionality constant is determined from the thermal equilibrium Boltzman polarization P_p of protons in a water sample,

$$P_p = \tanh\left(\frac{\mu_p B}{k_B T}\right), \tag{2.18}$$

where B equal 21.61 G for a proton 92 kHz resonance. The proton signals were a few μV, in comparison with 200 mV of ^{3}He. Many sweeps were needed to form an average. The typical water signal is shown in fig. 2.9. The



Figure 2.9: AFP-NMR signals of ^{3}He and water.

uncertainty in the height of that signal (1.8%) together with the uncertainty of the normalized ^{3}He density (1.6%) and the computed coil flux ratio (1.6%) were the dominant contributions to the total of 3.4% uncertainty [83] of the AFP-NMR polarization measurement.

The second method, the EPR polarimetry, was used to calibrate the AFP system. It is based on the shift of the rubidium Zeeman resonance due to the magnetic field created by the polarized ^{3}He . There are two contributions to the EPR resonance shift. The first one is proportional to the ^{3}He polarization coming from the polarization transfer in the Rb–^{3}He spin exchange interaction. The other is due to the classical magnetic field produced by the ^{3}He magnetization. The shifts are substantial, about 20 kHz out of 8 MHz, and easy to measure. By taking the difference in the EPR frequency between two measurements with opposite ^{3}He polarization we could isolate the part proportional to the ^{3}He polarization. The resonance was observed by monitoring the intensity of the fluorescence photons of the D_2 Rb line as a function of the RF frequency [88]. In the highly polarized Rb vapor the atoms are in the $F = 3$, $m = 3$ state (or $m = -3$ for the opposite polarization). The RF field at the EPR frequency corresponding to m transition $3 \rightarrow 2$ increases the $m = 2$ population capable of absorbing the laser light, and thus increases the fluorescence intensity. The typical EPR frequency measurement is shown in fig. 2.10. For the precise determination of the EPR frequency the magnetic field was measured by the Flux-Gate magnetometer and was kept stable at the 10^{-5} level. That enabled the frequency shift to be



Figure 2.10: EPR frequency measurement for both polarization directions (denoted by arrows) for the target cell Picard.

measured with 0.5% accuracy. The proportionality constant, $\kappa_\circ$, between the frequency shift and the ^{3}He polarization was measured [88] within 1%. Other uncertainties include the polarization gradient (1.5%) needed to connect the EPR measured polarization in the pumping cell with the AFP measurement in the target cell, the gas density (1.5%) and the magnetic field shift (1.3%). The total systematic error of the EPR method is 3.0%.

The AFP and EPR polarization measurements disagree by 1.2 σ (combined in quadrature). Assuming Gaussian distributions for the errors, that corresponds to 23% probability of that or a larger difference. We do not exclude the possibility of unknown systematic effects. Thus, rather that taking the average of the results, we conservatively increase the systematic uncertainty so it covers both measurements with their errors. We also include the total systematic drift of the AFP coils calibration constants, equal to 1.7%, of measurements before and after the runs. The final target polarization uncertainty is 4.8%.

2.5.5 Target polarization direction

The direction of the ^{3}He polarization with respect to the beam direction was determined from the orientation of the holding magnetic field with respect to the beam direction and the orientation of the magnetic field with respect to the polarization.

The direction of the magnetic field was determined in four ways: 1. using a compass. 2. Next the sign of the field was measured with the Flux-gate magnetometer. 3. Then with a Hall probe calibrated by the spectrometer magnets. 4. Finally, knowledge of the direction of the coil winding and the electric current also established the magnetic field orientation. All methods gave the same result.

The relative orientation of the polarization to the magnetic field was established in three ways. The first method uses the sign of the AFP signals. The magnetic moments of the protons and ^{3}He have opposite signs. If their spins are initially aligned, then the sweep with the same sign of dB/dt will produce the opposite signs of the NMR signal. That relates the direction of the ^{3}He spin to that of protons, which are aligned with the magnetic field. The second method established the direction of the ^{3}He magnetization due to polarization and thus the direction of the spin from the sign of the EPR shift. Finally, the observation of the masing effect resulting in the depolarization due to the coupling of the ^{3}He spins to the pickup coils gives the polarization orientation. All the methods were consistent.

2.6 Magnetic spectrometers

The new magnetic spectrometers were designed for the E154 and the following E155 experiments at End Station A [89, 90] with the 50 GeV electron beam. The goal was to obtain the highest possible counting rates consistent with the ability to cleanly identify and determine the energy of the scattered electrons in the presence of a large background of pions and low energy neutral particles. An important factor was the short SLED beam pulse width of about 240 ns resulting in 10 times higher instantaneous rates (with the same electrons per spill) in comparison with the previous ESA experiments, E142 and E143.

The system consisted of two independent magnetic spectrometers centered at 2.75° and 5.5° with respect to the incident 48.3 GeV electron beam. That allowed for the kinematic range of $0.014 \leq x \leq 0.8$ in the Bjorken x and $1\ \mathrm{GeV}^2 \leq \mathrm{Q}^2 \leq 17\ \mathrm{GeV}^2$ in the momentum transfer squared. The fixed angles correspond to x and Q^2 values shown in fig. 2.11. The lowest x value, corresponding to the lowest momentum of the scattered electrons, was determined by the tolerable level of the hadronic background and the requirement $y = (E - E')/E \leq 0.8$ dictated by the rapid increase of radiative corrections