

Measurement of the Electron Energy Spectrum and its Moments in Inclusive $B \rightarrow X e \nu$ Decays

- B. Aubert,¹ R. Barate,¹ D. Boutigny,¹ F. Couderc,¹ J.-M. Gaillard,¹ A. Hicheur,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Tisserand,¹ A. Zghiche,¹ A. Palano,² A. Pompili,² J. C. Chen,³ N. D. Qi,³ G. Rong,³ P. Wang,³ Y. S. Zhu,³ G. Eigen,⁴ I. Ofte,⁴ B. Stugu,⁴ G. S. Abrams,⁵ A. W. Borgland,⁵ A. B. Breon,⁵ D. N. Brown,⁵ J. Button-Shafer,⁵ R. N. Cahn,⁵ E. Charles,⁵ C. T. Day,⁵ M. S. Gill,⁵ A. V. Gritsan,⁵ Y. Groysman,⁵ R. G. Jacobsen,⁵ R. W. Kadel,⁵ J. Kadyk,⁵ L. T. Kerth,⁵ Yu. G. Kolomensky,⁵ G. Kukartsev,⁵ C. LeClerc,⁵ M. E. Levi,⁵ G. Lynch,⁵ L. M. Mir,⁵ P. J. Oddone,⁵ T. J. Orimoto,⁵ M. Pripstein,⁵ N. A. Roe,⁵ M. T. Ronan,⁵ V. G. Shelkov,⁵ A. V. Telnov,⁵ W. A. Wenzel,⁵ K. Ford,⁶ T. J. Harrison,⁶ C. M. Hawkes,⁶ S. E. Morgan,⁶ A. T. Watson,⁶ N. K. Watson,⁶ M. Fritsch,⁷ K. Goetzen,⁷ T. Held,⁷ H. Koch,⁷ B. Lewandowski,⁷ M. Pelizaeus,⁷ M. Steinke,⁷ J. T. Boyd,⁸ N. Chevalier,⁸ W. N. Cottingham,⁸ M. P. Kelly,⁸ T. E. Latham,⁸ F. F. Wilson,⁸ K. Abe,⁹ T. Cuhadar-Donszelmann,⁹ C. Hearty,⁹ T. S. Mattison,⁹ J. A. McKenna,⁹ D. Thiessen,⁹ P. Kyberd,¹⁰ L. Teodorescu,¹⁰ V. E. Blinov,¹¹ A. D. Bukin,¹¹ V. P. Druzhinin,¹¹ V. B. Golubev,¹¹ V. N. Ivanchenko,¹¹ E. A. Kravchenko,¹¹ A. P. Onuchin,¹¹ S. I. Serednyakov,¹¹ Yu. I. Skovpen,¹¹ E. P. Solodov,¹¹ A. N. Yushkov,¹¹ D. Best,¹² M. Bruinsma,¹² M. Chao,¹² I. Eschrich,¹² D. Kirkby,¹² A. J. Lankford,¹² M. Mandelkern,¹² R. K. Mommsen,¹² W. Roethel,¹² D. P. Stoker,¹² C. Buchanan,¹³ B. L. Hartfiel,¹³ J. W. Gary,¹⁴ B. C. Shen,¹⁴ K. Wang,¹⁴ D. del Re,¹⁵ H. K. Hadavand,¹⁵ E. J. Hill,¹⁵ D. B. MacFarlane,¹⁵ H. P. Paar,¹⁵ Sh. Rahatlou,¹⁵ V. Sharma,¹⁵ J. W. Berryhill,¹⁶ C. Campagnari,¹⁶ B. Dahmes,¹⁶ S. L. Levy,¹⁶ O. Long,¹⁶ A. Lu,¹⁶ M. A. Mazur,¹⁶ J. D. Richman,¹⁶ W. Verkerke,¹⁶ T. W. Beck,¹⁷ A. M. Eisner,¹⁷ C. A. Heusch,¹⁷ W. S. Lockman,¹⁷ T. Schalk,¹⁷ R. E. Schmitz,¹⁷ B. A. Schumm,¹⁷ A. Seiden,¹⁷ P. Spradlin,¹⁷ D. C. Williams,¹⁷ M. G. Wilson,¹⁷ J. Albert,¹⁸ E. Chen,¹⁸ G. P. Dubois-Felsmann,¹⁸ A. Dvoretskii,¹⁸ D. G. Hitlin,¹⁸ I. Narsky,¹⁸ T. Piatenko,¹⁸ F. C. Porter,¹⁸ A. Ryd,¹⁸ A. Samuel,¹⁸ S. Yang,¹⁸ S. Jayatilleke,¹⁹ G. Mancinelli,¹⁹ B. T. Meadows,¹⁹ M. D. Sokoloff,¹⁹ T. Abe,²⁰ F. Blanc,²⁰ P. Bloom,²⁰ S. Chen,²⁰ P. J. Clark,²⁰ W. T. Ford,²⁰ U. Nauenberg,²⁰ A. Olivas,²⁰ P. Rankin,²⁰ J. G. Smith,²⁰ W. C. van Hoek,²⁰ L. Zhang,²⁰ J. L. Barton,²¹ T. Hu,²¹ A. Soffer,²¹ W. H. Toki,²¹ R. J. Wilson,²¹ D. Altenburg,²² T. Brandt,²² J. Brosse,²² T. Colberg,²² M. Dickopp,²² E. Feltresi,²² A. Hauke,²² H. M. Lacker,²² E. Maly,²² R. Müller-Pfefferkorn,²² R. Nogowski,²² S. Otto,²² J. Schubert,²² K. R. Schubert,²² R. Schwierz,²² B. Spaan,²² D. Bernard,²³ G. R. Bonneauaud,²³ F. Brochard,²³ P. Grenier,²³ Ch. Thiebaux,²³ G. Vasileiadis,²³ M. Verderi,²³ D. J. Bard,²⁴ A. Khan,²⁴ D. Lavin,²⁴ F. Muheim,²⁴ S. Playfer,²⁴ M. Andreotti,²⁵ V. Azzolini,²⁵ D. Bettoni,²⁵ C. Bozzi,²⁵ R. Calabrese,²⁵ G. Cibinetto,²⁵ E. Luppi,²⁵ M. Negrini,²⁵ A. Sarti,²⁵ E. Treadwell,²⁶ R. Baldini-Ferroli,²⁷ A. Calcaterra,²⁷ R. de Sangro,²⁷ G. Finocchiaro,²⁷ P. Patteri,²⁷ M. Piccolo,²⁷ A. Zallo,²⁷ A. Buzzo,²⁸ R. Capra,²⁸ R. Contri,²⁸ G. Crosetti,²⁸ M. Lo Vetere,²⁸ M. Macri,²⁸ M. R. Monge,²⁸ S. Passaggio,²⁸ C. Patrignani,²⁸ E. Robutti,²⁸ A. Santroni,²⁸ S. Tosi,²⁸ S. Bailey,²⁹ G. Brandenburg,²⁹ M. Morii,²⁹ E. Won,²⁹ R. S. Dubitzky,³⁰ U. Langenegger,³⁰ W. Bhimji,³¹ D. A. Bowerman,³¹ P. D. Dauncey,³¹ U. Egede,³¹ J. R. Gaillard,³¹ G. W. Morton,³¹ J. A. Nash,³¹ G. P. Taylor,³¹ G. J. Grenier,³² S.-J. Lee,³² U. Mallik,³² J. Cochran,³³ H. B. Crawley,³³ J. Lamsa,³³ W. T. Meyer,³³ S. Prell,³³ E. I. Rosenberg,³³ J. Yi,³³ M. Davier,³⁴ G. Grosdidier,³⁴ A. Höcker,³⁴ S. Laplace,³⁴ F. Le Diberder,³⁴ V. Lepeltier,³⁴ A. M. Lutz,³⁴ T. C. Petersen,³⁴ S. Plaszczynski,³⁴ M. H. Schune,³⁴ L. Tantot,³⁴ G. Wormser,³⁴ C. H. Cheng,³⁵ D. J. Lange,³⁵ M. C. Simani,³⁵ D. M. Wright,³⁵ A. J. Bevan,³⁶ J. P. Coleman,³⁶ J. R. Fry,³⁶ E. Gabathuler,³⁶ R. Gamet,³⁶ M. Kay,³⁶ R. J. Parry,³⁶ D. J. Payne,³⁶ R. J. Sloane,³⁶ C. Touramanis,³⁶ J. J. Back,³⁷ P. F. Harrison,³⁷ G. B. Mohanty,³⁷ C. L. Brown,³⁸ G. Cowan,³⁸ R. L. Flack,³⁸ H. U. Flaecher,³⁸ S. George,³⁸ M. G. Green,³⁸ A. Kurup,³⁸ C. E. Marker,³⁸ T. R. McMahon,³⁸ S. Ricciardi,³⁸ F. Salvatore,³⁸ G. Vaitsas,³⁸ M. A. Winter,³⁸ D. Brown,³⁹ C. L. Davis,³⁹ J. Allison,⁴⁰ N. R. Barlow,⁴⁰ R. J. Barlow,⁴⁰ P. A. Hart,⁴⁰ M. C. Hodgkinson,⁴⁰ G. D. Lafferty,⁴⁰ A. J. Lyon,⁴⁰ J. C. Williams,⁴⁰ A. Farbin,⁴¹ W. D. Hulsbergen,⁴¹ A. Jawahery,⁴¹ D. Kovalskyi,⁴¹ C. K. Lae,⁴¹ V. Lillard,⁴¹ D. A. Roberts,⁴¹ G. Blaylock,⁴² C. Dallapiccola,⁴² K. T. Flood,⁴² S. S. Hertzbach,⁴² R. Kofler,⁴² V. B. Koptchev,⁴² T. B. Moore,⁴² S. Saremi,⁴² H. Staengle,⁴² S. Willocq,⁴² R. Cowan,⁴³ G. Sciolla,⁴³ F. Taylor,⁴³ R. K. Yamamoto,⁴³ D. J. J. Mangeol,⁴⁴ P. M. Patel,⁴⁴ S. H. Robertson,⁴⁴ A. Lazzaro,⁴⁵ F. Palombo,⁴⁵ J. M. Bauer,⁴⁶ L. Cremaldi,⁴⁶

V. Eschenburg,⁴⁶ R. Godang,⁴⁶ R. Kroeger,⁴⁶ J. Reidy,⁴⁶ D. A. Sanders,⁴⁶ D. J. Summers,⁴⁶ H. W. Zhao,⁴⁶ S. Brunet,⁴⁷ D. Côté,⁴⁷ P. Taras,⁴⁷ H. Nicholson,⁴⁸ C. Cartaro,⁴⁹ N. Cavallo,⁴⁹ F. Fabozzi,^{49,*} C. Gatto,⁴⁹ L. Lista,⁴⁹ D. Monorchio,⁴⁹ P. Paolucci,⁴⁹ D. Piccolo,⁴⁹ C. Sciacca,⁴⁹ M. Baak,⁵⁰ G. Raven,⁵⁰ L. Wilden,⁵⁰ C. P. Jessop,⁵¹ J. M. LoSecco,⁵¹ T. A. Gabriel,⁵² T. Allmendinger,⁵³ B. Brau,⁵³ K. K. Gan,⁵³ K. Honscheid,⁵³ D. Hufnagel,⁵³ H. Kagan,⁵³ R. Kass,⁵³ T. Pulliam,⁵³ R. Ter-Antonyan,⁵³ Q. K. Wong,⁵³ J. Brau,⁵⁴ R. Frey,⁵⁴ O. Igonkina,⁵⁴ C. T. Potter,⁵⁴ N. B. Sinev,⁵⁴ D. Strom,⁵⁴ E. Torrence,⁵⁴ F. Colecchia,⁵⁵ A. Dorigo,⁵⁵ F. Galeazzi,⁵⁵ M. Margoni,⁵⁵ M. Morandin,⁵⁵ M. Posocco,⁵⁵ M. Rotondo,⁵⁵ F. Simonetto,⁵⁵ R. Stroili,⁵⁵ G. Tiozzo,⁵⁵ C. Voci,⁵⁵ M. Benayoun,⁵⁶ H. Briand,⁵⁶ J. Chauveau,⁵⁶ P. David,⁵⁶ Ch. de la Vaissière,⁵⁶ L. Del Buono,⁵⁶ O. Hamon,⁵⁶ M. J. J. John,⁵⁶ Ph. Leruste,⁵⁶ J. Ocariz,⁵⁶ M. Pivk,⁵⁶ L. Roos,⁵⁶ S. T'Jampens,⁵⁶ G. Therin,⁵⁶ P. F. Manfredi,⁵⁷ V. Re,⁵⁷ P. K. Behera,⁵⁸ L. Gladney,⁵⁸ Q. H. Guo,⁵⁸ J. Panetta,⁵⁸ F. Anulli,^{27,59} M. Biasini,⁵⁹ I. M. Peruzzi,^{27,59} M. Pioppi,⁵⁹ C. Angelini,⁶⁰ G. Batignani,⁶⁰ S. Bettarini,⁶⁰ M. Bondioli,⁶⁰ F. Bucci,⁶⁰ G. Calderini,⁶⁰ M. Carpinelli,⁶⁰ V. Del Gamba,⁶⁰ F. Forti,⁶⁰ M. A. Giorgi,⁶⁰ A. Lusiani,⁶⁰ G. Marchiori,⁶⁰ F. Martinez-Vidal,^{60,†} M. Morganti,⁶⁰ N. Neri,⁶⁰ E. Paoloni,⁶⁰ M. Rama,⁶⁰ G. Rizzo,⁶⁰ F. Sandrelli,⁶⁰ J. Walsh,⁶⁰ M. Haire,⁶¹ D. Judd,⁶¹ K. Paick,⁶¹ D. E. Wagoner,⁶¹ N. Danielson,⁶² P. Elmer,⁶² C. Lu,⁶² V. Miftakov,⁶² J. Olsen,⁶² A. J. S. Smith,⁶² E. W. Varnes,⁶² F. Bellini,⁶³ G. Cavoto,^{62,63} R. Faccini,⁶³ F. Ferrarotto,⁶³ F. Ferroni,⁶³ M. Gaspero,⁶³ L. Li Gioi,⁶³ M. A. Mazzoni,⁶³ S. Morganti,⁶³ M. Pierini,⁶³ G. Piredda,⁶³ F. Safai Tehrani,⁶³ C. Voena,⁶³ S. Christ,⁶⁴ G. Wagner,⁶⁴ R. Waldi,⁶⁴ T. Adye,⁶⁵ N. De Groot,⁶⁵ B. Franek,⁶⁵ N. I. Geddes,⁶⁵ G. P. Gopal,⁶⁵ E. O. Olaiya,⁶⁵ S. M. Xella,⁶⁵ R. Aleksan,⁶⁶ S. Emery,⁶⁶ A. Gaidot,⁶⁶ S. F. Ganzhur,⁶⁶ P.-F. Giraud,⁶⁶ G. Hamel de Monchenault,⁶⁶ W. Kozanecki,⁶⁶ M. Langer,⁶⁶ M. Legendre,⁶⁶ G. W. London,⁶⁶ B. Mayer,⁶⁶ G. Schott,⁶⁶ G. Vasseur,⁶⁶ Ch. Yèche,⁶⁶ M. Zito,⁶⁶ M. V. Purohit,⁶⁷ A. W. Weidemann,⁶⁷ F. X. Yumiceva,⁶⁷ D. Aston,⁶⁸ R. Bartoldus,⁶⁸ N. Berger,⁶⁸ A. M. Boyarski,⁶⁸ O. L. Buchmueller,⁶⁸ M. R. Convery,⁶⁸ M. Cristinziani,⁶⁸ G. De Nardo,⁶⁸ D. Dong,⁶⁸ J. Dorfan,⁶⁸ D. Dujmic,⁶⁸ W. Dunwoodie,⁶⁸ E. E. Elsen,⁶⁸ R. C. Field,⁶⁸ T. Glanzman,⁶⁸ S. J. Gowdy,⁶⁸ T. Hadig,⁶⁸ V. Halyo,⁶⁸ T. Hryna'ova,⁶⁸ W. R. Innes,⁶⁸ M. H. Kelsey,⁶⁸ P. Kim,⁶⁸ M. L. Kocian,⁶⁸ D. W. G. S. Leith,⁶⁸ J. Libby,⁶⁸ S. Luitz,⁶⁸ V. Luth,⁶⁸ H. L. Lynch,⁶⁸ H. Marsiske,⁶⁸ R. Messner,⁶⁸ D. R. Muller,⁶⁸ C. P. O'Grady,⁶⁸ V. E. Ozcan,⁶⁸ A. Perazzo,⁶⁸ M. Perl,⁶⁸ S. Petrak,⁶⁸ B. N. Ratcliff,⁶⁸ A. Roodman,⁶⁸ A. A. Salnikov,⁶⁸ R. H. Schindler,⁶⁸ J. Schwiening,⁶⁸ G. Simi,⁶⁸ A. Snyder,⁶⁸ A. Soha,⁶⁸ J. Stelzer,⁶⁸ D. Su,⁶⁸ M. K. Sullivan,⁶⁸ J. Va'vra,⁶⁸ S. R. Wagner,⁶⁸ M. Weaver,⁶⁸ A. J. R. Weinstein,⁶⁸ W. J. Wisniewski,⁶⁸ M. Wittgen,⁶⁸ D. H. Wright,⁶⁸ C. C. Young,⁶⁸ P. R. Burchat,⁶⁹ A. J. Edwards,⁶⁹ T. I. Meyer,⁶⁹ B. A. Petersen,⁶⁹ C. Roat,⁶⁹ S. Ahmed,⁷⁰ M. S. Alam,⁷⁰ J. A. Ernst,⁷⁰ M. A. Saeed,⁷⁰ M. Saleem,⁷⁰ F. R. Wappler,⁷⁰ W. Bugg,⁷¹ M. Krishnamurthy,⁷¹ S. M. Spanier,⁷¹ R. Eckmann,⁷² H. Kim,⁷² J. L. Ritchie,⁷² A. Satpathy,⁷² R. F. Schwitters,⁷² J. M. Izen,⁷³ I. Kitayama,⁷³ X. C. Lou,⁷³ S. Ye,⁷³ F. Bianchi,⁷⁴ M. Bona,⁷⁴ F. Gallo,⁷⁴ D. Gamba,⁷⁴ C. Borean,⁷⁵ L. Bosisio,⁷⁵ F. Cossutti,⁷⁵ G. Della Ricca,⁷⁵ S. Dittongo,⁷⁵ S. Grancagnolo,⁷⁵ L. Lanceri,⁷⁵ P. Poropat,^{75,‡} L. Vitale,⁷⁵ G. Vuagnin,⁷⁵ R. S. Panvini,⁷⁶ Sw. Banerjee,⁷⁷ C. M. Brown,⁷⁷ D. Fortin,⁷⁷ P. D. Jackson,⁷⁷ R. Kowalewski,⁷⁷ J. M. Roney,⁷⁷ H. R. Band,⁷⁸ S. Dasu,⁷⁸ M. Datta,⁷⁸ A. M. Eichenbaum,⁷⁸ J. J. Hollar,⁷⁸ J. R. Johnson,⁷⁸ P. E. Kutter,⁷⁸ H. Li,⁷⁸ R. Liu,⁷⁸ F. Di Lodovico,⁷⁸ A. Mihalyi,⁷⁸ A. K. Mohapatra,⁷⁸ Y. Pan,⁷⁸ R. Prepost,⁷⁸ S. J. Sekula,⁷⁸ P. Tan,⁷⁸ J. H. von Wimmersperg-Toeller,⁷⁸ J. Wu,⁷⁸ S. L. Wu,⁷⁸ Z. Yu,⁷⁸ and H. Neal⁷⁹

(The BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

²Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

³Institute of High Energy Physics, Beijing 100039, China

⁴University of Bergen, Inst. of Physics, N-5007 Bergen, Norway

⁵Lawrence Berkeley National Laboratory and University of California, Berkeley, CA 94720, USA

⁶University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁷Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁸University of Bristol, Bristol BS8 1TL, United Kingdom

⁹University of British Columbia, Vancouver, BC, Canada V6T 1Z1

¹⁰Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹¹Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹²University of California at Irvine, Irvine, CA 92697, USA

¹³University of California at Los Angeles, Los Angeles, CA 90024, USA

¹⁴University of California at Riverside, Riverside, CA 92521, USA

¹⁵University of California at San Diego, La Jolla, CA 92093, USA

- ¹⁶*University of California at Santa Barbara, Santa Barbara, CA 93106, USA*
- ¹⁷*University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, CA 95064, USA*
- ¹⁸*California Institute of Technology, Pasadena, CA 91125, USA*
- ¹⁹*University of Cincinnati, Cincinnati, OH 45221, USA*
- ²⁰*University of Colorado, Boulder, CO 80309, USA*
- ²¹*Colorado State University, Fort Collins, CO 80523, USA*
- ²²*Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany*
- ²³*Ecole Polytechnique, LLR, F-91128 Palaiseau, France*
- ²⁴*University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom*
- ²⁵*Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy*
- ²⁶*Florida A&M University, Tallahassee, FL 32307, USA*
- ²⁷*Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy*
- ²⁸*Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy*
- ²⁹*Harvard University, Cambridge, MA 02138, USA*
- ³⁰*Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany*
- ³¹*Imperial College London, London, SW7 2AZ, United Kingdom*
- ³²*University of Iowa, Iowa City, IA 52242, USA*
- ³³*Iowa State University, Ames, IA 50011-3160, USA*
- ³⁴*Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France*
- ³⁵*Lawrence Livermore National Laboratory, Livermore, CA 94550, USA*
- ³⁶*University of Liverpool, Liverpool L69 7ZE, United Kingdom*
- ³⁷*Queen Mary, University of London, E1 4NS, United Kingdom*
- ³⁸*University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom*
- ³⁹*University of Louisville, Louisville, KY 40292, USA*
- ⁴⁰*University of Manchester, Manchester M13 9PL, United Kingdom*
- ⁴¹*University of Maryland, College Park, MD 20742, USA*
- ⁴²*University of Massachusetts, Amherst, MA 01003, USA*
- ⁴³*Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, MA 02139, USA*
- ⁴⁴*McGill University, Montréal, QC, Canada H3A 2T8*
- ⁴⁵*Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy*
- ⁴⁶*University of Mississippi, University, MS 38677, USA*
- ⁴⁷*Université de Montréal, Laboratoire René J. A. Lévesque, Montréal, QC, Canada H3C 3J7*
- ⁴⁸*Mount Holyoke College, South Hadley, MA 01075, USA*
- ⁴⁹*Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy*
- ⁵⁰*NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands*
- ⁵¹*University of Notre Dame, Notre Dame, IN 46556, USA*
- ⁵²*Oak Ridge National Laboratory, Oak Ridge, TN 37831, USA*
- ⁵³*Ohio State University, Columbus, OH 43210, USA*
- ⁵⁴*University of Oregon, Eugene, OR 97403, USA*
- ⁵⁵*Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy*
- ⁵⁶*Universités Paris VI et VII, Lab de Physique Nucléaire H. E., F-75252 Paris, France*
- ⁵⁷*Università di Pavia, Dipartimento di Elettronica and INFN, I-27100 Pavia, Italy*
- ⁵⁸*University of Pennsylvania, Philadelphia, PA 19104, USA*
- ⁵⁹*Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy*
- ⁶⁰*Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy*
- ⁶¹*Prairie View A&M University, Prairie View, TX 77446, USA*
- ⁶²*Princeton University, Princeton, NJ 08544, USA*
- ⁶³*Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy*
- ⁶⁴*Universität Rostock, D-18051 Rostock, Germany*
- ⁶⁵*Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom*
- ⁶⁶*DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France*
- ⁶⁷*University of South Carolina, Columbia, SC 29208, USA*
- ⁶⁸*Stanford Linear Accelerator Center, Stanford, CA 94309, USA*
- ⁶⁹*Stanford University, Stanford, CA 94305-4060, USA*
- ⁷⁰*State Univ. of New York, Albany, NY 12222, USA*
- ⁷¹*University of Tennessee, Knoxville, TN 37996, USA*
- ⁷²*University of Texas at Austin, Austin, TX 78712, USA*
- ⁷³*University of Texas at Dallas, Richardson, TX 75083, USA*
- ⁷⁴*Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy*
- ⁷⁵*Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy*
- ⁷⁶*Vanderbilt University, Nashville, TN 37235, USA*

⁷⁷*University of Victoria, Victoria, BC, Canada V8W 3P6*

⁷⁸*University of Wisconsin, Madison, WI 53706, USA*

⁷⁹*Yale University, New Haven, CT 06511, USA*

(Dated: March 15, 2003)

We report a measurement of the inclusive electron energy spectrum for semileptonic decays of B mesons in a data sample of 52 million $\Upsilon(4S) \rightarrow B\bar{B}$ decays collected with the *BABAR* detector at the PEP-II asymmetric-energy B -meson factory at SLAC. We determine the branching fraction, first, second, and third moments of the spectrum for lower cut-offs on the electron energy between 0.6 and 1.5 GeV. We measure the partial branching fraction to be $\mathcal{B}(B \rightarrow Xe\nu, E_e > 0.6 \text{ GeV}) = (10.36 \pm 0.06(\text{stat.}) \pm 0.23(\text{sys.}))\%$.

PACS numbers: 12.15.Hh, 11.30.Er, 13.25.Hw

The operator product expansion provides corrections to the relation between the semileptonic B decay rate and the magnitude of the Cabibbo-Kobayashi-Maskawa (CKM) [1] matrix element V_{cb} in the free-quark model [2]. The corrections are expressed in terms of non-perturbative quantities that can be extracted from moments of inclusive distributions. We plan to use the precision measurements of moments of the lepton energy spectra presented here and of hadron mass distributions [3] to determine those parameters and thereby to improve the determination of $|V_{cb}|$ [4].

In this paper, we present a new measurement of the inclusive electron energy spectrum from semileptonic B decays, averaged over charged and neutral B mesons produced at the $\Upsilon(4S)$ resonance. After correcting for charmless semileptonic decays, we derive from this spectrum several moments as a function of the minimum electron energy E_0 ranging from 0.6 GeV to 1.5 GeV, where the lower endpoint is set by the limits of electron identification and prevalence of background. We define $R_i(E_0, \mu)$ as $\int_{E_0}^{\infty} (E_e - \mu)^i (d\Gamma/dE_e) dE_e$, and measure the first moment $M_1(E_0) = R_1(E_0, 0)/R_0(E_0, 0)$, the central moments $M_n(E_0) = R_n(E_0, M_1(E_0))/R_0(E_0, 0)$ for $n = 2, 3$ and the partial branching fraction $\mathcal{B}(E_0) = \tau_B R_0(E_0, 0)$, where τ_B is the average lifetime of charged and neutral B mesons.

The measurements presented here are based on data collected by the *BABAR* detector [5] at the PEP-II asymmetric e^+e^- storage ring; they correspond to an integrated luminosity of 47.4 fb^{-1} on the $\Upsilon(4S)$ resonance and 9.1 fb^{-1} at an energy 40 MeV below the resonance (off-resonance). Where background and efficiency corrections cannot be measured directly from data, we use a full simulation of the detector based on GEANT4 [6].

This analysis is similar to the *BABAR* measurement of the semileptonic branching fraction [7], including use of the same electron identification criteria, but supersedes it by an order of magnitude in integrated luminosity. We identify $B\bar{B}$ events by observing an electron, e_{tag} , with a momentum of $1.4 < p^* < 2.3 \text{ GeV}/c$ in the $\Upsilon(4S)$ rest frame. These electrons make up the tagged

sample that is used as normalization for the branching fraction. A second electron, e_{sig} , for which we require $p^* > 0.5 \text{ GeV}/c$ is assigned either to the unlike-sign sample (charge $Q(e_{tag}) = -Q(e_{sig})$) or to the like-sign sample ($Q(e_{tag}) = Q(e_{sig})$). In events without $B^0\bar{B}^0$ mixing, primary electrons from semileptonic B decays belong to the unlike-sign sample while secondary electrons contribute to the like-sign sample.

Multi-hadron events are selected by either requiring a track multiplicity $N_{ch} \geq 5$, or $N_{ch} = 4$ plus at least two photon candidates with energies above 80 MeV. Track pairs from converted photons are not included in N_{ch} , but count as one photon. For further suppression of non- $B\bar{B}$ events we require $R_2 < 0.8$, where R_2 is the ratio of the second to the zeroth Fox-Wolfram moment [8].

Electrons originating from the same B meson as the tagged electron typically have opposite charge and direction. To reject them we require

$$\cos \alpha > 1.0 - p_e^*(\text{GeV}/c) \quad \text{and} \quad \cos \alpha > -0.2, \quad (1)$$

where α is the angle between the two electrons in the $\Upsilon(4S)$ rest frame. To suppress background contributions from $J/\psi \rightarrow e^+e^-$ decays to the tag-sample, we require the invariant mass M_{ee} of the tag electron, paired with any electron of opposite charge with $\cos \alpha < -0.2$, to be outside the interval $2.9 < M_{ee} < 3.15 \text{ GeV}/c^2$. The efficiencies of these selection criteria are estimated by Monte Carlo (MC) simulation.

Continuum background is subtracted from the tagged, like- and unlike-sign samples by scaling the off-resonance yields by the ratio of on- to off-resonance integrated luminosities, corrected for the energy dependence of the continuum cross section. In the off-resonance sample, the momenta are scaled by the ratio of the on- and off-resonance energies.

Electron spectra from photon conversions and Dalitz decays are extracted from data, taking into account the pair-reconstruction efficiencies from MC simulation. The relative uncertainty in these efficiencies is estimated to be 13% and 19% for conversion and Dalitz pairs, respectively.

The misidentification rates for pions, kaons, and protons are extracted from data control samples. They rise from 0.05% to 0.12% for pions and fall from 0.4% to 0.1% for kaons as p^* increases from 0.5 to 2.5 GeV/c. The systematic errors are estimated from the control sample purities and from the uncertainties in the π , K and p abundances. The resulting relative uncertainties are less than 40%.

There is a small residual background in the sample of unlike-sign pairs originating from the same B meson and fulfilling the requirement on the opening angle α from Eq. 1. It is estimated from a fit to the $\cos\alpha$ distribution, separately for each 50- MeV/c-wide bin in p^* . The distribution is flat for signal pairs, while for background pairs it is taken from MC simulation, with a maximum at $\cos\alpha = -1$ and gradually decreasing to 0 at $\cos\alpha = 1$.




FIG. 1: Measured momentum spectrum (points) and estimated backgrounds (histograms) for electron candidates in (a) the unlike-sign sample, and (b) the like-sign sample.

Fig. 1 shows the electron momentum spectra and the background contributions discussed so far. Further backgrounds arise from decays of τ leptons, charmed mesons produced in $b \rightarrow c\bar{s}$ decays and J/ψ or $\psi(2S) \rightarrow e^+e^-$ decays with only one detected e . These backgrounds are irreducible, and their contributions to the three electron samples are estimated from MC simulations, using the ISGW2 model [9] to describe semileptonic D and D_s meson decays. Assuming $\Gamma(D_s \rightarrow Xe\nu) = \Gamma(D \rightarrow Xe\nu)$, we obtain $\mathcal{B}(D_s \rightarrow Xe\nu) = (8.05 \pm 0.66)\%$. Using 0.84 ± 0.09 [10] for the measured fraction of $B \rightarrow D_sX$ decays where the D_s originates from fragmentation of

the W Boson, and $\mathcal{B}(B \rightarrow D_sX) = (10.5 \pm 2.6)\%$ [11] yields $\mathcal{B}(B^{0,+} \rightarrow D_s^+ \rightarrow e^+) = (0.71 \pm 0.20)\%$. Assuming equal production rates of D and D^* and using $\mathcal{B}(B \rightarrow \overline{D}D^{(*)}X) = (8.2 \pm 1.3)\%$ [10], we arrive at $\mathcal{B}(B^{0,+} \rightarrow D^{0,+} \rightarrow e^+) = (0.84 \pm 0.21)\%$. To estimate the contribution of electrons from τ decays, we consider the cascades $B \rightarrow \tau \rightarrow e$ and $B \rightarrow D_s \rightarrow \tau \rightarrow e$, with branching fractions taken from [11]. The rates for the decays $B \rightarrow J/\psi \rightarrow e^+e^-$ and $B \rightarrow \psi(2S) \rightarrow e^+e^-$ are also adjusted to [11].

These irreducible background spectra are subtracted from the like-sign and unlike-sign spectra after correction for electron identification efficiency. We determine this efficiency as a function of p^* and the polar angle θ^* using $e^+e^- \rightarrow e^+e^-\gamma$ events and then use MC simulation to estimate losses in hadronic events with higher multiplicities. For $p^* > 0.6$ GeV/c, the average efficiency is 91% with an uncertainty of 1.5% estimated from the size of the MC correction. A summary of the yields is given in Table I.

TABLE I: Unlike-sign and like-sign pair yields for $0.6 < p^* < 2.5$ GeV/c and their corrections with statistical and systematic errors. Numbers are quoted after all selection criteria.

| | e^+e^- sample | $e^\pm e^\pm$ sample |
|----------------------|---------------------------|---------------------------|
| All candidates | 183493 ± 434 | 133842 ± 371 |
| continuum bkgd. | 22922 ± 349 | 15758 ± 290 |
| conversion, Dalitz | $2978 \pm 286 \pm 327$ | $10730 \pm 502 \pm 1177$ |
| fake e | $885 \pm 63 \pm 423$ | $2229 \pm 182 \pm 966$ |
| e from same B | $3200 \pm 34 \pm 160$ | |
| e yield | $153508 \pm 630 \pm 558$ | $105126 \pm 712 \pm 1523$ |
| eff. corr. e yield | $169654 \pm 732 \pm 2235$ | $117192 \pm 803 \pm 2510$ |
| irreducible bkgd. | $13912 \pm 92 \pm 1341$ | $14512 \pm 97 \pm 2513$ |
| corr. e yield | $155742 \pm 738 \pm 2606$ | $102680 \pm 809 \pm 3551$ |

To account for $B^0\overline{B}^0$ mixing, we determine the number of primary electrons in the i -th p^* bin from the like-sign and unlike-sign pairs as

$$N_{b \rightarrow c,u}^i = \frac{1 - f_0 \chi_0}{1 - 2f_0 \chi_0} \frac{N_{e^+e^-}^i}{\epsilon_\alpha^i} - \frac{f_0 \chi_0}{1 - 2f_0 \chi_0} N_{e^\pm e^\pm}^i \quad (2)$$

where $\chi_0 = 0.186 \pm 0.004$ [11] is the $B^0\overline{B}^0$ mixing parameter and $f_0 = \mathcal{B}(\Upsilon(4S) \rightarrow B^0\overline{B}^0) = 0.490 \pm 0.018$ [11]. The parameter ϵ_α^i is the efficiency of the additional requirement for the unlike-sign sample as defined in Eq. 1.

The spectrum obtained from Eq. 2 is corrected for the effects of bremsstrahlung in the detector material using MC simulation. Fig. 2 shows the resulting spectrum of primary electrons. Since this correction significantly impacts the first moments, 3% for $E_0 = 0.6$ GeV and 0.5% for $E_0 = 1.5$ GeV, we have verified that the detector material is simulated to better than 3%.




FIG. 2: Electron momentum spectrum from $B \rightarrow X_e \nu(\gamma)$ decays in the $\Upsilon(4S)$ frame after correction for efficiencies and bremsstrahlung, with combined statistical and systematic errors.

Charmless semileptonic $B \rightarrow X_u e \nu$ decays are modeled as in [12] by a combination of semileptonic decays with resonant and non-resonant hadronic systems. Using $\mathcal{B}(B \rightarrow X_u e \nu) = (2.2 \pm 0.5) \times 10^{-3}$ [12] to correct for this background, we determine the moments $\tilde{M}_n = \sum_k p_k^n N_{b \rightarrow c}^k / \sum_k N_{b \rightarrow c}^k$ where k runs over all bins above the energy E_0 and p_k are the bin centers for $n = 1$ and the bin centers shifted by \tilde{M}_1 for $n = 2, 3$. These moments are then transformed into E_e moments M_n by correcting for the movement of the B mesons in the center-of-mass frame. Further biases due to the event selection criteria and binning are estimated from MC simulation. The spectra and moments presented are those of $B \rightarrow X_c e \nu(\gamma)$ decays with any number of photons. The moments as a function of E_0 are shown in Fig. 3 and Table II lists the principal systematic errors for $E_0 = 0.6$ and 1.5 GeV. Without subtraction of $B \rightarrow X_u e \nu$ decays, we measure $M_1^{b \rightarrow x}(1.5 \text{ GeV}) = (1779.0 \pm 1.9 \pm 0.7) \text{ MeV}$, which is consistent with a recent measurement by CLEO [13]. Measurements with $E_0 = 0$ GeV have been performed by DELPHI [14].

We determine the partial branching fraction as $(\sum_k N_{b \rightarrow c,u}^k) / (N_{tag} \epsilon_{evt} \epsilon_{cuts})$, where k runs over all bins with $E_e > E_0$, $N_{tag} = (3616.8 \pm 3.5(\text{stat.}) \pm 21.8(\text{syst.})) \times 10^3$ is the background-corrected number of tag electrons, $\epsilon_{evt} = (98.9 \pm 0.5)\%$ refers to the relative efficiency for selecting two-electron events compared to events with a single e_{tag} , and $\epsilon_{cuts} = (82.8 \pm 0.3)\%$ is the acceptance for the signal electron for $E_0 = 0.6$ GeV. The result,

$$\begin{aligned} \mathcal{B}(B \rightarrow X_e \nu(\gamma), E_e > 0.6 \text{ GeV}) \\ = (10.36 \pm 0.06(\text{stat.}) \pm 0.23(\text{syst.}))\%, \end{aligned}$$

is consistent with our previous measurement [7], with the overall error improved by 25%. The result can be extrapolated to $E_0 = 0$ as part of a combined fit of the Heavy Quark Effective Theory (HQET) parameters to




FIG. 3: Measured moments as a function of the cut-off energy, (a) M_1 and \mathcal{B} , (b) M_2 and M_3 .

the full set of moments [4].

Current theoretical predictions on the lepton energy moments do not incorporate photon emission. Therefore we use PHOTOS [15] to simulate QED radiation and correct the moments for its impact. We verify that radiation that is not included in PHOTOS, e.g. additional hard photons, have no significant effect on the moments. The radiatively corrected moments and the estimated PHOTOS uncertainty [16] are given in Table II. The complete listing of all moments and the full correlation matrix, with and without PHOTOS corrections can be found in Tables III-V. For fitting purposes, a set of tables and matrices with a precision of 5 significant digits can be obtained from the authors.

In summary, we report a measurement of the electron energy spectrum of the inclusive decay $B \rightarrow X_e \nu$ and its branching fraction for electron energies above 0.6 GeV, which supersedes our previous result [7]. We have also derived branching fractions, first, second, and third moments of electron energy spectrum from $B \rightarrow X_c e \nu$ decays for energy thresholds from 0.6 to 1.5 GeV. This set of moments combined with hadron mass moments [3] will be used for a significantly improved determination of HQET parameters and of $|V_{cb}|$ [4].

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from the A. P. Sloan Foundation, Research Corporation, and Alexander von Humboldt Foundation.

TABLE II: Results and breakdown of the systematic errors for $\mathcal{B} = \tau_B \int_{E_0}^{\infty} (d\Gamma/dE_e) dE_e$, and the moments M_1 , M_2 , and M_3 for $B \rightarrow X_c e \nu$ in the B -meson rest frame for two values of E_0 .

| E_0 [GeV] | \mathcal{B} [10 ⁻²] | | M_1 [MeV] | | M_2 [10 ⁻³ GeV ²] | | M_3 [10 ⁻³ GeV ³] | |
|-------------------------------|-----------------------------------|-------|-------------|--------|--|------|--|------|
| | 0.6 | 1.5 | 0.6 | 1.5 | 0.6 | 1.5 | 0.6 | 1.5 |
| conversion and Dalitz pairs | 0.029 | 0.001 | 1.6 | 0.02 | 0.6 | 0.00 | 0.06 | 0.00 |
| e identification efficiency | 0.151 | 0.044 | 2.5 | 0.30 | 0.6 | 0.07 | 0.29 | 0.08 |
| e from same B | 0.019 | 0.000 | 1.3 | 0.00 | 0.6 | 0.00 | 0.03 | 0.00 |
| irreducible background | 0.114 | 0.012 | 5.9 | 0.08 | 2.4 | 0.01 | 0.20 | 0.00 |
| $f_0 \chi_0$ | 0.034 | 0.021 | 0.8 | 0.01 | 0.3 | 0.00 | 0.03 | 0.00 |
| tracking efficiency | 0.084 | 0.033 | 1.0 | 0.06 | 0.3 | 0.02 | 0.07 | 0.00 |
| bremsstrahlung correction | 0.011 | 0.028 | 1.9 | 0.43 | 0.0 | 0.05 | 0.19 | 0.00 |
| event selection | 0.052 | 0.024 | 0.6 | 0.14 | 0.0 | 0.03 | 0.07 | 0.01 |
| $b \rightarrow u$ subtraction | 0.047 | 0.030 | 1.2 | 1.24 | 0.6 | 0.48 | 0.20 | 0.17 |
| B momentum correction | 0.000 | 0.005 | 0.0 | 0.19 | 0.1 | 0.10 | 0.04 | 0.02 |
| N_{tag} normalization | 0.068 | 0.030 | | | | | | |
| Moments | 10.17 | 4.54 | 1414.3 | 1769.2 | 148.5 | 29.8 | -9.97 | 2.11 |
| \pm (stat.) | 0.06 | 0.03 | 3.7 | 1.8 | 2.0 | 0.8 | 0.79 | 0.44 |
| \pm (sys.) | 0.23 | 0.08 | 7.4 | 1.4 | 2.7 | 0.5 | 0.48 | 0.20 |
| Moments with rad. correction | 10.30 | 4.79 | 1432.8 | 1774.3 | 148.0 | 30.3 | -12.05 | 2.12 |
| \pm (stat.) | 0.06 | 0.03 | 3.9 | 1.9 | 2.2 | 0.9 | 0.88 | 0.47 |
| \pm (sys.) | 0.24 | 0.09 | 7.8 | 1.4 | 3.1 | 0.5 | 0.46 | 0.20 |

TABLE III: Measured moments M_1 , M_2 , M_3 , and \mathcal{B} for five cut-off energies E_0 with first their statistical error and second their systematic error. The moments are given in the B -meson rest frame and are defined to include X_c hadronic states only and to include decays $B \rightarrow X_c e \nu \gamma$ with any number of photons.

| E_0 [GeV] | \mathcal{B} [10 ⁻²] | M_1 [MeV] | M_2 [10 ⁻³ GeV ²] | M_3 [10 ⁻³ GeV ³] |
|-------------|-----------------------------------|--------------------------|--|--|
| 0.6 | $10.17 \pm 0.06 \pm 0.23$ | $1414.3 \pm 3.7 \pm 7.4$ | $148.5 \pm 2.0 \pm 2.7$ | $-9.97 \pm 0.79 \pm 0.48$ |
| 0.8 | $9.43 \pm 0.05 \pm 0.19$ | $1469.8 \pm 2.4 \pm 3.6$ | $117.3 \pm 1.2 \pm 0.9$ | $-1.82 \pm 0.60 \pm 0.39$ |
| 1.0 | $8.42 \pm 0.04 \pm 0.16$ | $1537.2 \pm 1.9 \pm 2.1$ | $88.3 \pm 0.9 \pm 0.6$ | $1.95 \pm 0.53 \pm 0.30$ |
| 1.2 | $7.05 \pm 0.04 \pm 0.13$ | $1621.3 \pm 1.8 \pm 1.6$ | $61.2 \pm 0.9 \pm 0.6$ | $2.97 \pm 0.49 \pm 0.24$ |
| 1.5 | $4.54 \pm 0.03 \pm 0.08$ | $1769.2 \pm 1.8 \pm 1.4$ | $29.8 \pm 0.8 \pm 0.5$ | $2.11 \pm 0.44 \pm 0.20$ |

TABLE IV: Measured moments M_1 , M_2 , M_3 , and \mathcal{B} for five cut-off energies E_0 with first their statistical error and second their systematic error. The moments are given in the B -meson rest frame and are corrected for QED radiative effects.

| E_0 [GeV] | \mathcal{B} [10 ⁻²] | M_1 [MeV] | M_2 [10 ⁻³ GeV ²] | M_3 [10 ⁻³ GeV ³] |
|-------------|-----------------------------------|--------------------------|--|--|
| 0.6 | $10.30 \pm 0.06 \pm 0.24$ | $1432.8 \pm 3.9 \pm 7.8$ | $148.0 \pm 2.2 \pm 3.1$ | $-12.05 \pm 0.88 \pm 0.46$ |
| 0.8 | $9.61 \pm 0.05 \pm 0.20$ | $1484.8 \pm 2.6 \pm 3.7$ | $117.7 \pm 1.3 \pm 1.0$ | $-3.18 \pm 0.64 \pm 0.38$ |
| 1.0 | $8.65 \pm 0.04 \pm 0.17$ | $1548.7 \pm 2.0 \pm 2.2$ | $89.1 \pm 1.0 \pm 0.6$ | $1.19 \pm 0.57 \pm 0.30$ |
| 1.2 | $7.31 \pm 0.04 \pm 0.14$ | $1629.9 \pm 1.9 \pm 1.7$ | $62.1 \pm 0.9 \pm 0.6$ | $2.66 \pm 0.52 \pm 0.24$ |
| 1.5 | $4.79 \pm 0.03 \pm 0.09$ | $1774.3 \pm 1.9 \pm 1.4$ | $30.3 \pm 0.9 \pm 0.5$ | $2.12 \pm 0.47 \pm 0.20$ |

TABLE V: Correlation matrix of the 20 measured moments. Matrix elements are for the full errors (statistical and systematic added in quadrature) in percent. The superscript i refers to the energy thresholds $E_i = 0.6, 0.8, 1.0, 1.2, 1.5$ GeV for $i = 1, 2, 3, 4, 5$ respectively.

| | M_1^1 | M_1^2 | M_1^3 | M_1^4 | M_1^5 | M_2^1 | M_2^2 | M_2^3 | M_2^4 | M_2^5 | M_3^1 | M_3^2 | M_3^3 | M_3^4 | M_3^5 | \mathcal{B}^1 | \mathcal{B}^2 | \mathcal{B}^3 | \mathcal{B}^4 | \mathcal{B}^5 |
|-----------------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|-----------------|-----------------|-----------------|-----------------|-----------------|
| M_1^1 | 100.0 | 83.2 | 64.0 | 48.3 | 34.0 | -73.6 | -26.5 | 12.3 | 22.5 | 23.4 | 13.5 | -9.2 | 0.7 | 10.5 | 15.7 | -50.8 | -26.4 | -11.8 | -2.5 | 8.0 |
| M_1^2 | | 100.0 | 81.4 | 66.8 | 51.5 | -35.1 | -20.0 | 27.3 | 37.8 | 38.3 | -7.3 | -0.3 | 9.7 | 21.3 | 27.2 | -36.5 | -24.1 | -7.4 | 3.3 | 16.7 |
| M_1^3 | | | 100.0 | 84.0 | 68.5 | -3.5 | 26.6 | 40.4 | 53.9 | 54.5 | 5.5 | 6.6 | 21.0 | 34.3 | 41.1 | -16.7 | -7.2 | -0.8 | 12.0 | 27.8 |
| M_1^4 | | | | 100.0 | 79.1 | 15.5 | 49.0 | 69.8 | 61.9 | 65.7 | 23.8 | 28.0 | 33.9 | 43.2 | 51.6 | -0.3 | 6.7 | 12.1 | 15.9 | 35.1 |
| M_1^5 | | | | | 100.0 | 28.7 | 63.1 | 85.6 | 88.8 | 79.7 | 48.7 | 59.7 | 64.4 | 66.4 | 65.1 | 17.6 | 23.4 | 27.5 | 31.5 | 36.3 |
| M_2^1 | | | | | | 100.0 | 66.5 | 42.8 | 35.2 | 33.4 | 3.9 | 41.9 | 38.0 | 33.4 | 30.9 | 66.1 | 44.1 | 33.9 | 29.6 | 28.0 |
| M_2^2 | | | | | | | 100.0 | 75.2 | 68.9 | 67.3 | 59.0 | 61.0 | 63.1 | 61.4 | 60.6 | 53.9 | 51.4 | 40.3 | 38.0 | 41.1 |
| M_2^3 | | | | | | | | 100.0 | 90.9 | 88.6 | 66.7 | 81.3 | 79.7 | 78.9 | 78.8 | 36.3 | 39.4 | 41.9 | 39.3 | 45.1 |
| M_2^4 | | | | | | | | | 100.0 | 94.5 | 66.3 | 82.1 | 87.4 | 87.3 | 84.9 | 30.3 | 35.3 | 38.4 | 41.9 | 44.0 |
| M_2^5 | | | | | | | | | | 100.0 | 67.4 | 83.8 | 90.8 | 94.5 | 93.6 | 29.0 | 34.3 | 37.3 | 40.1 | 45.3 |
| M_3^1 | | | | | | | | | | | 100.0 | 79.0 | 76.8 | 72.2 | 68.2 | 31.4 | 43.7 | 41.6 | 38.2 | 34.4 |
| M_3^2 | | | | | | | | | | | | 100.0 | 95.9 | 90.5 | 85.2 | 47.5 | 48.3 | 48.5 | 44.5 | 39.5 |
| M_3^3 | | | | | | | | | | | | | 100.0 | 97.4 | 92.8 | 41.4 | 44.0 | 44.8 | 44.0 | 40.0 |
| M_3^4 | | | | | | | | | | | | | | 100.0 | 97.5 | 33.7 | 37.4 | 39.1 | 40.6 | 39.5 |
| M_3^5 | | | | | | | | | | | | | | | 100.0 | 28.3 | 32.4 | 34.5 | 36.1 | 39.3 |
| \mathcal{B}^1 | | | | | | | | | | | | | | | | 100.0 | 95.4 | 90.4 | 86.3 | 80.1 |
| \mathcal{B}^2 | | | | | | | | | | | | | | | | | 100.0 | 97.7 | 95.1 | 90.1 |
| \mathcal{B}^3 | | | | | | | | | | | | | | | | | | 100.0 | 98.3 | 94.3 |
| \mathcal{B}^4 | | | | | | | | | | | | | | | | | | | 100.0 | 96.5 |
| \mathcal{B}^5 | | | | | | | | | | | | | | | | | | | | 100.0 |

* Also with Università della Basilicata, Potenza, Italy
 † Also with IFIC, Instituto de Física Corpuscular, CSIC-Universidad de Valencia, Valencia, Spain
 ‡ Deceased

- [1] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [2] A. V. Manohar and M. B. Wise, Phys. Rev. **D49**, 1310 (1994).
- [3] B. Aubert *et al.* (BABAR Collaboration), *Measurement of Moments of the Hadronic Mass Distribution in Semileptonic B Decays*, to be submitted to Phys. Rev. **D** (2004).
- [4] B. Aubert *et al.* (BABAR Collaboration), *Determination of $|V_{cb}|$ using Lepton Energy and Hadronic Mass Moments in Semileptonic B Decays*, to be submitted to Phys. Rev. Lett. (2004).
- [5] B. Aubert *et al.* (BABAR Collaboration), Nucl. Instr. Meth. **A479**, 1 (2003.)
- [6] GEANT4 Collaboration, S. Agostinelli *et al.*, Nucl. Instr. Meth. **A506**, 250 (2003.)
- [7] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. **D67**, 031101 (2003).
- [8] G.C. Fox and S. Wolfram, Phys. Rev. Lett. **41**, 1581 (1978)
- [9] D. Scora and N. Isgur, Phys. Rev. **D52**, 2783 (1995).
- [10] The LEP Electroweak Working Group, hep-ex/0112028.
- [11] K. Hagiwara *et al.*, Phys. Rev. **D 66**, 010001 (2002) and 2003 off-year partial update for the 2004 edition (URL: <http://pdg.lbl.gov/>).
- [12] B. Aubert *et al.* (BABAR Collaboration), (2003).
- [13] A. H. Mahmood *et al.* (CLEO Collaboration), Phys. Rev. **D 67**, 072001 (2003).
- [14] M. Battaglia *et al.*, Phys. Lett. **B556** (2003) 41.
- [15] E. Barberio and Z. Was, Comp. Phys. Commun. **79**, 291 (1994).
- [16] E. Richter-Was, Phys. Lett. **B 303** (1993) 163.