## Measurement of the Ratio of Decay Amplitudes for $\overline{B}{}^0 \to J/\psi K^{*0}$ and $B^0 \to J/\psi K^{*0}$

B. Aubert,<sup>1</sup> R. Barate,<sup>1</sup> D. Boutigny,<sup>1</sup> F. Couderc,<sup>1</sup> J.-M. Gaillard,<sup>1</sup> A. Hicheur,<sup>1</sup> Y. Karyotakis,<sup>1</sup> J. P. Lees,<sup>1</sup> V. Tisserand,<sup>1</sup> A. Zghiche,<sup>1</sup> A. Palano,<sup>2</sup> A. Pompili,<sup>2</sup> J. C. Chen,<sup>3</sup> N. D. Qi,<sup>3</sup> G. Rong,<sup>3</sup> P. Wang,<sup>3</sup> Y. S. Zhu,<sup>3</sup> G. Eigen,<sup>4</sup> I. Ofte,<sup>4</sup> B. Stugu,<sup>4</sup> G. S. Abrams,<sup>5</sup> A. W. Borgland,<sup>5</sup> A. B. Breon,<sup>5</sup> D. N. Brown,<sup>5</sup> J. Button-Shafer,<sup>5</sup> R. N. Cahn,<sup>5</sup> E. Charles,<sup>5</sup> C. T. Day,<sup>5</sup> M. S. Gill,<sup>5</sup> A. V. Gritsan,<sup>5</sup> Y. Groysman,<sup>5</sup> R. G. Jacobsen,<sup>5</sup> R. W. Kadel,<sup>5</sup> J. Kadyk,<sup>5</sup> L. T. Kerth,<sup>5</sup> Yu. G. Kolomensky,<sup>5</sup> G. Kukartsev,<sup>5</sup> G. Lynch,<sup>5</sup> L. M. Mir,<sup>5</sup> P. J. Oddone,<sup>5</sup> T. J. Orimoto,<sup>5</sup> M. Pripstein,<sup>5</sup> N. A. Roe,<sup>5</sup> M. T. Ronan,<sup>5</sup> V. G. Shelkov,<sup>5</sup> W. A. Wenzel,<sup>5</sup> K. E. Ford,<sup>6</sup> T. J. Harrison,<sup>6</sup> C. M. Hawkes,<sup>6</sup> S. E. Morgan,<sup>6</sup> A. T. Watson,<sup>6</sup> M. Fritsch,<sup>7</sup> K. Goetzen,<sup>7</sup> T. Held,<sup>7</sup> H. Koch,<sup>7</sup> B. Lewandowski,<sup>7</sup> M. Pelizaeus,<sup>7</sup> M. Steinke,<sup>7</sup> J. T. Boyd,<sup>8</sup> N. Chevalier,<sup>8</sup> W. N. Cottingham,<sup>8</sup> M. P. Kelly,<sup>8</sup> T. E. Latham,<sup>8</sup> F. F. Wilson,<sup>8</sup> T. Cuhadar-Donszelmann,<sup>9</sup> C. Hearty,<sup>9</sup> N. S. Knecht,<sup>9</sup> T. S. Mattison,<sup>9</sup> J. A. McKenna,<sup>9</sup> D. Thiessen,<sup>9</sup> A. Khan,<sup>10</sup> P. Kyberd,<sup>10</sup> L. Teodorescu,<sup>10</sup> V. E. Blinov,<sup>11</sup> A. D. Bukin,<sup>11</sup> V. P. Druzhinin,<sup>11</sup> V. B. Golubev,<sup>11</sup> V. N. Ivanchenko,<sup>11</sup> E. A. Kravchenko,<sup>11</sup> A. P. Onuchin,<sup>11</sup> S. I. Serednyakov,<sup>11</sup> Yu. I. Skovpen,<sup>11</sup> E. P. Solodov,<sup>11</sup> A. N. Yushkov,<sup>11</sup> D. Best,<sup>12</sup> M. Bruinsma,<sup>12</sup> M. Chao,<sup>12</sup> I. Eschrich,<sup>12</sup> D. Kirkby,<sup>12</sup> A. J. Lankford,<sup>12</sup> M. Mandelkern,<sup>12</sup> R. K. Mommsen,<sup>12</sup> W. Roethel,<sup>12</sup> D. P. Stoker,<sup>12</sup> C. Buchanan,<sup>13</sup> B. L. Hartfiel,<sup>13</sup> J. W. Gary,<sup>14</sup> B. C. Shen,<sup>14</sup> K. Wang,<sup>14</sup> D. del Re,<sup>15</sup> H. K. Hadavand,<sup>15</sup> E. J. Hill,<sup>15</sup> D. B. MacFarlane,<sup>15</sup> H. P. Paar,<sup>15</sup> Sh. Rahatlou,<sup>15</sup> V. Sharma,<sup>15</sup> J. W. Berryhill,<sup>16</sup> C. Campagnari,<sup>16</sup> B. Dahmes,<sup>16</sup> S. L. Levy,<sup>16</sup> O. Long,<sup>16</sup> A. Lu,<sup>16</sup> M. A. Mazur,<sup>16</sup> J. D. Richman,<sup>16</sup> W. Verkerke,<sup>16</sup> T. W. Beck,<sup>17</sup> A. M. Eisner,<sup>17</sup> C. A. Heusch,<sup>17</sup> W. S. Lockman,<sup>17</sup> T. Schalk,<sup>17</sup> R. E. Schmitz,<sup>17</sup> B. A. Schumm,<sup>17</sup> A. Seiden,<sup>17</sup> P. Spradlin,<sup>17</sup> D. C. Williams,<sup>17</sup> M. G. Wilson,<sup>17</sup> J. Albert,<sup>18</sup> E. Chen,<sup>18</sup> G. P. Dubois-Felsmann,<sup>18</sup> A. Dvoretskii,<sup>18</sup> D. G. Hitlin,<sup>18</sup> I. Narsky,<sup>18</sup> T. Piatenko,<sup>18</sup> F. C. Porter,<sup>18</sup> A. Ryd,<sup>18</sup> A. Samuel,<sup>18</sup> S. Yang,<sup>18</sup> S. Jayatilleke,<sup>19</sup> G. Mancinelli,<sup>19</sup> B. T. Meadows,<sup>19</sup> M. D. Sokoloff,<sup>19</sup> T. Abe,<sup>20</sup> F. Blanc,<sup>20</sup> P. Bloom,<sup>20</sup> S. Chen,<sup>20</sup> W. T. Ford,<sup>20</sup> U. Nauenberg,<sup>20</sup> A. Olivas,<sup>20</sup> P. Rankin,<sup>20</sup> J. G. Smith,<sup>20</sup> J. Zhang,<sup>20</sup> L. Zhang,<sup>20</sup> A. Chen,<sup>21</sup> J. L. Harton,<sup>21</sup> A. Soffer,<sup>21</sup> W. H. Toki,<sup>21</sup> R. J. Wilson,<sup>21</sup> Q. L. Zeng,<sup>21</sup> D. Altenburg,<sup>22</sup> T. Brandt,<sup>22</sup> J. Brose,<sup>22</sup> T. Colberg,<sup>22</sup> M. Dickopp,<sup>22</sup> E. Feltresi,<sup>22</sup> A. Hauke,<sup>22</sup> H. M. Lacker,<sup>22</sup> E. Maly,<sup>22</sup> R. Müller-Pfefferkorn,<sup>22</sup> R. Nogowski,<sup>22</sup> S. Otto,<sup>22</sup> A. Petzold,<sup>22</sup> J. Schubert,<sup>22</sup> K. R. Schubert,<sup>22</sup> R. Schwierz,<sup>22</sup> B. Spaan,<sup>22</sup> J. E. Sundermann,<sup>22</sup> D. Bernard,<sup>23</sup> G. R. Bonneaud,<sup>23</sup> F. Brochard,<sup>23</sup> P. Grenier,<sup>23</sup> S. Schrenk,<sup>23</sup> Ch. Thiebaux,<sup>23</sup> G. Vasileiadis,<sup>23</sup> M. Verderi,<sup>23</sup> D. J. Bard,<sup>24</sup> P. J. Clark,<sup>24</sup> D. Lavin,<sup>24</sup> F. Muheim,<sup>24</sup> S. Playfer,<sup>24</sup> Y. Xie,<sup>24</sup> M. Andreotti,<sup>25</sup> V. Azzolini,<sup>25</sup> D. Bettoni,<sup>25</sup> C. Bozzi,<sup>25</sup> R. Calabrese,<sup>25</sup> G. Cibinetto,<sup>25</sup> E. Luppi,<sup>25</sup> M. Negrini,<sup>25</sup> L. Piemontese,<sup>25</sup> A. Sarti,<sup>25</sup> E. Treadwell,<sup>26</sup> R. Baldini-Ferroli,<sup>27</sup> A. Calcaterra,<sup>27</sup> R. de Sangro,<sup>27</sup> G. Finocchiaro,<sup>27</sup> P. Patteri,<sup>27</sup> M. Piccolo,<sup>27</sup> A. Zallo,<sup>27</sup> A. Buzzo,<sup>28</sup> R. Capra,<sup>28</sup> R. Contri,<sup>28</sup> G. Crosetti,<sup>28</sup> M. Lo Vetere,<sup>28</sup> M. Macri,<sup>28</sup> M. R. Monge,<sup>28</sup> S. Passaggio,<sup>28</sup> C. Patrignani,<sup>28</sup> E. Robutti,<sup>28</sup> A. Santroni,<sup>28</sup> S. Tosi,<sup>28</sup> S. Bailey,<sup>29</sup> G. Brandenburg,<sup>29</sup> M. Morii,<sup>29</sup> E. Won,<sup>29</sup> R. S. Dubitzky,<sup>30</sup> U. Langenegger,<sup>30</sup> W. Bhimji,<sup>31</sup> D. A. Bowerman,<sup>31</sup> P. D. Dauncey,<sup>31</sup> U. Egede,<sup>31</sup> J. R. Gaillard,<sup>31</sup> G. W. Morton,<sup>31</sup> J. A. Nash,<sup>31</sup> G. P. Taylor,<sup>31</sup> M. J. Charles,<sup>32</sup> G. J. Grenier,<sup>32</sup> U. Mallik,<sup>32</sup> J. Cochran,<sup>33</sup> H. B. Crawley,<sup>33</sup> J. Lamsa,<sup>33</sup> W. T. Meyer,<sup>33</sup> S. Prell,<sup>33</sup> E. I. Rosenberg,<sup>33</sup> J. Yi,<sup>33</sup> M. Davier,<sup>34</sup> G. Grosdidier,<sup>34</sup> A. Höcker,<sup>34</sup> S. Laplace,<sup>34</sup> F. Le Diberder,<sup>34</sup> V. Lepeltier,<sup>34</sup> A. M. Lutz,<sup>34</sup> T. C. Petersen,<sup>34</sup> S. Plaszczynski,<sup>34</sup> M. H. Schune,<sup>34</sup> L. Tantot,<sup>34</sup> G. Wormser,<sup>34</sup> C. H. Cheng,<sup>35</sup> D. J. Lange,<sup>35</sup> M. C. Simani,<sup>35</sup> D. M. Wright,<sup>35</sup> A. J. Bevan,<sup>36</sup> J. P. Coleman,<sup>36</sup> J. R. Fry,<sup>36</sup> E. Gabathuler,<sup>36</sup> R. Gamet,<sup>36</sup> R. J. Parry,<sup>36</sup> D. J. Payne,<sup>36</sup> R. J. Sloane,<sup>36</sup> C. Touramanis,<sup>36</sup> J. J. Back,<sup>37</sup> C. M. Cormack,<sup>37</sup> P. F. Harrison,<sup>37, \*</sup> G. B. Mohanty,<sup>37</sup> C. L. Brown,<sup>38</sup> G. Cowan,<sup>38</sup> R. L. Flack,<sup>38</sup> H. U. Flaecher,<sup>38</sup> M. G. Green,<sup>38</sup> C. E. Marker,<sup>38</sup> T. R. McMahon,<sup>38</sup> S. Ricciardi,<sup>38</sup> F. Salvatore,<sup>38</sup> G. Vaitsas,<sup>38</sup> M. A. Winter,<sup>38</sup> D. Brown,<sup>39</sup> C. L. Davis,<sup>39</sup> J. Allison,<sup>40</sup> N. R. Barlow,<sup>40</sup> R. J. Barlow,<sup>40</sup> P. A. Hart,<sup>40</sup> M. C. Hodgkinson,<sup>40</sup> G. D. Lafferty,<sup>40</sup> A. J. Lyon,<sup>40</sup> J. C. Williams,<sup>40</sup> A. Farbin,<sup>41</sup> W. D. Hulsbergen,<sup>41</sup> A. Jawahery,<sup>41</sup> D. Kovalskyi,<sup>41</sup> C. K. Lae,<sup>41</sup> V. Lillard,<sup>41</sup> D. A. Roberts,<sup>41</sup> G. Blaylock,<sup>42</sup> C. Dallapiccola,<sup>42</sup> K. T. Flood,<sup>42</sup> S. S. Hertzbach,<sup>42</sup> R. Kofler,<sup>42</sup> V. B. Koptchev,<sup>42</sup> T. B. Moore,<sup>42</sup> S. Saremi,<sup>42</sup> H. Staengle,<sup>42</sup> S. Willocq,<sup>42</sup> R. Cowan,<sup>43</sup> G. Sciolla,<sup>43</sup> F. Taylor,<sup>43</sup> R. K. Yamamoto,<sup>43</sup> D. J. J. Mangeol,<sup>44</sup> P. M. Patel,<sup>44</sup> S. H. Robertson,<sup>44</sup> A. Lazzaro,<sup>45</sup> F. Palombo,<sup>45</sup> J. M. Bauer,<sup>46</sup> L. Cremaldi,<sup>46</sup> V. Eschenburg,<sup>46</sup> R. Godang,<sup>46</sup> R. Kroeger,<sup>46</sup> J. Reidy,<sup>46</sup> D. A. Sanders,<sup>46</sup> D. J. Summers,<sup>46</sup> H. W. Zhao,<sup>46</sup> S. Brunet,<sup>47</sup> D. Côté,<sup>47</sup>

P. Taras,<sup>47</sup> H. Nicholson,<sup>48</sup> N. Cavallo,<sup>49</sup> F. Fabozzi,<sup>49,†</sup> C. Gatto,<sup>49</sup> L. Lista,<sup>49</sup> D. Monorchio,<sup>49</sup> P. Paolucci,<sup>49</sup> D. Piccolo,<sup>49</sup> C. Sciacca,<sup>49</sup> M. Baak,<sup>50</sup> H. Bulten,<sup>50</sup> G. Raven,<sup>50</sup> L. Wilden,<sup>50</sup> C. P. Jessop,<sup>51</sup> J. M. LoSecco,<sup>51</sup> T. A. Gabriel,<sup>52</sup> T. Allmendinger,<sup>53</sup> B. Brau,<sup>53</sup> K. K. Gan,<sup>53</sup> K. Honscheid,<sup>53</sup> D. Hufnagel,<sup>53</sup> H. Kagan,<sup>53</sup> R. Kass,<sup>53</sup> T. Pulliam,<sup>53</sup> A. M. Rahimi,<sup>53</sup> R. Ter-Antonyan,<sup>53</sup> Q. K. Wong,<sup>53</sup> J. Brau,<sup>54</sup> R. Frey,<sup>54</sup> O. Igonkina,<sup>54</sup> C. T. Potter,<sup>54</sup> N. B. Sinev,<sup>54</sup> D. Strom,<sup>54</sup> E. Torrence,<sup>54</sup> F. Colecchia,<sup>55</sup> A. Dorigo,<sup>55</sup> F. Galeazzi,<sup>55</sup> M. Margoni,<sup>55</sup> M. Morandin,<sup>55</sup> M. Posocco,<sup>55</sup> M. Rotondo,<sup>55</sup> F. Simonetto,<sup>55</sup> R. Stroili,<sup>55</sup> G. Tiozzo,<sup>55</sup> C. Voci,<sup>55</sup> M. Benavoun,<sup>56</sup> H. Briand,<sup>56</sup> J. Chauveau,<sup>56</sup> P. David,<sup>56</sup> Ch. de la Vaissière,<sup>56</sup> L. Del Buono,<sup>56</sup> O. Hamon,<sup>56</sup> M. J. J. John,<sup>56</sup> Ph. Leruste,<sup>56</sup> J. Malcles,<sup>56</sup> J. Ocariz,<sup>56</sup> M. Pivk,<sup>56</sup> L. Roos,<sup>56</sup> S. T'Jampens,<sup>56</sup> G. Therin,<sup>56</sup> P. F. Manfredi,<sup>57</sup> V. Re,<sup>57</sup> P. K. Behera,<sup>58</sup> L. Gladney,<sup>58</sup> Q. H. Guo,<sup>58</sup> J. Panetta,<sup>58</sup> F. Anulli,<sup>27,59</sup> M. Biasini,<sup>59</sup> I. M. Peruzzi,<sup>27,59</sup> M. Pioppi,<sup>59</sup> C. Angelini,<sup>60</sup> G. Batignani,<sup>60</sup> S. Bettarini,<sup>60</sup> M. Bondioli,<sup>60</sup> F. Bucci,<sup>60</sup> G. Calderini,<sup>60</sup> M. Carpinelli,<sup>60</sup> V. Del Gamba,<sup>60</sup> F. Forti,<sup>60</sup> M. A. Giorgi,<sup>60</sup> A. Lusiani,<sup>60</sup> G. Marchiori,<sup>60</sup> F. Martinez-Vidal,<sup>60,‡</sup> M. Morganti,<sup>60</sup> N. Neri,<sup>60</sup> E. Paoloni,<sup>60</sup> M. Rama,<sup>60</sup> G. Rizzo,<sup>60</sup> F. Sandrelli,<sup>60</sup> J. Walsh,<sup>60</sup> M. Haire,<sup>61</sup> D. Judd,<sup>61</sup> K. Paick,<sup>61</sup> D. E. Wagoner,<sup>61</sup> N. Danielson,<sup>62</sup> P. Elmer,<sup>62</sup> Y. P. Lau,<sup>62</sup> C. Lu,<sup>62</sup> V. Miftakov,<sup>62</sup> J. Olsen,<sup>62</sup> A. J. S. Smith,<sup>62</sup> A. V. Telnov,<sup>62</sup> F. Bellini,<sup>63</sup> G. Cavoto,<sup>62,63</sup> R. Faccini,<sup>63</sup> F. Ferrarotto,<sup>63</sup> F. Ferroni,<sup>63</sup> M. Gaspero,<sup>63</sup> L. Li Gioi,<sup>63</sup> M. A. Mazzoni,<sup>63</sup> S. Morganti,<sup>63</sup> M. Pierini,<sup>63</sup> G. Piredda,<sup>63</sup> F. Safai Tehrani,<sup>63</sup> C. Voena,<sup>63</sup> S. Christ,<sup>64</sup> G. Wagner,<sup>64</sup> R. Waldi,<sup>64</sup> T. Adye,<sup>65</sup> N. De Groot,<sup>65</sup> B. Franek,<sup>65</sup> N. I. Geddes,<sup>65</sup> G. P. Gopal,<sup>65</sup> E. O. Olaiya,<sup>65</sup> R. Aleksan,<sup>66</sup> S. Emery,<sup>66</sup> A. Gaidot,<sup>66</sup> S. F. Ganzhur,<sup>66</sup> P.-F. Giraud,<sup>66</sup> G. Hamel de Monchenault,<sup>66</sup> W. Kozanecki,<sup>66</sup> M. Langer,<sup>66</sup> M. Legendre,<sup>66</sup> G. W. London,<sup>66</sup> B. Mayer,<sup>66</sup> G. Schott,<sup>66</sup> G. Vasseur,<sup>66</sup> Ch. Yèche,<sup>66</sup> M. Zito,<sup>66</sup> M. V. Purohit,<sup>67</sup> A. W. Weidemann,<sup>67</sup> J. R. Wilson,<sup>67</sup> F. X. Yumiceva,<sup>67</sup> D. Aston,<sup>68</sup> R. Bartoldus,<sup>68</sup> N. Berger,<sup>68</sup> A. M. Boyarski,<sup>68</sup> O. L. Buchmueller,<sup>68</sup> M. R. Convery,<sup>68</sup> M. Cristinziani,<sup>68</sup> G. De Nardo,<sup>68</sup> D. Dong,<sup>68</sup> J. Dorfan,<sup>68</sup> D. Dujmic,<sup>68</sup> W. Dunwoodie,<sup>68</sup> E. E. Elsen,<sup>68</sup> S. Fan,<sup>68</sup> R. C. Field,<sup>68</sup> T. Glanzman,<sup>68</sup> S. J. Gowdy,<sup>68</sup> T. Hadig,<sup>68</sup> V. Halyo,<sup>68</sup> C. Hast,<sup>68</sup> T. Hryn'ova,<sup>68</sup> W. R. Innes,<sup>68</sup> M. H. Kelsey,<sup>68</sup> P. Kim,<sup>68</sup> M. L. Kocian,<sup>68</sup> D. W. G. S. Leith,<sup>68</sup> J. Libby,<sup>68</sup> S. Luitz,<sup>68</sup> V. Luth,<sup>68</sup> H. L. Lynch,<sup>68</sup> H. Marsiske,<sup>68</sup> R. Messner,<sup>68</sup> D. R. Muller,<sup>68</sup> C. P. O'Grady,<sup>68</sup> V. E. Ozcan,<sup>68</sup> A. Perazzo,<sup>68</sup> M. Perl,<sup>68</sup> S. Petrak,<sup>68</sup> B. N. Ratcliff,<sup>68</sup> A. Roodman,<sup>68</sup> A. A. Salnikov,<sup>68</sup> R. H. Schindler,<sup>68</sup> J. Schwiening,<sup>68</sup> G. Simi,<sup>68</sup> A. Snyder,<sup>68</sup> A. Soha,<sup>68</sup> J. Stelzer,<sup>68</sup> D. Su,<sup>68</sup> M. K. Sullivan,<sup>68</sup> J. Va'vra,<sup>68</sup> S. R. Wagner,<sup>68</sup> M. Weaver,<sup>68</sup> A. J. R. Weinstein,<sup>68</sup> W. J. Wisniewski,<sup>68</sup> M. Wittgen,<sup>68</sup> D. H. Wright,<sup>68</sup> A. K. Yarritu,<sup>68</sup> C. C. Young,<sup>68</sup> P. R. Burchat,<sup>69</sup> A. J. Edwards,<sup>69</sup> T. I. Meyer,<sup>69</sup> B. A. Petersen,<sup>69</sup> C. Roat,<sup>69</sup> S. Ahmed,<sup>70</sup> M. S. Alam,<sup>70</sup> J. A. Ernst,<sup>70</sup> M. A. Saeed,<sup>70</sup> M. Saleem,<sup>70</sup> F. R. Wappler,<sup>70</sup> W. Bugg,<sup>71</sup> M. Krishnamurthy,<sup>71</sup> S. M. Spanier,<sup>71</sup> R. Eckmann,<sup>72</sup> H. Kim,<sup>72</sup> J. L. Ritchie,<sup>72</sup> A. Satpathy,<sup>72</sup> R. F. Schwitters,<sup>72</sup> J. M. Izen,<sup>73</sup> I. Kitayama,<sup>73</sup> X. C. Lou,<sup>73</sup> S. Ye,<sup>73</sup> F. Bianchi,<sup>74</sup> M. Bona,<sup>74</sup> F. Gallo,<sup>74</sup> D. Gamba,<sup>74</sup> C. Borean,<sup>75</sup> L. Bosisio,<sup>75</sup> C. Cartaro,<sup>75</sup> F. Cossutti,<sup>75</sup> G. Della Ricca,<sup>75</sup> S. Dittongo,<sup>75</sup> S. Grancagnolo,<sup>75</sup> L. Lanceri,<sup>75</sup> P. Poropat,<sup>75, §</sup> L. Vitale,<sup>75</sup> G. Vuagnin,<sup>75</sup> R. S. Panvini,<sup>76</sup> Sw. Banerjee,<sup>77</sup> C. M. Brown,<sup>77</sup> D. Fortin,<sup>77</sup> P. D. Jackson,<sup>77</sup> R. Kowalewski,<sup>77</sup> J. M. Roney,<sup>77</sup> H. R. Band,<sup>78</sup> S. Dasu,<sup>78</sup> M. Datta,<sup>78</sup> A. M. Eichenbaum,<sup>78</sup> M. Graham,<sup>78</sup> J. J. Hollar,<sup>78</sup> J. R. Johnson,<sup>78</sup> P. E. Kutter,<sup>78</sup> H. Li,<sup>78</sup> R. Liu,<sup>78</sup> F. Di Lodovico,<sup>78</sup> A. Mihalvi,<sup>78</sup> A. K. Mohapatra,<sup>78</sup> Y. Pan,<sup>78</sup> R. Prepost,<sup>78</sup> A. E. Rubin,<sup>78</sup> S. J. Sekula,<sup>78</sup>

P. Tan,<sup>78</sup> J. H. von Wimmersperg-Toeller,<sup>78</sup> J. Wu,<sup>78</sup> S. L. Wu,<sup>78</sup> Z. Yu,<sup>78</sup> M. G. Greene,<sup>79</sup> and H. Neal<sup>79</sup>

(The BABAR Collaboration)

<sup>1</sup>Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

<sup>2</sup>Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

- <sup>3</sup>Institute of High Energy Physics, Beijing 100039, China
- <sup>4</sup>University of Bergen, Inst. of Physics, N-5007 Bergen, Norway

<sup>5</sup>Lawrence Berkeley National Laboratory and University of California, Berkeley, CA 94720, USA

<sup>6</sup>University of Birmingham, Birmingham, B15 2TT, United Kingdom

<sup>7</sup>Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

<sup>8</sup>University of Bristol, Bristol BS8 1TL, United Kingdom

<sup>9</sup>University of British Columbia, Vancouver, BC, Canada V6T 1Z1

<sup>10</sup>Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

<sup>11</sup>Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

<sup>12</sup>University of California at Irvine, Irvine, CA 92697, USA

<sup>13</sup>University of California at Los Angeles, Los Angeles, CA 90024, USA

<sup>14</sup>University of California at Riverside, Riverside, CA 92521, USA

<sup>15</sup>University of California at San Diego, La Jolla, CA 92093, USA

<sup>16</sup>University of California at Santa Barbara, Santa Barbara, CA 93106, USA

<sup>17</sup>University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, CA 95064, USA <sup>18</sup>California Institute of Technology, Pasadena, CA 91125, USA

Caujornia institute of Technology, Fasaaena, CA 91125, USA

<sup>19</sup>University of Cincinnati, Cincinnati, OH 45221, USA

<sup>20</sup>University of Colorado, Boulder, CO 80309, USA

<sup>21</sup>Colorado State University, Fort Collins, CO 80523, USA

<sup>22</sup> Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany

<sup>23</sup>Ecole Polytechnique, LLR, F-91128 Palaiseau, France

<sup>24</sup>University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom

<sup>25</sup>Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy

<sup>26</sup>Florida A&M University, Tallahassee, FL 32307, USA

<sup>27</sup>Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy

<sup>28</sup>Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy

<sup>29</sup>Harvard University. Cambridge. MA 02138. USA

<sup>30</sup> Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany

<sup>31</sup>Imperial College London, London, SW7 2AZ, United Kingdom

<sup>32</sup>University of Iowa, Iowa City, IA 52242, USA

<sup>33</sup>Iowa State University, Ames, IA 50011-3160, USA

<sup>34</sup>Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France

<sup>35</sup>Lawrence Livermore National Laboratory, Livermore, CA 94550, USA

<sup>36</sup>University of Liverpool, Liverpool L69 72E, United Kingdom

<sup>37</sup>Queen Mary, University of London, E1 4NS, United Kingdom

<sup>38</sup>University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom

<sup>39</sup>University of Louisville, Louisville, KY 40292, USA

<sup>40</sup>University of Manchester, Manchester M13 9PL, United Kingdom

<sup>41</sup>University of Maryland, College Park, MD 20742, USA

<sup>42</sup>University of Massachusetts, Amherst, MA 01003, USA

<sup>43</sup>Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, MA 02139, USA

44 McGill University, Montréal, QC, Canada H3A 2T8

<sup>45</sup>Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy

<sup>46</sup>University of Mississippi, University, MS 38677, USA

<sup>47</sup>Université de Montréal, Laboratoire René J. A. Lévesque, Montréal, QC, Canada H3C 3J7

<sup>48</sup>Mount Holyoke College, South Hadley, MA 01075, USA

<sup>49</sup>Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy

<sup>50</sup>NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands

<sup>51</sup>University of Notre Dame, Notre Dame, IN 46556, USA

<sup>52</sup>Oak Ridge National Laboratory, Oak Ridge, TN 37831, USA

<sup>53</sup>Ohio State University, Columbus, OH 43210, USA

<sup>54</sup>University of Oregon, Eugene, OR 97403, USA

<sup>55</sup>Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy

<sup>56</sup> Universités Paris VI et VII, Lab de Physique Nucléaire H. E., F-75252 Paris, France

<sup>57</sup>Università di Pavia, Dipartimento di Elettronica and INFN, I-27100 Pavia, Italy

<sup>58</sup>University of Pennsylvania, Philadelphia, PA 19104, USA

<sup>59</sup>Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy

<sup>60</sup>Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy <sup>61</sup>Prairie View A&M University, Prairie View, TX 77446, USA

<sup>62</sup>Princeton University, Princeton, NJ 08544, USA

<sup>63</sup>Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy

<sup>64</sup> Universität Rostock, D-18051 Rostock, Germany

<sup>65</sup>Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom

<sup>66</sup>DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France

<sup>67</sup>University of South Carolina, Columbia, SC 29208, USA

<sup>68</sup>Stanford Linear Accelerator Center, Stanford, CA 94309, USA

<sup>69</sup>Stanford University, Stanford, CA 94305-4060, USA

<sup>70</sup>State Univ. of New York, Albany, NY 12222, USA

<sup>71</sup>University of Tennessee, Knoxville, TN 37996, USA

<sup>72</sup>University of Texas at Austin, Austin, TX 78712, USA

<sup>73</sup>University of Texas at Dallas, Richardson, TX 75083, USA

<sup>74</sup> Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy

<sup>75</sup>Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy

<sup>76</sup> Vanderbilt University, Nashville, TN 37235, USA

<sup>77</sup>University of Victoria, Victoria, BC, Canada V8W 3P6

<sup>78</sup>University of Wisconsin, Madison, WI 53706, USA

79 Yale University, New Haven, CT 06511, USA

(Dated: April 3, 2004)

We have measured the time-dependent decay rate for the process  $B \to J/\psi K^{*0}(892)$  in a sample of about 88 million  $\Upsilon(4S) \to B\overline{B}$  decays collected with the *BABAR* detector at the PEP-II asymmetricenergy *B* Factory at SLAC. In this sample we study flavor-tagged events in which one neutral *B* meson is reconstructed in the  $J/\psi K^{*0}$  or  $J/\psi \overline{K}^{*0}$  final state. We measure the coefficients of the cosine and sine terms in the time-dependent asymmetries for  $J/\psi K^{*0}$  and  $J/\psi \overline{K}^{*0}$ , find them to be consistent with the Standard Model expectations, and set upper limits at 90% C.L. on the decay amplitude ratios  $|A(\overline{B}^0 \to J/\psi K^{*0})|/|A(B^0 \to J/\psi K^{*0})| < 0.26$  and  $|A(B^0 \to J/\psi \overline{K}^{*0})|/|A(\overline{B}^0 \to J/\psi \overline{K}^{*0})|$  $W \overline{K}^{*0}| < 0.32$ . For a single ratio of wrong-flavor to favored amplitudes for  $B^0$  and  $\overline{B}^0$  combined, we obtain an upper limit of 0.25 at 90% C.L.

PACS numbers: 13.25.Hw, 12.15.Hh, 11.30.Er

The Standard Model of electroweak interactions describes *CP* violation in weak interactions of quarks by the presence of a complex phase in the three-generation Cabibbo-Kobayashi-Maskawa (CKM) quark-mixing matrix [1]. In this framework, the CP asymmetries in the proper-time distributions of neutral B decays to  $J/\psi K_s^0$ and  $J/\psi K_r^0$  are directly related to the *CP*-violation parameter  $\sin 2\beta$  [2]. The time-dependent *CP* asymmetries for  $J/\psi K_s^0$  and  $J/\psi K_L^0$  are of opposite sign and, to a very good approximation, equal in magnitude [3]. The decay  $B^0 \to J/\psi K_s^0 \ (B^0 \to J/\psi K_L^0)$  proceeds through the CKMfavored, color-suppressed decay  $B^0 \rightarrow J/\psi K^0$  [4] followed by  $K^0 \to K^0_S (K^0 \to K^0_L)$ . The so-called wrong-flavor  $B^0$ decay amplitude to the opposite strangeness final state  $B^0 \rightarrow J/\psi \,\overline{K}{}^0$  is expected to be negligible in the Standard Model [3]. Interference between a wrong-flavor amplitude and the favored amplitude can alter the relation between the *CP* asymmetries,  $A_{CP}$ , for the  $J/\psi K_s^0$  and  $J/\psi K_L^0$  final states. In general, a difference between  $A_{CP}(J/\psi K_s^0)$ and  $-A_{CP}(J/\psi K_L^0)$  of more than a few times  $10^{-3}$  requires a wrong-flavor amplitude [3]. A limit on the CPodd part of the phase difference between the wrong-flavor amplitude and the favored amplitude can be derived from the measured values of  $\sin 2\beta$  from B decays to the  $J/\psi K_s^0$ and  $J/\psi K_L^0$  final states. No test of the modulus of the wrong-flavor amplitude currently exists.

The decay mode  $B^0 \to J/\psi K^{*0}$  proceeds via the same quark transition as  $B^0 \to J/\psi K^0$ . The matrix elements, and therefore the ratio of wrong-flavor to favored amplitudes, are expected to be similar for  $B^0 \to J/\psi K^{*0}$  and  $B^0 \to J/\psi K^0$  [3]. In this Letter we present a measurement of the ratio of wrong-flavor to favored amplitude for the decay  $B^0 \to J/\psi K^{*0}$ , from the time-dependent asymmetry, where we use  $K^{*0} \to K^+\pi^-$  to identify the strangeness of the final state. The data sample consists of about 88 million  $B\overline{B}$  pairs produced in  $e^+e^-$  interactions at the  $\Upsilon(4S)$  resonance, corresponding to an integrated luminosity of 82 fb<sup>-1</sup>, collected with the BABAR detector [5] at the PEP-II asymmetric-energy collider at SLAC.

Charged particles are detected, and their momenta measured, by a combination of a vertex tracker consisting of five layers of double-sided silicon microstrip detectors, and a 40-layer central drift chamber, both operating in the 1.5-T magnetic field of a superconducting solenoid. We identify photons and electrons using a CsI(Tl) electromagnetic calorimeter. Further charged particle identification is provided by the average energy loss (dE/dx) in the tracking devices and by an internally reflecting ring imaging Cherenkov detector covering the central region.

The analysis method is similar to that of other time-dependent mixing measurements performed at *BABAR* [6]. We use a sample of events  $(B_{J/\psi K\pi})$  in which one neutral *B* meson is reconstructed in the state  $J/\psi K^{*0}$ or  $J/\psi \overline{K}^{*0}$ . The  $J/\psi$  meson is reconstructed through its decay to  $e^+e^-$  or  $\mu^+\mu^-$ , and the  $K^{*0}$  ( $\overline{K}^{*0}$ ) meson through its decay to  $K^+\pi^-$  ( $K^-\pi^+$ ). We examine each event in this sample for evidence that the other *B* meson decayed either as a  $B^0$  or  $\overline{B}^0$  (flavor tag).

The pseudoscalar to vector-vector decay  $B^0 \rightarrow J/\psi K^{*0}(892)$  is described by three amplitudes  $A_0, A_{\parallel}$ , and  $A_{\perp}$ , for the longitudinal, parallel, and perpendicular transverse polarization [7], respectively, of the vector mesons. In the selection of  $B^0 \rightarrow J/\psi K^{*0}(892)$  there is a small contribution from  $B^0 \rightarrow J/\psi K^{*0}(1430)$ , whose decay amplitude is denoted with  $A_s$ . The favored decay amplitudes  $A_{\lambda}(B^0 \rightarrow J/\psi K^+\pi^-) = a_{\lambda}e^{i\delta_{\lambda}^a}e^{+i\phi^a}$  are described by the magnitudes  $a_{\lambda}$ , weak phase  $\phi^a$ , and strong phases  $\delta_{\lambda}^a$ , where  $\lambda = 0, \parallel, \perp, s$ . The amplitudes for the wrong-flavor decays are given by  $A_{\lambda}(\overline{B}^0 \rightarrow J/\psi K^+\pi^-) =$  $b_{\lambda}e^{i\delta_{\lambda}^b}e^{+i\phi^b}$ . The corresponding decay amplitudes for the charge-conjugate final state  $J/\psi K^-\pi^+$  are obtained by replacing  $\phi^a$  with  $-\bar{\phi}^a$ ,  $b_{\lambda}$  with  $\bar{b}_{\lambda}$ ,  $\delta_{\lambda}^b$  with  $\bar{\delta}_{\lambda}^b$ , and  $\phi^b$ with  $-\bar{\phi}^b$ . We assume  $a_{\lambda} = \bar{a}_{\lambda}$ .

The proper-time distributions of B meson decays to  $J/\psi K^+\pi^- (J/\psi K^-\pi^+)$ , having either a  $B^0$  or  $\overline{B}^0$  tag, can be expressed in terms of the  $B^0-\overline{B}^0$  oscillation amplitude and the amplitudes describing  $\overline{B}^0$  and  $B^0$  decays to this final state [8]. The angular-integrated decay rate  $f_+(f_-)$  to the final state  $J/\psi K^+\pi^-$  when the tagging meson is a  $B^0(\overline{B}^0)$  is given by

$$f_{\pm}(\Delta t) = \frac{e^{-|\Delta t|/\tau_{B^0}}}{4\tau_{B^0}} \bigg[ 1 \mp C \cos\left(\Delta m_d \Delta t\right) \\ \pm S \sin\left(\Delta m_d \Delta t\right) \bigg], \qquad (1)$$

where  $\Delta t \equiv t_{\rm rec} - t_{\rm tag}$  is the difference between the proper decay times of the reconstructed *B* meson (*B*<sub>rec</sub>) and the tagging B meson  $(B_{\text{tag}})$ ,  $\tau_{B^0}$  is the  $B^0$  lifetime, and  $\Delta m_d$ is the  $B^0$ - $\overline{B}^0$  oscillation frequency. The corresponding decay rates  $\overline{f}_+$  and  $\overline{f}_-$  for the charge-conjugate final state  $J/\psi K^-\pi^+$  are obtained by replacing C with  $-\overline{C}$  and Swith  $-\overline{S}$ .

The C and S coefficients are related to the wrong-flavor and favored amplitudes by

$$C = \frac{a^2 - b^2}{a^2 + b^2}, \text{ and } S = \frac{2\sum_{\lambda} \eta \, a_{\lambda} b_{\lambda} \sin(\phi + \delta_{\lambda})}{a^2 + b^2}, \quad (2)$$

with  $a^2 \equiv a_0^2 + a_{\parallel}^2 + a_{\perp}^2 + a_s^2$ ,  $b^2 \equiv b_0^2 + b_{\parallel}^2 + b_{\perp}^2 + b_s^2$ , and  $\eta = +1 \ (-1)$  for  $\lambda = 0, \parallel, s \ (\perp)$ . The strong and weak phase differences are given by  $\delta_{\lambda} = \delta_{\lambda}^b - \delta_{\lambda}^a$  and  $\phi = \arg(q/p) + (\phi_b - \phi_a)$ , respectively, where (q/p) contains the weak phase of  $B^0 - \overline{B}^0$  oscillations. The  $\overline{C}$  and  $\overline{S}$  coefficients are given by the same expressions, replacing  $b_{(\lambda)}$  with  $\overline{b}_{(\lambda)}, \ \delta_{\lambda}$  with  $\overline{\delta}_{\lambda}$ , and  $\phi$  with  $-\overline{\phi}$ .

In the  $B \to J/\psi K^{*0}$  selection, a  $J/\psi$  candidate must consist of two identified lepton tracks [5] that form a good vertex. The lepton-pair invariant mass must be in the range  $3.06-3.14 \text{ GeV}/c^2$  for muons and  $2.95-3.14 \text{ GeV}/c^2$ for electrons. This corresponds to a  $\pm 3\sigma$  interval for muons, and, for electrons, accommodates the remaining radiative tail after bremsstrahlung correction [6]. We form  $K^+\pi^-$  candidate pairs, where the track that is most consistent with being a kaon is assigned to be the kaon candidate. The  $K^+\pi^-$  pair must have an invariant mass within 100 MeV/ $c^2$  of the nominal  $K^{*0}(892)$  mass [9]. In the selected mass window the  $K_0^*(1430)$  contributes  $(7.3 \pm 1.6)\%$  of the  $K^+\pi^-$  events.

The *B*-meson candidates are formed from  $J/\psi$  and  $K^+\pi^-$  candidates with the requirement that the difference  $\Delta E = E_B^{\rm cm} - E_{\rm beam}^{\rm cm}$  between their energy and the beam energy in the center-of-mass frame be less than 30 MeV from zero. The beam-energy-substituted mass  $m_{\rm ES} = \sqrt{(E_{\rm beam}^{\rm cm})^2 - (p_B^{\rm cm})^2}$  must be greater than 5.2 GeV/ $c^2$ , where  $p_B^{\rm cm}$  is the measured *B* momentum in the center-of-mass frame. We define a signal region with  $m_{\rm ES} > 5.27 \ {\rm GeV}/c^2$  to determine event yields and purities, and a sideband region with  $m_{\rm ES} < 5.27 \ {\rm GeV}/c^2$  to study background properties. If several *B* candidates are found in an event, the one with the smallest  $|\Delta E|$  is retained.

A measurement of the asymmetry coefficients  $C, S, \overline{C}$ , and  $\overline{S}$  requires a determination of the experimental  $\Delta t$ resolution and the fraction w of events in which the flavor tag assignment is incorrect. This mistag fraction reduces the amplitudes of the observed asymmetries by a factor 1 - 2w. Mistag fractions and  $\Delta t$  resolution functions are determined from a sample of neutral B mesons that decay to final states with one charmed meson  $(B_{Dh})$ , and consists of the channels  $D^{(*)-}h^+$   $(h^+ = \pi^+, \rho^+, \text{ and } a_1^+)$ .

The algorithm for *B*-flavor tagging is explained in Ref. [10]. The total efficiency for assigning a reconstructed *B* candidate to one of four hierarchical, mutually exclusive tagging categories is  $(65.6 \pm 0.5)\%$ . Untagged events are excluded from further consideration. The effective tagging efficiency  $Q \equiv \sum_i \varepsilon_i (1 - 2w_i)^2$ , where  $\varepsilon_i$  and  $w_i$  are the efficiencies and mistag probabilities, for events tagged in category *i*, is measured to be  $(28.1 \pm 0.7)\%$ .

The time interval  $\Delta t$  between the two *B* decays is calculated from the measured separation  $\Delta z$  between the decay vertices of the  $B_{\rm rec}$  and  $B_{\rm tag}$  along the collision (z) axis [6]. We determine the *z* position of the  $B_{\rm rec}$  vertex from its charged tracks. The  $B_{\rm tag}$  vertex is determined by fitting tracks not belonging to the  $B_{\rm rec}$  candidate to a common vertex, employing constraints from the beam spot location and the  $B_{\rm rec}$  momentum [6]. We accept events with a  $\Delta t$  uncertainty of less than 2.5 ps and  $|\Delta t| < 20$  ps. The fraction of events satisfying these requirements is 95%.



FIG. 1: Distributions of  $m_{\rm ES}$  a) for  $J/\psi K^+\pi^-$  candidates and b) for  $J/\psi K^-\pi^+$  candidates satisfying the tagging and vertexing requirements. The fit is described in the text.

Figure 1 shows the  $m_{\rm ES}$  distributions of the  $J/\psi K^+\pi^$ and  $J/\psi K^-\pi^+$  candidates that satisfy the tagging and vertexing requirements. The  $m_{\rm ES}$  distributions are fit with the sum of a threshold function [11], which accounts for the background from random combinations of tracks in the event, and a Gaussian distribution describing the signal. In Table I we list the event yields and signal purities for the tagged  $B \rightarrow J/\psi K^+\pi^-$  and  $B \rightarrow J/\psi K^-\pi^+$ candidates. The fraction of events in the Gaussian component of the  $m_{\rm ES}$  fits due to other B decay modes is estimated to be  $(1.6 \pm 0.4)\%$  based on simulated events.

We determine the  $C, S, \overline{C}$ , and  $\overline{S}$  coefficients with a simultaneous unbinned maximum likelihood fit to the  $\Delta t$  distributions of the tagged  $B_{J/\psi K\pi}$  and  $B_{Dh}$  samples. In this fit the  $\Delta t$  distributions of the  $J/\psi K^+\pi^$ and  $J/\psi K^-\pi^+$  samples are described by Eq. (1). The  $\Delta t$  distributions of the  $B_{Dh}$  sample are described by the same equation with C = 1 and S = 0. The observed amplitudes for the time-dependent asymmetries in the  $B_{J/\psi K\pi}$  sample and for flavor oscillation in the  $B_{Dh}$  sample are reduced by the same factor, 1 - 2w, due to flavor mistags. Events are assigned signal and background probabilities based on the  $m_{\rm ES}$  distributions. The  $\Delta t$ distributions for the signal are convolved with a common resolution function, modeled by the sum of three Gaussians [6]. Backgrounds are incorporated by means of an empirical description of their  $\Delta t$  spectra, obtained from the  $m_{\rm ES}$ -sideband region, containing prompt and nonprompt components convolved with a resolution function [6] distinct from that of the signal.

There are 48 free parameters in the fit. The fit parameters that describe the signal  $\Delta t$  distributions are C,  $S, \overline{C}, \text{ and } \overline{S}$  (4), the average mistag fraction w, the difference  $\Delta w$  between  $B^0$  and  $\overline{B}^0$  mistag fractions, and the linear dependence of the mistag fraction on the  $\Delta t$ error for each tagging category (12), parameters for the signal  $\Delta t$  resolution (8), and parameters to account for differences in reconstruction and tagging efficiencies for  $B^0$  and  $\overline{B}^0$  mesons (5). The  $B_{J/\psi K\pi}$  and  $B_{Dh}$  background  $\Delta t$  distributions are described by parameters for the background time dependence (8),  $\Delta t$  resolution (3), and mistag fractions (8). We fix  $\tau_{B^0}$  at 1.542 ps and  $\Delta m_d$  at 0.489 ps<sup>-1</sup> [9]. The determination of the mistag fractions and  $\Delta t$  resolution function parameters for the signal is dominated by the large  $B_{Dh}$  sample. Background parameters are determined from events with  $m_{\rm ES} < 5.27 \, {\rm GeV}/c^2$ .

The fit to the  $B_{J/\psi K\pi}$  and  $B_{Dh}$  samples yields  $C = 1.045 \pm 0.058 \pm 0.035$ ,  $S = -0.024 \pm 0.095 \pm 0.041$ ,  $\overline{C} = 0.966 \pm 0.051 \pm 0.035$ , and  $\overline{S} = 0.004 \pm 0.090 \pm 0.041$ , where the first error is statistical and the second error is systematic. Figure 2 shows the  $\Delta t$  distributions and the asymmetries in yields between  $B^0$  tags and  $\overline{B}^0$  tags as a function of  $\Delta t$  for the  $J/\psi K^+\pi^-$  and  $J/\psi K^-\pi^+$  samples, overlaid with the projection of the likelihood fit result.

We estimate common systematic errors for C(S) and  $\overline{C}(\overline{S})$ . The dominant sources of systematic error are the uncertainties in the level, composition, and time-dependent asymmetry of the background in the selected  $B_{J/\psi K\pi}$  sample (0.016 for C, 0.017 for S), uncertainties in the beam spot location and the internal alignment of the vertex detector (0.016 for C, 0.021 for S), and the statistics of the simulated event sample (0.016 for C, 0.015 for S). Another significant contribution to the systematic uncertainty in the cosine coefficients comes from possible differences between the  $B_{Dh}$  and  $B_{J/\psi K\pi}$  mistag fractions (0.012). The uncertainty in the interference between the

TABLE I: Number of events,  $N_{\text{tag}}$ , and signal purity, P, in the signal region for the  $J/\psi K^+\pi^-$  and  $J/\psi K^-\pi^+$  samples, and for the  $B_{Dh}$  sample. Errors are statistical only.

510	1	5
Sample	$N_{\mathrm{tag}}$	P(%)
$J/\psi K^+\pi^-$ sample	860	$95.5\pm0.7$
$J/\psi K^{-}\pi^{+}$ sample	856	$96.5\pm0.6$
$B_{Dh}$ sample	25375	$84.9\pm0.2$



FIG. 2: Number of  $J/\psi K^+\pi^-$  and  $J/\psi K^-\pi^+$  candidates in the signal region a) with an opposite-flavor *B* tag,  $N_{OF}$ , b) with a same-flavor *B* tag,  $N_{SF}$ , and c) the observed asymmetry  $(N_{OF} - N_{SF})/(N_{OF} + N_{SF})$  as functions of  $\Delta t$ . In each figure the solid (dashed) curves represent the fit projection in  $\Delta t$  for  $J/\psi K^+\pi^- (J/\psi K^-\pi^+)$  candidates. The shaded regions in (a) and (b) represent the background contributions.

suppressed  $\bar{b} \rightarrow \bar{u}c\bar{d}$  amplitude with the favored  $b \rightarrow c\bar{u}d$ amplitude for the decay modes in the  $B_{Dh}$  sample and for certain tag-side *B* decays to hadronic final states [12] contributes to the systematic uncertainty in the sine coefficients (0.019). Finally, there are differences in the angular-integrated efficiency for the  $B \rightarrow J/\psi K^{*0}(892)$ helicity amplitudes and the  $B \rightarrow J/\psi K_0^*(1430)$  amplitude (0.007 for *C*, 0.016 for *S*). The total systematic errors for the cosine coefficients and sine coefficients are 0.035 and 0.041, respectively. Most systematic errors are determined with data and are expected to decrease with larger sample size.

The large  $J/\psi K^+\pi^-$  and  $J/\psi K^-\pi^+$  samples allow a number of consistency checks, including separation by data-taking period and tagging category. The results of fits to these subsamples are found to be statistically consistent.

The measured values of the cosine and sine coefficients are consistent with  $C = \overline{C} = 1$  and  $S = \overline{S} = 0$ , as expected for no contributions from the wrong-flavor decays  $B^0 \rightarrow J/\psi K^-\pi^+$  and  $\overline{B}^0 \rightarrow J/\psi K^+\pi^-$ . We use the measured cosine coefficients C and  $\overline{C}$  and assume |q/p| =1 [13] to calculate the wrong-flavor to favored decay rate ratios  $\Gamma(\overline{B}^0 \rightarrow J/\psi K^+\pi^-)/\Gamma(B^0 \rightarrow J/\psi K^+\pi^-) =$  $|b/a|^2 = -0.022 \pm 0.028$  (stat.)  $\pm 0.016$  (syst.) and  $\Gamma(B^0 \rightarrow J/\psi K^-\pi^+)/\Gamma(\overline{B}^0 \rightarrow J/\psi K^-\pi^+) = |\overline{b}/a|^2 =$  $0.017 \pm 0.026$  (stat.)  $\pm 0.016$  (syst.), where the negative central value occurs because C > 1. From these measurements the wrong-flavor to favored amplitude ratios for  $B \to J/\psi K^{*0}(892)$  and  $B \to J/\psi \overline{K}^{*0}(892)$  can be calculated. Using the measured fraction of  $B \to J/\psi K_0^*(1430)$  events contributing in the  $B \to J/\psi K^+\pi^-$  selection, the upper limits for the decay amplitude ratios at 90% confidence level (C.L.) are found to be  $|A(\overline{B}^0 \to J/\psi \overline{K}^{*0})|/|A(B^0 \to J/\psi \overline{K}^{*0})| < 0.26$  and  $|A(B^0 \to J/\psi \overline{K}^{*0})|/|A(\overline{B}^0 \to J/\psi \overline{K}^{*0})| < 0.32$ . For the single ratio of wrong-flavor to favored amplitude for  $B^0$  and  $\overline{B}^0$  combined, we determine an upper limit of 0.25 at 90% C.L.

In conclusion, we observe no evidence for the wrongflavor decays  $\overline{B}^0 \to J/\psi K^{*0}(892)$  and  $B^0 \to J/\psi \overline{K}^{*0}(892)$ . Together with theoretical information on the relation between the matrix elements for  $B^0 \to J/\psi K^0$  and  $B^0 \to J/\psi K^{*0}$  [3], the results presented here can be used to set a limit on the difference between  $A_{CP}(J/\psi K_S^0)$  and  $-A_{CP}(J/\psi K_S^0)$ .

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from the A. P. Sloan Foundation, Research Corporation, and Alexander von Humboldt Foundation.

- <sup>†</sup> Also with Università della Basilicata, Potenza, Italy
- <sup>‡</sup> Also with IFIC, Instituto de Física Corpuscular, CSIC-Universidad de Valencia, Valencia, Spain
   <sup>§</sup> Deceased
- Deceased
- N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [2] A.B. Carter and A.I. Sanda, Phys. Rev. D 23, 1567 (1981);
  I.I. Bigi and A.I. Sanda, Nucl. Phys. B193, 85 (1981).
- [3] Y. Grossman, A.L. Kagan, and Z. Ligeti, Phys. Lett. B 538, 327 (2002).
- [4] Charge conjugation is implied throughout this letter, unless explicitly stated otherwise.
- [5] BABAR Collaboration, B. Aubert *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A 479, 1 (2002).
- [6] BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. D 66, 032003 (2002).
- [7] A.S. Dighe, I. Dunietz, H.J. Lipkin, and J.L. Rosner, Phys. Lett. B 369, 144 (1996).
- [8] See, for example, L. Wolfenstein, in Phys. Rev. D 66, 010001 (2002).
- [9] Particle Data Group, K. Hagiwara *et al.*, Phys. Rev. D 66, 010001 (2002).
- [10] BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. 89, 201802 (2002).
- [11] ARGUS Collaboration, H. Albrecht *et al.*, Z. Phys. C 48, 543 (1990).
- [12] O. Long, M. Baak, R.N. Cahn, and D. Kirkby, Phys. Rev. D 68, 034010 (2003).
- [13] BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. 88, 231801 (2002).

<sup>\*</sup> Now at Department of Physics, University of Warwick, Coventry, United Kingdom