Accretion of Ghost Condensate by Black Holes

Andrei V. Frolov^{*} KIPAC/ITP, Stanford University Stanford, CA, 94305-4060 (Dated: April 27, 2004)

The intent of this letter is to point out that the accretion of a ghost condensate by black holes could be extremely efficient. We analyze steady-state spherically symmetric flows of the ghost fluid in the gravitational field of a Schwarzschild black hole and calculate the accretion rate. Unlike minimally coupled scalar field or quintessence, the accretion rate is set not by the cosmological energy density of the field, but by the energy scale of the ghost condensate theory. If hydrodynamical flow is established, it could be as high as tenth of a solar mass per second for 10MeV-scale ghost condensate accreting onto a stellar-sized black hole, which puts serious constraints on the parameters of the ghost condensate model.

I. INTRODUCTION

Prompted by an increasingly precise experimental measurements of cosmological parameters, and in particular detection of acceleration of the universe due to an unknown source which looks like a cosmological constant, in the recent years there has been a wide discussion in the literature about modifications of Einstein gravity on cosmological scales as a possible alternative to dark matter and/or energy. However, finding a self-consistent and well-motivated theory which agrees with all the observations is proving to be quite a challenge.

Recently, Arkani-Hamed *et. al.* proposed a model [1], dubbed *a ghost condensation*, which they argued to be consistent with all experimental observations and provide an interesting modification of gravity in the infrared, with potential applications to inflation [2] and cosmological constant problem. It involves an introduction of a scalar field which develops a non-zero expectation value of its (timelike) gradient in vacuum, due to non-trivial kinetic term in the action. Such modifications of the scalar field kinetic term were considered earlier on phenomenological grounds in the model known as *k*-inflation [3, 4].

The ghost condensate model has been studied on perturbative level in effective field theory [1], which already leads to interesting consequences such as star trails [5, 6]. We look at the ghost condensate from a slightly different perspective, namely we would like to investigate its behaviour in the strong gravitational field, for instance, near a Schwarzschild black hole

$$ds^{2} = -f(r)dt^{2} + \frac{dr^{2}}{f(r)} + r^{2}d\Omega_{n}^{2}, \qquad (1)$$

where $f(r) = 1 - r_g/r$, and $r_g = 2Gm$ is a gravitational radius of Schwarzschild black hole of mass m. The problem is similar to interaction of a cosmological scalar field with a black hole [7], so one would expect ghost condensate to be accreted by a black hole. Accretion of fluid onto a black hole has long been an important problem in astrophysics. Spherically symmetric steady-state fluid accretion onto a Schwarzschild black hole was derived in Ref. [8]. Minimally coupled scalar field [9] and quintessence [7, 10] accrete onto black holes as well, although the accretion rate is limited by the cosmological density of the field [7]. Accretion of exotic matter fields can lead to unusual results. For instance, accretion of a phantom energy (which violates energy dominance conditions) *decreases* the black hole size [11].

In this letter, we calculate the steady-state accretion rate of the ghost condensate by a black hole, and point out that it could be extremely efficient. This puts serious constraints on the parameters of the ghost condensate model.

II. GHOST CONDENSATE AS A FLUID

Ghost condensate model adds a non-minimally coupled scalar field to Einstein theory of gravity. However, instead of the usual kinetic term

$$X = -(\nabla \phi)^2, \tag{2}$$

the action is assumed to involve a more complicated function of the field gradient squared

$$S = \int \left\{ \frac{R}{16\pi G} + M^4 P(X) \right\} \sqrt{-g} \, d^4 x, \qquad (3)$$

as well as higher-derivative terms. We will ignore higherderivative terms in what follows. They complicate calculations significantly, and we are concerned with large scale flows, while one would expect higher-derivative terms to be important on short scales.

As the ghost field ϕ is not directly coupled to other fields, its dimensionality and normalization are arbitrary. If the field ϕ is chosen to have dimension of length, the field gradient X and the function P are dimensionless, and the only dimensional quantity in the ghost sector is M, which sets the overall energy scale of the ghost condensate. The specific form of the function P(X) is not rigidly fixed, although the defining feature of the

^{*}Electronic address: afrolov@stanford.edu



FIG. 1: Ghost condensate kinetic term (top) and equivalent fluid description (bottom). Equation of state w and sound speed c_s^2 are shown by dashed and solid curves, respectively.

ghost condensate model is that P has a minimum at nonvanishing (timelike) value of the field gradient. Because of that, the ghost field rolls even in its vacuum state. The simplest choice for P with this property is

$$P(X) = \frac{1}{2} (X - A)^2, \qquad (4)$$

illustrated in Fig. 1. One could also add a cosmological constant term Λ , but since it is not accreted by a black hole, we will not discuss it further.

Variation of the action (3) with respect to the ghost field ϕ yields equation of motion

$$\nabla_{\mu} \left[P'(X) \nabla^{\mu} \phi \right] \equiv \frac{1}{\sqrt{-g}} \partial_{\mu} \left[\sqrt{-g} P'(X) \partial^{\mu} \phi \right] = 0.$$
 (5)

The equation of motion is implied by conservation of the stress-energy tensor, which for the ghost condensate is

$$T_{\mu\nu} = 2M^4 P'(X)\phi_{;\mu}\phi_{;\nu} + M^4 P(X)g_{\mu\nu}.$$
 (6)

Configurations with $P'(X) \equiv 0$ solve the equation of motion identically for any spacetime metric. However, such configurations are indistinguishable gravitationally from a purely Einstein theory, as the stress-energy tensor becomes trivial as well.

The stress-energy tensor of the ghost condensate (6) can be transformed into that of a perfect fluid

$$T_{\mu\nu} = (\rho + p)u_{\mu}u_{\nu} + pg_{\mu\nu} \tag{7}$$

by a formal identification

$$\rho = M^4 (2XP' - P), \quad p = M^4 P, \quad u_\mu = \frac{\phi_{;\mu}}{\sqrt{X}}.$$
(8)

The fluid analogy is very useful in understanding the physics behind the solutions of the ghost equation of motion (5), although it is not an exact correspondence. Unlike ordinary fluids, ghost condensate is *irrotational*, that is, the vorticity tensor of the flow u^{μ} vanishes identically

$$\omega_{\alpha\beta} = \frac{1}{2} \left(u_{\alpha;\mu} q^{\mu}_{\ \beta} - u_{\beta;\mu} q^{\mu}_{\ \alpha} \right) \equiv 0, \tag{9}$$

where $q_{\mu\nu} = g_{\mu\nu} + u_{\mu}u_{\nu}$. This is a direct consequence of the vector flow u^{μ} being derived from a scalar.

Important parameters of the fluid are its equation of state and sound speed

$$w \equiv \frac{p}{\rho}, \quad c_s^2 \equiv \frac{dp}{d\rho} = \frac{p'}{\rho'}.$$
 (10)

For the ghost condensate with kinetic term (4), they are

$$w = \frac{X - A}{3X + A}, \quad c_s^2 = \frac{X - A}{3X - A}.$$
 (11)

The equation of state and the sound speed change from dust-like in the minimum X = A to radiation-like for large displacements $X \gg A$, as shown in Fig. 1. Configurations with X < A are unstable, as the sound speed squared becomes negative.

III. STEADY-STATE ACCRETION

Steady-state accretion means that the flow of the field (i.e. its gradient) does not change with time, that is $\pounds_{\partial_t}(\nabla_{\mu}\phi) = \partial_{\mu}\partial_t\phi = 0$, which in turn implies that $\partial_t\phi$ is constant (and can be set to one by a choice of the field normalization). Therefore, a general steady-state spherically symmetric field configuration is of the form

$$\phi = t + \psi(r), \tag{12}$$

and, in Schwarzschild spacetime (1), has a gradient

$$X = \frac{1 - (\partial_r^* \psi)^2}{f(r)},$$
(13)

where we introduced a "tortoise" derivative $\partial_r^* \equiv f(r)\partial_r$.

For steady-state accretion of the spherically symmetric ghost condensate profile (12) onto a Schwarzschild black hole (1), the equation of motion (5) becomes

$$\partial_r^* (r^2 P' \partial_r^* \psi) = 0, \qquad (14)$$

which can be immediately integrated to yield the flow conservation equation

$$P'\partial_r^*\psi = \alpha \frac{r_g^2}{r^2}.$$
 (15)

The meaning of the dimensionless constant of integration α becomes clear if one looks at the accretion rate

$$\dot{m} = 4\pi r^2 T_t^r = 2\alpha \cdot 4\pi r_g^2 M^4, \qquad (16)$$



FIG. 2: Flow diagrams v(x) of the ghost condensate accretion onto a black hole for A = 3/4 (left) and A = 1 (right). Inflow, no-flow, and (unstable) outflow branches are shown by solid, dot-dash, and dotted lines correspondingly. Flow trajectories passing through the critical point are emphasized by thicker lines. Negative v region corresponds to reversed flow direction, and is not shown.

which does not depend on r and describes a steady-state transfer of mass from infinity into a black hole. The numerical value of the coefficient α is picked by the solution of the flow equation (15) that is regular at the horizon and becomes homogeneous far from the black hole.

The flow equation (15) is actually algebraic in $\partial_r^* \psi$, and could be analyzed for an arbitrary function P. We will restrict our discussion to the ghost condensate with kinetic term (4) and further assume $A \leq 1$, as the choice A > 1places the solution (12) on the unstable branch of the kinetic term far from a black hole and is not physically relevant. Introducing the short-hand notation $v \equiv \partial_r^* \psi$ and $x \equiv f(r)$, the flow equation (15) can be written as

$$\left(\frac{1-v^2}{x} - A\right)\frac{v}{(1-x)^2} = \alpha.$$
 (17)

Solutions v(x) for various values of α are shown in Fig. 2. Although cubic equation (17) can be directly solved in radicals, the flow is more readily analyzed using standard phase space diagram techniques.

Both at the horizon (x = 0) and infinity (x = 1), all flow trajectories converge to one of three roots

$$\begin{aligned} x &= 0: \quad v_0 = 0, \pm 1 \\ x &= 1: \quad v_1 = 0, \pm \sqrt{1 - A} . \end{aligned}$$
 (18)

All flow trajectories must start and end at these roots, and they do not intersect except at the critical points. The critical points are defined as the points where the full differential of (17),

$$-\frac{3v^2 - 1 + Ax}{x(1-x)^2} \, dv - \frac{(1-v^2)(1-3x) + 2Ax^2}{x^2(1-x)^3} \, v \, dx = 0,$$
(19)

becomes degenerate, i.e. when coefficients in front of dvand dx both vanish. In the positive v region, there is (at most) one critical point

$$v_*^2 = \frac{A + \sqrt{A^2 - 36A + 36}}{18}, \quad x_* = \frac{1 - 3v_*^2}{A}.$$
 (20)

Regularity at the horizon for ingoing flow demands that $v_0 = 1$, while proper fall-off at infinity requires $v_1 = 0$. For A < 1, the only flow trajectory that connects the two is the one that passes through the critical point (20), as it is clear from the left panel of Fig. 2. The flow starts out subsonic at infinity, and turns supersonic at the critical point. As it is easy to see from equation (17), the field gradient v falls off at infinity as r^{-2} , and so the field profile levels off as r^{-1} far from the black hole.

Thus, the coefficient α for the steady-state accretion flow is calculated by evaluating equation (17) at the critical point (20). The resulting expression is straightforward, but cumbersome for arbitrary A, so we will not write it down here. Instead, we will show the graph of α as a function of A in Fig. 3. The coefficient α decreases monotonically from $3\sqrt{3}/2$ at A = 0 to 1 at A = 1.

The case of A = 1 is special, and is shown on the right panel of Fig. 2. The three roots at infinity merge into one triple root at $v_1 = 0$, and one can get from infinity



FIG. 3: Dependence of the accretion rate coefficient α on A.

to horizon without going through a critical point. These solutions correspond to a dust-like flow with $0 \le \alpha < 1$, and are always supersonic. However, their gradient v falls off at infinity only as $r^{-\frac{1}{2}}$, which means that the ghost field does not become homogeneous far from a black hole, but in fact grows as $r^{\frac{1}{2}}$. In particular, the trivial solution $(P' \equiv 0, \alpha = 0)$ is

$$\phi = t + 2r_g^{\frac{1}{2}} \left[r^{\frac{1}{2}} - r_g^{\frac{1}{2}} \operatorname{arctanh} \sqrt{\frac{r_g}{r}} \right].$$
(21)

The likely reason behind the change in field asymptotic is that spherically symmetric dust accretion is not steadystate. The accretion rate is ever growing, as the dust from larger and larger volume falls inside the black hole.

For A = 1, the flow trajectory which passes through a critical point ($\alpha = 1$) is simply $v = 1 - x = r_g/r$. The corresponding field profile is

$$\phi = t + r_g \ln\left[\frac{r}{r_g} - 1\right]. \tag{22}$$

Its asymptotic at infinity is also non-homogeneous, but the growth is only logarithmic.

IV. DISCUSSION

In the last section, we calculated the steady-state accretion rate of the ghost condensate by a black hole for spherically symmetric flows. The most important result of the calculation is that the dimensionless coefficient α , which determines the accretion rate of the flow, is bounded below by 1 even as density of the ghost condensate far from a black hole becomes vanishingly small.

This means that it is the energy scale M of the ghost condensate theory that sets the accretion rate, and not the cosmological abundance of the ghost condensate field as one might have naively expected.

Up to a numerical coefficient of order one, the accretion rate is equal to the energy density M^4 falling down through the horizon area $4\pi r_g^2$ at the speed of light. The top value of 10MeV for the ghost energy scale quoted in [1] corresponds to a rather high density

$$(10 \text{MeV})^4 = \frac{(10 \text{MV} \cdot e)^4}{\hbar^3 c^5} = 2.32 \cdot 10^{12} \frac{\text{kg}}{\text{m}^3}.$$
 (23)

If the steady-state flow of the kind we considered is established, the accretion rate of a 10MeV-scale ghost condensate by an astrophysical black hole would be enormous

$$\dot{m} = 0.08 \,\alpha \,\frac{M_{\odot}}{\mathrm{s}} \left(\frac{r_g}{\mathrm{3km}}\right)^2 \left(\frac{M}{10 \mathrm{MeV}}\right)^4. \tag{24}$$

To avoid rapid black hole growth and its astrophysical consequences, energy scale M of the ghost condensate should be significantly less than 10MeV. Stellar-size black hole would double in size over the lifetime of the universe (roughly 14Gyrs, or $4 \cdot 10^{17}$ s) for the ghost energy scale of order 1keV. This estimate goes down to 10eV for supermassive $(10^9 M_{\odot})$ black holes.

Spherically symmetric steady-state accretion is an idealized situation, of course. We have not considered timescale required to establish such flows, the role of initial conditions, motion of the black hole with respect to the condensate, or what happens if the ghost field becomes highly inhomogeneous. All of these are much harder problems, and it might turn out that some factors prevent the accretion from settling into an efficient steady-state regime. Still, having ghost condensate capable of such high accretion rates is alarming, and the issue should be further addressed by the ghost condensate scenario.

Acknowledgments

I am grateful to Shinji Mukohyama for sharing his insights into the ghost condensate dynamics. I would like to thank Renata Kallosh, Lev Kofman, and Marco Peloso for their helpful comments.

- N. Arkani-Hamed, H. C. Cheng, M. A. Luty and S. Mukohyama, Ghost condensation and a consistent infrared modification of gravity, hep-th/0312099.
- [2] N. Arkani-Hamed, P. Creminelli, S. Mukohyama and M. Zaldarriaga, *Ghost inflation*, JCAP **0404**, 001 (2004) [hep-th/0312100].
- [3] C. Armendariz-Picon, T. Damour and V. Mukhanov, *k-inflation*, Phys. Lett. B 458, 209 (1999) [hep-th/9904075].
- [4] J. Garriga and V. F. Mukhanov, *Perturbations* in k-inflation, Phys. Lett. B **458**, 219 (1999) [hep-th/9904176].

- [5] S. L. Dubovsky, Star tracks in the ghost condensate, hep-ph/0403308.
- [6] M. Peloso and L. Sorbo, Moving sources in a ghost condensate, hep-th/0404005.
- [7] A. Frolov and L. Kofman, Inflation and de Sitter thermodynamics, JCAP 0305, 009 (2003) [hep-th/0212327].
- [8] F. C. Michel, Accretion of matter by condensed objects, Astrophysics and Space Science 15, 153 (1972).
- [9] T. Jacobson, Primordial black hole evolution in ten-

sor scalar cosmology, Phys. Rev. Lett. 83, 2699 (1999) [astro-ph/9905303].

- [10] R. Bean and J. Magueijo, Could supermassive black holes be quintessential primordial black holes?, Phys. Rev. D 66, 063505 (2002) [astro-ph/0204486].
- [11] E. Babichev, V. Dokuchaev and Y. Eroshenko, Black hole mass decreasing due to phantom energy accretion, gr-qc/0402089.