

Measurement of branching fractions and mass spectra of $B \rightarrow K\pi\pi\gamma$

B. Aubert,¹ R. Barate,¹ D. Boutigny,¹ F. Couderc,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Poireau,¹ V. Tisserand,¹
 A. Zghiche,¹ E. Grauges,² A. Palano,³ M. Pappagallo,³ A. Pompili,³ J. C. Chen,⁴ N. D. Qi,⁴ G. Rong,⁴ P. Wang,⁴
 Y. S. Zhu,⁴ G. Eigen,⁵ I. Ofte,⁵ B. Stugu,⁵ G. S. Abrams,⁶ M. Battaglia,⁶ A. B. Breon,⁶ D. N. Brown,⁶
 J. Button-Shafer,⁶ R. N. Cahn,⁶ E. Charles,⁶ C. T. Day,⁶ M. S. Gill,⁶ A. V. Gritsan,⁶ Y. Groysman,⁶
 R. G. Jacobsen,⁶ R. W. Kadel,⁶ J. Kadyk,⁶ L. T. Kerth,⁶ Yu. G. Kolomensky,⁶ G. Kukartsev,⁶ G. Lynch,⁶
 L. M. Mir,⁶ P. J. Oddone,⁶ T. J. Orimoto,⁶ M. Pripstein,⁶ N. A. Roe,⁶ M. T. Ronan,⁶ W. A. Wenzel,⁶ M. Barrett,⁷
 K. E. Ford,⁷ T. J. Harrison,⁷ A. J. Hart,⁷ C. M. Hawkes,⁷ S. E. Morgan,⁷ A. T. Watson,⁷ M. Fritsch,⁸ K. Goetzen,⁸
 T. Held,⁸ H. Koch,⁸ B. Lewandowski,⁸ M. Pelizaeus,⁸ K. Peters,⁸ T. Schroeder,⁸ M. Steinke,⁸ J. T. Boyd,⁹
 J. P. Burke,⁹ N. Chevalier,⁹ W. N. Cottingham,⁹ M. P. Kelly,⁹ T. Cuhadar-Donszelmann,¹⁰ B. G. Fulsom,¹⁰
 C. Hearty,¹⁰ N. S. Knecht,¹⁰ T. S. Mattison,¹⁰ J. A. McKenna,¹⁰ A. Khan,¹¹ P. Kyberd,¹¹ M. Saleem,¹¹
 L. Teodorescu,¹¹ A. E. Blinov,¹² V. E. Blinov,¹² A. D. Bukin,¹² V. P. Druzhinin,¹² V. B. Golubev,¹²
 E. A. Kravchenko,¹² A. P. Onuchin,¹² S. I. Serednyakov,¹² Yu. I. Skoppen,¹² E. P. Solodov,¹² A. N. Yushkov,¹²
 D. Best,¹³ M. Bondioli,¹³ M. Bruinsma,¹³ M. Chao,¹³ I. Eschrich,¹³ D. Kirkby,¹³ A. J. Lankford,¹³
 M. Mandelkern,¹³ R. K. Mommsen,¹³ W. Roethel,¹³ D. P. Stoker,¹³ C. Buchanan,¹⁴ B. L. Hartfiel,¹⁴
 S. D. Foulkes,¹⁵ J. W. Gary,¹⁵ O. Long,¹⁵ B. C. Shen,¹⁵ K. Wang,¹⁵ L. Zhang,¹⁵ D. del Re,¹⁶ H. K. Hadavand,¹⁶
 E. J. Hill,¹⁶ D. B. MacFarlane,¹⁶ H. P. Paar,¹⁶ S. Rahatlou,¹⁶ V. Sharma,¹⁶ J. W. Berryhill,¹⁷ C. Campagnari,¹⁷
 A. Cunha,¹⁷ B. Dahmes,¹⁷ T. M. Hong,¹⁷ M. A. Mazur,¹⁷ J. D. Richman,¹⁷ W. Verkerke,¹⁷ T. W. Beck,¹⁸
 A. M. Eisner,¹⁸ C. J. Flacco,¹⁸ C. A. Heusch,¹⁸ J. Kroeseberg,¹⁸ W. S. Lockman,¹⁸ G. Nesom,¹⁸ T. Schalk,¹⁸
 B. A. Schumm,¹⁸ A. Seiden,¹⁸ P. Spradlin,¹⁸ D. C. Williams,¹⁸ M. G. Wilson,¹⁸ J. Albert,¹⁹ E. Chen,¹⁹
 G. P. Dubois-Felsmann,¹⁹ A. Dvoretskii,¹⁹ D. G. Hitlin,¹⁹ I. Narsky,¹⁹ T. Piatenko,¹⁹ F. C. Porter,¹⁹ A. Ryd,¹⁹
 A. Samuel,¹⁹ R. Andreassen,²⁰ S. Jayatilleke,²⁰ G. Mancinelli,²⁰ B. T. Meadows,²⁰ M. D. Sokoloff,²⁰ F. Blanc,²¹
 P. Bloom,²¹ S. Chen,²¹ W. T. Ford,²¹ U. Nauenberg,²¹ A. Olivas,²¹ P. Rankin,²¹ W. O. Ruddick,²¹ J. G. Smith,²¹
 K. A. Ulmer,²¹ S. R. Wagner,²¹ J. Zhang,²¹ A. Chen,²² E. A. Eckhart,²² A. Soffer,²² W. H. Toki,²² R. J. Wilson,²²
 Q. Zeng,²² D. Altenburg,²³ E. Feltresi,²³ A. Hauke,²³ B. Spaan,²³ T. Brandt,²⁴ J. Brose,²⁴ M. Dickopp,²⁴
 V. Klose,²⁴ H. M. Lacker,²⁴ R. Nogowski,²⁴ S. Otto,²⁴ A. Petzold,²⁴ G. Schott,²⁴ J. Schubert,²⁴ K. R. Schubert,²⁴
 R. Schwierz,²⁴ J. E. Sundermann,²⁴ D. Bernard,²⁵ G. R. Bonneauaud,²⁵ P. Grenier,²⁵ S. Schrenk,²⁵ Ch. Thiebaut,²⁵
 G. Vasileiadis,²⁵ M. Verderi,²⁵ D. J. Bard,²⁶ P. J. Clark,²⁶ W. Gradl,²⁶ F. Muheim,²⁶ S. Playfer,²⁶ Y. Xie,²⁶
 M. Andreotti,²⁷ V. Azzolini,²⁷ D. Bettoni,²⁷ C. Bozzi,²⁷ R. Calabrese,²⁷ G. Cibinetto,²⁷ E. Luppi,²⁷ M. Negrini,²⁷
 L. Piemontese,²⁷ F. Anulli,²⁸ R. Baldini-Ferroli,²⁸ A. Calcaterra,²⁸ R. de Sangro,²⁸ G. Finocchiaro,²⁸ P. Patteri,²⁸
 I. M. Peruzzi,²⁸, * M. Piccolo,²⁸ A. Zallo,²⁸ A. Buzzo,²⁹ R. Capra,²⁹ R. Contri,²⁹ M. Lo Vetere,²⁹ M. Macri,²⁹
 M. R. Monge,²⁹ S. Passaggio,²⁹ C. Patrignani,²⁹ E. Robutti,²⁹ A. Santroni,²⁹ S. Tosi,²⁹ S. Bailey,³⁰
 G. Brandenburg,³⁰ K. S. Chaisanguanthum,³⁰ M. Morii,³⁰ E. Won,³⁰ J. Wu,³⁰ R. S. Dubitzky,³¹ U. Langenegger,³¹
 J. Marks,³¹ S. Schenk,³¹ U. Uwer,³¹ W. Bhimji,³² D. A. Bowerman,³² P. D. Dauncey,³² U. Egede,³² R. L. Flack,³²
 J. R. Gaillard,³² G. W. Morton,³² J. A. Nash,³² M. B. Nikolich,³² G. P. Taylor,³² W. P. Vazquez,³² M. J. Charles,³³
 W. F. Mader,³³ U. Mallik,³³ A. K. Mohapatra,³³ J. Cochran,³⁴ H. B. Crawley,³⁴ V. Eges,³⁴ W. T. Meyer,³⁴
 S. Prell,³⁴ E. I. Rosenberg,³⁴ A. E. Rubin,³⁴ J. Yi,³⁴ N. Arnaud,³⁵ M. Davier,³⁵ X. Giroux,³⁵ G. Grosdidier,³⁵
 A. Höcker,³⁵ F. Le Diberder,³⁵ V. Lepeltier,³⁵ A. M. Lutz,³⁵ A. Oyanguren,³⁵ T. C. Petersen,³⁵ M. Pierini,³⁵
 S. Plaszczynski,³⁵ S. Rodier,³⁵ P. Roudeau,³⁵ M. H. Schune,³⁵ A. Stocchi,³⁵ G. Wormser,³⁵ C. H. Cheng,³⁶
 D. J. Lange,³⁶ M. C. Simani,³⁶ D. M. Wright,³⁶ A. J. Bevan,³⁷ C. A. Chavez,³⁷ J. P. Coleman,³⁷ I. J. Forster,³⁷
 J. R. Fry,³⁷ E. Gabathuler,³⁷ R. Gamet,³⁷ K. A. George,³⁷ D. E. Hutchcroft,³⁷ R. J. Parry,³⁷ D. J. Payne,³⁷
 K. C. Schofield,³⁷ C. Touramanis,³⁷ C. M. Cormack,³⁸ F. Di Lodovico,³⁸ R. Sacco,³⁸ C. L. Brown,³⁹ G. Cowan,³⁹
 H. U. Flaecher,³⁹ M. G. Green,³⁹ D. A. Hopkins,³⁹ P. S. Jackson,³⁹ T. R. McMahon,³⁹ S. Ricciardi,³⁹ F. Salvatore,³⁹
 D. Brown,⁴⁰ C. L. Davis,⁴⁰ J. Allison,⁴¹ N. R. Barlow,⁴¹ R. J. Barlow,⁴¹ M. C. Hodgkinson,⁴¹ G. D. Lafferty,⁴¹

M. T. Naisbit,⁴¹ J. C. Williams,⁴¹ C. Chen,⁴² A. Farbin,⁴² W. D. Hulsbergen,⁴² A. Jawahery,⁴² D. Kovalskyi,⁴² C. K. Lae,⁴² V. Lillard,⁴² D. A. Roberts,⁴² G. Simi,⁴² G. Blaylock,⁴³ C. Dallapiccola,⁴³ S. S. Hertzbach,⁴³ R. Kofler,⁴³ V. B. Koptchev,⁴³ X. Li,⁴³ T. B. Moore,⁴³ S. Saremi,⁴³ H. Staengle,⁴³ S. Willocq,⁴³ R. Cowan,⁴⁴ K. Koeneke,⁴⁴ G. Sciolla,⁴⁴ S. J. Sekula,⁴⁴ M. Spitznagel,⁴⁴ F. Taylor,⁴⁴ R. K. Yamamoto,⁴⁴ H. Kim,⁴⁵ P. M. Patel,⁴⁵ S. H. Robertson,⁴⁵ A. Lazzaro,⁴⁶ V. Lombardo,⁴⁶ F. Palombo,⁴⁶ J. M. Bauer,⁴⁷ L. Cremaldi,⁴⁷ V. Eschenburg,⁴⁷ R. Godang,⁴⁷ R. Kroeger,⁴⁷ J. Reidy,⁴⁷ D. A. Sanders,⁴⁷ D. J. Summers,⁴⁷ H. W. Zhao,⁴⁷ S. Brunet,⁴⁸ D. Côté,⁴⁸ P. Taras,⁴⁸ B. Viaud,⁴⁸ H. Nicholson,⁴⁹ N. Cavallo,^{50,†} G. De Nardo,⁵⁰ F. Fabozzi,^{50,†} C. Gatto,⁵⁰ L. Lista,⁵⁰ D. Monorchio,⁵⁰ P. Paolucci,⁵⁰ D. Piccolo,⁵⁰ C. Sciacca,⁵⁰ M. Baak,⁵¹ H. Bulten,⁵¹ G. Raven,⁵¹ H. L. Snoek,⁵¹ L. Wilden,⁵¹ C. P. Jessop,⁵² J. M. LoSecco,⁵² T. Allmendinger,⁵³ G. Benelli,⁵³ K. K. Gan,⁵³ K. Honscheid,⁵³ D. Hufnagel,⁵³ P. D. Jackson,⁵³ H. Kagan,⁵³ R. Kass,⁵³ T. Pulliam,⁵³ A. M. Rahimi,⁵³ R. Ter-Antonyan,⁵³ Q. K. Wong,⁵³ J. Brau,⁵⁴ R. Frey,⁵⁴ O. Igonkina,⁵⁴ M. Lu,⁵⁴ C. T. Potter,⁵⁴ N. B. Sinev,⁵⁴ D. Strom,⁵⁴ J. Strube,⁵⁴ E. Torrence,⁵⁴ A. Dorigo,⁵⁵ F. Galeazzi,⁵⁵ M. Margoni,⁵⁵ M. Morandin,⁵⁵ M. Posocco,⁵⁵ M. Rotondo,⁵⁵ F. Simonetto,⁵⁵ R. Stroili,⁵⁵ C. Voci,⁵⁵ M. Benayoun,⁵⁶ H. Briand,⁵⁶ J. Chauveau,⁵⁶ P. David,⁵⁶ L. Del Buono,⁵⁶ Ch. de la Vaissière,⁵⁶ O. Hamon,⁵⁶ M. J. J. John,⁵⁶ Ph. Leruste,⁵⁶ J. Malclès,⁵⁶ J. Ocariz,⁵⁶ L. Roos,⁵⁶ G. Therin,⁵⁶ P. K. Behera,⁵⁷ L. Gladney,⁵⁷ Q. H. Guo,⁵⁷ J. Panetta,⁵⁷ M. Biasini,⁵⁸ R. Covarelli,⁵⁸ S. Pacetti,⁵⁸ M. Pioppi,⁵⁸ C. Angelini,⁵⁹ G. Batignani,⁵⁹ S. Bettarini,⁵⁹ F. Bucci,⁵⁹ G. Calderini,⁵⁹ M. Carpinelli,⁵⁹ R. Cenci,⁵⁹ F. Forti,⁵⁹ M. A. Giorgi,⁵⁹ A. Lusiani,⁵⁹ G. Marchiori,⁵⁹ M. Morganti,⁵⁹ N. Neri,⁵⁹ E. Paoloni,⁵⁹ M. Rama,⁵⁹ G. Rizzo,⁵⁹ J. Walsh,⁵⁹ M. Haire,⁶⁰ D. Judd,⁶⁰ D. E. Wagoner,⁶⁰ J. Biesiada,⁶¹ N. Danielson,⁶¹ P. Elmer,⁶¹ Y. P. Lau,⁶¹ C. Lu,⁶¹ J. Olsen,⁶¹ A. J. S. Smith,⁶¹ A. V. Telnov,⁶¹ F. Bellini,⁶² G. Cavoto,⁶² A. D'Orazio,⁶² E. Di Marco,⁶² R. Faccini,⁶² F. Ferrarotto,⁶² F. Ferroni,⁶² M. Gaspero,⁶² L. Li Gioi,⁶² M. A. Mazzoni,⁶² S. Morganti,⁶² G. Piredda,⁶² F. Polci,⁶² F. Safai Tehrani,⁶² C. Voena,⁶² H. Schröder,⁶³ G. Wagner,⁶³ R. Waldi,⁶³ T. Adye,⁶⁴ N. De Groot,⁶⁴ B. Franek,⁶⁴ G. P. Gopal,⁶⁴ E. O. Olaiya,⁶⁴ F. F. Wilson,⁶⁴ R. Aleksan,⁶⁵ S. Emery,⁶⁵ A. Gaidot,⁶⁵ S. F. Ganzhur,⁶⁵ P.-F. Giraud,⁶⁵ G. Graziani,⁶⁵ G. Hamel de Monchenault,⁶⁵ W. Kozanecki,⁶⁵ M. Legendre,⁶⁵ G. W. London,⁶⁵ B. Mayer,⁶⁵ G. Vasseur,⁶⁵ Ch. Yèche,⁶⁵ M. Zito,⁶⁵ M. V. Purohit,⁶⁶ A. W. Weidemann,⁶⁶ J. R. Wilson,⁶⁶ F. X. Yumiceva,⁶⁶ T. Abe,⁶⁷ M. T. Allen,⁶⁷ D. Aston,⁶⁷ R. Bartoldus,⁶⁷ N. Berger,⁶⁷ A. M. Boyarski,⁶⁷ O. L. Buchmueller,⁶⁷ R. Claus,⁶⁷ M. R. Convery,⁶⁷ M. Cristinziani,⁶⁷ J. C. Dingfelder,⁶⁷ D. Dong,⁶⁷ J. Dorfan,⁶⁷ D. Dujmic,⁶⁷ W. Dunwoodie,⁶⁷ S. Fan,⁶⁷ R. C. Field,⁶⁷ T. Glanzman,⁶⁷ S. J. Gowdy,⁶⁷ T. Hadig,⁶⁷ V. Halyo,⁶⁷ C. Hast,⁶⁷ T. Hryna'ova,⁶⁷ W. R. Innes,⁶⁷ M. H. Kelsey,⁶⁷ P. Kim,⁶⁷ M. L. Kocian,⁶⁷ D. W. G. S. Leith,⁶⁷ J. Libby,⁶⁷ S. Luitz,⁶⁷ V. Luth,⁶⁷ H. L. Lynch,⁶⁷ H. Marsiske,⁶⁷ R. Messner,⁶⁷ D. R. Muller,⁶⁷ C. P. O'Grady,⁶⁷ V. E. Ozcan,⁶⁷ A. Perazzo,⁶⁷ M. Perl,⁶⁷ B. N. Ratcliff,⁶⁷ A. Roodman,⁶⁷ A. A. Salnikov,⁶⁷ R. H. Schindler,⁶⁷ J. Schwiening,⁶⁷ A. Snyder,⁶⁷ J. Stelzer,⁶⁷ D. Su,⁶⁷ M. K. Sullivan,⁶⁷ K. Suzuki,⁶⁷ S. Swain,⁶⁷ J. M. Thompson,⁶⁷ J. Va'vra,⁶⁷ M. Weaver,⁶⁷ A. J. R. Weinstein,⁶⁷ W. J. Wisniewski,⁶⁷ M. Wittgen,⁶⁷ D. H. Wright,⁶⁷ A. K. Yarritu,⁶⁷ K. Yi,⁶⁷ C. C. Young,⁶⁷ P. R. Burchat,⁶⁸ A. J. Edwards,⁶⁸ S. A. Majewski,⁶⁸ B. A. Petersen,⁶⁸ C. Roat,⁶⁸ M. Ahmed,⁶⁹ S. Ahmed,⁶⁹ M. S. Alam,⁶⁹ J. A. Ernst,⁶⁹ M. A. Saeed,⁶⁹ F. R. Wappler,⁶⁹ S. B. Zain,⁶⁹ W. Bugg,⁷⁰ M. Krishnamurthy,⁷⁰ S. M. Spanier,⁷⁰ R. Eckmann,⁷¹ J. L. Ritchie,⁷¹ A. Satpathy,⁷¹ R. F. Schwitters,⁷¹ J. M. Izen,⁷² I. Kitayama,⁷² X. C. Lou,⁷² S. Ye,⁷² F. Bianchi,⁷³ M. Bona,⁷³ F. Gallo,⁷³ D. Gamba,⁷³ M. Bomben,⁷⁴ L. Bosisio,⁷⁴ C. Cartaro,⁷⁴ F. Cossutti,⁷⁴ G. Della Ricca,⁷⁴ S. Dittongo,⁷⁴ S. Grancagnolo,⁷⁴ L. Lanceri,⁷⁴ L. Vitale,⁷⁴ F. Martinez-Vidal,⁷⁵ R. S. Panvini,^{76,†} Sw. Banerjee,⁷⁷ B. Bhuyan,⁷⁷ C. M. Brown,⁷⁷ D. Fortin,⁷⁷ K. Hamano,⁷⁷ R. Kowalewski,⁷⁷ J. M. Roney,⁷⁷ R. J. Sobie,⁷⁷ J. J. Back,⁷⁸ P. F. Harrison,⁷⁸ T. E. Latham,⁷⁸ G. B. Mohanty,⁷⁸ H. R. Band,⁷⁹ X. Chen,⁷⁹ B. Cheng,⁷⁹ S. Dasu,⁷⁹ M. Datta,⁷⁹ A. M. Eichenbaum,⁷⁹ K. T. Flood,⁷⁹ M. Graham,⁷⁹ J. J. Hollar,⁷⁹ J. R. Johnson,⁷⁹ P. E. Kutter,⁷⁹ H. Li,⁷⁹ R. Liu,⁷⁹ B. Mellado,⁷⁹ A. Mihalyi,⁷⁹ Y. Pan,⁷⁹ R. Prepost,⁷⁹ P. Tan,⁷⁹ J. H. von Wimmersperg-Toeller,⁷⁹ S. L. Wu,⁷⁹ Z. Yu,⁷⁹ and H. Neal⁸⁰

(The BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

²IFAE, Universitat Autònoma de Barcelona, E-08193 Bellaterra, Barcelona, Spain

³Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

⁴Institute of High Energy Physics, Beijing 100039, China

⁵University of Bergen, Inst. of Physics, N-50007 Bergen, Norway

⁶Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

⁷University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁸Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁹University of Bristol, Bristol BS8 1TL, United Kingdom

¹⁰University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

- ¹¹Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom
¹²Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia
¹³University of California at Irvine, Irvine, California 92697, USA
¹⁴University of California at Los Angeles, Los Angeles, California 90024, USA
¹⁵University of California at Riverside, Riverside, California 92521, USA
¹⁶University of California at San Diego, La Jolla, California 92093, USA
¹⁷University of California at Santa Barbara, Santa Barbara, California 93106, USA
¹⁸University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA
¹⁹California Institute of Technology, Pasadena, California 91125, USA
²⁰University of Cincinnati, Cincinnati, Ohio 45221, USA
²¹University of Colorado, Boulder, Colorado 80309, USA
²²Colorado State University, Fort Collins, Colorado 80523, USA
²³Universität Dortmund, Institut für Physik, D-44221 Dortmund, Germany
²⁴Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany
²⁵Ecole Polytechnique, LLR, F-91128 Palaiseau, France
²⁶University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom
²⁷Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy
²⁸Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy
²⁹Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy
³⁰Harvard University, Cambridge, Massachusetts 02138, USA
³¹Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany
³²Imperial College London, London, SW7 2AZ, United Kingdom
³³University of Iowa, Iowa City, Iowa 52242, USA
³⁴Iowa State University, Ames, Iowa 50011-3160, USA
³⁵Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France
³⁶Lawrence Livermore National Laboratory, Livermore, California 94550, USA
³⁷University of Liverpool, Liverpool L69 7ZE, United Kingdom
³⁸Queen Mary, University of London, E1 4NS, United Kingdom
³⁹University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom
⁴⁰University of Louisville, Louisville, Kentucky 40292, USA
⁴¹University of Manchester, Manchester M13 9PL, United Kingdom
⁴²University of Maryland, College Park, Maryland 20742, USA
⁴³University of Massachusetts, Amherst, Massachusetts 01003, USA
⁴⁴Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA
⁴⁵McGill University, Montréal, Quebec, Canada H3A 2T8
⁴⁶Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy
⁴⁷University of Mississippi, University, Mississippi 38677, USA
⁴⁸Université de Montréal, Laboratoire René J. A. Lévesque, Montréal, Quebec, Canada H3C 3J7
⁴⁹Mount Holyoke College, South Hadley, Massachusetts 01075, USA
⁵⁰Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy
⁵¹NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands
⁵²University of Notre Dame, Notre Dame, Indiana 46556, USA
⁵³Ohio State University, Columbus, Ohio 43210, USA
⁵⁴University of Oregon, Eugene, Oregon 97403, USA
⁵⁵Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy
⁵⁶Universités Paris VI et VII, Laboratoire de Physique Nucléaire et de Hautes Energies, F-75252 Paris, France
⁵⁷University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
⁵⁸Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy
⁵⁹Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy
⁶⁰Prairie View A&M University, Prairie View, Texas 77446, USA
⁶¹Princeton University, Princeton, New Jersey 08544, USA
⁶²Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy
⁶³Universität Rostock, D-18051 Rostock, Germany
⁶⁴Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom
⁶⁵DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France
⁶⁶University of South Carolina, Columbia, South Carolina 29208, USA
⁶⁷Stanford Linear Accelerator Center, Stanford, California 94309, USA
⁶⁸Stanford University, Stanford, California 94305-4060, USA
⁶⁹State University of New York, Albany, New York 12222, USA
⁷⁰University of Tennessee, Knoxville, Tennessee 37996, USA
⁷¹University of Texas at Austin, Austin, Texas 78712, USA
⁷²University of Texas at Dallas, Richardson, Texas 75083, USA
⁷³Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy
⁷⁴Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy

⁷⁵IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain

⁷⁶Vanderbilt University, Nashville, Tennessee 37235, USA

⁷⁷University of Victoria, Victoria, British Columbia, Canada V8W 3P6

⁷⁸Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom

⁷⁹University of Wisconsin, Madison, Wisconsin 53706, USA

⁸⁰Yale University, New Haven, Connecticut 06511, USA

(Dated: April 24, 2008)

We present a measurement of the partial branching fractions and mass spectra of the exclusive radiative penguin processes $B \rightarrow K\pi\pi\gamma$ in the range $m_{K\pi\pi} < 1.8 \text{ GeV}/c^2$. We reconstruct four final states: $K^+\pi^-\pi^+\gamma$, $K^+\pi^-\pi^0\gamma$, $K_S^0\pi^-\pi^+\gamma$, and $K_S^0\pi^+\pi^0\gamma$, where $K_S^0 \rightarrow \pi^+\pi^-$. Using 232 million $e^+e^- \rightarrow B\bar{B}$ events recorded by the BABAR experiment at the PEP-II asymmetric-energy storage ring, we measure the branching fractions $\mathcal{B}(B^+ \rightarrow K^+\pi^-\pi^+\gamma) = (2.95 \pm 0.13 \text{ (stat.)} \pm 0.20 \text{ (syst.)}) \times 10^{-5}$, $\mathcal{B}(B^0 \rightarrow K^+\pi^-\pi^0\gamma) = (4.07 \pm 0.22 \text{ (stat.)} \pm 0.31 \text{ (syst.)}) \times 10^{-5}$, $\mathcal{B}(B^0 \rightarrow K^0\pi^+\pi^-\gamma) = (1.85 \pm 0.21 \text{ (stat.)} \pm 0.12 \text{ (syst.)}) \times 10^{-5}$, and $\mathcal{B}(B^+ \rightarrow K^0\pi^+\pi^0\gamma) = (4.56 \pm 0.42 \text{ (stat.)} \pm 0.31 \text{ (syst.)}) \times 10^{-5}$.

PACS numbers: 13.25.Hw, 12.15.Hh, 11.30.Er

In the standard model (SM) the radiative penguin decay $B \rightarrow X_s\gamma$, where X_s is a hadronic system with unit strangeness, proceeds via weak-interaction loop diagrams. New physics, beyond the SM, may also contribute to the loop amplitude, and lead to differences from the SM. This possibility has been pursued in inclusive measurements, which are theoretically clean but experimentally challenging, and in exclusive measurements, such as $B \rightarrow K\pi\gamma$. We report measurements of the branching fractions and mass spectra for the decays $B \rightarrow K\pi\pi\gamma$ in four channels. SM predictions of the rates and resonance structure of these decays have large uncertainties [1]. The $K^+\pi^+\pi^-\gamma$ and $K^0\pi^+\pi^-\gamma$ decay channels have previously been observed [2]. Throughout this Letter, stated decays include charge conjugate modes.

The decays $B \rightarrow K\pi^+\pi^0\gamma$, which have not previously been observed, are of particular interest because these three-body hadronic states permit the measurement, given sufficient statistics, of the photon polarization [3]. The polarization measurement depends on the interference between processes such as $(K\pi^+)\pi^0\gamma$ and $(K\pi^0)\pi^+\gamma$, where () indicates resonant substructure. This measurement may be compared with the SM prediction of nearly complete left-handed polarization.

We use a sample of $(232 \pm 1.5) \times 10^6$ $B\bar{B}$ pairs in a 210.9 fb^{-1} dataset collected at the $\Upsilon(4S)$ resonance with the BABAR detector at the PEP-II asymmetric-energy e^+e^- collider. For background studies, we also use a 21.7 fb^{-1} sample collected below the $B\bar{B}$ threshold. The measurement procedure was designed using simulated signal and background events, data in sideband kinematic regions, and reconstructed $B \rightarrow D\pi^+$, $D \rightarrow K\pi\pi$ decays. Only after we established the selection and fit procedures did we examine signal candidates in the data sample.

A description of the detector exists elsewhere [4]. For this measurement, the most important detector elements are the five-layer silicon microstrip tracking detector (SVT) and the forty-layer drift chamber (DCH), situated in a 1.5 T solenoidal magnetic field, which measure charged particle momenta; the CsI(Tl) electromagnetic

calorimeter (EMC), which measures the energies and directions of the photons; and the detector of internally reflected Cherenkov light (DIRC). The DIRC response and energy loss (dE/dx) measured in the SVT and DCH are used to identify charged kaons and pions.

We reconstruct the photon candidate in the $K\pi\pi\gamma$ decay from an EMC shower not associated with a charged track. The photon must be in the fiducial region of the EMC, have a shower-profile consistent with a single photon, and be well-separated from other showers. To remove photons from π^0 (η) decays, we combine the candidate with other photons having energies of at least 50 (250) MeV/c^2 , and reject it if the invariant mass of any combination is within 25 (40) MeV/c^2 of the π^0 (η) mass.

We select K^\pm and π^\pm candidates from charged tracks consistent with a kaon or pion mass hypothesis in the DIRC and in the dE/dx in the SVT and DCH. We reconstruct K_S^0 candidates from pairs of oppositely-charged tracks, and determine the decay vertex with a fit. We require that the invariant mass falls within $11 \text{ MeV}/c^2$ of the K_S^0 mass; that the distance between the B decay vertex and the K_S^0 vertex exceeds 5 times the uncertainty on the distance; and that the angle between the K_S^0 trajectory and its momentum is less than 100 mrad. We reconstruct π^0 candidates from pairs of EMC showers each with energy $> 50 \text{ MeV}$. We require the invariant mass to be within $16 \text{ MeV}/c^2$ of the π^0 mass, and that the energy of each pair in the $\Upsilon(4S)$ center of mass (CM) frame exceeds 450 MeV; this last selection is about 83% efficient.

The dominant source of background is continuum production of light quark-antiquark pairs, in which a high-energy photon typically is produced either by initial state radiation, or from the decay of a π^0 or η in which one photon is not detected. To reject these backgrounds, we construct a Fisher discriminant [5] from the polar angle of the B candidate in the CM frame, the angle between the thrust axis of the B and the thrust axis of the remaining charged and neutral particles, and the ratio of the second to zeroth angular moments of the remaining charged and neutral particles around the thrust axis of the B . We op-

imize the coefficients independently in each channel to discriminate between simulated signal and continuum.

We perform a geometric fit to the reconstructed B candidate, with production vertex constrained to the nominal beam spot, rejecting the candidate if the final state is inconsistent with decay from a single vertex. We define $\Delta E^* \equiv E_B^* - E_{\text{beam}}^*$ and $m_{\text{ES}} \equiv \sqrt{E_{\text{beam}}^{*2} - \mathbf{p}_B^{*2}}$, where E_B^* and \mathbf{p}_B^* are the CM energy and momentum of the B candidate, and E_{beam}^* is the CM energy of each beam. We require $m_{\text{ES}} > 5.2 \text{ GeV}/c^2$ and $|\Delta E^*| < 0.15 \text{ GeV}$. We also require that the invariant mass of the $K\pi\pi$ system, $m_{K\pi\pi}$, fall below $1.8 \text{ GeV}/c^2$; this eliminates much of a rising continuum background with very little expected signal loss. It also removes $K\pi\pi$ combinations from D decays, in $B \rightarrow D\pi^0$ and $B \rightarrow D\eta$ where the π^0 or η are mis-reconstructed as a photon. In an event in which we reconstruct multiple candidates in one channel that pass the selection requirements (occurring in 11–27% of selected signal events, depending on the channel), we keep the candidate with the largest vertex probability (with the best π^0 mass in the $K_s^0\pi^+\pi^0\gamma$ channel; with the π^0 mass as a tie breaker in the $K^+\pi^-\pi^0\gamma$ channel) and reject the others. Candidates reconstructed in different channels are allowed in the same event. The dependence of the efficiency of our selection requirements on intermediate resonance and on $m_{K\pi\pi}$ has been checked and found to be small; systematic uncertainties are discussed later.

The dominant backgrounds from $B\bar{B}$ events after the selection criteria have been applied are $b \rightarrow s\gamma$ processes. We categorize these backgrounds: (i) “crossfeed” from mis-reconstructed $K\pi\pi\gamma$ decays, such as by choosing incorrectly a particle from the other B ; (ii) $B \rightarrow K\pi\gamma$ decays that combine with a track from the other B to form a $K\pi\pi\gamma$ candidate; and (iii) backgrounds from all other $b \rightarrow s\gamma$ decays. A crossfeed candidate may be reconstructed in the same decay channel in which it is produced, or in a different channel, and can also be produced in a $B \rightarrow K\pi\pi\gamma$ decay that is not used in this analysis (such as $B^+ \rightarrow K^+\pi^0\pi^0\gamma$). We model our signal as well as crossfeed backgrounds with simulated $B \rightarrow K_X\gamma$ decays, where K_X is any of the five lowest-lying $J > 0$ kaon resonances above the $K^*(892)$. We study backgrounds from $K\pi\gamma$ using simulated $B \rightarrow K^*(892)\gamma$ and $B \rightarrow K_2^*(1430)\gamma$ decays. We study backgrounds from other $b \rightarrow s\gamma$ decays using an inclusive simulation according to the model of Kagan and Neubert [6] with $m_b = 4.8 \text{ GeV}/c^2$, tuned to match multiplicity distributions measured in inclusive $b \rightarrow s\gamma$ decays [7]. The largest final background contributions from $b \rightarrow s\gamma$ processes are crossfeed backgrounds, for which we obtain yields ranging from 55% to 95% of the signal yields.

We estimate other sources of background candidates from B decays other than $b \rightarrow s\gamma$ processes by simulating generic B decays. We pay special attention to B decays with $K\pi\pi^0$ and $K\pi\pi\eta$ final states; if the π^0 or η decays asymmetrically and we don’t detect the lower-

energy photon, the kinematic properties of the resulting B candidate may resemble a signal candidate. We study these decays using high-statistics simulated samples, and look for signal candidates that are reconstructed from a single $B \rightarrow K\pi\pi^0$ or $B \rightarrow K\pi\pi\eta$ decay. We expect to reconstruct fewer than two such candidates per channel.

We perform a maximum likelihood fit to the joint $m_{\text{ES}}-\Delta E^*$ distribution of our selected candidates. We fit all four channels simultaneously to account for crossfeed backgrounds between channels. The likelihood function contains terms for correctly reconstructed signal candidates, crossfeed background candidates between all 16 combinations of the production and reconstruction channels, backgrounds from $B \rightarrow K\pi\gamma$ and from other $b \rightarrow s\gamma$ decays, and backgrounds from continuum events. We have determined from simulations that the dominant continuum background component adequately accounts for combinatoric backgrounds from other $B\bar{B}$ decays, which do not show strong peaks in m_{ES} and ΔE^* .

The likelihood function for a candidate reconstructed in decay channel i with kinematic variables $y \equiv (m_{\text{ES}}, \Delta E^*)$ is given by,

$$\mathcal{L}^i(y) = N_{B\bar{B}} \left(\mathcal{B}^i \epsilon_s^i f_s^i(y) + \sum_j \mathcal{B}^j \epsilon_x^{ji} f_x^{ji}(y) \right) \\ + n_c^i f_c^i(y) + n_b^i f_b^i(y),$$

where $N_{B\bar{B}}$ is the number of $B\bar{B}$ pairs in our dataset; \mathcal{B}^i is the branching fraction for decay channel i ; ϵ_s^i and f_s^i are the efficiency and probability density function (PDF) for correctly reconstructed signal candidates in decay channel i ; ϵ_x^{ji} and f_x^{ji} are the efficiency and PDF for crossfeed background candidates produced in channel j and reconstructed in channel i ; n_c^i and f_c^i are the yield and PDF for backgrounds from continuum and generic $B\bar{B}$ decay events in channel i ; and n_b^i and f_b^i are the yield and PDF for backgrounds from other $b \rightarrow s\gamma$ processes in channel i . We further parameterize the likelihood function by the four data-taking runs during which data were collected, accounting for slight changes in experimental conditions.

The branching fractions \mathcal{B}^i , yields n_c^i , and shape parameters of f_c^i are varied in the fit; other efficiencies, yields, and PDF shapes are fixed from simulation studies. We parameterize f_s^i as the product of Crystal Ball functions [8] of m_{ES} and of ΔE^* , f_x^i as the product of a Crystal Ball function of m_{ES} and a linear function of ΔE^* , and f_b^i as the product of an Argus function [9] of m_{ES} and an exponential function of ΔE^* . We use a binned parameterization for f_b^i . As the signal and crossfeed terms are both scaled by the parameters \mathcal{B}^i , the crossfeed background yields vary with the signal branching fractions, and we measure the branching fractions from yields of both signal and crossfeed candidates.

Table I shows the fit results. Projections in m_{ES} , along with the fit results, are displayed in Fig. 1. The fit proba-




FIG. 1: Distributions of m_{ES} (points). Projected m_{ES} distributions from the fit are shown as cumulative curves: continuum and generic $B\bar{B}$ component (dashed), $b \rightarrow s\gamma$ component (dotted, includes cross-feed), and signal (solid). The small oscillation in the dotted and solid curves is due to the use of binned distributions to model the $b \rightarrow s\gamma$ component.

bility (P -value) is evaluated with a likelihood ratio statistic [10], assuming Poisson-distributed bin contents, to be 10%. The distribution of the test statistic under the null hypothesis is evaluated by simulation.

Figure 2 shows background-subtracted $m_{K\pi\pi}$ mass spectra. Background subtraction is achieved using the results of the fits to calculate event-by-event weights to extract the signal component [11]. We present branching fractions in bins of $m_{K\pi\pi}$, which are largely model-independent, instead of extracting $B \rightarrow K_X\gamma$ branching fractions for specific K_X resonances. Disentangling the

TABLE I: Results of the fit for $B \rightarrow K\pi\pi\gamma$, for $m_{K\pi\pi} < 1.8$ GeV/ c^2 . The first error is statistical, the second systematic. The yields do not include the channel crossfeeds, which are included in the fit to obtain the branching fractions.

Channel	Yield	Branching Fraction (10^{-5})
$K^+\pi^-\pi^+\gamma$	899 ± 38	$2.95 \pm 0.13 \pm 0.20$
$K^+\pi^-\pi^0\gamma$	572 ± 31	$4.07 \pm 0.22 \pm 0.31$
$K^0\pi^+\pi^-\gamma$	176 ± 20	$1.85 \pm 0.21 \pm 0.12$
$K^0\pi^+\pi^0\gamma$	164 ± 15	$4.56 \pm 0.42 \pm 0.31$




FIG. 2: Background-subtracted $m_{K\pi\pi}$ spectra. The branching fraction in each bin is computed from the weighted event yield. Error bars show statistical uncertainties; the systematic uncertainties due to $b \rightarrow s\gamma$ model assumptions are small.

resonance structure requires careful modeling of amplitudes and relative phases of interfering processes, including in the decays of the K_X resonances, not all of which are well measured. A partial wave analysis to extract the resonance structure and measure the photon polarization should be possible with future datasets.

We validate the procedure for extracting branching fractions and $m_{K\pi\pi}$ distributions using fits to simulated samples. We verify that the branching fractions and mass spectra obtained from these toy fits reproduce on average the simulation inputs. We use the same procedure to extract the $m_{K\pi\pi}$ distributions for continuum and generic $B\bar{B}$ backgrounds and for backgrounds from $b \rightarrow s\gamma$ decays; these are consistent with the expected distributions.

Systematic uncertainties arise from various sources, shown in Table II. The largest sources are: (i) The $\Upsilon(4S)$ branching fractions to $B^+\bar{B}^-$ and $B^0\bar{B}^0$ are each assumed to be 0.5. We assign a 2.6% systematic uncertainty to this, based on current information [12]. (ii) The uncertainty on the photon selection efficiency determined from simulated events is estimated to be 2.7%. (iii) From studies of $B \rightarrow D\pi^\pm$, $D \rightarrow K\pi\pi$ events, we assign an uncertainty of 4.2% to the charged kaon identification efficiency. (iv) The uncertainty of the π^0 selection efficiency is estimated at 3.0%. (v) There is considerable uncertainty in the models we use to estimate backgrounds,

including cross-feed dependence, from $b \rightarrow s\gamma$ processes. We estimate the effect of this uncertainty on both the branching fractions and mass spectra by simulating these backgrounds with substantially different models. The largest effect is in the $K_s^0\pi^-\pi^+\gamma$ channel, where the uncertainty is 4.0%. (vi) We measure a shift in the beam energy in $B \rightarrow D\pi^\pm$ decays, on average 0.6 MeV; we estimate the effect of this on our fits. (vii) We estimate bias in the fit due to uncertain parameterization of the signal and background PDFs. The largest effect is in the $K_s^0\pi^+\pi^0\gamma$ channel, where the uncertainty is 3.5%.

We have measured branching fractions for $B \rightarrow K\pi\pi\gamma$ in four decay channels for $m_{K\pi\pi} < 1.8 \text{ GeV}/c^2$. The $K\pi^+\pi^-$ channels are consistent with the previous measurement [2]. We present first observations of decays in the $K\pi^+\pi^0\gamma$ channels that are important to measuring the photon polarization. The branching fractions are relatively large in the context of $B \rightarrow X_s\gamma$ decays, providing encouragement that a polarization measurement may be possible with future datasets. Mass spectra for the $K\pi\pi$ system are also presented. We observe an enhancement near $1.3 \text{ GeV}/c^2$ and substantial branching fractions at higher masses. Untangling the resonant contributions presents a challenge for the polarization measurement.

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), CEA and CNRS-IN2P3

TABLE II: Estimated systematic uncertainties in the branching fractions, in percent, by source and decay channel.

Source	$K^+\pi^-\pi^+$	$K^+\pi^-\pi^0$	$K_s^0\pi^-\pi^+$	$K_s^0\pi^+\pi^0$
$B\bar{B}$ count	1.1	1.1	1.1	1.1
$\Upsilon(4S)$ BF	2.6	2.6	2.6	2.6
Efficiencies:				
Photon selection	2.7	2.7	2.7	2.7
π^0 and η veto	1.0	1.0	1.0	1.0
Tracking	1.4	1.1	1.1	0.8
Kaon selection	4.2	4.2	1.6	1.6
π^\pm selection	1.4	1.0	1.4	1.0
π^0 selection		3.0		3.0
Fisher cut	1.0	1.0	1.0	1.0
Vertex probability	0.7	0.7	0.7	0.7
MC statistics	0.5	0.7	0.8	1.1
Backgrounds:				
$b \rightarrow s\gamma$ model	1.4	1.0	4.0	1.3
$B \rightarrow K\pi\pi\pi^0/\eta$	0.2	0.1	0.0	0.6
Beam energy shift	1.0	0.5	1.6	0.6
PDF shape	0.1	2.9	0.9	0.2
Fit bias	1.6	1.3	1.3	3.5
Total	6.7	7.6	6.7	6.8

(France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MES (Russia), MEC (Spain), and STFC (United Kingdom). Individuals have received support from the Marie Curie EIF (European Union) and the A. P. Sloan Foundation.

* Also with Università di Perugia, Dipartimento di Fisica, Perugia, Italy

† Also with Università della Basilicata, Potenza, Italy

‡ Deceased

- [1] For example, H. Cheng and C. Chua, Phys. Rev. D **69**, 094007 (2004); D. Ebert et al., Phys. Rev. D **64**, 054001 (2001); A. S. Safir, Eur. Phys. J. C **3**, 15 (2001); S. Veseli and M. G. Olsson, Phys. Lett. B **367**, 309 (1996).
- [2] Belle Collaboration, H. Yang *et al.*, Phys. Rev. Lett. **94**, 111802 (2005).
- [3] M. Gronau, *et al.*, Phys. Rev. Lett. **88**, 051802 (2002); M. Gronau and D. Pirjol, Phys. Rev. D **66**, 054008 (2002).
- [4] *BABAR* Collaboration, B. Aubert *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 1 (2002).
- [5] R. A. Fisher, Annals Eugen. **7**, 179 (1936).
- [6] A. Kagan and M. Neubert, Eur. Phys. J. C **7**, 5 (1999).
- [7] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **93**, 021804 (2004).
- [8] T. Skwarnicki, DESY F31-86-02 (thesis, unpublished) (1986).
- [9] ARGUS Collaboration, H. Albrecht *et al.*, Phys. Lett. **B185**, 218 (1975).
- [10] W. T. Eadie *et al.*, "Statistical Methods in Experimental Physics", North-Holland (1971).
- [11] M. Pivk and F. R. LeDiberder, physics/0402083 (2004).
- [12] *BABAR* Collaboration, B. Aubert *et al.*, hep-ex/0504001, to be published in Phys. Rev. Lett. (2005).