

# Improved Measurement of $CP$ Asymmetries in $B^0 \rightarrow (c\bar{c})K^{(*)0}$ Decays

The *BABAR* Collaboration

July 29, 2006

## Abstract

We present an updated measurements of time-dependent  $CP$  asymmetries in fully reconstructed neutral  $B$  decays to several  $CP$  eigenstates containing a charmonium meson. The measurements use a data sample of  $(347.5 \pm 3.8) \times 10^6 \Upsilon(4S) \rightarrow B\bar{B}$  decays collected with the *BABAR* detector at the PEP-II  $B$  factory between 1999 and 2006. We determine  $\sin 2\beta = 0.710 \pm 0.034(\text{stat}) \pm 0.019(\text{syst})$  and  $|\lambda| = 0.932 \pm 0.026(\text{stat}) \pm 0.017(\text{syst})$ . Both of these results are preliminary.

Submitted to the 33<sup>rd</sup> International Conference on High-Energy Physics, ICHEP 06,  
26 July—2 August 2006, Moscow, Russia.

---

*Stanford Linear Accelerator Center, Stanford University, Stanford, CA 94309*

Work supported in part by Department of Energy contract DE-AC02-76SF00515.

The BABAR Collaboration,

B. Aubert, R. Barate, M. Bona, D. Boutigny, F. Couderc, Y. Karyotakis, J. P. Lees, V. Poireau,  
V. Tisserand, A. Zghiche

*Laboratoire de Physique des Particules, IN2P3/CNRS et Université de Savoie, F-74941 Annecy-Le-Vieux,  
France*

E. Grauges

*Universitat de Barcelona, Facultat de Física, Departament ECM, E-08028 Barcelona, Spain*

A. Palano

*Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy*

J. C. Chen, N. D. Qi, G. Rong, P. Wang, Y. S. Zhu

*Institute of High Energy Physics, Beijing 100039, China*

G. Eigen, I. Ofte, B. Stugu

*University of Bergen, Institute of Physics, N-5007 Bergen, Norway*

G. S. Abrams, M. Battaglia, D. N. Brown, J. Button-Shafer, R. N. Cahn, E. Charles, M. S. Gill,  
Y. Groysman, R. G. Jacobsen, J. A. Kadyk, L. T. Kerth, Yu. G. Kolomensky, G. Kukartsev, G. Lynch,  
L. M. Mir, T. J. Orimoto, M. Pripstein, N. A. Roe, M. T. Ronan, W. A. Wenzel

*Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA*

P. del Amo Sanchez, M. Barrett, K. E. Ford, A. J. Hart, T. J. Harrison, C. M. Hawkes, S. E. Morgan,  
A. T. Watson

*University of Birmingham, Birmingham, B15 2TT, United Kingdom*

T. Held, H. Koch, B. Lewandowski, M. Pelizaeus, K. Peters, T. Schroeder, M. Steinke  
*Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany*

J. T. Boyd, J. P. Burke, W. N. Cottingham, D. Walker

*University of Bristol, Bristol BS8 1TL, United Kingdom*

D. J. Asgeirsson, T. Cuhadar-Donszelmann, B. G. Fulsom, C. Hearty, N. S. Knecht, T. S. Mattison,  
J. A. McKenna

*University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1*

A. Khan, P. Kyberd, M. Saleem, D. J. Sherwood, L. Teodorescu

*Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom*

V. E. Blinov, A. D. Bukin, V. P. Druzhinin, V. B. Golubev, A. P. Onuchin, S. I. Serednyakov,  
Yu. I. Skovpen, E. P. Solodov, K. Yu Todyshev

*Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia*

D. S. Best, M. Bondioli, M. Bruinsma, M. Chao, S. Curry, I. Eschrich, D. Kirkby, A. J. Lankford, P. Lund,  
M. Mandelkern, E. Martin, R. K. Mommsen, W. Roethel, D. P. Stoker

*University of California at Irvine, Irvine, California 92697, USA*

S. Abachi, C. Buchanan

*University of California at Los Angeles, Los Angeles, California 90024, USA*

S. D. Foulkes, J. W. Gary, O. Long, B. C. Shen, K. Wang, L. Zhang  
*University of California at Riverside, Riverside, California 92521, USA*

H. K. Hadavand, E. J. Hill, H. P. Paar, S. Rahatlou, V. Sharma  
*University of California at San Diego, La Jolla, California 92093, USA*

J. W. Berryhill, C. Campagnari, A. Cunha, B. Dahmes, T. M. Hong, D. Kovalskyi, J. D. Richman  
*University of California at Santa Barbara, Santa Barbara, California 93106, USA*

T. W. Beck, A. M. Eisner, C. J. Flacco, C. A. Heusch, J. Kroseberg, W. S. Lockman, G. Nesom, T. Schalk,  
B. A. Schumm, A. Seiden, P. Spradlin, D. C. Williams, M. G. Wilson  
*University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA*

J. Albert, E. Chen, A. Dvoretzkii, F. Fang, D. G. Hitlin, I. Narsky, T. Piatenko, F. C. Porter, A. Ryd,  
A. Samuel  
*California Institute of Technology, Pasadena, California 91125, USA*

G. Mancinelli, B. T. Meadows, K. Mishra, M. D. Sokoloff  
*University of Cincinnati, Cincinnati, Ohio 45221, USA*

F. Blanc, P. C. Bloom, S. Chen, W. T. Ford, J. F. Hirschauer, A. Kreisel, M. Nagel, U. Nauenberg,  
A. Olivas, W. O. Ruddick, J. G. Smith, K. A. Ulmer, S. R. Wagner, J. Zhang  
*University of Colorado, Boulder, Colorado 80309, USA*

A. Chen, E. A. Eckhart, A. Soffer, W. H. Toki, R. J. Wilson, F. Winklmeier, Q. Zeng  
*Colorado State University, Fort Collins, Colorado 80523, USA*

D. D. Altenburg, E. Feltresi, A. Hauke, H. Jasper, J. Merkel, A. Petzold, B. Spaan  
*Universität Dortmund, Institut für Physik, D-44221 Dortmund, Germany*

T. Brandt, V. Klose, H. M. Lacker, W. F. Mader, R. Nogowski, J. Schubert, K. R. Schubert, R. Schwierz,  
J. E. Sundermann, A. Volk  
*Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany*

D. Bernard, G. R. Bonneaud, E. Latour, Ch. Thiebaux, M. Verderi  
*Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, F-91128 Palaiseau, France*

P. J. Clark, W. Gradl, F. Muheim, S. Playfer, A. I. Robertson, Y. Xie  
*University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom*

M. Andreotti, D. Bettoni, C. Bozzi, R. Calabrese, G. Cibinetto, E. Luppi, M. Negrini, A. Petrella,  
L. Piemontese, E. Prencipe  
*Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy*

F. Anulli, R. Baldini-Ferrolì, A. Calcaterra, R. de Sangro, G. Finocchiaro, S. Pacetti, P. Patteri,  
I. M. Peruzzi,<sup>1</sup> M. Piccolo, M. Rama, A. Zallo  
*Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy*

---

<sup>1</sup>Also with Università di Perugia, Dipartimento di Fisica, Perugia, Italy

A. Buzzo, R. Capra, R. Contri, M. Lo Vetere, M. M. Macri, M. R. Monge, S. Passaggio, C. Patrignani,  
E. Robutti, A. Santroni, S. Tosi

*Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy*

G. Brandenburg, K. S. Chaisanguanthum, M. Morii, J. Wu

*Harvard University, Cambridge, Massachusetts 02138, USA*

R. S. Dubitzky, J. Marks, S. Schenk, U. Uwer

*Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany*

D. J. Bard, W. Bhimji, D. A. Bowerman, P. D. Dauncey, U. Egede, R. L. Flack, J. A. Nash,  
M. B. Nikolich, W. Panduro Vazquez

*Imperial College London, London, SW7 2AZ, United Kingdom*

P. K. Behera, X. Chai, M. J. Charles, U. Mallik, N. T. Meyer, V. Ziegler

*University of Iowa, Iowa City, Iowa 52242, USA*

J. Cochran, H. B. Crawley, L. Dong, V. Eyges, W. T. Meyer, S. Prell, E. I. Rosenberg, A. E. Rubin

*Iowa State University, Ames, Iowa 50011-3160, USA*

A. V. Gritsan

*Johns Hopkins University, Baltimore, Maryland 21218, USA*

A. G. Denig, M. Fritsch, G. Schott

*Universität Karlsruhe, Institut für Experimentelle Kernphysik, D-76021 Karlsruhe, Germany*

N. Arnaud, M. Davier, G. Grosdidier, A. Höcker, F. Le Diberder, V. Lepeltier, A. M. Lutz, A. Oyanguren,  
S. Pruvot, S. Rodier, P. Roudeau, M. H. Schune, A. Stocchi, W. F. Wang, G. Wormser

*Laboratoire de l'Accélérateur Linéaire, IN2P3/CNRS et Université Paris-Sud 11, Centre Scientifique  
d'Orsay, B.P. 34, F-91898 ORSAY Cedex, France*

C. H. Cheng, D. J. Lange, D. M. Wright

*Lawrence Livermore National Laboratory, Livermore, California 94550, USA*

C. A. Chavez, I. J. Forster, J. R. Fry, E. Gabathuler, R. Gamet, K. A. George, D. E. Hutchcroft,  
D. J. Payne, K. C. Schofield, C. Touramanis

*University of Liverpool, Liverpool L69 7ZE, United Kingdom*

A. J. Bevan, F. Di Lodovico, W. Menges, R. Sacco

*Queen Mary, University of London, E1 4NS, United Kingdom*

G. Cowan, H. U. Flaecher, D. A. Hopkins, P. S. Jackson, T. R. McMahon, S. Ricciardi, F. Salvatore,  
A. C. Wren

*University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United  
Kingdom*

D. N. Brown, C. L. Davis

*University of Louisville, Louisville, Kentucky 40292, USA*

J. Allison, N. R. Barlow, R. J. Barlow, Y. M. Chia, C. L. Edgar, G. D. Lafferty, M. T. Naisbit,  
J. C. Williams, J. I. Yi

*University of Manchester, Manchester M13 9PL, United Kingdom*

C. Chen, W. D. Hulsbergen, A. Jawahery, C. K. Lae, D. A. Roberts, G. Simi

*University of Maryland, College Park, Maryland 20742, USA*

G. Blaylock, C. Dallapiccola, S. S. Hertzbach, X. Li, T. B. Moore, S. Saremi, H. Staengle

*University of Massachusetts, Amherst, Massachusetts 01003, USA*

R. Cowan, G. Sciolla, S. J. Sekula, M. Spitznagel, F. Taylor, R. K. Yamamoto

*Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139,  
USA*

H. Kim, S. E. McLachlin, P. M. Patel, S. H. Robertson

*McGill University, Montréal, Québec, Canada H3A 2T8*

A. Lazzaro, V. Lombardo, F. Palombo

*Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy*

J. M. Bauer, L. Cremaldi, V. Eschenburg, R. Godang, R. Kroeger, D. A. Sanders, D. J. Summers,  
H. W. Zhao

*University of Mississippi, University, Mississippi 38677, USA*

S. Brunet, D. Côté, M. Simard, P. Taras, F. B. Viaud

*Université de Montréal, Physique des Particules, Montréal, Québec, Canada H3C 3J7*

H. Nicholson

*Mount Holyoke College, South Hadley, Massachusetts 01075, USA*

N. Cavallo,<sup>2</sup> G. De Nardo, F. Fabozzi,<sup>3</sup> C. Gatto, L. Lista, D. Monorchio, P. Paolucci, D. Piccolo,  
C. Sciacca

*Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy*

M. A. Baak, G. Raven, H. L. Snoek

*NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The  
Netherlands*

C. P. Jessop, J. M. LoSecco

*University of Notre Dame, Notre Dame, Indiana 46556, USA*

T. Allmendinger, G. Benelli, L. A. Corwin, K. K. Gan, K. Honscheid, D. Hufnagel, P. D. Jackson,  
H. Kagan, R. Kass, A. M. Rahimi, J. J. Regensburger, R. Ter-Antonyan, Q. K. Wong

*Ohio State University, Columbus, Ohio 43210, USA*

N. L. Blount, J. Brau, R. Frey, O. Igonkina, J. A. Kolb, M. Lu, R. Rahmat, N. B. Sinev, D. Strom,  
J. Strube, E. Torrence

*University of Oregon, Eugene, Oregon 97403, USA*

---

<sup>2</sup>Also with Università della Basilicata, Potenza, Italy

<sup>3</sup>Also with Università della Basilicata, Potenza, Italy

A. Gaz, M. Margoni, M. Morandin, A. Pompili, M. Posocco, M. Rotondo, F. Simonetto, R. Stroili, C. Voci  
*Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy*

M. Benayoun, H. Briand, J. Chauveau, P. David, L. Del Buono, Ch. de la Vaissière, O. Hamon,  
B. L. Hartfiel, M. J. J. John, Ph. Leruste, J. Malcès, J. Ocariz, L. Roos, G. Therin  
*Laboratoire de Physique Nucléaire et de Hautes Energies, IN2P3/CNRS, Université Pierre et Marie  
Curie-Paris6, Université Denis Diderot-Paris7, F-75252 Paris, France*

L. Gladney, J. Panetta  
*University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*

M. Biasini, R. Covarelli, E. Manoni  
*Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy*

C. Angelini, G. Batignani, S. Bettarini, F. Bucci, G. Calderini, M. Carpinelli, R. Cenci, F. Forti,  
M. A. Giorgi, A. Lusiani, G. Marchiori, M. A. Mazur, M. Morganti, N. Neri, E. Paoloni, G. Rizzo,  
J. J. Walsh  
*Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy*

M. Haire, D. Judd, D. E. Wagoner  
*Prairie View A&M University, Prairie View, Texas 77446, USA*

J. Biesiada, N. Danielson, P. Elmer, Y. P. Lau, C. Lu, J. Olsen, A. J. S. Smith, A. V. Telnov  
*Princeton University, Princeton, New Jersey 08544, USA*

F. Bellini, G. Cavoto, A. D’Orazio, D. del Re, E. Di Marco, R. Faccini, F. Ferrarotto, F. Ferroni,  
M. Gaspero, L. Li Gioi, M. A. Mazzoni, S. Morganti, G. Piredda, F. Polci, F. Safai Tehrani, C. Voena  
*Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy*

M. Ebert, H. Schröder, R. Waldi  
*Universität Rostock, D-18051 Rostock, Germany*

T. Adye, N. De Groot, B. Franek, E. O. Olaiya, F. F. Wilson  
*Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom*

R. Aleksan, S. Emery, A. Gaidot, S. F. Ganzhur, G. Hamel de Monchenault, W. Kozanecki, M. Legendre,  
G. Vasseur, Ch. Yèche, M. Zito  
*DSM/Daphnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France*

X. R. Chen, H. Liu, W. Park, M. V. Purohit, J. R. Wilson  
*University of South Carolina, Columbia, South Carolina 29208, USA*

M. T. Allen, D. Aston, R. Bartoldus, P. Bechtle, N. Berger, R. Claus, J. P. Coleman, M. R. Convery,  
M. Cristinziani, J. C. Dingfelder, J. Dorfan, G. P. Dubois-Felsmann, D. Dujmic, W. Dunwoodie,  
R. C. Field, T. Glanzman, S. J. Gowdy, M. T. Graham, P. Grenier,<sup>4</sup> V. Halyo, C. Hast, T. Hryn’ova,  
W. R. Innes, M. H. Kelsey, P. Kim, D. W. G. S. Leith, S. Li, S. Luitz, V. Luth, H. L. Lynch,  
D. B. MacFarlane, H. Marsiske, R. Messner, D. R. Muller, C. P. O’Grady, V. E. Ozcan, A. Perazzo,  
M. Perl, T. Pulliam, B. N. Ratcliff, A. Roodman, A. A. Salnikov, R. H. Schindler, J. Schwiening,  
A. Snyder, J. Stelzer, D. Su, M. K. Sullivan, K. Suzuki, S. K. Swain, J. M. Thompson, J. Va’vra, N. van

---

<sup>4</sup>Also at Laboratoire de Physique Corpusculaire, Clermont-Ferrand, France

Bakel, M. Weaver, A. J. R. Weinstein, W. J. Wisniewski, M. Wittgen, D. H. Wright, A. K. Yarritu, K. Yi,  
C. C. Young

*Stanford Linear Accelerator Center, Stanford, California 94309, USA*

P. R. Burchat, A. J. Edwards, S. A. Majewski, B. A. Petersen, C. Roat, L. Wilden

*Stanford University, Stanford, California 94305-4060, USA*

S. Ahmed, M. S. Alam, R. Bula, J. A. Ernst, V. Jain, B. Pan, M. A. Saeed, F. R. Wappler, S. B. Zain

*State University of New York, Albany, New York 12222, USA*

W. Bugg, M. Krishnamurthy, S. M. Spanier

*University of Tennessee, Knoxville, Tennessee 37996, USA*

R. Eckmann, J. L. Ritchie, A. Satpathy, C. J. Schilling, R. F. Schwitters

*University of Texas at Austin, Austin, Texas 78712, USA*

J. M. Izen, X. C. Lou, S. Ye

*University of Texas at Dallas, Richardson, Texas 75083, USA*

F. Bianchi, F. Gallo, D. Gamba

*Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy*

M. Bomben, L. Bosisio, C. Cartaro, F. Cossutti, G. Della Ricca, S. Dittongo, L. Lanceri, L. Vitale

*Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy*

V. Azzolini, N. Lopez-March, F. Martinez-Vidal

*IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain*

Sw. Banerjee, B. Bhuyan, C. M. Brown, D. Fortin, K. Hamano, R. Kowalewski, I. M. Nugent, J. M. Roney,  
R. J. Sobie

*University of Victoria, Victoria, British Columbia, Canada V8W 3P6*

J. J. Back, P. F. Harrison, T. E. Latham, G. B. Mohanty, M. Pappagallo

*Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom*

H. R. Band, X. Chen, B. Cheng, S. Dasu, M. Datta, K. T. Flood, J. J. Hollar, P. E. Kutter, B. Mellado,  
A. Mihalyi, Y. Pan, M. Pierini, R. Prepost, S. L. Wu, Z. Yu

*University of Wisconsin, Madison, Wisconsin 53706, USA*

H. Neal

*Yale University, New Haven, Connecticut 06511, USA*

# 1 INTRODUCTION

Charge conjugation-parity ( $CP$ ) violation in the  $B$  meson system has been established by the *BABAR* [1] and Belle [2] collaborations. The Standard Model (SM) of electroweak interactions describes  $CP$  violation as a consequence of a complex phase in the three-generation Cabibbo-Kobayashi-Maskawa (CKM) quark-mixing matrix [3]. In this framework, measurements of  $CP$  asymmetries in the proper-time distribution of neutral  $B$  decays to  $CP$  eigenstates containing a charmonium and  $K^0$  meson provide a direct measurement of  $\sin 2\beta$  [4]. The angle  $\beta$  is defined as  $\arg[-(V_{cd}V_{cb}^*)/(V_{td}V_{tb}^*)]$ , where the  $V_{ij}$  are CKM matrix elements.

In this paper, we report on an updated measurement of  $\sin 2\beta$  in  $(347.5 \pm 3.8) \times 10^6 \Upsilon(4S) \rightarrow B\bar{B}$  decays collected with the *BABAR* detector using  $B^0$  decays to the final states  $J/\psi K_S^0$ ,  $J/\psi K_L^0$ ,  $\psi(2S)K_S^0$ ,  $\chi_{c1}K_S^0$ ,  $\eta_c K_S^0$ , and  $J/\psi K^{*0}(K^{*0} \rightarrow K_S^0\pi^0)$ . Since our previous measurement [5], we have added a sample of  $120 \times 10^6 \Upsilon(4S) \rightarrow B\bar{B}$  decays, and applied an improved event reconstruction to the complete dataset. A new  $\eta_c K_S^0$  event selection has been developed based on the Dalitz structure of the  $\eta_c \rightarrow K_S^0 K^+\pi^-$  decay. We have also performed a more detailed study of the  $CP$  properties of our background events resulting in a reduced systematic error.

## 2 THE *BABAR* DETECTOR AND DATASET

The data used in this analysis were collected with the *BABAR* detector at the PEP-II asymmetric  $e^+e^-$  storage ring from 1999 to 2006. This represents a total integrated luminosity of  $(316.2 \pm 3.5) \text{ fb}^{-1}$  taken at the  $\Upsilon(4S)$  resonance (onpeak), corresponding to a sample of  $(347.5 \pm 3.8) \times 10^6 \Upsilon(4S) \rightarrow B\bar{B}$  decays.

The *BABAR* detector is described in detail elsewhere [6]. Charged particles are selected and their momenta are measured by a combination of a vertex tracker consisting of five layers of double-sided silicon microstrip detectors and a 40-layer central drift chamber, both operating in the 1.5 T magnetic field of a superconducting solenoid. We identify photons and electrons using a CsI(Tl) electromagnetic calorimeter (EMC). Charged particle identification (PID) is provided by an internally reflected ring imaging Cherenkov detector (DIRC) covering the central region of the detector, the average energy loss ( $dE/dx$ ) in the tracking devices, and by the EMC. In addition, the instrumented flux return (IFR) containing resistive plate chambers are used for muon and long-lived neutral hadron identifications. We use the GEANT4 [7] software to simulate interactions of particles traversing the *BABAR* detector.

## 3 ANALYSIS METHOD

We use information from the other  $B$  meson,  $B_{\text{tag}}$ , in the event to tag the initial flavor of the fully reconstructed  $B$  candidate. The decay rate  $f_+(f_-)$  for a neutral  $B$  meson decaying to a  $CP$  eigenstate accompanied by a  $B^0(\bar{B}^0)$  tag can be expressed in terms of a complex parameter  $\lambda$  [8] as

$$f_{\pm}(\Delta t) = \frac{e^{-|\Delta t|/\tau_{B^0}}}{4\tau_{B^0}} \left\{ (1 \mp \Delta\omega) \pm (1 - 2\omega) \times \left[ \frac{2\text{Im}\lambda}{1 + |\lambda|^2} \sin(\Delta m_d \Delta t) - \frac{1 - |\lambda|^2}{1 + |\lambda|^2} \cos(\Delta m_d \Delta t) \right] \right\}, \quad (1)$$

where  $\Delta t = t_{\text{rec}} - t_{\text{tag}}$  is the difference between the proper decay times of the reconstructed ( $B_{\text{rec}}$ ) and tagged ( $B_{\text{tag}}$ )  $B$  mesons.  $\tau_{B^0}$  is the  $B^0$  lifetime and  $\Delta m_d$  is the mass difference determined



from  $B^0$ - $\bar{B}^0$  oscillations [9]. The average mistag probability  $\omega$  describes the effect of incorrect tags, and  $\Delta\omega$  is the difference between the mistag rate for  $B^0$  and  $\bar{B}^0$ . Here we assume that the decay width difference  $\Delta\Gamma_d$  between the neutral  $B$  mass eigenstates is zero. The sine term in Eq. 1 is due to the interference between the direct decay and the decay after  $B^0$ - $\bar{B}^0$  oscillation. A non-zero cosine term arises from the interference between decay amplitudes with different weak and strong phases (direct  $CP$  violation) or from  $CP$  violation in  $B^0$ - $\bar{B}^0$  mixing.

In the SM,  $CP$  violation in mixing is negligible, as is direct  $CP$  violation for  $b \rightarrow c\bar{c}s$  decays that contain a charmonium meson [8]. Under these assumptions,  $\lambda = \eta_f e^{-2i\beta}$  where  $\eta_f = \pm 1$  is the  $CP$  eigenvalue of the final state  $f$ . Thus, the time-dependent  $CP$ -violating asymmetry is

$$A_{CP}(\Delta t) \equiv \frac{f_+(\Delta t) - f_-(\Delta t)}{f_+(\Delta t) + f_-(\Delta t)} \propto -(1 - 2\omega)\eta_f \sin 2\beta \sin(\Delta m_d \Delta t). \quad (2)$$

We reconstruct a sample of neutral  $B$  mesons,  $B_{CP}$ , decaying to the  $\eta_f = -1$  final states [10]  $J/\psi K_S^0$ ,  $\psi(2S)K_S^0$ ,  $\chi_{c1}K_S^0$  and  $\eta_c K_S^0$ , and the  $\eta_f = +1$  final state  $J/\psi K_L^0$ . The  $J/\psi$  and  $\psi(2S)$  mesons are reconstructed through their decays to  $e^+e^-$  and  $\mu^+\mu^-$ ; the  $\psi(2S)$  is also reconstructed through its decay to  $J/\psi\pi^+\pi^-$ . The  $\chi_{c1}$  meson is reconstructed in the decay mode  $J/\psi\gamma$ . The  $\eta_c$  meson is reconstructed in the decay mode  $K_S^0 K^+\pi^-$ . We also reconstruct the  $J/\psi K^{*0}$  ( $K^{*0} \rightarrow K_S^0\pi^0$ ) final state which can be  $CP$ -even or  $CP$ -odd due to the presence of even ( $L=0, 2$ ) and odd ( $L=1$ ) orbital angular momenta contributions. Ignoring the angular information in  $J/\psi K^{*0}$  results in a reduction of the measured  $CP$  asymmetry by a factor  $|1 - 2R_\perp|$ , where  $R_\perp$  is the fraction of the  $L=1$  contribution. We have measured  $R_\perp = 0.233 \pm 0.010(\text{stat}) \pm 0.005(\text{syst})$  [11], which gives an effective  $\eta_f = 0.504 \pm 0.033$ , after acceptance corrections.

In addition to the  $CP$  modes described above, a large sample  $B_{\text{flav}}$  of  $B^0$  decays to the flavor eigenstates  $D^{(*)-}h^+$  ( $h^+ = \pi^+, \rho^+$ , and  $a_1^+$ ) and  $J/\psi K^{*0}$  ( $K^{*0} \rightarrow K^+\pi^-$ ) is used for calibrating the flavor tagging performance and  $\Delta t$  resolution. We perform studies to measure apparent  $CP$  violation from unphysical sources using a control sample of  $B^+$  mesons decaying to the final states  $J/\psi K^{(*)+}$ ,  $\psi(2S)K^+$ ,  $\chi_{c1}K^+$ , and  $\eta_c K^+$ . Since the previous *BABAR* analyses, we apply an improved event reconstruction to all the events and we find that on a subset of the data the new and old event reconstruction algorithms give consistent results. The event selection and candidate reconstruction are unchanged from those described in Refs. [5, 12, 13] with the exceptions described below. As in Ref. [5] we reconstruct the  $B^0 \rightarrow \eta_c K_S^0$  and  $B^\pm \rightarrow \eta_c K^\pm$  modes using only the  $\eta_c \rightarrow K_S^0 K^+\pi^-$  decay with  $2.91 < m_{K_S^0 K^+\pi^-} < 3.05 \text{ GeV}/c^2$ . We now exploit the fact that the  $\eta_c$  decays predominantly through a  $K\pi$  resonance at around  $1430 \text{ MeV}/c^2$ , and apply the selection criteria  $1.26 \text{ GeV}/c^2 < m(K_S^0\pi^-) < 1.63 \text{ GeV}/c^2$  and  $1.26 \text{ GeV}/c^2 < m(K^+\pi^-) < 1.63 \text{ GeV}/c^2$ .

We calculate the time interval  $\Delta t$  between the two  $B$  decays from the measured separation  $\Delta z$  between the decay vertices of  $B_{\text{rec}}$  and  $B_{\text{tag}}$  along the collision ( $z$ ) axis [12]. The  $z$  position of the  $B_{\text{rec}}$  vertex is determined from the charged daughter tracks. The  $B_{\text{tag}}$  decay vertex is determined by fitting tracks not belonging to the  $B_{\text{rec}}$  candidate to a common vertex, employing constraints from the beam spot location and the  $B_{\text{rec}}$  momentum [12]. Events are accepted if the calculated  $\Delta t$  uncertainty is less than 2.5 ps and  $|\Delta t|$  is less than 20 ps. The fraction of events satisfying these requirements is 95%. The r.m.s.  $\Delta t$  resolution is 1.1 ps for the 99.7% of events that are not outliers.

Multivariate algorithms are used to identify signatures of  $B$  decays that determine (“tag”) the flavor of the  $B_{\text{tag}}$  at decay to be either a  $B^0$  or  $\bar{B}^0$  candidate. Primary leptons from semileptonic  $B$  decays are selected from identified electrons and muons as well as isolated energetic tracks. The charges of identified kaon candidates are used in a kaon tag. Low momentum pions from  $D^{*+}$  decays are selected on the basis of their momentum and direction with respect to the thrust axis

of  $B_{\text{tag}}$ . These algorithms are combined to account for correlations among different sources of flavor information and to provide an estimate of the mistag probability for each event. Each event whose estimated mistag probability is less than 45% is assigned to one of six tagging categories. The **Lepton** category contains events with an identified lepton; the remaining events are divided into the **Kaon I**, **Kaon II**, **Kaon-Pion**, **Pion**, or **Other** categories based on the estimated mistag probability. For each category  $i$ , the tagging efficiency  $\varepsilon_i$  and fraction  $w_i$  of events having the wrong tag assignment are measured from data (Table 1). The figure of merit for tagging is the effective tagging efficiency  $Q \equiv \sum_i \varepsilon_i(1 - 2w_i)^2 = (30.4 \pm 0.3)\%$ , where the error shown is statistical only.

Table 1: Efficiencies  $\varepsilon_i$ , average mistag fractions  $w_i$ , mistag fraction differences  $\Delta w_i \equiv w_i(B^0) - w_i(\bar{B}^0)$ , and effective tagging efficiency  $Q_i$  extracted for each tagging category  $i$  from the  $B_{\text{flav}}$  sample.

Category	$\varepsilon$ (%)	$w$ (%)	$\Delta w$ (%)	$Q$ (%)
<b>Lepton</b>	$8.67 \pm 0.08$	$3.0 \pm 0.3$	$-0.2 \pm 0.6$	$7.67 \pm 0.13$
<b>Kaon I</b>	$10.96 \pm 0.09$	$5.3 \pm 0.4$	$-0.6 \pm 0.7$	$8.74 \pm 0.16$
<b>Kaon II</b>	$17.21 \pm 0.11$	$15.5 \pm 0.4$	$-0.4 \pm 0.7$	$8.21 \pm 0.19$
<b>Kaon-Pion</b>	$13.77 \pm 0.10$	$23.5 \pm 0.5$	$-2.4 \pm 0.8$	$3.87 \pm 0.14$
<b>Pion</b>	$14.38 \pm 0.10$	$33.0 \pm 0.5$	$5.2 \pm 0.8$	$1.67 \pm 0.10$
<b>Other</b>	$9.61 \pm 0.08$	$41.9 \pm 0.6$	$4.6 \pm 0.9$	$0.25 \pm 0.04$
<b>All</b>	$74.60 \pm 0.12$			$30.4 \pm 0.3$

With the exception of the  $J/\psi K_L^0$  mode, we use the beam-energy substituted mass  $m_{\text{ES}} = \sqrt{(E_{\text{beam}}^*)^2 - (p_B^*)^2}$  to determine the composition of our final sample, where  $E_{\text{beam}}^*$  and  $p_B^*$  are the beam energy and  $B$  momentum in the  $e^+e^-$  center-of-mass frame. For the  $J/\psi K_L^0$  mode we use the difference  $\Delta E$  between the candidate center-of-mass energy and  $E_{\text{beam}}^*$ . The composition of our final sample is shown in Fig. 1. We use events with  $m_{\text{ES}} > 5.2 \text{ GeV}/c^2$  ( $\Delta E < 80 \text{ MeV}$  for  $J/\psi K_L^0$ ) in order to determine the properties of the background contributions. We define a signal region  $5.27 < m_{\text{ES}} < 5.29 \text{ GeV}/c^2$  ( $|\Delta E| < 10 \text{ MeV}$  for  $J/\psi K_L^0$ ) that contains  $CP$  candidate events that satisfy the tagging and vertexing requirements as listed in Table 2.

For all modes except  $\eta_c K_S^0$  and  $J/\psi K_L^0$  we use simulated events to estimate the fractions of events that peak in the  $m_{\text{ES}}$  signal region due to cross-feed from other decay modes (Peaking background). For the  $\eta_c K_S^0$  mode the cross-feed fraction is determined from a fit to the  $m_{KK\pi}$  and  $m_{\text{ES}}$  distributions in data. For the  $J/\psi K_L^0$  decay mode, the sample composition, effective  $\eta_f$ , and  $\Delta E$  distribution of the individual background sources are determined either from simulation (for  $B \rightarrow J/\psi X$ ) or from the  $m_{\ell^+\ell^-}$  sidebands in data (for fake  $J/\psi \rightarrow \ell^+\ell^-$ ).

We determine  $\sin 2\beta$  with a simultaneous maximum likelihood fit to the  $\Delta t$  distribution of the tagged  $B_{CP}$  and  $B_{\text{flav}}$  samples. The  $\Delta t$  distributions of the  $B_{CP}$  sample are modeled by Eq. 1 with  $|\lambda| = 1$ . Those of the  $B_{\text{flav}}$  sample evolve according to the known frequency for flavor oscillation in  $B^0$  mesons. We assume that the observed amplitudes for the  $CP$  asymmetry in the  $B_{CP}$  sample and for flavor oscillation in the  $B_{\text{flav}}$  sample are reduced by the same factor  $1 - 2w$  due to flavor mistags. The  $\Delta t$  distributions for the signal are convolved with a resolution function common to both the




Figure 1: Distributions for  $B_{CP}$  and  $B_{\text{flav}}$  candidates satisfying the tagging and vertexing requirements: a)  $m_{\text{ES}}$  for the final states  $J/\psi K_S^0$ ,  $\psi(2S)K_S^0$ ,  $\chi_{c1}K_S^0$ , and  $\eta_c K_S^0$ , b)  $\Delta E$  for the final state  $J/\psi K_L^0$ , c)  $m_{\text{ES}}$  for  $J/\psi K^{*0}$  ( $K^{*0} \rightarrow K_S^0 \pi^0$ ), and d)  $m_{\text{ES}}$  for the  $B_{\text{flav}}$  sample. In each plot, the shaded region is the estimated background contribution.

$B_{\text{flav}}$  and  $B_{CP}$  samples, modeled by the sum of three Gaussians [12]. Backgrounds are incorporated with an empirical description of their  $\Delta t$  spectra, containing zero and non-zero lifetime components convolved with a resolution function [12] distinct from that of the signal.

There are 65 free parameters in the fit:  $\sin 2\beta$  (1), the average mistag fractions  $w$  and the differences  $\Delta w$  between  $B^0$  and  $\bar{B}^0$  mistag fractions for each tagging category (12), parameters for the signal  $\Delta t$  resolution (7), parameters for  $CP$  background time dependence (8), and the difference between  $B^0$  and  $\bar{B}^0$  reconstruction and tagging efficiencies (7); for  $B_{\text{flav}}$  background, time dependence (3),  $\Delta t$  resolution (3), and mistag fractions (24). For the  $CP$  modes (except for  $J/\psi K_L^0$ ), the apparent  $CP$  asymmetry of the non-peaking background in each tagging category is allowed to be a free parameter in the fit.

We fix  $\tau_{B^0} = 1.530$  ps,  $\Delta m_d = 0.507$  ps $^{-1}$  [9],  $|\lambda| = 1$ , and  $\Delta\Gamma_d = 0$ . The determination of the mistag fractions and  $\Delta t$  resolution function parameters for the signal is dominated by the large  $B_{\text{flav}}$  sample. We determine background parameters mainly from events outside the peaks in the

Table 2: Number of events  $N_{\text{tag}}$  in the signal region after tagging and vertexing requirements, signal purity  $P$  including the contribution from peaking background, and results of fitting for  $CP$  asymmetries in the  $B_{CP}$  sample and various subsamples. In addition, results on the  $B_{\text{flav}}$  and charged  $B$  control samples test that no artificial  $CP$  asymmetry is found where we expect no  $CP$  violation ( $\sin 2\beta = 0$ ). Errors are statistical only. The signal region is  $5.27 < m_{\text{ES}} < 5.29 \text{ GeV}/c^2$  ( $|\Delta E| < 10 \text{ MeV}$  for  $J/\psi K_L^0$ ).

Sample	$N_{\text{tag}}$	$P(\%)$	$\sin 2\beta$
Full $CP$ sample	11496	76	$0.710 \pm 0.034$
$J/\psi K_S^0, \psi(2S)K_S^0, \chi_{c1}K_S^0, \eta_c K_S^0$	6028	92	$0.713 \pm 0.038$
$J/\psi K_L^0$	4323	55	$0.716 \pm 0.080$
$J/\psi K^{*0} (K^{*0} \rightarrow K_S^0 \pi^0)$	965	68	$0.526 \pm 0.284$
1999-2002 data	3084	79	$0.755 \pm 0.067$
2003-2004 data	4850	77	$0.724 \pm 0.052$
2005-2006 data	3562	74	$0.663 \pm 0.062$
<hr/>			
$J/\psi K_S^0, \psi(2S)K_S^0, \chi_{c1}K_S^0, \eta_c K_S^0$ only ( $\eta_f = -1$ )			
$J/\psi K_S^0 (K_S^0 \rightarrow \pi^+ \pi^-)$	4076	96	$0.715 \pm 0.044$
$J/\psi K_S^0 (K_S^0 \rightarrow \pi^0 \pi^0)$	988	88	$0.581 \pm 0.105$
$\psi(2S)K_S^0 (K_S^0 \rightarrow \pi^+ \pi^-)$	622	83	$0.892 \pm 0.120$
$\chi_{c1}K_S^0$	279	89	$0.709 \pm 0.174$
$\eta_c K_S^0$	243	75	$0.717 \pm 0.229$
Lepton category	703	97	$0.754 \pm 0.068$
Kaon I category	900	93	$0.713 \pm 0.066$
Kaon II category	1437	91	$0.711 \pm 0.075$
Kaon-Pion category	1107	89	$0.635 \pm 0.117$
Pion category	1238	91	$0.587 \pm 0.175$
Other category	823	89	$0.454 \pm 0.469$
<hr/>			
$B_{\text{flav}}$ sample	112878	83	$0.016 \pm 0.011$
$B^+$ sample	27775	93	$0.008 \pm 0.017$

$m_{\text{ES}}$  and  $\Delta E$  distributions, as shown in Fig. 1.

The fit to the  $B_{CP}$  and  $B_{\text{flav}}$  samples yields

$$\sin 2\beta = 0.710 \pm 0.034(\text{stat}) \pm 0.019(\text{syst}).$$

Figure 2 shows the  $\Delta t$  distributions and asymmetries in yields between events with  $B^0$  tags and  $\bar{B}^0$  tags for the  $\eta_f = -1$  and  $\eta_f = +1$  samples as a function of  $\Delta t$ , overlaid with the projection of the likelihood fit result. We perform a separate fit with only the cleanest  $\eta_f = -1$  sample, in which we treat both  $|\lambda|$  and  $\sin 2\beta$  as free parameters. We do not use the modes  $J/\psi K^{*0}$  and  $J/\psi K_L^0$  to minimize the dependence of the results on the background parametrization. We obtain  $|\lambda| = 0.932 \pm 0.026(\text{stat}) \pm 0.017(\text{syst})$ . The correlation between the coefficients multiplying the  $\sin(\Delta m_d \Delta t)$  and  $\cos(\Delta m_d \Delta t)$  terms in Eq. 1 is  $-1.2\%$ .




Figure 2: a) Number of  $\eta_f = -1$  candidates ( $J/\psi K_S^0$ ,  $\psi(2S)K_S^0$ ,  $\chi_{c1}K_S^0$ , and  $\eta_c K_S^0$ ) in the signal region with a  $B^0$  tag ( $N_{B^0}$ ) and with a  $\bar{B}^0$  tag ( $N_{\bar{B}^0}$ ), and b) the raw asymmetry  $(N_{B^0} - N_{\bar{B}^0}) / (N_{B^0} + N_{\bar{B}^0})$ , as functions of  $\Delta t$ . Figures c) and d) are the corresponding distributions for the  $\eta_f = +1$  mode  $J/\psi K_L^0$ . All distributions exclude `Other`-tagged events. The solid (dashed) curves represent the fit projections in  $\Delta t$  for  $B^0$  ( $\bar{B}^0$ ) tags. The shaded regions represent the estimated background contributions.

## 4 SYSTEMATIC UNCERTAINTIES

The systematic uncertainties on  $\sin 2\beta$  and  $|\lambda|$  are summarized in Table 3. These include the uncertainties in the level and  $CP$  asymmetry of the peaking background, the assumed parameterization of the  $\Delta t$  resolution function, possible differences between the  $B_{\text{flav}}$  and  $B_{CP}$  tagging performances and  $\Delta t$  resolution functions, knowledge of the event-by-event beam spot position, and the possible interference between the suppressed  $\bar{b} \rightarrow \bar{u}c\bar{d}$  amplitude with the favored  $b \rightarrow c\bar{u}d$  amplitude for

some tag-side  $B$  decays [14]. In addition, we include the variation due to the assumed values of  $\Delta m_d$  and  $\tau_B$  [9]. We also assign the change in the measured  $\sin 2\beta$  as the corresponding systematic uncertainties when we let  $|\lambda|$  to be a free parameter in the fit and when we set  $\Delta\Gamma_d/\Gamma_d = \pm 0.02$ , the latter being considerably larger than SM estimates [15]. The total systematic error on  $\sin 2\beta$  ( $|\lambda|$ ) is 0.019 (0.017).

Table 3: Systematic uncertainties on  $\sin 2\beta$  and  $|\lambda|$ .

Source	$\sigma(\sin 2\beta)$	$\sigma( \lambda )$
$CP$ backgrounds	0.007	0.002
$\Delta t$ resolution function	0.008	0.002
$J/\psi K_L^0$ backgrounds	0.007	N/A
Mistag fraction differences	0.009	0.007
Beam spot	0.008	0.004
$\Delta m_d, \tau_B, \Delta\Gamma_d/\Gamma_d,  \lambda $	0.003	0.001
Tag-side interference	0.002	0.014
MC statistics	0.003	0.005
Total systematic error	0.019	0.017

The large  $B_{CP}$  sample allows a number of consistency checks, including separation of the data by decay mode and tagging category. The results of those checks are listed in Table 2. We observe no statistically significant asymmetry from fits to the control samples of non- $CP$  decay modes.

## 5 SUMMARY

In summary, we report on improved measurements of  $\sin 2\beta$  and  $|\lambda|$  that supersede our previous result [5]. We measure  $\sin 2\beta = 0.710 \pm 0.034(\text{stat}) \pm 0.019(\text{syst})$  and  $|\lambda| = 0.932 \pm 0.026(\text{stat}) \pm 0.017(\text{syst})$ . The updated value of  $\sin 2\beta$  is consistent with the current world average [16] and the theoretical estimates of the magnitudes of CKM matrix elements in the context of the SM [17]. The theoretical uncertainty on the interpretation of the measurement of  $\sin 2\beta$  in these modes is approximately 0.01 [18].

## 6 ACKNOWLEDGMENTS

We are grateful for the extraordinary contributions of our PEP-II colleagues in achieving the excellent luminosity and machine conditions that have made this work possible. The success of this project also relies critically on the expertise and dedication of the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and the kind hospitality extended to them. This work is supported by the US Department of Energy and National Science Foundation, the Natural Sciences and Engineering Research Council (Canada), Institute of High Energy Physics (China), the Commissariat à l’Energie Atomique and Institut National de Physique Nucléaire et de Physique des Particules (France), the Bundesministerium für Bildung und Forschung and Deutsche Forschungsgemeinschaft (Germany), the Istituto Nazionale di

Fisica Nucleare (Italy), the Foundation for Fundamental Research on Matter (The Netherlands), the Research Council of Norway, the Ministry of Science and Technology of the Russian Federation, and the Particle Physics and Astronomy Research Council (United Kingdom). Individuals have received support from the Marie-Curie IEF program (European Union) and the A. P. Sloan Foundation.

## References

- [1] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **89**, 201802 (2002).
- [2] BELLE Collaboration, K. Abe *et al.*, Phys. Rev. D **66**, 071102 (2002).
- [3] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963);  
M. Kobayashi and T. Maskawa, Prog. Th. Phys. **49**, 652 (1973).
- [4] A.B. Carter and A.I. Sanda, Phys. Rev. D **23**, 1567 (1981);  
I.I. Bigi and A.I. Sanda, Nucl. Phys. **193**, 85 (1981).
- [5] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **94**, 161803 (2005).
- [6] *BABAR* Collaboration, B. Aubert *et al.*, Nucl. Instr. Methods Phys. Res., Sect. A **479**, 1 (2002).
- [7] Geant4 Collaboration, S. Agostinelli *et al.*, Nucl. Instr. Methods Phys. Res., Sect. A **506**, 250 (2003).
- [8] See, for example, D. Kirkby, and Y. Nir in W.-M. Yao *et al.*, J. Phys. **G33**, 1 (2006).
- [9] W.-M. Yao *et al.*, [Particle Data Group], J. Phys. **G33**, 1 (2006).
- [10] Charge-conjugate reactions are included implicitly unless otherwise specified.
- [11] *BABAR* Collaboration, B. Aubert *et al.*, hep-ex/0607081
- [12] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. D **66**, 032003 (2002).
- [13] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. D **70**, 011101 (2004).
- [14] O. Long, M. Baak, R. N. Cahn, D. Kirkby, Phys. Rev. D **68**, 034010 (2003).
- [15] A. S. Dighe *et al.*, Nucl. Phys. **B624**, 377 (2002);  
M. Ciuchini *et al.* JHEP **0308**, 031 (2003).
- [16] E. Barberio *et al.*, hep-ex/0603003 and online update at  
<http://www.slac.stanford.edu/xorg/hfag>.
- [17] See, for example, A. Ceccucci, Z. Ligeti, and Y. Sakai, in Ref. [9].
- [18] M. Ciuchini, M. Pierini and L. Silvestrini, Phys. Rev. Lett. **95**, 221804 (2005).