

Search for the Rare Decay $B \rightarrow \pi \ell^+ \ell^-$

- B. Aubert,¹ M. Bona,¹ D. Boutigny,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Poireau,¹ X. Prudent,¹ V. Tisserand,¹
 A. Zghiche,¹ J. Garra Tico,² E. Grauges,² L. Lopez,³ A. Palano,³ G. Eigen,⁴ I. Ofte,⁴ B. Stugu,⁴ L. Sun,⁴
 G. S. Abrams,⁵ M. Battaglia,⁵ D. N. Brown,⁵ J. Button-Shafer,⁵ R. N. Cahn,⁵ Y. Groysman,⁵ R. G. Jacobsen,⁵
 J. A. Kadyk,⁵ L. T. Kerth,⁵ Yu. G. Kolomensky,⁵ G. Kukartsev,⁵ D. Lopes Pegna,⁵ G. Lynch,⁵ L. M. Mir,⁵
 T. J. Orimoto,⁵ M. Pripstein,⁵ N. A. Roe,⁵ M. T. Ronan,^{5,*} K. Tackmann,⁵ W. A. Wenzel,⁵ P. del Amo Sanchez,⁶
 C. M. Hawkes,⁶ A. T. Watson,⁶ T. Held,⁷ H. Koch,⁷ B. Lewandowski,⁷ M. Pelizaeus,⁷ T. Schroeder,⁷ M. Steinke,⁷
 J. T. Boyd,⁸ J. P. Burke,⁸ W. N. Cottingham,⁸ D. Walker,⁸ D. J. Asgeirsson,⁹ T. Cuhadar-Donszelmann,⁹
 B. G. Fulsom,⁹ C. Hearty,⁹ N. S. Knecht,⁹ T. S. Mattison,⁹ J. A. McKenna,⁹ A. Khan,¹⁰ M. Saleem,¹⁰
 L. Teodorescu,¹⁰ V. E. Blinov,¹¹ A. D. Bukin,¹¹ V. P. Druzhinin,¹¹ V. B. Golubev,¹¹ A. P. Onuchin,¹¹
 S. I. Serednyakov,¹¹ Yu. I. Skovpen,¹¹ E. P. Solodov,¹¹ K. Yu. Todyshev,¹¹ M. Bondioli,¹² M. Bruinsma,¹²
 S. Curry,¹² I. Eschrich,¹² D. Kirkby,¹² A. J. Lankford,¹² P. Lund,¹² M. Mandelkern,¹² E. C. Martin,¹² D. P. Stoker,¹²
 S. Abachi,¹³ C. Buchanan,¹³ S. D. Foulkes,¹⁴ J. W. Gary,¹⁴ F. Liu,¹⁴ O. Long,¹⁴ B. C. Shen,¹⁴ L. Zhang,¹⁴
 H. P. Paar,¹⁵ S. Rahatlou,¹⁵ V. Sharma,¹⁵ J. W. Berryhill,¹⁶ C. Campagnari,¹⁶ A. Cunha,¹⁶ B. Dahmes,¹⁶
 T. M. Hong,¹⁶ D. Kovalskyi,¹⁶ J. D. Richman,¹⁶ T. W. Beck,¹⁷ A. M. Eisner,¹⁷ C. J. Flacco,¹⁷ C. A. Heusch,¹⁷
 J. Kroeseberg,¹⁷ W. S. Lockman,¹⁷ T. Schalk,¹⁷ B. A. Schumm,¹⁷ A. Seiden,¹⁷ D. C. Williams,¹⁷ M. G. Wilson,¹⁷
 L. O. Winstrom,¹⁷ E. Chen,¹⁸ C. H. Cheng,¹⁸ A. Dvoretskii,¹⁸ F. Fang,¹⁸ D. G. Hitlin,¹⁸ I. Narsky,¹⁸ T. Piatenko,¹⁸
 F. C. Porter,¹⁸ G. Mancinelli,¹⁹ B. T. Meadows,¹⁹ K. Mishra,¹⁹ M. D. Sokoloff,¹⁹ F. Blanc,²⁰ P. C. Bloom,²⁰
 S. Chen,²⁰ W. T. Ford,²⁰ J. F. Hirschauer,²⁰ A. Kreisel,²⁰ M. Nagel,²⁰ U. Nauenberg,²⁰ A. Olivas,²⁰ J. G. Smith,²⁰
 K. A. Ulmer,²⁰ S. R. Wagner,²⁰ J. Zhang,²⁰ A. Chen,²¹ E. A. Eckhart,²¹ A. Soffer,²¹ W. H. Toki,²¹ R. J. Wilson,²¹
 F. Winklmeier,²¹ Q. Zeng,²¹ D. D. Altenburg,²² E. Feltresi,²² A. Hauke,²² H. Jasper,²² J. Merkel,²² A. Petzold,²²
 B. Spaan,²² K. Wacker,²² T. Brandt,²³ V. Klose,²³ H. M. Lacker,²³ W. F. Mader,²³ R. Nogowski,²³ J. Schubert,²³
 K. R. Schubert,²³ R. Schwierz,²³ J. E. Sundermann,²³ A. Volk,²³ D. Bernard,²⁴ G. R. Bonneau,²⁴ E. Latour,²⁴
 Ch. Thiebaux,²⁴ M. Verderi,²⁴ P. J. Clark,²⁵ W. Gradl,²⁵ F. Muheim,²⁵ S. Playfer,²⁵ A. I. Robertson,²⁵ Y. Xie,²⁵
 M. Andreotti,²⁶ D. Bettoni,²⁶ C. Bozzi,²⁶ R. Calabrese,²⁶ A. Cecchi,²⁶ G. Cibinetto,²⁶ P. Franchini,²⁶ E. Luppi,²⁶
 M. Negrini,²⁶ A. Petrella,²⁶ L. Piemontese,²⁶ E. Prencipe,²⁶ V. Santoro,²⁶ F. Anulli,²⁷ R. Baldini-Ferroli,²⁷
 A. Calcaterra,²⁷ R. de Sangro,²⁷ G. Finocchiaro,²⁷ S. Pacetti,²⁷ P. Patteri,²⁷ I. M. Peruzzi,^{27,†} M. Piccolo,²⁷
 M. Rama,²⁷ A. Zallo,²⁷ A. Buzzo,²⁸ R. Contri,²⁸ M. Lo Vetere,²⁸ M. M. Macri,²⁸ M. R. Monge,²⁸ S. Passaggio,²⁸
 C. Patrignani,²⁸ E. Robutti,²⁸ A. Santroni,²⁸ S. Tosi,²⁸ K. S. Chaisanguanthum,²⁹ M. Morii,²⁹ J. Wu,²⁹
 R. S. Dubitzky,³⁰ J. Marks,³⁰ S. Schenk,³⁰ U. Uwer,³⁰ D. J. Bard,³¹ P. D. Dauncey,³¹ R. L. Flack,³¹ J. A. Nash,³¹
 M. B. Nikolich,³¹ W. Panduro Vazquez,³¹ P. K. Behera,³² X. Chai,³² M. J. Charles,³² U. Mallik,³² N. T. Meyer,³²
 V. Ziegler,³² J. Cochran,³³ H. B. Crawley,³³ L. Dong,³³ V. Eyges,³³ W. T. Meyer,³³ S. Prell,³³ E. I. Rosenberg,³³
 A. E. Rubin,³³ A. V. Gritsan,³⁴ C. K. Lae,³⁴ A. G. Denig,³⁵ M. Fritsch,³⁵ G. Schott,³⁵ N. Arnaud,³⁶
 J. Béquilleux,³⁶ M. Davier,³⁶ G. Grosdidier,³⁶ A. Höcker,³⁶ V. Lepeltier,³⁶ F. Le Diberder,³⁶ A. M. Lutz,³⁶
 S. Pruvot,³⁶ S. Rodier,³⁶ P. Roudeau,³⁶ M. H. Schune,³⁶ J. Serrano,³⁶ V. Sordini,³⁶ A. Stocchi,³⁶ W. F. Wang,³⁶
 G. Wormser,³⁶ D. J. Lange,³⁷ D. M. Wright,³⁷ C. A. Chavez,³⁸ I. J. Forster,³⁸ J. R. Fry,³⁸ E. Gabathuler,³⁸
 R. Gamet,³⁸ D. E. Hutchcroft,³⁸ D. J. Payne,³⁸ K. C. Schofield,³⁸ C. Touramanis,³⁸ A. J. Bevan,³⁹ K. A. George,³⁹
 F. Di Lodovico,³⁹ W. Menges,³⁹ R. Sacco,³⁹ G. Cowan,⁴⁰ H. U. Flaecher,⁴⁰ D. A. Hopkins,⁴⁰ P. S. Jackson,⁴⁰
 T. R. McMahon,⁴⁰ F. Salvatore,⁴⁰ A. C. Wren,⁴⁰ D. N. Brown,⁴¹ C. L. Davis,⁴¹ J. Allison,⁴² N. R. Barlow,⁴²
 R. J. Barlow,⁴² Y. M. Chia,⁴² C. L. Edgar,⁴² G. D. Lafferty,⁴² T. J. West,⁴² J. I. Yi,⁴² J. Anderson,⁴³ C. Chen,⁴³
 A. Jawahery,⁴³ D. A. Roberts,⁴³ G. Simi,⁴³ J. M. Tuggle,⁴³ G. Blaylock,⁴⁴ C. Dallapiccola,⁴⁴ S. S. Hertzbach,⁴⁴
 X. Li,⁴⁴ T. B. Moore,⁴⁴ E. Salvati,⁴⁴ S. Saremi,⁴⁴ R. Cowan,⁴⁵ P. H. Fisher,⁴⁵ G. Sciolla,⁴⁵ S. J. Sekula,⁴⁵
 M. Spitznagel,⁴⁵ F. Taylor,⁴⁵ R. K. Yamamoto,⁴⁵ H. Kim,⁴⁶ S. E. McLachlin,⁴⁶ P. M. Patel,⁴⁶ S. H. Robertson,⁴⁶
 A. Lazzaro,⁴⁷ V. Lombardo,⁴⁷ F. Palombo,⁴⁷ J. M. Bauer,⁴⁸ L. Cremaldi,⁴⁸ V. Eschenburg,⁴⁸ R. Godang,⁴⁸
 R. Kroeger,⁴⁸ D. A. Sanders,⁴⁸ D. J. Summers,⁴⁸ H. W. Zhao,⁴⁸ S. Brunet,⁴⁹ D. Côté,⁴⁹ M. Simard,⁴⁹ P. Taras,⁴⁹

F. B. Viaud,⁴⁹ H. Nicholson,⁵⁰ G. De Nardo,⁵¹ F. Fabozzi,^{51,‡} L. Lista,⁵¹ D. Monorchio,⁵¹ C. Sciacca,⁵¹ M. A. Baak,⁵² G. Raven,⁵² H. L. Snoek,⁵² C. P. Jessop,⁵³ J. M. LoSecco,⁵³ G. Benelli,⁵⁴ L. A. Corwin,⁵⁴ K. K. Gan,⁵⁴ K. Honscheid,⁵⁴ D. Hufnagel,⁵⁴ H. Kagan,⁵⁴ R. Kass,⁵⁴ J. P. Morris,⁵⁴ A. M. Rahimi,⁵⁴ J. J. Regensburger,⁵⁴ R. Ter-Antonyan,⁵⁴ Q. K. Wong,⁵⁴ N. L. Blount,⁵⁵ J. Brau,⁵⁵ R. Frey,⁵⁵ O. Igonkina,⁵⁵ J. A. Kolb,⁵⁵ M. Lu,⁵⁵ R. Rahmat,⁵⁵ N. B. Sinev,⁵⁵ D. Strom,⁵⁵ J. Strube,⁵⁵ E. Torrence,⁵⁵ N. Gagliardi,⁵⁶ A. Gaz,⁵⁶ M. Margoni,⁵⁶ M. Morandin,⁵⁶ A. Pompili,⁵⁶ M. Posocco,⁵⁶ M. Rotondo,⁵⁶ F. Simonetto,⁵⁶ R. Stroili,⁵⁶ C. Voci,⁵⁶ E. Ben-Haim,⁵⁷ H. Briand,⁵⁷ J. Chauveau,⁵⁷ P. David,⁵⁷ L. Del Buono,⁵⁷ Ch. de la Vaissière,⁵⁷ O. Hamon,⁵⁷ B. L. Hartfiel,⁵⁷ Ph. Leruste,⁵⁷ J. Malclès,⁵⁷ J. Ocariz,⁵⁷ A. Perez,⁵⁷ L. Gladney,⁵⁸ M. Biasini,⁵⁹ R. Covarelli,⁵⁹ E. Manoni,⁵⁹ C. Angelini,⁶⁰ G. Batignani,⁶⁰ S. Bettarini,⁶⁰ G. Calderini,⁶⁰ M. Carpinelli,⁶⁰ R. Cenci,⁶⁰ F. Forti,⁶⁰ M. A. Giorgi,⁶⁰ A. Lusiani,⁶⁰ G. Marchiori,⁶⁰ M. A. Mazur,⁶⁰ M. Morganti,⁶⁰ N. Neri,⁶⁰ E. Paoloni,⁶⁰ G. Rizzo,⁶⁰ J. J. Walsh,⁶⁰ M. Haire,⁶¹ J. Biesiada,⁶² P. Elmer,⁶² Y. P. Lau,⁶² C. Lu,⁶² J. Olsen,⁶² A. J. S. Smith,⁶² A. V. Telnov,⁶² E. Baracchini,⁶³ F. Bellini,⁶³ G. Cavoto,⁶³ A. D'Orazio,⁶³ D. del Re,⁶³ E. Di Marco,⁶³ R. Faccini,⁶³ F. Ferrarotto,⁶³ F. Ferroni,⁶³ M. Gaspero,⁶³ P. D. Jackson,⁶³ L. Li Gioi,⁶³ M. A. Mazzoni,⁶³ S. Morganti,⁶³ G. Piredda,⁶³ F. Polci,⁶³ F. Renga,⁶³ C. Voena,⁶³ M. Ebert,⁶⁴ H. Schröder,⁶⁴ R. Waldi,⁶⁴ T. Adye,⁶⁵ G. Castelli,⁶⁵ B. Franek,⁶⁵ E. O. Olaiya,⁶⁵ S. Ricciardi,⁶⁵ W. Roethel,⁶⁵ F. F. Wilson,⁶⁵ R. Aleksan,⁶⁶ S. Emery,⁶⁶ M. Escalier,⁶⁶ A. Gaidot,⁶⁶ S. F. Ganzhur,⁶⁶ G. Hamel de Monchenault,⁶⁶ W. Kozanecki,⁶⁶ M. Legendre,⁶⁶ G. Vasseur,⁶⁶ Ch. Yéche,⁶⁶ M. Zito,⁶⁶ X. R. Chen,⁶⁷ H. Liu,⁶⁷ W. Park,⁶⁷ M. V. Purohit,⁶⁷ J. R. Wilson,⁶⁷ M. T. Allen,⁶⁸ D. Aston,⁶⁸ R. Bartoldus,⁶⁸ P. Bechtle,⁶⁸ N. Berger,⁶⁸ R. Claus,⁶⁸ J. P. Coleman,⁶⁸ M. R. Convery,⁶⁸ J. C. Dingfelder,⁶⁸ J. Dorfan,⁶⁸ G. P. Dubois-Felsmann,⁶⁸ D. Dujmic,⁶⁸ W. Dunwoodie,⁶⁸ R. C. Field,⁶⁸ T. Glanzman,⁶⁸ S. J. Gowdy,⁶⁸ M. T. Graham,⁶⁸ P. Grenier,⁶⁸ V. Halyo,⁶⁸ C. Hast,⁶⁸ T. Hryna'ova,⁶⁸ W. R. Innes,⁶⁸ M. H. Kelsey,⁶⁸ P. Kim,⁶⁸ D. W. G. S. Leith,⁶⁸ S. Li,⁶⁸ S. Luitz,⁶⁸ V. Luth,⁶⁸ H. L. Lynch,⁶⁸ D. B. MacFarlane,⁶⁸ H. Marsiske,⁶⁸ R. Messner,⁶⁸ D. R. Muller,⁶⁸ C. P. O'Grady,⁶⁸ V. E. Ozcan,⁶⁸ A. Perazzo,⁶⁸ M. Perl,⁶⁸ T. Pulliam,⁶⁸ B. N. Ratcliff,⁶⁸ A. Roodman,⁶⁸ A. A. Salnikov,⁶⁸ R. H. Schindler,⁶⁸ J. Schwiening,⁶⁸ A. Snyder,⁶⁸ J. Stelzer,⁶⁸ D. Su,⁶⁸ M. K. Sullivan,⁶⁸ K. Suzuki,⁶⁸ S. K. Swain,⁶⁸ J. M. Thompson,⁶⁸ J. Va'vra,⁶⁸ N. van Bakel,⁶⁸ A. P. Wagner,⁶⁸ M. Weaver,⁶⁸ W. J. Wisniewski,⁶⁸ M. Wittgen,⁶⁸ D. H. Wright,⁶⁸ A. K. Yarritu,⁶⁸ K. Yi,⁶⁸ C. C. Young,⁶⁸ P. R. Burchat,⁶⁹ A. J. Edwards,⁶⁹ S. A. Majewski,⁶⁹ B. A. Petersen,⁶⁹ L. Wilden,⁶⁹ S. Ahmed,⁷⁰ M. S. Alam,⁷⁰ R. Bula,⁷⁰ J. A. Ernst,⁷⁰ V. Jain,⁷⁰ B. Pan,⁷⁰ M. A. Saeed,⁷⁰ F. R. Wappler,⁷⁰ S. B. Zain,⁷⁰ W. Bugg,⁷¹ M. Krishnamurthy,⁷¹ S. M. Spanier,⁷¹ R. Eckmann,⁷² J. L. Ritchie,⁷² A. M. Ruland,⁷² C. J. Schilling,⁷² R. F. Schwitters,⁷² J. M. Izen,⁷³ X. C. Lou,⁷³ S. Ye,⁷³ F. Bianchi,⁷⁴ F. Gallo,⁷⁴ D. Gamba,⁷⁴ M. Pelliccioni,⁷⁴ M. Bomben,⁷⁵ L. Bosisio,⁷⁵ C. Cartaro,⁷⁵ F. Cossutti,⁷⁵ G. Della Ricca,⁷⁵ L. Lanceri,⁷⁵ L. Vitale,⁷⁵ V. Azzolini,⁷⁶ N. Lopez-March,⁷⁶ F. Martinez-Vidal,⁷⁶ D. A. Milanes,⁷⁶ A. Oyanguren,⁷⁶ J. Albert,⁷⁷ Sw. Banerjee,⁷⁷ B. Bhuyan,⁷⁷ K. Hamano,⁷⁷ R. Kowalewski,⁷⁷ I. M. Nugent,⁷⁷ J. M. Roney,⁷⁷ R. J. Sobie,⁷⁷ J. J. Back,⁷⁸ P. F. Harrison,⁷⁸ T. E. Latham,⁷⁸ G. B. Mohanty,⁷⁸ M. Pappagallo,^{78,§} H. R. Band,⁷⁹ X. Chen,⁷⁹ S. Dasu,⁷⁹ K. T. Flood,⁷⁹ J. J. Hollar,⁷⁹ P. E. Kutter,⁷⁹ Y. Pan,⁷⁹ M. Pierini,⁷⁹ R. Prepost,⁷⁹ S. L. Wu,⁷⁹ Z. Yu,⁷⁹ and H. Neal⁸⁰

(The BABAR Collaboration)

¹Laboratoire de Physique des Particules, IN2P3/CNRS et Université de Savoie, F-74941 Annecy-Le-Vieux, France

²Universitat de Barcelona, Facultat de Fisica, Departament ECM, E-08028 Barcelona, Spain

³Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

⁴University of Bergen, Institute of Physics, N-5007 Bergen, Norway

⁵Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

⁶University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁷Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁸University of Bristol, Bristol BS8 1TL, United Kingdom

⁹University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

¹⁰Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹¹Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹²University of California at Irvine, Irvine, California 92697, USA

¹³University of California at Los Angeles, Los Angeles, California 90024, USA

¹⁴University of California at Riverside, Riverside, California 92521, USA

¹⁵University of California at San Diego, La Jolla, California 92093, USA

¹⁶University of California at Santa Barbara, Santa Barbara, California 93106, USA

¹⁷University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

¹⁸California Institute of Technology, Pasadena, California 91125, USA

¹⁹University of Cincinnati, Cincinnati, Ohio 45221, USA

²⁰University of Colorado, Boulder, Colorado 80309, USA

- ²¹Colorado State University, Fort Collins, Colorado 80523, USA
²²Universität Dortmund, Institut für Physik, D-44221 Dortmund, Germany
²³Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany
²⁴Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, F-91128 Palaiseau, France
²⁵University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom
²⁶Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy
²⁷Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy
²⁸Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy
²⁹Harvard University, Cambridge, Massachusetts 02138, USA
³⁰Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany
³¹Imperial College London, London, SW7 2AZ, United Kingdom
³²University of Iowa, Iowa City, Iowa 52242, USA
³³Iowa State University, Ames, Iowa 50011-3160, USA
³⁴Johns Hopkins University, Baltimore, Maryland 21218, USA
³⁵Universität Karlsruhe, Institut für Experimentelle Kernphysik, D-76021 Karlsruhe, Germany
³⁶Laboratoire de l'Accélérateur Linéaire, IN2P3/CNRS et Université Paris-Sud 11,
Centre Scientifique d'Orsay, B. P. 34, F-91898 ORSAY Cedex, France
³⁷Lawrence Livermore National Laboratory, Livermore, California 94550, USA
³⁸University of Liverpool, Liverpool L69 7ZE, United Kingdom
³⁹Queen Mary, University of London, E1 4NS, United Kingdom
⁴⁰University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom
⁴¹University of Louisville, Louisville, Kentucky 40292, USA
⁴²University of Manchester, Manchester M13 9PL, United Kingdom
⁴³University of Maryland, College Park, Maryland 20742, USA
⁴⁴University of Massachusetts, Amherst, Massachusetts 01003, USA
⁴⁵Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA
⁴⁶McGill University, Montréal, Québec, Canada H3A 2T8
⁴⁷Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy
⁴⁸University of Mississippi, University, Mississippi 38677, USA
⁴⁹Université de Montréal, Physique des Particules, Montréal, Québec, Canada H3C 3J7
⁵⁰Mount Holyoke College, South Hadley, Massachusetts 01075, USA
⁵¹Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy
⁵²NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands
⁵³University of Notre Dame, Notre Dame, Indiana 46556, USA
⁵⁴Ohio State University, Columbus, Ohio 43210, USA
⁵⁵University of Oregon, Eugene, Oregon 97403, USA
⁵⁶Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy
⁵⁷Laboratoire de Physique Nucléaire et de Hautes Energies,
IN2P3/CNRS, Université Pierre et Marie Curie-Paris6,
Université Denis Diderot-Paris7, F-75252 Paris, France
⁵⁸University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
⁵⁹Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy
⁶⁰Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy
⁶¹Prairie View A&M University, Prairie View, Texas 77446, USA
⁶²Princeton University, Princeton, New Jersey 08544, USA
⁶³Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy
⁶⁴Universität Rostock, D-18051 Rostock, Germany
⁶⁵Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom
⁶⁶DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France
⁶⁷University of South Carolina, Columbia, South Carolina 29208, USA
⁶⁸Stanford Linear Accelerator Center, Stanford, California 94309, USA
⁶⁹Stanford University, Stanford, California 94305-4060, USA
⁷⁰State University of New York, Albany, New York 12222, USA
⁷¹University of Tennessee, Knoxville, Tennessee 37996, USA
⁷²University of Texas at Austin, Austin, Texas 78712, USA
⁷³University of Texas at Dallas, Richardson, Texas 75083, USA
⁷⁴Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy
⁷⁵Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy
⁷⁶IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain
⁷⁷University of Victoria, Victoria, British Columbia, Canada V8W 3P6
⁷⁸Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom
⁷⁹University of Wisconsin, Madison, Wisconsin 53706, USA
⁸⁰Yale University, New Haven, Connecticut 06511, USA

(Dated: August 6, 2007)

We have performed a search for the flavor-changing neutral-current decays $B \rightarrow \pi\ell^+\ell^-$, where $\ell^+\ell^-$ is either e^+e^- or $\mu^+\mu^-$, using a sample of $230 \times 10^6 \Upsilon(4S) \rightarrow B\bar{B}$ decays collected with the *BABAR* detector. We observe no evidence of a signal and measure the upper limit on the isospin-averaged branching fraction to be $\mathcal{B}(B \rightarrow \pi\ell^+\ell^-) < 9.1 \times 10^{-8}$ at 90% confidence level. We also search for the lepton-flavor-violating decays $B \rightarrow \pi e^\pm \mu^\mp$ and measure an upper limit on the isospin-averaged branching fraction of $\mathcal{B}(B \rightarrow \pi e^\pm \mu^\mp) < 9.2 \times 10^{-8}$ at 90% confidence level.

PACS numbers: 13.20 He

In the Standard Model (SM), the decays $B \rightarrow \pi\ell^+\ell^-$, where $\ell^+\ell^-$ is either e^+e^- or $\mu^+\mu^-$, proceed through $b \rightarrow d\ell^+\ell^-$ flavor-changing neutral-current processes (FCNC) that do not occur at tree level. Three amplitudes contribute at leading order: a photon penguin, a Z penguin, and a W^+W^- box diagram. With highly suppressed SM rates, predicted to be $(3.3 \pm 1.0) \times 10^{-8}$ [1], these decays provide a promising means to search for effects of new flavor-changing interactions. Such effects are predicted in a wide variety of models, usually in the context of $b \rightarrow s\ell^+\ell^-$ decays [2–4]. The $b \rightarrow d\ell^+\ell^-$ decay involves quark-flavor transitions different from $b \rightarrow s\ell^+\ell^-$ and thus its measurement constitutes an independent test for new flavor-changing interactions. An experimentally similar but otherwise unrelated process, the lepton-flavor-violating (LFV) decay $B \rightarrow \pi e^\pm \mu^\mp$, is forbidden in the SM, but can occur in some models beyond the SM, such as theories involving leptoquarks [4]. Earlier searches by other experiments [5] have reached branching fraction upper limits at the 10^{-3} level for the FCNC decay and the 10^{-6} level for the LFV decays.

In this Letter we report the findings of a search for the decays $B \rightarrow \pi\ell^+\ell^-$ and $B \rightarrow \pi e^\pm \mu^\mp$ in 208.9 fb^{-1} of data recorded at the $\Upsilon(4S)$ resonance, corresponding to $(230.1 \pm 2.5) \times 10^6 B\bar{B}$ decays. The data were collected with the *BABAR* detector [6] at the PEP-II storage ring located at the Stanford Linear Accelerator Center. The event selection criteria are optimized using simulated data and data samples independent of those selected as signal. The signal model used for efficiency evaluation of the $\ell^+\ell^-$ modes uses form-factors from [7] and amplitudes from [3]. Calculations of the same type have previously been shown by [8] to describe the kinematic distributions of $B \rightarrow K\ell^+\ell^-$ well. The efficiency of the $e\mu$ event selection is estimated using a 3-body phase space model with QED photon radiative corrections.

We reconstruct signal events by combining two oppositely charged leptons (e^+e^- , $\mu^+\mu^-$ or $e^\pm\mu^\mp$) with a pion (π^\pm or π^0). Electron (muon) candidates are required to have a momentum larger than 0.3 (0.7) GeV/c . We suppress backgrounds due to photon conversions in the $B \rightarrow \pi e^+e^-$ channels by removing e^+e^- pairs with invariant mass less than $30 \text{ MeV}/c^2$. Bremsstrahlung photons from electrons are recovered if the photon has an energy of $E > 30 \text{ MeV}$ and a direction within a small angular region around the initial electron momentum vector. The identification of electrons (muons) is about 92%

(68%) efficient on average with a hadron misidentification rate of less than 1% (4%). Charged pion identification is more than 85% efficient and has a kaon misidentification rate of less than 5%. Neutral pions are identified as pairs of photons, each having an energy of at least 50 MeV. The invariant mass of the pair is required to satisfy $115 < m_{\gamma\gamma} < 150 \text{ MeV}/c^2$.

Correctly reconstructed B decays produce narrow peaks in the distributions of two kinematic variables: the beam-energy substituted mass, $m_{\text{ES}} = \sqrt{E_b^{*2} - |\mathbf{p}_B^*|^2}$, and $\Delta E = E_B^* - E_b^*$. Here, E_b^* is the beam energy and E_B^* (\mathbf{p}_B^*) is the energy (momentum) of the reconstructed B meson, evaluated in the center-of-mass (c.m.) frame. For signal events the m_{ES} distribution is centered at the B -meson mass and the ΔE distribution is centered at zero. The mean and width of these distributions are determined from smearing and shifting the values from simulated signal events according to studies of $B^+ \rightarrow J/\psi K^+$ and $B^0 \rightarrow J/\psi \pi^0$ events in data control samples and simulations. We find the width of m_{ES} to be 2.5 (1.8) MeV/c^2 for the π^\pm (π^0) modes and widths of ΔE to be 23 , 50 , 20 and 39 MeV for the $\pi^\pm e^+e^-$, $\pi^0 e^+e^-$, $\pi^\pm \mu^+\mu^-$ and $\pi^0 \mu^+\mu^-$ final states, respectively. For events reconstructed as $e^\pm\mu^\mp$, we assume the same mean and width as for the corresponding e^+e^- modes.

The primary sources of background are random combinations of particles from $e^+e^- \rightarrow q\bar{q}$ ($q = u, d, s, c$) and from $\Upsilon(4S) \rightarrow B\bar{B}$ decays. These combinatorial backgrounds typically arise from pairs of semileptonic decays of B and $D^{(*)}$ mesons. Additionally, there is background from events that are peaking in m_{ES} and ΔE as they have the same topology as signal events. These events include $B \rightarrow J/\psi \pi$ ($J/\psi \rightarrow \ell^+\ell^-$), $B^\pm \rightarrow J/\psi K^\pm$ or $B^\pm \rightarrow K^\pm \ell^+\ell^-$ (with K^\pm misidentified as π^\pm), and $B \rightarrow \pi hh$ (with two hadrons $h = K^\pm, \pi^\pm$ misidentified as muons).

Contributions from $e^+e^- \rightarrow q\bar{q}$ processes are reduced by exploiting the difference between the spherical track distribution in $B\bar{B}$ events and the jetlike structure of $e^+e^- \rightarrow q\bar{q}$ events. We consider events for which the ratio of second to zeroth Fox-Wolfram moments R_2 is less than 0.5. Further suppression by a factor of ~ 45 is obtained by constructing a Fisher discriminant from the following four quantities [9] defined in the center-of-mass frame: R_2 , $|\cos \theta_{thr}|$ where θ_{thr} is the angle between the thrust axis of the signal particles and that of the remaining particles in the event, $|\cos \theta_B|$ where θ_B is the

angle of the B candidate's momentum vector with respect to the beam axis, and the ratio of second- to zeroth-order Legendre moments [10].

Combinatorial background from $B\bar{B}$ events is reduced by a factor of ~ 3 by using a likelihood ratio composed of [9]: the missing energy of the event (computed from all charged tracks and neutral energy clusters), the vertex fit probability of all tracks from the B candidate, the vertex fit probability of the two leptons, and $\cos\theta_B$. Missing energy provides the strongest suppression of these events, which typically contain energetic neutrinos from at least two semileptonic B or $D^{(*)}$ meson decays.

We veto events that have a dilepton invariant mass consistent with the J/ψ resonance ($2.90 < m_{e^+e^-} < 3.20 \text{ GeV}/c^2$ and $3.00 < m_{\mu^+\mu^-} < 3.20 \text{ GeV}/c^2$) or with the $\psi(2S)$ resonance ($3.60 < m_{\ell^+\ell^-} < 3.75 \text{ GeV}/c^2$). For electron modes, the vetoes are applied to $m_{\ell^+\ell^-}$ computed both with and without bremsstrahlung recovery. When a lepton radiates or is mismeasured, $m_{\ell^+\ell^-}$ may shift to values below the charmonium mass, with ΔE shifting downward accordingly. Therefore, we veto events that lie in linearly dependent ΔE - $m_{\ell\ell}$ bands, whose widths are determined from simulation, similar to the technique applied in [8]. For $e^\pm\mu^\mp$ modes, we use the same vetoes as for the e^+e^- modes. In modes with muons, in order to veto events with tracks that are consistent with hadronic decays $D \rightarrow K\pi$ or $D \rightarrow \pi\pi$, we require $m_{\ell\ell}$ and $m_{\pi\ell}$ to lie outside the range $1.84 - 1.89 \text{ GeV}/c^2$ when the ℓ is assigned the mass of a π or K . For the π^0 modes, the range for $m_{\pi\ell}$ is increased to $1.79 - 1.94 \text{ GeV}/c^2$.

The events removed by the charmonium vetoes are kinematically similar to signal events and serve as large control samples for studying signal shapes, selection efficiencies, and systematic errors. The branching fraction of $B \rightarrow J/\psi\pi$ is also extracted from the control sample and found to be in agreement with the current world average [11]. We also select a control sample of $B^+ \rightarrow J/\psi K^+$ events to measure the efficiencies and systematic uncertainties of lepton identification and the Fisher and likelihood selection.

We extract the signal yield by counting events within a signal region defined as $\pm 2\sigma$ around the mean values of the m_{ES} and ΔE distributions expected for signal events, and comparing observed event counts with estimations of the remaining background in the same region, summarized in Table I.

To determine the peaking background from hadronic $B \rightarrow \pi\pi$ or $B \rightarrow K\pi\pi$ events, we select a control sample where one track is required to pass hadron identification in place of muon identification. This selects hadronic B -decays where the remaining track that passes the muon selection is a mis-identified hadron. Each event is further weighted by the probability that one more hadron is mis-identified as a muon. The expected contribution to the $B \rightarrow \pi\ell^+\ell^-$ signal region is extracted from a one-

dimensional fit to the distribution of m_{ES} for events that pass the ΔE signal selection.

Backgrounds from B decays to final states with real leptons are estimated from high-statistics samples of simulated events. $B \rightarrow K\ell^+\ell^-$ events are the largest peaking background component for the $\pi^+e^+e^-$ mode, but are shifted towards lower ΔE than signal and fall outside the signal region. $B \rightarrow \rho\ell^+\ell^-$ events contribute even less, since the reconstructed B mesons are missing a pion. Background from charmonium resonances are found to be negligible.

The expected number of combinatorial background events is extracted from a two-dimensional, unbinned maximum-likelihood fit to m_{ES} and ΔE in a sideband defined by $5.2 < m_{\text{ES}} < 5.2724 \text{ GeV}/c^2$ and $|\Delta E| < 0.25 \text{ GeV}$, i.e., below the m_{ES} value expected for signal B events. The signal-region yield is obtained from extrapolation of this fit into the signal region. This procedure has been validated by studies of simulated background events and data events in the $e\mu$ channel where no signal-like events are expected. The background probability distribution function (PDF) is modeled as the product of an ARGUS function [12] for m_{ES} and an exponential function for ΔE . The slopes and normalization are floating in the fit. Average biases in the background central value and its uncertainty were corrected for, based on a study of a large ensemble of simulated experiments generated from the background PDF obtained from data. The corrections amount to 35% in the low-statistics $B^0 \rightarrow \pi^0\mu^+\mu^-$ channel and $< 10\%$ in all others.

Systematic uncertainties due to the background estimates are summarized in Table I. The uncertainty in the combinatorial-background estimate is determined by varying the fit parameters by $\pm 1\sigma$ of the best fit. We also consider the effect of using alternative PDF parameterizations on the background estimates, and use the computed differences to bound the systematic uncertainty. Alternatives considered include a PDF that is correlated in m_{ES} and ΔE via a linear ΔE dependence in the m_{ES} slope parameter, and PDFs for which the ΔE shape is a linear or quadratic polynomial. For peaking background with real leptons the uncertainty is dominated by limited knowledge [11] of the branching fractions for these processes, and for hadronic B peaking background the uncertainty is dominated by the control sample statistics from which it is derived.

Systematic uncertainties due to the signal efficiency include: charged-particle tracking efficiency (0.8% per lepton, 1.4% per charged hadron) and identification (0.7% per electron pair, 1.9% per muon pair, 0.5% per pion), neutral pion efficiency (3%), the Fisher and likelihood selection (1.4% for all modes involving electrons, 1.7% for $B^+ \rightarrow \pi^+\mu^+\mu^-$ and 1.9% for $B^0 \rightarrow \pi^0\mu^+\mu^-$), and signal simulation statistics (0.1%). A systematic uncertainty in signal-region selection efficiency arises from the uncertainty in the mean and width of the m_{ES} and

TABLE I: Number of background events with associated systematic uncertainties expected in the signal region.

	$\pi^+ e^+ e^-$	$\pi^0 e^+ e^-$	$\pi^+ \mu^+ \mu^-$	$\pi^0 \mu^+ \mu^-$	$\pi^+ e\mu$	$\pi^0 e\mu$
$m_{ES}-\Delta E$ fit	0.84 ± 0.24	0.43 ± 0.23	0.90 ± 0.25	0.23 ± 0.20	1.55 ± 0.34	1.22 ± 0.43
$m_{ES}-\Delta E$ correlations	± 0.02	± 0.03	± 0.06	± 0.03	± 0.17	± 0.05
ΔE shape	± 0.02	± 0.01	± 0.12	± 0.02	± 0.31	± 0.24
Peaking ($\ell^+\ell^-$)	0.057 ± 0.016	0.009 ± 0.003	0.032 ± 0.008	0.005 ± 0.001	0.0 ± 0.001	0.0 ± 0.001
Peaking (hadronic)	< 0.001	< 0.001	0.027 ± 0.033	0.035 ± 0.022	0.0 ± 0.02	0.0 ± 0.02
Total	0.90 ± 0.24	0.44 ± 0.23	0.96 ± 0.29	0.27 ± 0.20	1.55 ± 0.49	1.22 ± 0.50

ΔE distributions determined from charmonium control samples. This contributes a total uncertainty of 0.7% for charged modes for which a high-statistics sample of $B^+ \rightarrow J/\psi K^+$ events is used, and a total of 7% uncertainty for neutral modes for which a small statistics sample of $B^0 \rightarrow J/\psi \pi^0$ events is used. For the electron modes, we vary the amount of the bremsstrahlung tail in the ΔE distribution, introducing a systematic uncertainty of 1-1.4%. The number of $B\bar{B}$ events in the data sample is known to a precision of 1.1%. Additional systematic uncertainties for the efficiency result from the choice of the form factor model and the relative magnitudes of the $b \rightarrow d\ell^+\ell^-$ amplitudes, which affect the distribution of four-momentum transfer $q^2 = m_{\ell^+\ell^-}^2$ of the signal. We evaluate these systematics from the spread in efficiencies when using alternative form-factor models [13], and when varying the Wilson coefficients in the amplitudes by a factor of ± 2 . The former uncertainty varies from 1.1% for $B^+ \rightarrow \pi^+ e^+ e^-$ to 7.3% for $B^0 \rightarrow \pi^0 \mu^+ \mu^-$; the latter uncertainty varies from 0.3% for $B^0 \rightarrow \pi^0 \mu^+ \mu^-$ to 1.2% for $B^+ \rightarrow \pi^+ e^+ e^-$. For the $e\mu$ modes we use the spread in efficiency when applying two alternative theoretical models for these decays, which amounts to 17% (19%) for the $\pi^{\pm(0)}$ mode. The total systematic uncertainty of the signal efficiencies are 4% (9%), 6% (11%) and 17% (21%) for $\pi^{\pm(0)} ee$, $\pi^{\pm(0)} \mu^+ \mu^-$ and $\pi^{\pm(0)} e\mu$ modes, respectively.


Figure 1 shows the distribution of events from data in the $m_{ES}-\Delta E$ plane. The rectangles in the plots indicate the signal regions. Three $B \rightarrow \pi\ell^+\ell^-$ candidates and one $B \rightarrow \pi e^\pm \mu^\mp$ candidate are observed in the signal regions, which is consistent with the expected background. In Table II we calculate the branching fraction upper limits at 90% confidence level (C.L.) using a frequentist method that takes systematic uncertainties and their correlations into account. We follow the algorithm of [14], but differ from it in that we assume Gaussian distribution truncated at zero for the systematic uncertainties in signal sensitivity and background expectation. We combine modes and determine the $e\mu$ -averaged branching fractions to be $\mathcal{B}(B^+ \rightarrow \pi^+ \ell^+\ell^-) < 1.2 \times 10^{-7}$ and $\mathcal{B}(B^0 \rightarrow \pi^0 \ell^+\ell^-) < 1.2 \times 10^{-7}$ at 90% C.L., where charged conjugate modes are implied. Defining the isospin averaged branching fraction $\mathcal{B}(B \rightarrow \pi\ell^+\ell^-) \equiv$

$\mathcal{B}(B^+ \rightarrow \pi^+ \ell^+\ell^-) = 2 \times \frac{\tau_{B^+}}{\tau_{B^0}} \mathcal{B}(B^0 \rightarrow \pi^0 \ell^+\ell^-)$, where the different B -meson lifetimes τ_B [11] are taken into account, we find the combined upper limit

$$\mathcal{B}(B \rightarrow \pi\ell^+\ell^-) < 9.1 \times 10^{-8} \text{ at 90\% C.L.}$$

This is about a factor three above the nominal SM prediction [1]. We similarly compute the combined limit for the $e\mu$ modes of

$$\mathcal{B}(B \rightarrow \pi e^\pm \mu^\mp) < 9.2 \times 10^{-8} \text{ at 90\% C.L.}$$

FIG. 1: $m_{ES}-\Delta E$ distributions for events selected in each mode. The rectangles indicate the signal regions.

In conclusion, we have presented the result of a search for $B \rightarrow \pi\ell^+\ell^-$ using a sample of $(230.1 \pm 2.5) \times 10^6$ $B\bar{B}$ pairs produced at the $\Upsilon(4S)$ resonance. No excess of events is observed in the signal regions, and at 90% confidence limit we measure the upper limit of $\mathcal{B}(B \rightarrow \pi\ell^+\ell^-) < 9.1 \times 10^{-8}$, which is within a factor three of SM expectations. We also measure the upper limit of the lepton-flavor-violating branching fractions to be $\mathcal{B}(B \rightarrow \pi e^\pm \mu^\mp) < 9.2 \times 10^{-8}$.

TABLE II: The observed yields, number of expected background events, signal efficiency, and branching fraction (\mathcal{B}) upper limit (U.L.) at 90% C.L. in units of 10^{-7} . The upper limits for combined modes are also given.

Mode	Observed events	Expected background	Signal efficiency	\mathcal{B} U.L. 90% C.L.
$B^+ \rightarrow \pi^+ e^+ e^-$	1	0.90 ± 0.24	(7.1 ± 0.3)%	1.8
$B^0 \rightarrow \pi^0 e^+ e^-$	0	0.44 ± 0.23	(5.7 ± 0.5)%	1.4
$B^+ \rightarrow \pi^+ \mu^+ \mu^-$	1	0.96 ± 0.29	(4.7 ± 0.3)%	2.8
$B^0 \rightarrow \pi^0 \mu^+ \mu^-$	1	0.27 ± 0.20	(3.1 ± 0.3)%	5.1
$B^+ \rightarrow \pi^+ e^\pm \mu^\mp$	1	1.55 ± 0.49	(6.3 ± 1.1)%	1.7
$B^0 \rightarrow \pi^0 e^\pm \mu^\mp$	0	1.22 ± 0.50	(3.7 ± 0.8)%	1.4
$B^+ \rightarrow \pi^+ \ell^+ \ell^-$				1.2
$B^0 \rightarrow \pi^0 \ell^+ \ell^-$				1.2
$B \rightarrow \pi \ell^+ \ell^-$				0.91
$B \rightarrow \pi e^\pm \mu^\mp$				0.92

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), MEC (Spain), and STFC (United Kingdom). Individuals have received support from the Marie Curie EIF (European Union) and the A. P. Sloan Foundation.

Perugia, Italy

[‡] Also with Università della Basilicata, Potenza, Italy

[§] Also with IPPP, Physics Department, Durham University, Durham DH1 3LE, United Kingdom

- [1] T. M. Aliev and M. Savci, Phys. Rev. D **60**, 014005 (1999).
- [2] G. Burdman, Phys. Rev. D **52**, 6400 (1995); J. L. Hewett and J. D. Wells, Phys. Rev. D **55**, 5549 (1997); G. Eilam, J. L. Hewett, and T. G. Rizzo, Phys. Rev. D **34**, 2773 (1986); T. M. Aliev, A. Ozpineci, and M. Savci, Eur. Phys. J. C **29**, 265 (2003).
- [3] A. Ali, E. Lunghi, C. Greub, and G. Hiller, Phys. Rev. D **66**, 034002 (2002).
- [4] S. Davidson, D. C. Bailey, and B. A. Campbell, Z. Phys. C **61**, 613 (1994).
- [5] A. J. Weir *et al.* (Mark II Collaboration), Phys. Rev. D **41**, 1384 (1990); K. W. Edwards *et al.* (CLEO Collaboration), Phys. Rev. D **65**, 111102 (2002).
- [6] B. Aubert *et al.* (*BABAR* Collaboration), Nucl. Instrum. Methods A **479**, 1 (2002).
- [7] P. Ball and R. Zwicky, Phys. Rev. D **71**, 014015 (2005).
- [8] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. D **73**, 092001 (2006).
- [9] Distributions can be seen in I. Ofte, Ph.D thesis, University of Bergen, Bergen, Norway, 2006, ISBN 82-308-0272-6
- [10] The i^{th} moment L_i is defined by $L_i = \sum_j |p_j^*| |\cos(\theta_j^*)|^i$, where the p_j^* are the c.m. momenta of all particles not used in reconstructing the signal B candidate, and the angle θ_j^* is between the particle's momentum and the thrust axis of the signal B .
- [11] W.-M. Yao *et al.* (Particle Data Group), J. Phys. G **33**, 1 (2006).
- [12] The function is $f(x) \propto x\sqrt{1-x^2} \exp[-\zeta(1-x^2)]$, where the slope ζ is a fit parameter and $x = m_{\text{ES}}/E_b^*$; H. Albrecht *et al.* (ARGUS Collaboration), Z. Phys. C **48**, 543 (1990).
- [13] D. Melikhov and N. Nikitin, hep-ph/9609503; D. Melikhov, Phys. Lett. B **380**, 363 (1996); D. Melikhov, Phys. Rev. D **53**, 2460 (1996); D. Melikhov and B. Stech, Phys. Rev. D **62**, 014006 (2000).
- [14] R. Barlow, Comput. Phys. Commun. **149**, 97 (2002).

^{*} Deceased

[†] Also with Università di Perugia, Dipartimento di Fisica,