

# The Determination of the Helicity of $W'$ Boson Couplings at the LHC \* †

Thomas G. Rizzo<sup>a</sup>

*Stanford Linear Accelerator Center, 2575 Sand Hill Rd., Menlo Park, CA, 94025*

## Abstract

Apart from its mass and width, the most important property of a new charged gauge boson,  $W'$ , is the helicity of its couplings to the SM fermions. Such particles are expected to exist in many extensions of the Standard Model. In this paper we explore the capability of the LHC to determine the  $W'$  coupling helicity at low integrated luminosities in the  $\ell + E_T^{miss}$  discovery channel. We find that measurements of the transverse mass distribution, reconstructed from this final state in the  $W - W'$  interference region, provides the best determination of this quantity. To make such measurements requires integrated luminosities of  $\sim 10(60) \text{ fb}^{-1}$  assuming  $M_{W'} = 1.5(2.5) \text{ TeV}$  and provided that the  $W'$  couplings have Standard Model magnitude. This helicity determination can be further strengthened by the use of various discovery channel leptonic asymmetries, also measured in the same interference regime, but with higher integrated luminosities.

---

\*Work supported in part by the Department of Energy, Contract DE-AC02-76SF00515

†e-mail: <sup>a</sup>rizzo@slac.stanford.edu

# 1 Introduction

The ATLAS and CMS experiments at the LHC will begin taking data in a few months and it is widely believed that new physics beyond the Standard Model(SM) will be discovered in the coming years. There are many expectations as to what this new physics may be and in what form it will manifest itself, but it is likely that we will be in for a surprise. Once this new physics is discovered our primary goal will be to understand its essential nature and how the specific discoveries, such as the production and observed properties of new particles, fit into a broader theoretical framework.

The existence of a new charged gauge boson,  $W'$ , or a  $W'$ -like object, is now a relatively common prediction which results from many new physics scenarios. These possibilities include the Little Higgs(LH) model[1], the Randall-Sundrum(RS)[2] model with bulk gauge fields[3], Universal Extra Dimensions(UED)[4], TeV scale extra dimensions[5, 6, 7], as well as many different extended electroweak gauge models, such as the prototypical Left-Right Symmetric Model(LRM)[8, 9]. Although the physics of a new  $Z'$  has gotten much attention in the literature[10], the detailed study of a possible  $W'$  has fared somewhat less well[11]. Perhaps the most important property of a  $W'$ , apart from its mass and width, is the helicity of its couplings to the fermions in the SM. For all of the models discussed in the literature above, these couplings are either purely left- or right-handed, apart from some possible small mixing effects. Determining the helicity of the couplings of a newly discovered  $W'$  is thus the first major step in opening up the underlying physics as it is an order one discriminator between different classes of models.<sup>‡</sup>

As will be discussed below, there have been many suggestions over the last 20-plus years as to how to measure the helicity of  $W'$  couplings, all of which have their own strengths and weaknesses. These analyses have generally relied upon the use of the narrow width approximation. However, in employing this approximation much valuable information about the properties of the  $W'$  can be lost, in particular, that obtained from  $W - W'$  interference. The goal of this paper will

---

<sup>‡</sup>This is similar in nature to determining whether the known light neutrinos are Dirac or Majorana particles.

be to explore the effects of this interference on the transverse mass dependent distributions of the  $W'$ . As we will see the rather straightforward measurement of the transverse mass distribution itself will allow us obtain the necessary  $W'$  helicity information. Furthermore, we will demonstrate that such measurements will require only relatively low integrated luminosities for  $W'$  masses which are not too large, and will employ the traditional  $\ell + E_T^{miss}$   $W'$  discovery channel.

Section II of the paper contains some background material and a historically-oriented overview of previous ideas that have been suggested to address the  $W'$  helicity issue including a discussion of their various strengths and weaknesses. Section III will present an analysis of the  $W'$  transverse mass distribution and its helicity dependence for a range of  $W'$  masses, coupling strengths and LHC integrated luminosities. The use of various asymmetries evaluated in the  $W - W'$  interference region in order to assist with the  $W'$  helicity determination will also be discussed. Section IV contains a final summary and discussion of our results.

## 2 Background and History

Let us begin by establishing some notation; since much of this should be fairly familiar we will be rather sketchy and refer the interested reader to Ref.[10] for details.

We denote the couplings of the SM fermions to the  $W_i = (W = W_{SM}, W')$  as

$$\left(\frac{G_F M_W^2}{\sqrt{2}}\right)^{1/2} V_{ff'} C_i^{\ell,q} \bar{f} \gamma_\mu (1 - h_i \gamma_5) f' W_i^\mu + h.c., \quad (1)$$

where for the case of  $W_i = W_{SM}$ , the coupling strength(for leptons and quarks, respectively) and helicity factors are given by  $C_i^{\ell,q}, h_i = 1$  and  $V_{ff'}$  is the CKM(unit) matrix when  $f, f'$  are quarks(leptons); note that the helicity structure for both leptons and quarks is assumed to be the same as in all the model cases above.<sup>§</sup> Following the notation given in Ref.[10], with some obvious

---

<sup>§</sup>For simplicity in what follows we will further assume that the corresponding RH and LH CKM matrices are identical up to phases and we will generally neglect any possible small effects arising from  $W - W'$  mixing. In the case of RH couplings, we will further assume that the SM neutrinos are Dirac fields.

modifications, the inclusive  $pp \rightarrow W_i^+ \rightarrow \ell^+ \nu + X$  differential cross section can be written as

$$\frac{d\sigma}{d\tau dy dz} = K \frac{G_F^2 M_W^4}{48\pi} \sum_{qq'} |V_{qq'}|^2 \left[ SG_{qq'}^+(1+z^2) + 2AG_{qq'}^- z \right], \quad (2)$$

where  $K$  is a kinematic/numerical factor that accounts for NLO and NNLO QCD corrections[12] as well as leading electroweak corrections[13] and is roughly of order  $\simeq 1.3$  for suitably defined couplings,  $\tau = M^2/s$  ( $\sqrt{s} = 14$  TeV at the LHC) with  $M^2$  being the lepton pair invariant mass. Furthermore,

$$\begin{aligned} S &= \sum_{ij} P_{ij} (C_i C_j)^\ell (C_i C_j)^q (1 + h_i h_j)^2 \\ A &= \sum_{ij} P_{ij} (C_i C_j)^\ell (C_i C_j)^q (h_i + h_j)^2, \end{aligned} \quad (3)$$

where the sums extend over all of the exchanged particles in the  $s$ -channel. Here

$$P_{ij} = \hat{s} \frac{(\hat{s} - M_i^2)(\hat{s} - M_j^2) + \Gamma_i \Gamma_j M_i M_j}{[(\hat{s} - M_i^2)^2 + \Gamma_i^2 M_i^2][i \rightarrow j]}, \quad (4)$$

with  $\hat{s} = M^2$  being the square of the total collision energy and  $\Gamma_i$  the total widths of the exchanged  $W_i$  particles. Note that we have employed  $z = \cos \theta$ , the scattering angle in the CM frame defined as that between the incoming  $u$ -type quark and the outgoing neutrino (both being fermions as opposed to being one fermion and one anti-fermion). Furthermore, the following combinations of parton distribution functions appear:

$$G_{qq'}^\pm = \left[ q(x_a, M^2) \bar{q}'(x_b, M^2) \pm q(x_b, M^2) \bar{q}'(x_a, M^2) \right], \quad (5)$$

where  $q(q')$  is a  $u(d)$ -type quark and  $x_{a,b} = \sqrt{\tau} e^{\pm y}$  are the corresponding parton momentum fractions. Analogous expressions can also be written in the case of  $W_i^-$  exchange by taking  $z \rightarrow -z$  and interchanging initial state quarks and anti-quarks.

In most cases of interest one usually converts the distribution over  $z$  above into one over the transverse mass,  $M_T$ , formed from the final state lepton and the missing transverse energy associated with the neutrino; at fixed  $M$ , one has  $z = (1 - M_T^2/M^2)^{1/2}$ . The resulting transverse mass distribution can then be written as

$$\frac{d\sigma}{dM_T} = \int_{M_T^2/s}^1 d\tau \int_{-Y}^Y dy J(z \rightarrow M_T) \frac{d\sigma}{d\tau dy dz}, \quad (6)$$

where  $Y = \min(y_{cut}, -1/2 \log \tau)$  allows for a rapidity cut on the outgoing leptons and  $J(z \rightarrow M_T)$  is the appropriate Jacobian factor[15]. In practice,  $y_{cut} \simeq 2.5$  for the two LHC detectors. Note that  $\frac{d\sigma}{dM_T}$  will only pick out the  $z$ -even part of  $\frac{d\sigma}{d\tau dy dz}$  as well as the even combination of terms in the product of the parton densities,  $G_{qq'}^+$ . In the usual analogous fashion to the  $Z'$  case[10], as we will see in our discussion below, one can also define the forward-backward asymmetry as a function of the transverse mass, in principle prior to integration over the rapidity  $y$ ,  $A_{FB}(M_T, y)$ , whose numerator now picks out the  $z$ -odd terms in  $\frac{d\sigma}{d\tau dy dz}$  as well as the odd combination of terms in the parton densities  $G_{qq'}^-$ .

To be complete, we note that historically when discussing new gauge boson production, particularly when dealing with states which are weakly coupled as will be the case in what follows, use is often made of the narrow width approximation(NWA). In the  $W'$  case of relevance here, the NWA essentially replaces the integration over  $d\tau \sim dM$  by a  $\delta$  function, *i.e.*, the  $W'$  is assumed to be produced on-shell. Thus, for any smooth function  $f(M)$ , essentially,  $\int dM f(M) \rightarrow \int dM f(M) \frac{\pi}{2} \Gamma_{W'} \delta(M - M_{W'}) \rightarrow \frac{\pi}{2} \Gamma_{W'} f(M_{W'})$ , apart from some overall factors. Note that use of the NWA implies that we evaluate quantities on the ‘peak’ of the  $W'$  mass distribution, *i.e.*, at  $M = M_{W'}$ . This approximation is usually claimed to be valid up to  $O(\Gamma_{W'}/M_{W'})$  corrections(at worst), but there are occasions, *e.g.*, when  $W - W'$  interference is important, when its use can lead to a loss of valuable information and may even lead to wrong conclusions[16]. Unfortunately, in the  $W'$  case, the quantity  $M$  itself is not a true observable due to the missing longitudinal momentum

of the neutrino.

Given this background, let us now turn to an historical discussion of the determination of the  $W'$  coupling helicity. To be concrete, we will consider two different  $W'$  models; we will assume for simplicity that  $C_{W'}^{\ell,q} = 1$  in both cases and that only the value of  $h_{W'} = \pm 1$  distinguishes them. In this situation, employing the NWA, the cross section for on-shell  $W'$  production (followed by its leptonic decay) is proportional to  $\sim (1 + h_{W'}^2)$  and is trivially seen to be independent of the helicity of the couplings. We would thus conclude that cross section measurements are not useful helicity discriminants. More interestingly, as was noted long ago[17], we find that the rapidity integrated value of  $A_{FB}$ , given in the NWA by

$$A_{FB} \sim \frac{h_{W'}^2}{(1 + h_{W'}^2)^2}, \quad (7)$$

also has the *same value* for either purely LH or RH couplings<sup>¶</sup>. Thus, in the NWA,  $A_{FB}$  provides no help in determining the  $W'$  coupling helicity structure for the cases we consider here. However, we note that if the quark and leptonic coupling helicities of the  $W'$  are *opposite*, then the value of  $A_{FB}$  will flip sign in comparison to the above expectation.

It is apparent from this result that some other observable(s) must be used to distinguish these two cases. Keeping the NWA assumption, the first suggestion[18] along these lines was to examine the polarization of  $\tau$ 's originating in the decay  $W' \rightarrow \tau\nu$ . In that paper it was explicitly shown that the the energy spectrum of the final state particle in the decay  $\tau \rightarrow \ell, \pi$  or  $\rho$  (in the  $\tau$  rest frame) was reasonably sensitive to the original  $W'$  helicity since the  $\tau$  itself effectively decays only through the SM LH couplings of the  $W$ (provided the  $W'$  is sufficiently massive as we will assume here). The difficulty with this method is that the observation of this decay mode at the LHC is not all that straightforward and even the corresponding  $Z' \rightarrow \tau\tau$  mode, which is somewhat easier to observe, is just beginning to be studied by the LHC experimental collaborations[19].

---

<sup>¶</sup>This follows immediately from the fact that we have assumed that both the hadronic and leptonic couplings of the  $W'$  have to have the *same* helicity.

Clearly, measuring the polarization of the  $\tau$ 's in  $W' \rightarrow \tau\nu$  will be reasonably difficult in the LHC detector environment and may, at the very least, require large integrated luminosities even for a relatively light  $W'$ . The results of detailed studies by the LHC collaborations to address this issue are anxiously awaited.

In the early 90's, two important NWA-based methods for probing the helicity of the  $W'$  were suggested[20]. The first of these is an examination of the rare decay mode  $W' \rightarrow \ell^+\ell^-W$  (with the  $W$  decaying into jets); in particular, one makes a measurement of the ratio of branching fractions

$$R_W = \frac{B(W' \rightarrow \ell^+\ell^-W)}{B(W' \rightarrow \ell\nu)}, \quad (8)$$

obtained by employing the NWA.  $R_W$  is expected to be roughly  $\sim O(0.01)$  or so after suitable cuts. One of the main SM backgrounds, *i.e.*,  $WZ$  production, can essentially be removed by demanding that the dileptons do not form a  $Z$ , demanding that the mass of the  $jj\ell\ell$  system be not far from the (already known) value of  $M_{W'}$  and that of the dijets reconstructs to the  $W$  mass. Even after these requirements, however, some background from the continuum would remain. Furthermore, as the energy of the final state  $W$  increases it is more likely that the resulting dijets will coalesce into a single jet depending on the jet cone definition which is employed. In this case, at the very least, a very large additional background from single jets may appear; it is also possible that the events with a final state  $W$  would be completely lost without the dijet mass reconstruction. The  $3\ell + E_T^{miss}$  final state, with suitable cuts, would be obviously cleaner and would avoid some of these issues but at the price of an overall suppression due to ratio of branching fractions of  $\simeq 1/3$  thus reducing the mass range over which this process would be useful.

In a general gauge model, the amplitude for this process is the sum of two graphs. In the first graph,  $W'^- \rightarrow \ell^- \bar{\nu}^*$ , *i.e.*, the production of a virtual neutrino followed by the ‘decay’  $\bar{\nu}^* \rightarrow \ell^+ W^-$ . Clearly, if the  $W'$  couples in a purely RH manner to the SM leptons then this graph will vanish in the limit of massless neutrinos due to the presence of two opposite helicity projection

operators. This graph will, of course, be non-zero only if the  $W'$  couples in an at least partially LH manner. The second graph involves the presence of the trilinear couplings  $W'ZW$  and  $W'Z'W$ ; recall that in any model with a  $W'$ , a  $Z'$  will also appear just based on gauge invariance. In this case, the decay proceeds as  $W' \rightarrow WZ/Z'^* \rightarrow W\ell^+\ell^-$ , noting that the on-shell SM  $Z$  contribution can be removed by a suitable cut on the dilepton invariant mass. The main issue is the size of the  $W'Z'W$  (and  $W'ZW$ ) couplings and this can involve such things such as, *e.g.*, the detailed electroweak symmetry breaking patterns of the given model under study. *Generically* in extra dimensional models[3, 4, 5, 6, 7], these couplings are absent in the limit of small mixing due the orthogonality of the Kaluza-Klein wavefunctions of the states. In models where the SM  $SU(2)_L$  arises from a diagonal breaking of the form  $G_1 \otimes G_2 \rightarrow SU(2)_{Diag}$ , such as in LH models[1], the  $W'Z'W$  coupling is of order the SM weak coupling,  $g$ , while the  $W'ZW$  coupling is either of order  $g$  or can be mixing angle suppressed. In other cases, such as in the LRM[8], where  $SU(2)_L \otimes SU(2)_R$  just breaks to  $SU(2)_L$ , the  $W'ZW, WZ'W$  couplings are only generated by mixings and for the diagrams of interest are not longitudinally enhanced. Since the amplitude associated with the pure leptonic graphs are absent in this case, the entire amplitude is mixing angle suppressed so that this process has an unobservably small rate. In fact, there are no known models where the  $W'$  helicity is RH and the  $W'ZW, WZ'W$  couplings are not mixing angle suppressed<sup>||</sup>. Thus, *based on known models*, it appears that the observation of the rare decay  $W' \rightarrow \ell^+\ell^-W$  would be a compelling indication that the  $W'$  is at least partially coupled in a LH manner with apparently most of the serious SM backgrounds being removable by conventional cuts. However, in making a truly model-independent analysis one must exercise care in the use of this result. A detailed analysis of the signal and backgrounds, including that for the  $jj\ell^+\ell^-$  final state, for such decays including realistic detector effects would be very useful in addressing all these issues and should be performed. However, it also seems clear that is unlikely that a reliable measurement of  $R_W$  can be made with relatively low integrated luminosities.

---

<sup>||</sup>In a *fundamental* UV complete theory, this may follow directly from arguments based solely on gauge invariance and the requirement of high energy unitarity.



A second, imaginative possibility is to observe  $WW'$  associated production[20] with  $W \rightarrow jj$  for the same reasons as above. Many of the arguments made in the previous paragraph will also apply in this case as well since the diagrams responsible for this process are quite similar to previously discussed. Essentially these graphs are obtained by crossing, with the final state leptons now replaced by an initial state  $q\bar{q}$ . In this case one looks for the  $jj\ell E_T^{miss}$  final state with the  $\ell E_T^{miss}$  transverse mass peaking near  $M_{W'}$ . One would anticipate this cross section to be of order  $\sim 0.01$  of that of the  $W'$  discovery channel. The main issues here are, as above, the SM backgrounds and the nature of the triple gauge vertices. It is not likely that a reliable measurement of this cross section will be performed with low luminosities that could be interpreted in a model-independent way until all of the background and detector issues are dealt with. Again, a detailed analysis including detector effects should be performed.

### 3 $W - W'$ Interference as a Function of $M_T$

What we have learned from the previous discussion is that tools which employ the NWA are not particularly useful when we are trying to determine the  $W'$  coupling helicity with relatively low luminosities in an easily examined final state. One of the key reasons for this is that the use of NWA does not allow us to examine the influence of  $W - W'$  interference to which we now turn[21]\*\*.

The most obvious distribution to examine first is  $\frac{d\sigma}{dM_T}$  itself; for the moment let us restrict ourselves to the two cases where  $C_{W'}^{\ell,q} = 1$  and  $h_{W'} = \pm 1$ . Fig. 1 shows this distribution for a large integrated luminosity, assuming  $M_{W'} = 1.5$  TeV[22], as well as the SM continuum background<sup>††</sup>. Several things are immediately clear: (i) In the region near the Jacobian peak both distributions are quite similar; this is not surprising as this is the region where the NWA is most applicable since

---

\*\*We note in passing that the usual experimental analyses at LHC[23] performed by both the ATLAS and CMS collaborations ignore the effects of  $W - W'$  interference since these contributions are absent from default versions of stand-alone PYTHIA[24].

<sup>††</sup>Note that we would expect to see many excess events for such  $W'$  masses as only  $\simeq 25$   $pb^{-1}$  of luminosity would be needed to discover( $5\sigma$ ) such as state at the LHC.




Figure 1: Transverse mass distribution for the production of a 1.5 TeV  $W'$  including interference effects at the LHC displayed on both log and linear scales assuming an integrated luminosity of  $300 \text{ fb}^{-1}$ . The lowest histogram is the SM continuum background. The upper blue(middle red) histogram at  $M_T = 600$  GeV corresponds to the case of  $h_{W'} = -1(1)$ .




Figure 2: Same as in the previous figure but now on a linear scale with lower luminosities and smeared by the detector resolution. In the top(bottom) panel an integrated luminosity of  $30(10) \text{ fb}^{-1}$  has been assumed. Detector smearing has now been included assuming  $\delta M_T/M_T = 2\%$ .

now  $M_T \simeq M$  and  $W - W'$  interference is minimal. In this limit we would indeed recover our earlier result that the cross section is helicity independent. (ii) In the lower  $M_T$  region where interference effects are important the two models lead to quite different distributions. In particular, for the LH case with  $h_{W'} = 1$ , we observe a destructive interference with the SM amplitude producing a distribution that lies below that of the pure SM continuum background. (This is not surprising as the overall signs of the  $W$  and  $W'$  contributions are the same but we are at values of  $\sqrt{\hat{s}}$  that are above  $M_W$  yet below  $M_{W'}$  so that the relevant propagators have opposite signs.) However, for the RH case with  $h_{W'} = -1$ , there is no such interference and therefore the resulting distribution always lies above the SM background. It is fairly obvious that these two distributions are trivially distinguishable at these large integrated luminosities.




Figure 3: High luminosity plot of the transverse mass distribution assuming  $M_{W'} = 2.5(3.5)$  for the upper(lower) pair of histograms along with the SM continuum background. In the interference region near  $\simeq 0.5M_{W'}$  the upper(lower) member of the pair corresponds to the case of  $h_{W'} = -1(1)$ . Detector smearing has now been included assuming  $\delta M_T/M_T = 2\%$ .

Fig. 2 shows the same  $\frac{d\sigma}{dM_T}$  distribution on a linear scale but now for far smaller integrated luminosities that may be obtained during early LHC running; here we include the effects of detector smearing, with  $\delta M_T/M_T \simeq 2\%$ , which is somewhat less important in the very large statistics sample

cases shown above. It is immediately apparent that even with only  $\sim 10 \text{ fb}^{-1}$  of luminosity the two cases remain quite distinct; however, it also appears unlikely that much smaller luminosities would be very useful in this regard. This result is a significant improvement over previous attempts to determine the  $W'$  coupling helicity with low luminosities in clean channels.

At this point there are several important questions one might ask: (i) What happens for a more massive  $W'$ , *i.e.*, how much luminosity will be needed in such cases to distinguish  $W'$  couplings of opposite helicities? (ii) What if the  $W'$  couplings are weaker than our canonical choice above? (iii) Do other observables, *e.g.*,  $A_{FB}$ , measured in the interference region below the Jacobian peak assist us in model separation? (iv) What if the  $W'$  couplings are not purely chiral and are an admixture of LH and RH helicities? It is to these issues that we now turn.

Fig. 3 provides us with a high luminosity overview for the more massive cases where  $M_{W'} = 2.5$  or  $3.5 \text{ TeV}$ . In the  $M_{W'} = 2.5 \text{ TeV}$  case, Fig. 4 demonstrates that the full  $300 \text{ fb}^{-1}$  luminosity is not required to distinguish the two possibly helicities;  $\sim 60 \text{ fb}^{-1}$  seems to be the approximate minimum luminosity that appears to be necessary. For higher masses, distinguishing the two cases becomes far more difficult due to the smaller production cross section as we see from Fig. 5 for the case of  $M_{W'} = 3.5 \text{ TeV}$  assuming a luminosity of  $300 \text{ fb}^{-1}$ ; essentially the full luminosity is required for model distinction in this case.

What if the  $W'$  couplings are weaker? Clearly if they are too weak there will be insufficient statistics to discriminate the two possible coupling helicity assignments for any fixed value of  $M_{W'}$ . In order to examine a realistic example of this situation, we consider the case of the second  $W$  KK excitation in the UED model[4, 25] with a conserved KK-parity. In such a scenario the LH couplings of this field to SM fermions vanish at tree level but are induced by one loop effects. In this case one finds that the effective values of  $C^{\ell,q}$  are distinct but are qualitatively of order  $\sim 0.05$  though we employ the specific values obtained in Ref.[4, 25] below in the actual calculations. Fig. 6 shows the transverse mass distributions in this case assuming that  $M_{W'} = 1 \text{ TeV}$  for the second level




Figure 4: Transverse mass distribution assuming a mass of 2.5 TeV for the  $W'$  along with the SM continuum background; the upper(lower) panel corresponds to a luminosity of 300(75)  $fb^{-1}$ . In the interference region near  $\simeq 0.5M_{W'}$ , the upper(lower) histogram corresponds to the case of  $h_{W'} = -1(1)$ .




Figure 5: High luminosity plot of the transverse mass distribution assuming  $M_{W'} = 3.5$  TeV along with the SM continuum background. In the interference region near  $\simeq 0.5M_{W'}$  the upper(lower) histogram corresponds to the case of  $h_{W'} = -1(1)$ .

KK state. The signal for this  $W$  KK state is clearly visible above the SM background. However, we also see that for even for these high luminosities and low masses the two helicity choices are not distinguishable. Clearly, one cannot determine the  $W'$  coupling helicity for such very weak interaction strengths. Semi-quantitatively, we find that that this breakdown in the discriminating power occurs when  $(C^\ell C^q)^{1/2} \sim 0.1$  at these luminosities and masses.

We now turn to the next question we need to address: can asymmetries be useful in strengthening our ability to determine the  $W'$  coupling helicity? We know from the discussion above that the answer is apparently ‘no’ in the NWA limit, *i.e.*, when  $M_T \simeq M$ . Thus we must focus our attention on the  $M_T$  region below the peak where  $W - W'$  interference is strongest or, more generally, examine the asymmetries’  $M_T$ -dependence directly. The most obvious quantity to begin with is the  $y$ -integrated value of  $A_{FB}$  for both  $W'^{\pm}$  channels. To make such a measurement, we need to know several things in addition to the sign of the lepton (which we assume can be done with  $\simeq 100\%$  efficiency). At the parton level, in the case of  $W'^{-}$  for example, the relevant angle used to define  $A_{FB}$  lies between the incoming  $d$ -type quark and the outgoing  $\ell^-$ . Reconstructing this direction




Figure 6: High luminosity plot of the transverse mass distribution assuming  $M_{W'} = 1$  TeV for the second  $W$  KK level in the UED model smeared by detector resolution as above. As usual the lower histogram is the SM background while the other two correspond to the signal cases with  $h_{W'} = -1(1)$  and are essentially indistinguishable.

presents us with two problems: first, since the longitudinal momentum of the  $\nu$  is unknown there is an, in principle, two-fold ambiguity in the motion of the center of mass in the lab frame; this can cause a serious dilution of the observed asymmetry but can be corrected for statistically using Monte Carlo once the  $W'$  mass is known. Second, even when it is determined, the direction of motion of the center of mass is not necessarily that of the  $d$ -type quark though it is likely to be so when the boost of the center of mass frame is large. The later problem also arises for the case of a  $Z'$  and has also been shown to be mostly correctable in detailed Monte Carlo studies[26]. For the moment, let us forget these issues and ask what the  $y$ -integrated  $A_{FB}(M_T)$  looks like in both  $\ell^\pm$  channels; the results are shown in Fig. 7 assuming high luminosities and  $M_{W'} = 1.5$  TeV. Here we see that these integrated quantities, even for luminosities of  $300 \text{ fb}^{-1}$ , are essentially useless in distinguishing the two coupling helicity cases. Furthermore, we also see that the two coupling helicities lead to essentially identical results when  $M_T \simeq M_{W'}$  as would be expected based on the NWA. A short analysis indicates that approximately ten times more integrated luminosity would be required before some separation in the two cases becomes possible[27]. Clearly this situation



would only become worse if we were to raise the mass of the  $W'$  or reduce its coupling strength.

It is perhaps possible that some information is lost by only using the integrated quantity  $A_{FB}$  and we need to consider instead  $A_{FB}(y_W)$ , where  $y_W$  is the rapidity of the center of mass frame. This distribution is odd under the interchange  $y_W \rightarrow -y_W$  at the LHC so we can simply fold this distribution over the  $y_W = 0$  boundary to double the statistics. Furthermore, by integrating over a wide  $M_T$  range in the interference region below the  $W'$  peak, *e.g.*,  $0.4 \leq M_T \leq 1$  TeV in the case of a 1.5 TeV  $W'$ , further statistics can be gained. Fig 8 shows the resulting  $A_{FB}(y_W)$  distributions for a  $W'^{\pm}$  with mass of 1.5 TeV assuming a luminosity of  $300 \text{ fb}^{-1}$  for  $h_{W'} = \pm 1$ . At these large luminosities, the  $A_{FB}(y_W)$  distributions for the two helicity choices are clearly distinguishable but this will certainly become more difficult for lower luminosities or for larger masses. We find that we essentially lose all coupling helicity information when the luminosity falls much below  $\simeq 100 \text{ fb}^{-1}$  for this  $W'$  mass.

The next observable we consider is the charge asymmetry,  $A_{WQ}(y_W)$ :

$$A_{WQ}(y_W) = \frac{N_+(y_W) - N_-(y_W)}{N_+(y_W) + N_-(y_W)}, \quad (9)$$

where  $N_{\pm}(y_W)$  are the number of events with charged leptons of sign  $\pm$  in a given bin of rapidity. Note that at the LHC,  $A_{WQ}(y_W)$  is symmetric under  $y_W \rightarrow -y_W$  so that we can again fold the distribution around  $y_W = 0$ . Fig. 9 shows this distribution, integrated over the interference region  $0.4 \leq M_T \leq 1$  TeV, assuming  $M_{W'} = 1.5$  TeV and a luminosity of  $300 \text{ fb}^{-1}$ . It is clear that at this level of integrated luminosity the two distributions are reasonably distinguishable. However, as we lower the luminosity or raise the mass of the  $W'$  the quality of the separation degrades significantly. Certainly for luminosities less than  $\simeq 100 \text{ fb}^{-1}$ , this asymmetry measurement would not be very helpful. Thus  $A_{WQ}(y_W)$  is not a very useful tool for coupling helicity determination until high luminosities become available.

A last asymmetry possibility to consider is the rapidity asymmetry for the final state charged




Figure 7: The  $y$ -integrated value of  $A_{FB}$ , as a function of the transverse mass, assuming a mass of 1.5 TeV for the  $W'^+(W'^-)$  in the top(bottom) panel. Here an integrated luminosity of  $300 \text{ fb}^{-1}$  has been assumed. The two essentially indistinguishable histograms correspond to the two possible choices of the helicity,  $h_{W'} = \pm 1$ .




Figure 8: The value of  $A_{FB}$  as a function the center of mass rapidity,  $y_W$ , integrated over the transverse mass bin 400-1000 GeV assuming a mass of 1.5 TeV for the  $W'^+(W'^-)$  in the top(bottom) panel. An integrated luminosity of  $300 \text{ fb}^{-1}$  has been assumed and the distribution has been folded around  $y_W = 0$ . The upper(lower) set of data points in the top(lower) panel for small values of  $y_W$  corresponds to the choice of  $h_{W'} = -1$ . Note that we have chosen signs to make the ranges of  $A_{FB}$  comparable in both cases.




Figure 9: The  $W - W'$  induced charge asymmetry, assuming  $M_{W'} = 1.5$  TeV, as a function the center of mass rapidity,  $y_W$ , integrated over the transverse mass bin 400-1000 GeV. An integrated luminosity of  $300 \text{ fb}^{-1}$  has been assumed and the distribution has been folded around  $y_W = 0$ . The upper set of data points at low values of  $y_W$  corresponds to the choice of  $h_{W'} = 1$ .

leptons themselves,  $A_\ell(y_\ell)$ :

$$A_\ell(y_\ell) = \frac{N_+(y_\ell) - N_-(y_\ell)}{N_+(y_\ell) + N_-(y_\ell)}, \quad (10)$$

which is also an even function of  $y_\ell$  so the distribution can again be folded around  $y_\ell = 0$ . The resulting distribution can be seen in Fig. 10 for large integrated luminosities. Here we again see reasonable model differentiation at low values of  $y_\ell \lesssim 1$  but this fades in utility as integrated luminosities drop much below  $\simeq 100 \text{ fb}^{-1}$  as the two curves are generally rather close.

From this general discussion of possibly asymmetries that one can form employing this final state we can thus conclude that their usefulness in coupling helicity determination will require  $\simeq 100 \text{ fb}^{-1}$ .

Lastly, and to be more general, we must at least consider possible scenarios where the couplings of the  $W'$  to SM fermions are a substantial admixture of both LH and RH helicities, though obvious examples of such kinds of models are apparently absent from the existing literature.




Figure 10: The  $W - W'$  induced lepton asymmetry, assuming  $M_{W'} = 1.5$  TeV, as a function the lepton's rapidity,  $y_\ell$ , integrated over the transverse mass bin 400-1100 GeV. An integrated luminosity of  $300 \text{ fb}^{-1}$  has been assumed and the distribution has been folded around  $y_\ell = 0$ . The upper set of data points at low values of  $y_\ell$  corresponds to the choice of  $h_{W'} = 1$ .

To get a feel for such a possibility, we perform two analyses: first, we set  $C^{\ell,q} = 1$  as before and vary the values of  $h_{W'}$  between pairs of positive and negative values. As we do this, the helicity of the couplings of the  $W'$  will vary as will its total decay width which behaves as  $\sim 1 + h_{W'}^2$ . In a second analysis, we can rescale the values of the  $C^{\ell,q}$  so that the  $W'$  width is held fixed. In this case, as we will see, the resulting histograms for the transverse mass distribution lie especially close to one another. The results of these two sets of calculations are shown in Fig. 11 in the case of large integrated luminosities assuming the default value of  $M_{W'} = 1.5$  TeV. In the first analysis shown in the top panel, we see that at these assumed luminosities all of the different histograms are distinguishable and not just the two pairs of cases with opposite helicities. This result generally remains true down to luminosities  $\sim 75 \text{ fb}^{-1}$  or so. If we are *only* interested in separating opposite helicity pairs then we find that the cases  $h_{W'} = \pm 0.8(0.6, 0.4, 0.2)$  can be distinguished down to luminosities of order  $\sim 10(25, 50, 75) \text{ fb}^{-1}$ , respectively.

In the second analysis, as seen in the lower panel of the figure, the histograms for  $h_{W'} =$

0.8, 0.6 and 0.4 (as well as for their corresponding opposite helicity partners) are very close to one another and are essentially inseparable even at these high luminosities. However, the two sets of opposite helicity histograms remain distinguishable and this will remain true down to luminosities of order  $30 - 75 \text{ fb}^{-1}$ . It would seem from these analyses that the transverse mass distribution will play the dominant role in  $W'$  coupling helicity determination in all possible cases although somewhat higher integrated luminosities may be required in some scenarios.

## 4 Summary and Conclusions

Apart from its mass and width, the most important property of a new charged gauge boson,  $W'$ , is the helicity of its couplings to the SM fermions. Such particles are predicted to exist in the TeV mass range in many new physics models and this coupling helicity is an order one discriminator between the various classes of models. The main difficulties with the existing techniques for determining this helicity are potentially threefold: (i) they require rather high integrated luminosities even for a relatively light  $W'$ , and/or (ii) they are sufficiently intricate as to require a detailed background and detector study to determine their feasibility, and/or (iii) they make use of more complex final states other than the standard  $\ell + E_T^{\text{miss}}$  discovery channel. Some of these techniques also suffer from employing the narrow width approximation which can result in loss of valuable information regarding the effects of  $W - W'$  interference. In this paper we propose a simple technique for making this helicity determination at the LHC. In order to attempt to circumvent all of these difficulties, we have examined the  $W - W'$  interference region of the transverse mass distribution for the  $\ell + E_T^{\text{miss}}$  discovery mode. We have found that this distribution is particularly sensitive to the helicity of the  $W'$  couplings. In particular, using this technique we have shown that such helicity differentiation requires only  $\sim 10(60, 300) \text{ fb}^{-1}$  assuming  $M_{W'} = 1.5(2.5, 3.5) \text{ TeV}$  and provided that the  $W'$  has Standard Model strength couplings. This helicity determination can be further strengthened by the use of various discovery channel leptonic asymmetries also measured




Figure 11: Same as the linear plot shown in Fig.1, but now for other values of the coupling helicities. From top to bottom the pairs of histograms in the upper panel correspond to  $h(W') = \pm 0.8, \pm 0.6, \pm 0.4$  and  $\pm 0.2$ , respectively. The next lowest single histogram corresponds to the case of pure vector couplings, *i.e.*,  $h(W') = 0$ . In producing these results we have assumed that the values of the  $C^{\ell,q}=1$ . In the lower panel, we show the same result now but with the overall couplings rescaled so as to keep the  $W'$  width a constant.

in the same interference regime once higher integrated luminosities are available as well as by the more traditional approaches. Hopefully the LHC will observe a  $W'$  so that this approach can be employed.

## Acknowledgments

The author would like to thank A. De Roeck, S. Godfrey and J. Hewett for input and discussions related to this paper.

## References

- [1] N. Arkani-Hamed, A. G. Cohen and H. Georgi, Phys. Lett. B **513**, 232 (2001) [arXiv:hep-ph/0105239]; N. Arkani-Hamed, A. G. Cohen, E. Katz and A. E. Nelson, JHEP **0207**, 034 (2002) [arXiv:hep-ph/0206021].
- [2] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999) [arXiv:hep-ph/9905221].
- [3] H. Davoudiasl, J. L. Hewett and T. G. Rizzo, Phys. Lett. B **473**, 43 (2000) [arXiv:hep-ph/9911262]; A. Pomarol, Phys. Lett. B **486**, 153 (2000) [arXiv:hep-ph/9911294]; Y. Grossman and M. Neubert, Phys. Lett. B **474**, 361 (2000) [arXiv:hep-ph/9912408]; H. Davoudiasl, J. L. Hewett and T. G. Rizzo, Phys. Rev. D **63**, 075004 (2001) [arXiv:hep-ph/0006041]; T. Gherghetta and A. Pomarol, Nucl. Phys. B **586**, 141 (2000) [arXiv:hep-ph/0003129]; S. Chang, J. Hisano, H. Nakano, N. Okada and M. Yamaguchi, Phys. Rev. D **62**, 084025 (2000) [arXiv:hep-ph/9912498]; S. J. Huber and Q. Shafi, Phys. Lett. B **498**, 256 (2001) [arXiv:hep-ph/0010195] and Phys. Rev. D **63**, 045010 (2001) [arXiv:hep-ph/0005286]; R. Kitano, Phys. Lett. B **481**, 39 (2000) [arXiv:hep-ph/0002279]; J. L. Hewett, F. J. Petriello and T. G. Rizzo, JHEP **0209**, 030 (2002) [arXiv:hep-ph/0203091].
- [4] T. Appelquist, H. C. Cheng and B. A. Dobrescu, Phys. Rev. D **64**, 035002 (2001) [arXiv:hep-ph/0012100]; H. C. Cheng, K. T. Matchev and M. Schmaltz, Phys. Rev. D **66**, 056006 (2002)



- [arXiv:hep-ph/0205314] and Phys. Rev. D **66**, 036005 (2002) [arXiv:hep-ph/0204342].
- [5] I. Antoniadis, Phys. Lett. B **246**, 377 (1990);
- [6] T. G. Rizzo and J. D. Wells, Phys. Rev. D **61**, 016007 (2000) [arXiv:hep-ph/9906234];  
 K. m. Cheung and G. Landsberg, Phys. Rev. D **65**, 076003 (2002) [arXiv:hep-ph/0110346];  
 A. Strumia, Phys. Lett. B **466**, 107 (1999) [arXiv:hep-ph/9906266]; A. Delgado, A. Pomarol  
 and M. Quiros, JHEP **0001**, 030 (2000) [arXiv:hep-ph/9911252]; F. Cornet, M. Relano and  
 J. Rico, Phys. Rev. D **61**, 037701 (2000) [arXiv:hep-ph/9908299]; C. D. Carone, Phys. Rev.  
 D **61**, 015008 (2000) [arXiv:hep-ph/9907362]; P. Nath and M. Yamaguchi, Phys. Rev. D **60**,  
 116004 (1999) [arXiv:hep-ph/9902323]; A. Muck, A. Pilaftsis and R. Ruckl, Phys. Rev. D **65**,  
 085037 (2002) [arXiv:hep-ph/0110391]; G. Polesello and M. Prata, Eur. Phys. J. C **32S2**, 55  
 (2003); P. Nath, Y. Yamada and M. Yamaguchi, Phys. Lett. B **466**, 100 (1999) [arXiv:hep-  
 ph/9905415];
- [7] H. Davoudiasl and T. G. Rizzo, arXiv:hep-ph/0702078.
- [8] For a classic review and original references, see R.N. Mohapatra, *Unification and Supersymmetry*, (Springer, New York, 1986).
- [9] See, for example, K. R. Lynch, E. H. Simmons, M. Narain and S. Mrenna, Phys. Rev. D **63**,  
 035006 (2001) [arXiv:hep-ph/0007286]; H. Georgi, E. Jenkins and E. H. Simmons, Nucl. Phys.  
 B **331**, 541 (1990); A. Bagneid, T. K. Kuo and N. Nakagawa, Int. J. Mod. Phys. A **2**, 1351  
 (1987).
- [10] For classic reviews of  $Z'$  physics, see A. Leike, Phys. Rept. **317**, 143 (1999) [arXiv:hep-  
 ph/9805494]; J. L. Hewett and T. G. Rizzo, Phys. Rept. **183**, 193 (1989); M. Cvetič and  
 S. Godfrey, arXiv:hep-ph/9504216; T. G. Rizzo, “Extended gauge sectors at future colliders:  
 Report of the new gauge boson subgroup,” eConf **C960625**, NEW136 (1996) [arXiv:hep-  
 ph/9612440] and arXiv:hep-ph/0610104.

- [11] See, however, S. Godfrey, P. Kalyniak, B. Kamal, M. A. Doncheski and A. Leike, *Int. J. Mod. Phys. A* **16S1B**, 879 (2001) [arXiv:hep-ph/0009325]; S. Godfrey, P. Kalyniak, B. Kamal and A. Leike, *Phys. Rev. D* **61**, 113009 (2000) [arXiv:hep-ph/0001074].
- [12] For a recent analysis and original references, see K. Melnikov and F. Petriello, arXiv:hep-ph/0609070.
- [13] U. Baur and D. Wackeroth, *Nucl. Phys. Proc. Suppl.* **116**, 159 (2003) [arXiv:hep-ph/0211089];
- [14] V. A. Zykunov, arXiv:hep-ph/0509315.
- [15] See, for example, V. Barger and R. J. N. Phillips in *Collider Physics*, Frontiers in Physics Series Vol.71, 1996.
- [16] D. Berdine, N. Kauer and D. Rainwater, arXiv:hep-ph/0703058.
- [17] H. E. Haber, “Signals Of New W’s And Z’s,” SLAC-PUB-3456 *Presented at 1984 Summer Study on the Design and Utilization of the Superconducting Super Collider, Snowmass, CO, Jun 23 - Jul 23, 1984*
- [18] H. E. Haber, “Taus: A Probe Of New W And Z Couplings,” *Presented at 1984 Summer Study on the Design and Utilization of the Superconducting Super Collider, Snowmass, CO, Jun 23 - Jul 23, 1984*.
- [19] See the talk given by T. Vickey, at the ATLAS Exotics Working Group Meeting, 2/21/07.
- [20] M. Cvetič, P. Langacker and B. Kayser, *Phys. Rev. Lett.* **68**, 2871 (1992); M. Cvetič, P. Langacker and J. Liu, *Phys. Rev. D* **49**, 2405 (1994) [arXiv:hep-ph/9308251]; M. Cvetič and P. Langacker, *Phys. Rev. D* **46**, 4943 (1992) [Erratum-ibid. *D* **48**, 4484 (1993)] [arXiv:hep-ph/9207216].
- [21] The possibility of probing  $W - W'$  interference in the  $t\bar{b}$  channel has recently been discussed in E. Boos, V. Bunichev, L. Dudko and M. Perfilov, arXiv:hep-ph/0610080.

- [22] The direct search lower limit on the mass of a  $W'$  with such couplings is approaching 1 TeV from Run II data at the Tevatron; see, for example, P. Savard, “Searches for Extra Dimensions and New Gauge Bosons at the Tevatron,” talk given at the *XXXIII International Conference on High Energy Physics*, 26 July-2 August 2006, Moscow, Russia; T. Adams, “Searches for New Phenomena with Lepton Final States at the Tevatron,” talk given at Rencontres de Moriond Electroweak Interactions and Unified Theories 2007, La Thuile, Italy 10-17 March 2007; A. Abulencia *et al.* [CDF Collaboration], arXiv:hep-ex/0611022. In the case of the LRM, the lower bound from indirect measurements may be somewhat larger: P. Langacker and S. Uma Sankar, Phys. Rev. D **40**, 1569 (1989).
- [23] See, for example, G. Azuelos *et al.*, Eur. Phys. J. C **39**S2, 13 (2005) [arXiv:hep-ph/0402037]; C. Hof, T. Hebbeker and K. Hoepfner,
- [24] T. Sjostrand, S. Mrenna and P. Skands, JHEP **0605**, 026 (2006) [arXiv:hep-ph/0603175].
- [25] A. Datta, K. Kong and K. T. Matchev, Phys. Rev. D **72**, 096006 (2005) [Erratum-ibid. D **72**, 119901 (2005)] [arXiv:hep-ph/0509246].
- [26] See, for example, R. Cousins, J. Mumford and V. Valuev, CMS Note 2005/022; I. Golutin *et al.*, CMS AN-2007/003.
- [27] F. Gianotti *et al.*, Eur. Phys. J. C **39**, 293 (2005) [arXiv:hep-ph/0204087].