

Production and Decay of Ω_c^0

B. Aubert,¹ M. Bona,¹ D. Boutigny,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Poireau,¹ X. Prudent,¹ V. Tisserand,¹
 A. Zghiche,¹ J. Garra Tico,² E. Grauges,² L. Lopez,³ A. Palano,³ G. Eigen,⁴ I. Ofte,⁴ B. Stugu,⁴ L. Sun,⁴
 G. S. Abrams,⁵ M. Battaglia,⁵ D. N. Brown,⁵ J. Button-Shafer,⁵ R. N. Cahn,⁵ Y. Groysman,⁵ R. G. Jacobsen,⁵
 J. A. Kadyk,⁵ L. T. Kerth,⁵ Yu. G. Kolomensky,⁵ G. Kukartsev,⁵ D. Lopes Pegna,⁵ G. Lynch,⁵ L. M. Mir,⁵
 T. J. Orimoto,⁵ M. Pripstein,⁵ N. A. Roe,⁵ M. T. Ronan,^{5,*} K. Tackmann,⁵ W. A. Wenzel,⁵ P. del Amo Sanchez,⁶
 C. M. Hawkes,⁶ A. T. Watson,⁶ T. Held,⁷ H. Koch,⁷ B. Lewandowski,⁷ M. Pelizaeus,⁷ T. Schroeder,⁷
 M. Steinke,⁷ W. N. Cottingham,⁸ D. Walker,⁸ D. J. Asgeirsson,⁹ T. Cuhadar-Donszelmann,⁹ B. G. Fulsom,⁹
 C. Hearty,⁹ N. S. Knecht,⁹ T. S. Mattison,⁹ J. A. McKenna,⁹ A. Khan,¹⁰ M. Saleem,¹⁰ L. Teodorescu,¹⁰
 V. E. Blinov,¹¹ A. D. Bukin,¹¹ V. P. Druzhinin,¹¹ V. B. Golubev,¹¹ A. P. Onuchin,¹¹ S. I. Serednyakov,¹¹
 Yu. I. Skovpen,¹¹ E. P. Solodov,¹¹ K. Yu Todyshev,¹¹ M. Bondioli,¹² S. Curry,¹² I. Eschrich,¹² D. Kirkby,¹²
 A. J. Lankford,¹² P. Lund,¹² M. Mandelkern,¹² E. C. Martin,¹² D. P. Stoker,¹² S. Abachi,¹³ C. Buchanan,¹³
 S. D. Foulkes,¹⁴ J. W. Gary,¹⁴ F. Liu,¹⁴ O. Long,¹⁴ B. C. Shen,¹⁴ L. Zhang,¹⁴ H. P. Paar,¹⁵ S. Rahatlou,¹⁵
 V. Sharma,¹⁵ J. W. Berryhill,¹⁶ C. Campagnari,¹⁶ A. Cunha,¹⁶ B. Dahmes,¹⁶ T. M. Hong,¹⁶ D. Kovalskyi,¹⁶
 J. D. Richman,¹⁶ T. W. Beck,¹⁷ A. M. Eisner,¹⁷ C. J. Flacco,¹⁷ C. A. Heusch,¹⁷ J. Kroseberg,¹⁷ W. S. Lockman,¹⁷
 T. Schalk,¹⁷ B. A. Schumm,¹⁷ A. Seiden,¹⁷ D. C. Williams,¹⁷ M. G. Wilson,¹⁷ L. O. Winstrom,¹⁷ E. Chen,¹⁸
 C. H. Cheng,¹⁸ A. Dvoretzki,¹⁸ F. Fang,¹⁸ D. G. Hitlin,¹⁸ I. Narsky,¹⁸ T. Piatenko,¹⁸ F. C. Porter,¹⁸
 G. Mancinelli,¹⁹ B. T. Meadows,¹⁹ K. Mishra,¹⁹ M. D. Sokoloff,¹⁹ F. Blanc,²⁰ P. C. Bloom,²⁰ S. Chen,²⁰
 W. T. Ford,²⁰ J. F. Hirschauer,²⁰ A. Kreisel,²⁰ M. Nagel,²⁰ U. Nauenberg,²⁰ A. Olivas,²⁰ J. G. Smith,²⁰
 K. A. Ulmer,²⁰ S. R. Wagner,²⁰ J. Zhang,²⁰ A. M. Gabareen,²¹ A. Soffer,²¹ W. H. Toki,²¹ R. J. Wilson,²¹
 F. Winklmeier,²¹ Q. Zeng,²¹ D. D. Altenburg,²² E. Feltresi,²² A. Hauke,²² H. Jasper,²² J. Merkel,²² A. Petzold,²²
 B. Spaan,²² K. Wacker,²² T. Brandt,²³ V. Klose,²³ H. M. Lacker,²³ W. F. Mader,²³ R. Nogowski,²³ J. Schubert,²³
 K. R. Schubert,²³ R. Schwierz,²³ J. E. Sundermann,²³ A. Volk,²³ D. Bernard,²⁴ G. R. Bonneaud,²⁴ E. Latour,²⁴
 V. Lombardo,²⁴ Ch. Thiebaut,²⁴ M. Verderi,²⁴ P. J. Clark,²⁵ W. Gradl,²⁵ F. Muheim,²⁵ S. Playfer,²⁵
 A. I. Robertson,²⁵ Y. Xie,²⁵ M. Andreotti,²⁶ D. Bettoni,²⁶ C. Bozzi,²⁶ R. Calabrese,²⁶ A. Cecchi,²⁶ G. Cibinetto,²⁶
 P. Franchini,²⁶ E. Luppi,²⁶ M. Negrini,²⁶ A. Petrella,²⁶ L. Piemontese,²⁶ E. Prencipe,²⁶ V. Santoro,²⁶ F. Anulli,²⁷
 R. Baldini-Ferroli,²⁷ A. Calcaterra,²⁷ R. de Sangro,²⁷ G. Finocchiaro,²⁷ S. Pacetti,²⁷ P. Patteri,²⁷ I. M. Peruzzi,^{27,†}
 M. Piccolo,²⁷ M. Rama,²⁷ A. Zallo,²⁷ A. Buzzo,²⁸ R. Contri,²⁸ M. Lo Vetere,²⁸ M. M. Macri,²⁸ M. R. Monge,²⁸
 S. Passaggio,²⁸ C. Patrignani,²⁸ E. Robutti,²⁸ A. Santroni,²⁸ S. Tosi,²⁸ K. S. Chaisanguanthum,²⁹ M. Morii,²⁹
 J. Wu,²⁹ R. S. Dubitzky,³⁰ J. Marks,³⁰ S. Schenk,³⁰ U. Uwer,³⁰ D. J. Bard,³¹ P. D. Dauncey,³¹ R. L. Flack,³¹
 J. A. Nash,³¹ M. B. Nikolich,³¹ W. Panduro Vazquez,³¹ P. K. Behera,³² X. Chai,³² M. J. Charles,³² U. Mallik,³²
 N. T. Meyer,³² V. Ziegler,³² J. Cochran,³³ H. B. Crawley,³³ L. Dong,³³ V. Eyges,³³ W. T. Meyer,³³ S. Prell,³³
 E. I. Rosenberg,³³ A. E. Rubin,³³ A. V. Gritsan,³⁴ Z. J. Guo,³⁴ C. K. Lae,³⁴ A. G. Denig,³⁵ M. Fritsch,³⁵
 G. Schott,³⁵ N. Arnaud,³⁶ J. Béquilleux,³⁶ M. Davier,³⁶ G. Grosdidier,³⁶ A. Höcker,³⁶ V. Lepeltier,³⁶
 F. Le Diberder,³⁶ A. M. Lutz,³⁶ S. Pruvot,³⁶ S. Rodier,³⁶ P. Roudeau,³⁶ M. H. Schune,³⁶ J. Serrano,³⁶ V. Sordini,³⁶
 A. Stocchi,³⁶ W. F. Wang,³⁶ G. Wormser,³⁶ D. J. Lange,³⁷ D. M. Wright,³⁷ C. A. Chavez,³⁸ I. J. Forster,³⁸
 J. R. Fry,³⁸ E. Gabathuler,³⁸ R. Gamet,³⁸ D. E. Hutchcroft,³⁸ D. J. Payne,³⁸ K. C. Schofield,³⁸ C. Touramanis,³⁸
 A. J. Bevan,³⁹ K. A. George,³⁹ F. Di Lodovico,³⁹ W. Menges,³⁹ R. Sacco,³⁹ G. Cowan,⁴⁰ H. U. Flaecher,⁴⁰
 D. A. Hopkins,⁴⁰ P. S. Jackson,⁴⁰ T. R. McMahon,⁴⁰ F. Salvatore,⁴⁰ A. C. Wren,⁴⁰ D. N. Brown,⁴¹ C. L. Davis,⁴¹
 J. Allison,⁴² N. R. Barlow,⁴² R. J. Barlow,⁴² Y. M. Chia,⁴² C. L. Edgar,⁴² G. D. Lafferty,⁴² T. J. West,⁴²
 J. I. Yi,⁴² J. Anderson,⁴³ C. Chen,⁴³ A. Jawahery,⁴³ D. A. Roberts,⁴³ G. Simi,⁴³ J. M. Tuggle,⁴³ G. Blaylock,⁴⁴
 C. Dallapiccola,⁴⁴ S. S. Hertzbach,⁴⁴ X. Li,⁴⁴ T. B. Moore,⁴⁴ E. Salvati,⁴⁴ S. Saremi,⁴⁴ R. Cowan,⁴⁵ P. H. Fisher,⁴⁵
 G. Sciolla,⁴⁵ S. J. Sekula,⁴⁵ M. Spitznagel,⁴⁵ F. Taylor,⁴⁵ R. K. Yamamoto,⁴⁵ S. E. Mclachlin,⁴⁶ P. M. Patel,⁴⁶
 S. H. Robertson,⁴⁶ A. Lazzaro,⁴⁷ F. Palombo,⁴⁷ J. M. Bauer,⁴⁸ L. Cremaldi,⁴⁸ V. Eschenburg,⁴⁸ R. Godang,⁴⁸
 R. Kroeger,⁴⁸ D. A. Sanders,⁴⁸ D. J. Summers,⁴⁸ H. W. Zhao,⁴⁸ S. Brunet,⁴⁹ D. Côté,⁴⁹ M. Simard,⁴⁹ P. Taras,⁴⁹
 F. B. Viaud,⁴⁹ H. Nicholson,⁵⁰ G. De Nardo,⁵¹ F. Fabozzi,^{51,‡} L. Lista,⁵¹ D. Monorchio,⁵¹ C. Sciacca,⁵¹
 M. A. Baak,⁵² G. Raven,⁵² H. L. Snoek,⁵² C. P. Jessop,⁵³ J. M. LoSecco,⁵³ G. Benelli,⁵⁴ L. A. Corwin,⁵⁴
 K. K. Gan,⁵⁴ K. Honscheid,⁵⁴ D. Hufnagel,⁵⁴ H. Kagan,⁵⁴ R. Kass,⁵⁴ J. P. Morris,⁵⁴ A. M. Rahimi,⁵⁴
 J. J. Regensburger,⁵⁴ R. Ter-Antonyan,⁵⁴ Q. K. Wong,⁵⁴ N. L. Blount,⁵⁵ J. Brau,⁵⁵ R. Frey,⁵⁵ O. Igonkina,⁵⁵
 J. A. Kolb,⁵⁵ M. Lu,⁵⁵ R. Rahmat,⁵⁵ N. B. Sinev,⁵⁵ D. Strom,⁵⁵ J. Strube,⁵⁵ E. Torrence,⁵⁵ N. Gagliardi,⁵⁶ A. Gaz,⁵⁶
 M. Margoni,⁵⁶ M. Morandin,⁵⁶ A. Pompili,⁵⁶ M. Posocco,⁵⁶ M. Rotondo,⁵⁶ F. Simonetto,⁵⁶ R. Stroili,⁵⁶ C. Voci,⁵⁶

E. Ben-Haim,⁵⁷ H. Briand,⁵⁷ J. Chauveau,⁵⁷ P. David,⁵⁷ L. Del Buono,⁵⁷ Ch. de la Vaissière,⁵⁷ O. Hamon,⁵⁷
 B. L. Hartfiel,⁵⁷ Ph. Leruste,⁵⁷ J. Malclès,⁵⁷ J. Ocariz,⁵⁷ A. Perez,⁵⁷ L. Gladney,⁵⁸ M. Biasini,⁵⁹ R. Covarelli,⁵⁹
 E. Manoni,⁵⁹ C. Angelini,⁶⁰ G. Batignani,⁶⁰ S. Bettarini,⁶⁰ G. Calderini,⁶⁰ M. Carpinelli,⁶⁰ R. Cenci,⁶⁰
 A. Cervelli,⁶⁰ F. Forti,⁶⁰ M. A. Giorgi,⁶⁰ A. Lusiani,⁶⁰ G. Marchiori,⁶⁰ M. A. Mazur,⁶⁰ M. Morganti,⁶⁰ N. Neri,⁶⁰
 E. Paoloni,⁶⁰ G. Rizzo,⁶⁰ J. J. Walsh,⁶⁰ M. Haire,⁶¹ J. Biesiada,⁶² P. Elmer,⁶² Y. P. Lau,⁶² C. Lu,⁶² J. Olsen,⁶²
 A. J. S. Smith,⁶² A. V. Telnov,⁶² E. Baracchini,⁶³ F. Bellini,⁶³ G. Cavoto,⁶³ A. D’Orazio,⁶³ D. del Re,⁶³ E. Di
 Marco,⁶³ R. Faccini,⁶³ F. Ferrarotto,⁶³ F. Ferroni,⁶³ M. Gaspero,⁶³ P. D. Jackson,⁶³ L. Li Gioi,⁶³ M. A. Mazzoni,⁶³
 S. Morganti,⁶³ G. Piredda,⁶³ F. Polci,⁶³ F. Renga,⁶³ C. Voena,⁶³ M. Ebert,⁶⁴ H. Schröder,⁶⁴ R. Waldi,⁶⁴ T. Adye,⁶⁵
 G. Castelli,⁶⁵ B. Franek,⁶⁵ E. O. Olaiya,⁶⁵ S. Ricciardi,⁶⁵ W. Roethel,⁶⁵ F. F. Wilson,⁶⁵ R. Aleksan,⁶⁶ S. Emery,⁶⁶
 M. Escalier,⁶⁶ A. Gaidot,⁶⁶ S. F. Ganzhur,⁶⁶ G. Hamel de Monchenault,⁶⁶ W. Kozanecki,⁶⁶ M. Legendre,⁶⁶
 G. Vasseur,⁶⁶ Ch. Yèche,⁶⁶ M. Zito,⁶⁶ X. R. Chen,⁶⁷ H. Liu,⁶⁷ W. Park,⁶⁷ M. V. Purohit,⁶⁷ J. R. Wilson,⁶⁷
 M. T. Allen,⁶⁸ D. Aston,⁶⁸ R. Bartoldus,⁶⁸ P. Bechtel,⁶⁸ N. Berger,⁶⁸ R. Claus,⁶⁸ J. P. Coleman,⁶⁸ M. R. Convery,⁶⁸
 J. C. Dingfelder,⁶⁸ J. Dorfan,⁶⁸ G. P. Dubois-Felsmann,⁶⁸ D. Dujmic,⁶⁸ W. Dunwoodie,⁶⁸ R. C. Field,⁶⁸
 T. Glanzman,⁶⁸ S. J. Gowdy,⁶⁸ M. T. Graham,⁶⁸ P. Grenier,⁶⁸ C. Hast,⁶⁸ T. Hryn’ova,⁶⁸ W. R. Innes,⁶⁸
 M. H. Kelsey,⁶⁸ H. Kim,⁶⁸ P. Kim,⁶⁸ D. W. G. S. Leith,⁶⁸ S. Li,⁶⁸ S. Luitz,⁶⁸ V. Luth,⁶⁸ H. L. Lynch,⁶⁸
 D. B. MacFarlane,⁶⁸ H. Marsiske,⁶⁸ R. Messner,⁶⁸ D. R. Muller,⁶⁸ C. P. O’Grady,⁶⁸ A. Perazzo,⁶⁸ M. Perl,⁶⁸
 T. Pulliam,⁶⁸ B. N. Ratcliff,⁶⁸ A. Roodman,⁶⁸ A. A. Salnikov,⁶⁸ R. H. Schindler,⁶⁸ J. Schwiening,⁶⁸ A. Snyder,⁶⁸
 J. Stelzer,⁶⁸ D. Su,⁶⁸ M. K. Sullivan,⁶⁸ K. Suzuki,⁶⁸ S. K. Swain,⁶⁸ J. M. Thompson,⁶⁸ J. Va’vra,⁶⁸ N. van Bakel,⁶⁸
 A. P. Wagner,⁶⁸ M. Weaver,⁶⁸ W. J. Wisniewski,⁶⁸ M. Wittgen,⁶⁸ D. H. Wright,⁶⁸ A. K. Yarritu,⁶⁸ K. Yi,⁶⁸
 C. C. Young,⁶⁸ P. R. Burchat,⁶⁹ A. J. Edwards,⁶⁹ S. A. Majewski,⁶⁹ B. A. Petersen,⁶⁹ L. Wilden,⁶⁹ S. Ahmed,⁷⁰
 M. S. Alam,⁷⁰ R. Bula,⁷⁰ J. A. Ernst,⁷⁰ V. Jain,⁷⁰ B. Pan,⁷⁰ M. A. Saeed,⁷⁰ F. R. Wappler,⁷⁰ S. B. Zain,⁷⁰
 W. Bugg,⁷¹ M. Krishnamurthy,⁷¹ S. M. Spanier,⁷¹ R. Eckmann,⁷² J. L. Ritchie,⁷² A. M. Ruland,⁷² C. J. Schilling,⁷²
 R. F. Schwitters,⁷² J. M. Izen,⁷³ X. C. Lou,⁷³ S. Ye,⁷³ F. Bianchi,⁷⁴ F. Gallo,⁷⁴ D. Gamba,⁷⁴ M. Pelliccioni,⁷⁴
 M. Bomben,⁷⁵ L. Bosisio,⁷⁵ C. Cartaro,⁷⁵ F. Cossutti,⁷⁵ G. Della Ricca,⁷⁵ L. Lanceri,⁷⁵ L. Vitale,⁷⁵ V. Azzolini,⁷⁶
 N. Lopez-March,⁷⁶ F. Martinez-Vidal,⁷⁶ D. A. Milanese,⁷⁶ A. Oyangueren,⁷⁶ J. Albert,⁷⁷ Sw. Banerjee,⁷⁷ B. Bhuyan,⁷⁷
 K. Hamano,⁷⁷ R. Kowalewski,⁷⁷ I. M. Nugent,⁷⁷ J. M. Roney,⁷⁷ R. J. Sobie,⁷⁷ J. J. Back,⁷⁸ P. F. Harrison,⁷⁸
 T. E. Latham,⁷⁸ G. B. Mohanty,⁷⁸ M. Pappagallo,^{78,§} H. R. Band,⁷⁹ X. Chen,⁷⁹ S. Dasu,⁷⁹ K. T. Flood,⁷⁹
 J. J. Hollar,⁷⁹ P. E. Kutter,⁷⁹ Y. Pan,⁷⁹ M. Pierini,⁷⁹ R. Prepost,⁷⁹ S. L. Wu,⁷⁹ Z. Yu,⁷⁹ and H. Neal⁸⁰

(The BABAR Collaboration)

¹Laboratoire de Physique des Particules, IN2P3/CNRS et Université de Savoie, F-74941 Annecy-Le-Vieux, France

²Universitat de Barcelona, Facultat de Física, Departament ECM, E-08028 Barcelona, Spain

³Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

⁴University of Bergen, Institute of Physics, N-5007 Bergen, Norway

⁵Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

⁶University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁷Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁸University of Bristol, Bristol BS8 1TL, United Kingdom

⁹University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

¹⁰Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹¹Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹²University of California at Irvine, Irvine, California 92697, USA

¹³University of California at Los Angeles, Los Angeles, California 90024, USA

¹⁴University of California at Riverside, Riverside, California 92521, USA

¹⁵University of California at San Diego, La Jolla, California 92093, USA

¹⁶University of California at Santa Barbara, Santa Barbara, California 93106, USA

¹⁷University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

¹⁸California Institute of Technology, Pasadena, California 91125, USA

¹⁹University of Cincinnati, Cincinnati, Ohio 45221, USA

²⁰University of Colorado, Boulder, Colorado 80309, USA

²¹Colorado State University, Fort Collins, Colorado 80523, USA

²²Universität Dortmund, Institut für Physik, D-44221 Dortmund, Germany

²³Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany

²⁴Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, F-91128 Palaiseau, France

²⁵University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom

²⁶Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy

²⁷Laboratori Nazionali di Frascati dell’INFN, I-00044 Frascati, Italy

- ²⁸ *Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy*
- ²⁹ *Harvard University, Cambridge, Massachusetts 02138, USA*
- ³⁰ *Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany*
- ³¹ *Imperial College London, London, SW7 2AZ, United Kingdom*
- ³² *University of Iowa, Iowa City, Iowa 52242, USA*
- ³³ *Iowa State University, Ames, Iowa 50011-3160, USA*
- ³⁴ *Johns Hopkins University, Baltimore, Maryland 21218, USA*
- ³⁵ *Universität Karlsruhe, Institut für Experimentelle Kernphysik, D-76021 Karlsruhe, Germany*
- ³⁶ *Laboratoire de l'Accélérateur Linéaire, IN2P3/CNRS et Université Paris-Sud 11, Centre Scientifique d'Orsay, B. P. 34, F-91898 ORSAY Cedex, France*
- ³⁷ *Lawrence Livermore National Laboratory, Livermore, California 94550, USA*
- ³⁸ *University of Liverpool, Liverpool L69 7ZE, United Kingdom*
- ³⁹ *Queen Mary, University of London, E1 4NS, United Kingdom*
- ⁴⁰ *University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom*
- ⁴¹ *University of Louisville, Louisville, Kentucky 40292, USA*
- ⁴² *University of Manchester, Manchester M13 9PL, United Kingdom*
- ⁴³ *University of Maryland, College Park, Maryland 20742, USA*
- ⁴⁴ *University of Massachusetts, Amherst, Massachusetts 01003, USA*
- ⁴⁵ *Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA*
- ⁴⁶ *McGill University, Montréal, Québec, Canada H3A 2T8*
- ⁴⁷ *Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy*
- ⁴⁸ *University of Mississippi, University, Mississippi 38677, USA*
- ⁴⁹ *Université de Montréal, Physique des Particules, Montréal, Québec, Canada H3C 3J7*
- ⁵⁰ *Mount Holyoke College, South Hadley, Massachusetts 01075, USA*
- ⁵¹ *Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy*
- ⁵² *NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands*
- ⁵³ *University of Notre Dame, Notre Dame, Indiana 46556, USA*
- ⁵⁴ *Ohio State University, Columbus, Ohio 43210, USA*
- ⁵⁵ *University of Oregon, Eugene, Oregon 97403, USA*
- ⁵⁶ *Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy*
- ⁵⁷ *Laboratoire de Physique Nucléaire et de Hautes Energies, IN2P3/CNRS, Université Pierre et Marie Curie-Paris6, Université Denis Diderot-Paris7, F-75252 Paris, France*
- ⁵⁸ *University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*
- ⁵⁹ *Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy*
- ⁶⁰ *Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy*
- ⁶¹ *Prairie View A&M University, Prairie View, Texas 77446, USA*
- ⁶² *Princeton University, Princeton, New Jersey 08544, USA*
- ⁶³ *Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy*
- ⁶⁴ *Universität Rostock, D-18051 Rostock, Germany*
- ⁶⁵ *Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom*
- ⁶⁶ *DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France*
- ⁶⁷ *University of South Carolina, Columbia, South Carolina 29208, USA*
- ⁶⁸ *Stanford Linear Accelerator Center, Stanford, California 94309, USA*
- ⁶⁹ *Stanford University, Stanford, California 94305-4060, USA*
- ⁷⁰ *State University of New York, Albany, New York 12222, USA*
- ⁷¹ *University of Tennessee, Knoxville, Tennessee 37996, USA*
- ⁷² *University of Texas at Austin, Austin, Texas 78712, USA*
- ⁷³ *University of Texas at Dallas, Richardson, Texas 75083, USA*
- ⁷⁴ *Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy*
- ⁷⁵ *Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy*
- ⁷⁶ *IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain*
- ⁷⁷ *University of Victoria, Victoria, British Columbia, Canada V8W 3P6*
- ⁷⁸ *Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom*
- ⁷⁹ *University of Wisconsin, Madison, Wisconsin 53706, USA*
- ⁸⁰ *Yale University, New Haven, Connecticut 06511, USA*

(Dated: August 7, 2007)

We present an analysis of inclusive Ω_c^0 baryon production and decays in 230.5 fb^{-1} of data recorded with the BABAR detector. Ω_c^0 baryons are reconstructed in four final states ($\Omega^- \pi^+$, $\Omega^- \pi^+ \pi^0$, $\Omega^- \pi^+ \pi^+ \pi^-$, $\Xi^- K^- \pi^+ \pi^+$) and the corresponding ratios of branching fractions are measured. We also measure the momentum spectrum in the e^+e^- center-of-mass frame. From the spectrum, we observe Ω_c^0 production from B decays and in $c\bar{c}$ events, and extract the two rates of production.

The Ω_c^0 ($c\bar{s}s$) is the heaviest weakly-decaying singly-charmed baryon. It has been observed independently in several decay modes by different experiments [1] and in a variety of production environments, including e^+e^- colliders operating at the $\Upsilon(4S)$ resonance [2–4], photoproduction [5–7], and hyperon beams [8]. So far, B meson decays to Ω_c^0 have not been observed. Several different mechanisms could contribute, principally weak decays of the following forms: $b \rightarrow c\bar{c}s$ (e.g., $B^- \rightarrow \Omega_c^0\Xi_c^-$); $b \rightarrow c\bar{u}s$ (e.g., $B^- \rightarrow \Omega_c^0\bar{\Sigma}^-$); and $b \rightarrow c\bar{u}d$ (e.g., $B^- \rightarrow \Omega_c^0\Xi_c^0\pi^-$). Beyond the requirement to produce at least one $s\bar{s}$ pair during fragmentation, we would expect these three types of decays to be further suppressed by the limited phase space, by $|V_{us}|^2$, and by needing to produce a second $s\bar{s}$ pair, respectively. Theoretical predictions for branching fractions of individual two-body contributions vary from $\mathcal{O}(10^{-5})$ to $\mathcal{O}(10^{-3})$ [9–11].

In this letter, we present a study of the Ω_c^0 baryon, reconstructed in four decay modes: $\Omega^- \pi^+$, $\Omega^- \pi^+ \pi^0$, $\Omega^- \pi^+ \pi^+ \pi^-$, and $\Xi^- K^- \pi^+ \pi^+$ [12]. We measure the ratios of branching fractions for these modes, normalizing to $\mathcal{B}(\Omega_c^0 \rightarrow \Omega^- \pi^+)$. The previous most precise measurements of these ratios are from an analysis of approximately 45 events from six Ω_c^0 decay modes [3]. We then measure the spectrum of the Ω_c^0 momentum in the e^+e^- center-of-mass frame (p^*) and observe significant production of Ω_c^0 baryons in the decays of B mesons.

The data for this analysis were recorded with the BABAR detector at the Stanford Linear Accelerator Center PEP-II asymmetric-energy e^+e^- collider. The detector is described in detail elsewhere [13]. A total integrated luminosity of 230.5 fb^{-1} is used, of which 208.9 fb^{-1} were collected at the $\Upsilon(4S)$ resonance (corresponding to 232 million $B\bar{B}$ pairs) and 21.6 fb^{-1} were collected 40 MeV below the $B\bar{B}$ production threshold.

Simulated events with the Ω_c^0 decaying into the relevant final states are generated for the processes $e^+e^- \rightarrow c\bar{c} \rightarrow \Omega_c^0 X$ and $e^+e^- \rightarrow \Upsilon(4S) \rightarrow B\bar{B} \rightarrow \Omega_c^0 X$, where X represents the rest of the event. The PYTHIA simulation package [14] is used for the $c\bar{c}$ fragmentation and for B decays to Ω_c^0 , and the GEANT4 [15] package is used to simulate the detector response. To investigate possible background contributions, additional samples of generic Monte Carlo (MC) events are used, equivalent to 990 fb^{-1} for $\Upsilon(4S)$ events ($e^+e^- \rightarrow \Upsilon(4S) \rightarrow B\bar{B}$), plus 320 fb^{-1} for $c\bar{c}$ continuum events ($e^+e^- \rightarrow c\bar{c}$) and 340 fb^{-1} for light quark continuum events ($e^+e^- \rightarrow q\bar{q}$, $q = u, d, s$).

The reconstruction of an Ω_c^0 candidate begins by identifying a proton, combining it with an oppositely charged track interpreted as a π^- , and fitting the tracks to a common vertex to form a Λ candidate. The Λ is then combined with a negatively charged track interpreted as a

K^- (π^-) and fit to a common vertex to form an Ω^- (Ξ^-) candidate. For each intermediate hyperon (Λ , Ξ^- , Ω^-) we require the invariant mass to be within $4.5 \text{ MeV}/c^2$ of its nominal value (corresponding to approximately 4, 3, and 3 times the detector resolution, respectively). We form π^0 candidates from pairs of photons in the electromagnetic calorimeter, requiring the energy of each photon to be above 80 MeV and the combined energy to be above 200 MeV. We require the invariant mass of the π^0 candidate, computed at the event primary vertex, to be in the range 120–150 MeV/c^2 .

Each Ω^- (Ξ^-) candidate that passes the requirements is then combined with one or three additional tracks that are identified as pions or kaons as appropriate. For the $\Omega^- \pi^+ \pi^0$ final state, we also combine the hyperon and π^+ with a π^0 . The Ω_c^0 candidate daughters are refit to a common vertex with their masses constrained to the nominal values. From this fit we extract the decay vertices and associated uncertainties of the Ω_c^0 and the intermediate hyperons, the four-momenta of the particles, and the Ω_c^0 candidate mass. For each intermediate hyperon we require a positive scalar product of the momentum vector in the laboratory frame and the displacement vector from its production vertex to its decay vertex.

To further suppress the background, we compute the likelihood ratio $\mathcal{L} = \prod_i p_i^S(x_i) / [\prod_i p_i^S(x_i) + \prod_i p_i^B(x_i)]$ for each Ω_c^0 candidate, where the index i refers to the likelihood variables x_i , and $p_i(x_i)$ are the probability density functions for signal (S) and background (B). For a given Ω_c^0 candidate \mathcal{L} has a value between 0 and 1. The likelihood variables x_i are the logarithm of the Ω^- or Ξ^- decay length significance, which is defined as the distance between the production and decay vertices divided by the uncertainty on that distance; the momentum of the Ω^- or Ξ^- in the e^+e^- rest frame; the total momentum of the mesons recoiling against the Ω^- or Ξ^- in the e^+e^- rest frame; and, for the $\Omega^- \pi^+ \pi^0$ mode, the π^0 momentum in the laboratory frame. These variables (particularly the decay length significance) cover the expected range effectively with a limited number of bins. The distributions of these variables for the signal hypothesis are derived from signal MC simulations, and for the background hypothesis from generic MC events in which contributions from real Ω_c^0 are excluded. Separate distributions are used for each final state when measuring ratios of branching fractions, and for each momentum range when measuring the momentum spectrum.

To measure the ratios of branching fractions, we require that $p^* > 2.4 \text{ GeV}/c$ in order to suppress combinatoric background. Since the kinematic limit for Ω_c^0 produced in B decays at BABAR is $p_{\text{max}}^* = 2.02 \text{ GeV}/c$, only Ω_c^0 produced in the $c\bar{c}$ continuum are retained. We also require that the value of \mathcal{L} for each candidate is greater

than a threshold \mathcal{L}_0 , chosen to maximize the expected signal significance for a given final state based on simulated events. We perform an unbinned maximum likelihood fit to the mass distributions shown in Fig. 1. The signal lineshape is parameterized as the sum of two Gaussian functions with a common mean; the background is parameterized as a first-order polynomial. In the fits to the data, the signal yield is a free parameter; the widths and relative amplitudes of the two Gaussian functions are fixed to values determined from a fit to simulated signal events. The mean mass is also a free parameter, except for the $\Xi^- K^- \pi^+ \pi^+$ final state where we fix it to the central value obtained in $\Omega_c^0 \rightarrow \Omega^- \pi^+$ in order to ensure proper fit convergence. The masses are found to be consistent with one another and with the current world average [1] within uncertainties.

The numbers of signal events are 177 ± 16 , 64 ± 15 , 25 ± 8 , and 45 ± 12 (statistical uncertainties only) for the final states $\Omega^- \pi^+$, $\Omega^- \pi^+ \pi^0$, $\Omega^- \pi^+ \pi^+ \pi^-$, and $\Xi^- K^- \pi^+ \pi^+$, respectively. These correspond to statistical significances of 18, 5.1, 4.2, and 4.3 standard deviations, respectively, where the significance is defined as $\sqrt{2\Delta\ell}$ and $\Delta\ell$ is the change in the logarithm of the likelihood between the fits with and without an Ω_c^0 signal component. The fitted yields are then corrected for efficiency, which is defined as the fraction of simulated signal events, generated in the appropriate p^* range, that are reconstructed and pass all selection criteria. Including the loss of efficiency due to the Λ and Ω^- branching fractions, we obtain efficiencies of $(8.6 \pm 0.6)\%$, $(2.5 \pm 0.3)\%$, $(4.3 \pm 0.4)\%$, and $(4.7 \pm 0.5)\%$ for the four final states, where the uncertainties include systematic effects and are partially correlated. The systematic uncertainties on, and corrections to, the ratios of branching fractions are listed in Table I and discussed further later. We measure the ratios to be

$$\frac{\mathcal{B}(\Omega_c^0 \rightarrow \Omega^- \pi^+ \pi^0)}{\mathcal{B}(\Omega_c^0 \rightarrow \Omega^- \pi^+)} = 1.27 \pm 0.31(\text{stat}) \pm 0.11(\text{syst}),$$

$$\frac{\mathcal{B}(\Omega_c^0 \rightarrow \Omega^- \pi^+ \pi^+ \pi^-)}{\mathcal{B}(\Omega_c^0 \rightarrow \Omega^- \pi^+)} = 0.28 \pm 0.09(\text{stat}) \pm 0.01(\text{syst}),$$

$$\frac{\mathcal{B}(\Omega_c^0 \rightarrow \Xi^- K^- \pi^+ \pi^+)}{\mathcal{B}(\Omega_c^0 \rightarrow \Omega^- \pi^+)} = 0.46 \pm 0.13(\text{stat}) \pm 0.03(\text{syst}).$$

We also measure the p^* spectrum of Ω_c^0 in order to study the production rates in both $c\bar{c}$ and $B\bar{B}$ events. Only the $\Omega^- \pi^+$ final state is used. The same reconstruction, optimization of selection criteria, and fitting procedures described above are applied, except that no requirement on p^* is made. Instead, the Ω_c^0 candidates are divided into nine equal intervals of p^* covering the range 0.0–4.5 GeV/c. We again require $\mathcal{L} > \mathcal{L}_0$ and compute the efficiency in each p^* interval as before with simulated signal events. In the numerator of the efficiency we count events with measured p^* in the appropriate interval, and in the denominator we count events with generated p^* in that interval: this definition removes the slight broaden-




FIG. 1: The invariant mass spectra for candidates passing the selection criteria. The data are fit with a double Gaussian lineshape on a linear background.

TABLE I: Systematic uncertainties on the ratios of branching fractions, where $R_1 \equiv \mathcal{B}(\Omega_c^0 \rightarrow \Omega^- \pi^+ \pi^0)/\mathcal{B}(\Omega_c^0 \rightarrow \Omega^- \pi^+)$, $R_2 \equiv \mathcal{B}(\Omega_c^0 \rightarrow \Omega^- \pi^+ \pi^+ \pi^-)/\mathcal{B}(\Omega_c^0 \rightarrow \Omega^- \pi^+)$, and $R_3 \equiv \mathcal{B}(\Omega_c^0 \rightarrow \Xi^- K^- \pi^+ \pi^+)/\mathcal{B}(\Omega_c^0 \rightarrow \Omega^- \pi^+)$.

Effect	R_1	R_2	R_3
Finite MC sample size	0.7%	0.7%	0.8%
Intermediate resonances in Ω_c^0 decay	1.3%	2.6%	3.7%
Signal lineshape	1.0%	1.0%	1.0%
Dependence on the fit procedure	1.5%	1.5%	1.5%
Hyperon branching fractions	—	—	1.0%
Particle identification efficiency	1.0%	1.0%	1.0%
Tracking efficiency	0.0%	2.8% ^a	2.8% ^a
p^* spectrum mismodeling	1.5%	0.6%	3.5%
π^0 fitting and efficiency	7.8% ^b	—	—
Total systematic uncertainty	8.3%	4.4%	6.3%

^aA relative correction of +0.5% applies to R_2 and R_3 .

^bA relative correction of +1.1% applies to R_1 .




FIG. 2: The background-subtracted and efficiency-corrected $\Omega_c^0 p^*$ spectrum. The black points represent the data, with vertical error bars giving the sum in quadrature of statistical and uncorrelated systematic uncertainties. The solid histogram shows the Bowler fragmentation function, binned and fit to the data for $p^* > 2$ GeV/c (vertical, dashed line).

ing effect of the detector momentum resolution. We also take into account a small difference in efficiency between $c\bar{c}$ and $B\bar{B}$ events. The efficiency-corrected yield in each p^* interval is shown in Fig. 2.

The systematic uncertainties are divided into two categories: normalization effects, which are treated as fully correlated between all p^* intervals, and shape effects, which are treated as uncorrelated between different p^* intervals. The normalization uncertainties are due to the mass resolution, which is determined from the MC and checked with studies of the control modes $\Xi_c^0 \rightarrow \Xi^- \pi^+$ and $\Xi_c^+ \rightarrow \Xi^- \pi^+ \pi^+$ (2.4%); the Λ and Ω^- branching fractions [1] (1.3%); and the tracking efficiency, which is corrected for data/MC discrepancies with control samples of τ decays (5.9%). The shape uncertainties are due to the limited size of MC samples ($< 1\%$); dependence on the fit procedure (1.5%); modeling of the p^* spectrum, which can affect the weighted average efficiency within a p^* bin (0–6%); the signal lineshape parameterization (1.0%); and the particle identification efficiency (2.0%). When fitting fragmentation functions (see below), we consider only the statistical and shape uncertainties, added in quadrature. When quoting total yields and rates, we include the normalization uncertainties, along with a relative correction of +1.0% due to a known data/MC discrepancy in tracking efficiency.

The double-peak structure seen in the p^* spectrum is due to two production mechanisms: the peak at lower p^* is due to Ω_c^0 production in B meson decays and the peak at higher p^* is due to Ω_c^0 production from the $c\bar{c}$ continuum. This is consistent with the pattern observed in Λ_c^+ and Ξ_c^0 spectra measured for e^+e^- annihilation at $\sqrt{s} = 10.6$ GeV [16–18]. We fit the p^* spectrum with the

Bowler fragmentation function [19] for $p^* > 2$ GeV/c. We then extract the continuum yield as the sum of the data points above 2 GeV/c plus the integral of the extrapolated function below 2 GeV/c. Similarly, the yield from B decays is the sum of the data points below 2 GeV/c minus the integral of the extrapolated function below 2 GeV/c. Note that we do not fit a fragmentation function to the data below 2 GeV/c. We obtain yields of 2583 ± 289 and 2426 ± 414 for Ω_c^0 produced in the continuum and in B decays, respectively, where the uncertainty includes all statistical and experimental effects. An additional model uncertainty arises from the extrapolation of the continuum tail for $p^* < 2$ GeV/c. To estimate this, we repeat the p^* spectrum fit and yield measurement with other fragmentation functions: Collins and Spiller (CS) [20], two versions of the phenomenological model of Kartvelishvili *et al.* (KLP-M and KLP-B) [21, 22] and the Peterson model [23]. The CS and KLP-M fits are inconsistent with the data for $p^* > 2$ GeV/c. The RMS of the yields from the three other fits is 240 events and is taken as the model uncertainty for the B and continuum Ω_c^0 yields. Dividing the Ω_c^0 yield in B decays by the total number of B mesons in the data sample, we obtain the branching fraction product $\mathcal{B}(B \rightarrow \Omega_c^0 X) \mathcal{B}(\Omega_c^0 \rightarrow \Omega^- \pi^+) = [5.2 \pm 0.9 (\text{exp}) \pm 0.5 (\text{model})] \times 10^{-6}$, where X represents the rest of the B meson decay products. Dividing the Ω_c^0 yield from the continuum by the integrated luminosity and correcting for the small variation in cross-section with \sqrt{s} , we obtain the cross-section product at $\sqrt{s} = 10.58$ GeV: $\sigma(e^+e^- \rightarrow \Omega_c^0 X) \mathcal{B}(\Omega_c^0 \rightarrow \Omega^- \pi^+) = [11.2 \pm 1.3 (\text{exp}) \pm 1.0 (\text{model})]$ fb, where X represents the rest of the event. As a cross-check, we also make model-independent estimates of the yields from the continuum and from B decays by subtracting the data below the $\Upsilon(4S)$ threshold. Within large uncertainties, these are consistent with the yields measured above.

It is thus clear that decays of B mesons to Ω_c^0 occur at a significant rate. Assuming the absolute branching fraction $\mathcal{B}(\Omega_c^0 \rightarrow \Omega^- \pi^+) \sim 1\%$, we conclude that $\mathcal{B}(B \rightarrow \Omega_c^0 X) \sim \text{few} \times 10^{-4}$. This is substantially lower than the inclusive B meson branching fractions to the charmed baryons Λ_c^+ and Ξ_c , which are $\sim \text{few} \times 10^{-2}$ [16–18]. One possible explanation for this is that both Λ_c^+ and Ξ_c can be produced in a $b \rightarrow c\bar{c}s$ transition without creating an $s\bar{s}$ pair from the vacuum, whereas at least one $s\bar{s}$ pair must be created for Ω_c^0 production. It is also possible that phase space suppression in B decays to baryons becomes significant when very close to threshold.

In conclusion, we have studied the Ω_c^0 baryon at *BABAR* through four hadronic decay modes, using 230.5 fb^{-1} of data. We measure the ratios of branching fractions for four modes, significantly improving upon the previous values [3]. We have also measured the p^* spectrum and found comparable production rates of Ω_c^0 baryons from the continuum and from B meson decays. The inclusive B branching fraction to Ω_c^0 is found to be substantially

lower than those to Ξ_c^0 and Λ_c^+ baryons, assuming the relevant baryon weak decay branching fractions are of the same order of magnitude.

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), MEC (Spain), and PPARC (United Kingdom). Individuals have received support from the Marie Curie EIF (European Union) and the A. P. Sloan Foundation.

* Deceased

† Also with Università di Perugia, Dipartimento di Fisica, Perugia, Italy

‡ Also with Università della Basilicata, Potenza, Italy

§ Also with IPPP, Physics Department, Durham University, Durham DH1 3LE, United Kingdom

- [1] Particle Data Group, W.-M. Yao *et al.*, *J. Phys. G* **33**, 1 (2006).
- [2] ARGUS Collaboration, H. Albrecht *et al.*, *Phys. Lett. B* **288**, 367 (1992).
- [3] CLEO Collaboration, D. Cronin-Hennessy *et al.*, *Phys. Rev. Lett.* **86**, 3730 (2001).
- [4] CLEO Collaboration, R. Ammar *et al.*, *Phys. Rev. Lett.* **89**, 171803 (2002).
- [5] E687 Collaboration, P. L. Frabetti *et al.*, *Phys. Lett. B* **300**, 190 (1993).
- [6] E687 Collaboration, P. L. Frabetti *et al.*, *Phys. Lett. B* **338**, 106 (1994).
- [7] FOCUS Collaboration, J. M. Link *et al.*, *Phys. Lett. B* **561**, 41 (2003).
- [8] WA89 Collaboration, M. I. Adamovich *et al.*, *Phys. Lett. B* **358**, 151 (1995).
- [9] P. Ball and H. G. Dosch, *Z. Phys. C* **51**, 445 (1991).
- [10] S. M. Sheikholeslami and M. P. Khanna, *Phys. Rev. D* **44**, 770 (1991).
- [11] V. L. Chernyak and I. R. Zhitnitsky, *Nucl. Phys.* **B345**, 137 (1990).
- [12] The use of charge conjugate modes is implied throughout.
- [13] *BABAR* Collaboration, B. Aubert *et al.*, *Nucl. Instr. Methods Phys. Res., Sect. A* **479**, 1 (2002).
- [14] T. Sjöstrand *et al.*, *Comput. Phys. Commun.* **135**, 238 (2001).
- [15] S. Agostinelli *et al.*, *Nucl. Instr. Methods Phys. Res., Sect. A* **506**, 250 (2003).
- [16] Belle Collaboration, R. Seuster *et al.*, *Phys. Rev. D* **73**, 032002 (2006).
- [17] *BABAR* Collaboration, B. Aubert *et al.*, *Phys. Rev. D* **75**, 012003 (2007).
- [18] *BABAR* Collaboration, B. Aubert *et al.*, *Phys. Rev. Lett.* **95**, 142003 (2005).
- [19] M.G. Bowler, *Z. Phys.* **C11**, 169 (1981).
- [20] P.D.B. Collins and T.P. Spiller, *J. Phys. G* **11**, 1289 (1985).
- [21] V.G. Kartvelishvili, A.K. Likhoded, and V.A. Petrov, *Phys. Lett. B* **78**, 615 (1978).
- [22] V.G. Kartvelishvili and A.K. Likhoded, *Sov. J. Nucl. Phys.* **29**, 390 (1979).
- [23] C. Peterson, D. Schlatter, I. Schmitt, and P.M. Zerwas, *Phys. Rev. D* **27**, 105 (1983).