

SLAC-PUB-1397
(T)
March 1974

REMARKS ON THE PHYSICAL DEGREES OF FREEDOM IN
TWO-DIMENSIONAL ELECTRODYNAMICS *

Jorge F. Willemsen
Stanford Linear Accelerator Center
Stanford University, Stanford, Calif. 94305

ABSTRACT

A simple argument is advanced for how it happens that two-dimensional electrodynamics is a theory of massive spinless bosons.

(Submitted to Phys. Rev. D. Comments and Addenda)

* Work supported by the U.S. Atomic Energy Commission.

It is well-known that, once the computational dust has settled, two dimensional quantum electrodynamics (TDED) collapses to a theory of a massive spinless non-interacting Bose field. Sophisticated arguments for why this occurs have been presented by Lowenstein and Swieca.¹ The purpose of this note is to supply a simple way of seeing why it is so.

The first observation required is that, in the gauge $A_1(x, t) = 0$, the interaction reduces to a self-interaction of the charge density via the (two-dimensional) Coulomb potential,²

$$\mathcal{L} = i \bar{\psi} \not{\partial} \psi + \frac{e^2}{4} j^0(x, t) \int dy |x - y| j^0(y, t) \quad (1)$$

Next, as is known from work on the Thirring model,³ the free Dirac theory in two dimensions may equivalently be discussed in terms of the associated vector current

$$j^\mu(x, t) = : \bar{\psi} \gamma^\mu \psi : \quad ; \quad (2)$$

and the symmetric, traceless, conserved tensor operator constructed from the current,

$$T^{\mu\nu}(x, t) = \frac{\pi}{2} \left(\left| j^\mu, j^\nu \right| - g^{\mu\nu} j_\lambda j^\lambda \right) . \quad (3)$$

That is, $H = \int dx T^{00}$ generates the free field equation of motion for ψ , given the anti-commutator $\left\{ \psi(x, t), \psi^\dagger(y, t) \right\} = \delta(x - y)$, and the definitions Eqs. (2) and (3). The result is not obvious, however, and depends for its demonstration on the operator equation³

$$\partial_x \psi(x, z) = \frac{i\pi}{2} \left[j^1 + \gamma^5 j^0, \psi \right] , \quad (4)$$

which is true in two dimensions.

It is possible to proceed by postulating a set of commutation relations satisfied by j^μ with itself, and with ψ .^{4,5} For the free theory, this is unnecessary. The required "current algebra" can be derived, and is

$$\left[j_0(x, t), j_0(y, t) \right] = 0 \quad ; \quad (5a)$$

$$\left[j_1(x, t), j_1(y, t) \right] = 0 \quad ; \quad (5b)$$

$$\left[j_0(x, t), j_1(y, t) \right] = \frac{i}{\pi} \partial_x \delta(x - y) \quad . \quad (5c)$$

This current algebra is solved by setting⁶

$$\sqrt{\pi} \ j_\mu(x, t) = \epsilon_{\mu\nu} \partial^\nu \tilde{\Phi}(x, t) \quad , \quad (6)$$

where $\tilde{\Phi}$ is a canonical (pseudo) scalar field. Then $T^{\mu\nu}$ is the canonical energy momentum tensor for this massless field. A consistent choice for an associated Lagrange density is

$$\mathcal{L}^{(0)} = 1/2 (\partial_\mu \tilde{\Phi}) (\partial^\mu \tilde{\Phi}) \quad (7)$$

[Note $j_\mu = \partial_\mu \Phi$ is also a possible choice. However, current conservation demands Φ is a massless free field, but places no constraints on $\tilde{\Phi}$.]

The result essential to our argument is that, using Eqs. (4) and (6), one can show

$$H_0 = -i \bar{\psi}_0 \gamma^1 \partial_1 \psi_1 = \frac{1}{2} \left[(\partial_0 \tilde{\Phi})^2 + (\partial_1 \tilde{\Phi})^2 \right] \quad , \quad (8)$$

where ψ_0 is the free Dirac field in two dimensions. This amazing relation is expected to be true only in two dimensions, and is a result of the fact Eq. (4)

reduces trilinears in ψ to a single ψ . Using this equation, in interaction representation Eq. (1) becomes

$$\begin{aligned} \mathcal{L} &= 1/2 (\partial_\mu \tilde{\Phi}) (\partial^\mu \tilde{\Phi}) + \frac{e^2}{4\pi} \int dy |x - y| \partial_x \tilde{\Phi}(x, t) \partial_y \tilde{\Phi}(y, t) \\ &\longrightarrow 1/2 (\partial_\mu \tilde{\Phi}) (\partial^\mu \tilde{\Phi}) - \frac{e^2}{2\pi} \tilde{\Phi}^2 . \end{aligned} \quad (9)$$

But this just describes a massive (pseudo) scalar field, of mass $\mu^2 = e^2/\pi$ and nothing else; this was the desired result.

Admittedly, we have seemed cavalier in obtaining this result, paying little attention to defining the various operators we have introduced with any rigour, and substituting free theory equations into the interacting theory. In fact, however, a rigorous momentum space analysis can be carried out. This analysis verifies the conclusions stated above.

As an example, consider TDED in the finite spatial interval $[0, \pi]$. The use of a finite interval may be viewed as an alternative to Klaiber's procedure for regularizing the bad infrared behaviour of the theory.^{1, 7, 8} The particular interval chosen is a matter of convenience. Any interval of arbitrary length L may be adopted. Taking the limit $L \rightarrow \infty$ at the end of the calculation turns momentum sums into momentum integrals.

It has been shown recently⁹ that

$$\begin{aligned} H &= \int_0^\pi dx \left[\psi^\dagger \partial_0 \psi - \mathcal{L} \right] \\ &= \sum_{n=1}^{\infty} (n - 1/2) (b_n^\dagger b_n + c_n^\dagger c_n) + (\mu^2/4) \sum_{p=1}^{\infty} p^{-1} \left[\rho^\dagger \rho^\dagger + \rho \rho + 2\rho^\dagger \rho \right]_p , \end{aligned} \quad (10)$$

where \mathcal{L} is given by Eq. (1) ; and

$$\rho(p) = p^{-1/2} \int_0^\pi dx \left[j^0(x, 0) \cos px - i j^1(x, 0) \sin px \right] , \quad (11)$$

satisfy Bose commutation relations $[\rho(p), \rho^+(q)] = \delta_{p,q}$. This Hamiltonian may be diagonalized by means of a Bogoliubov transformation,

$$\tilde{H} = e^{iS} H e^{-iS} \quad (12)$$

$$= (H_0 - T) + \sum_{p=1}^{\infty} E(p) \rho^+(p) \rho(p) + E_0 ,$$

where H_0 is the free Dirac Hamiltonian, and

$$S = \left(-\frac{i}{2}\right) \sum_{p=1}^{\infty} \left[\tanh^{-1} \left(\frac{\mu^2}{\mu^2 + p^2} \right) \right] \left[\rho^+ \rho^+ - \rho \rho \right]_p ; \quad (13a)$$

$$T = \sum_{p=1}^{\infty} p \rho^+(p) \rho(p) ; \quad (13b)$$

$$E(p) = \sqrt{\mu^2 + p^2} ; \quad (13c)$$

$$E_0 = \frac{1}{2} \sum_{p=1}^{\infty} \left[E(p) - p - \left(\frac{\mu^2}{2p} \right) \right] . \quad (13d)$$

The analogue of Eq. (8) in this case is¹⁰

$$H_0 = \frac{Q^2}{2} + T , \quad (14)$$

where the conserved charge $Q = \int dx j^0$. This equation may be verified by explicit, though tedious calculation, using identities which follow from Eq. (4).

Fourier analyzing Eq. (4), one obtains

$$Q b_k + \sum_{p=1}^{\infty} p \rho_p^+ b_{k+p} + \sum_{p=1}^{\infty} \sqrt{p+k-1} c_p^+ \rho_{p+k-1} + \sum_{p=1}^{k-1} \sqrt{p} \rho_p b_{k-p} = 0 ; \quad (15)$$

and an accompanying equation for c obtained by interchanging $b \leftrightarrow c$ where they appear explicitly in Eq. (15).

The validity of Eq. (15) as an operator equation may be verified explicitly by brute force. As mentioned earlier, the key to why this relation is possible is that ρ and ρ^+ are bilinear in fermion operators, hence many terms anti-commute to zero under summation.⁸

Implicit in Eq. (10) was the consistency requirement for TDED in axial gauge,

$$Q |\Phi_{\text{phys}}\rangle = 0 \quad . \quad (16)$$

This constraint follows, essentially, because the vector current is conserved, but the divergence of the axial current contains an anomaly. However, these currents are related by $a^\mu = \epsilon^{\mu\nu} j_\nu$. Consistency demands (in this gauge) that Eq. (16) be true.¹¹

Combining Eqs. (12), (14), and (16)

$$\tilde{H} = E_0 + \sum_{p=1}^{\infty} E(p) \rho^+(p) \rho(p) \quad . \quad (17)$$

All reference to the fermions has disappeared. This is a property of the solution independent of our choice of interval, and of boundary conditions in the interval.¹² As the size of the system $L \rightarrow \infty$, E_0 diverges logarithmically. This is the only remnant of the infrared problem.

To summarize, one first casts TDED as a theory of self-interacting fermions, by choosing a convenient gauge. This done, one attempts to represent as much of the theory as possible in terms of the currents, based on the experience with the Thirring model that these are the only genuine observables.^{3,4,5} Unlike the Thirring model, however, in which ψ preserves a role as an intertwining

operator between inequivalent irreducible representations of the current algebra, the vanishing of the charge in TDED deprives ψ of even this role. Indeed, the presence of massive excitations follows trivially from $(\square + \mu^2) j^\mu = 0$. The point being made is that ψ can be eliminated entirely from the problem. In the language of Ref. 12, there are no "quasi-particles", only plasmons.

I thank Professor T. M. Yan for comments on the manuscript.

REFERENCES

1. J. H. Lowenstein and J. A. Swieca, Ann. Phys. (N.Y.) 68, 172 (1971).
2. L. S. Brown, Nuovo Cimento 29, 617 (1963).
3. C. M. Sommerfield, Phys. Rev. 176, 2019 (1968).
4. K. Johnson, Nuovo Cimento 20, 773 (1961).
5. G. F. Dell' Antonio, Y. Frishman, and D. Zwanziger, Phys. Rev. D6, 988 (1972).
6. C. G. Callan, R. F. Dashen, and D. H. Sharp, Phys. Rev. 165 1883 (1968).
7. B. Klaiber, Lectures in Theoretical Physics, eds. A. O. Barnt and W. E. Brittin (Gordon and Breach, N. Y., 1968), Vol. XA, p. 141.
8. R. Pettorino and J. Willemsen, (unpublished).
9. C. E. Carlson, L. N. Chang, F. Mansouri, and J. F. Willemsen, SLAC PUB 1342 (to be published in Phys. Lett. B).
10. This result is not unexpected. It is known that $[H_0 - T]$ commutes with the current's Fourier components, ρ and ρ^+ (Ref. 12). It is then argued on general grounds ("irreducibility assumption") that $[H_0 - T]$ must be a function of Q only (Ref. 5). Presumably dimensional arguments and a few low order matrix elements can determine the precise form.
11. B. Zumino, Phys. Lett. 10, 224 (1964).

12. D. Mattis and E. Lieb, J. Math. Phys. 6 (1965) 304.

This paper does not explicitly verify the statement referred to, but provides the methods with which it may be verified.