

August 2008
BABAR-PROC-08/023
SLAC-PUB-13240
arXiv:0808.1011[hep-ex]

HADRONIC B DECAYS AT *BABAR* AND BELLE

VINCENZO LOMBARDO

(Representing the *BABAR* and BELLE Collaborations)

Dipartimento di Fisica and INFN Milano, via Celoria 16, Milano (Italy)

E-mail: vincenzo.lombardo@mi.infn.it

We review recent results of the *BABAR* and Belle Collaborations on the the α and γ angles of the unitarity triangle, on the $B \rightarrow K\pi\pi$ Dalitz-plot analyses, and on the searches for baryonic B decays and for $B \rightarrow D\bar{D}$ decays.

1 Introduction

In this review we present recent results from the *BABAR* and Belle Collaborations on hadronic B decays. The results are organized in 4 sections according to their theoretical interest. In section 2 we describe two recent results related to the angle α of the unitarity triangle (UT) of the Cabibbo-Kobayashi-Maskawa (CKM) quark mixing matrix¹, in section 3 two updated $B \rightarrow K\pi\pi$ Dalitz-plot analyses, in section 4 several new or updated measurements related to the CKM angle γ , and in section 5 two searches for baryonic B decays and for $B \rightarrow D^+D^-$ and \bar{D}^0D^0 decays.

2 Measurements related to the CKM angle α

Hadronic B decays from $\bar{b} \rightarrow \bar{u}u\bar{d}$ transitions provide the most direct information about the weak phase α governing the interference between $B^0 - \bar{B}^0$ mixing and B decay amplitudes. The difficulty in extracting α in these B decay modes is due to the presence of subleading penguin amplitudes with a different weak phase than that of the dominant tree amplitudes. For this reason the measured angle α_{eff} , obtained from time-dependent CP analyses, would differ from the UT angle α . This difficulty can be overcome by using symmetries, either isospin or approximate flavour $SU(3)$ ^{2,3,4}. In the first approach isospin relations can be used to interpret

Contributed to the Rencontres de Moriond QCD and High Energy Interactions, March 8-15, 2008, La Thuile, Italy

Work supported in part by US Department of Energy contract DE-AC02-76SF00515

results in terms of a constraint on α . The second approach can be used to extract an upper bound on $|\alpha - \alpha_{eff}|$ in terms of branching fractions of a complete set of SU(3) related processes where penguin amplitudes are expected to be enhanced.

The Belle experiment has recently presented at “Les Rencontres de Physique de la Valle d’Aoste” the analysis of neutral B mesons decays to $\rho^0\rho^0$ using 657 million $\bar{B}B$ events. The measurements of branching fraction and CP violating asymmetries of $B^0 \rightarrow \rho^0\rho^0$ are needed to complete the $B \rightarrow \rho\rho$ isospin relations which allow to constraint the value of α as measured in $B^0 \rightarrow \rho^+\rho^-$. The measured upper limit (UL) at 90% confidence level for the branching fraction of $B^0 \rightarrow \rho^0\rho^0$ is 1.0×10^{-6} . With no significant signal, this mode was assumed to be longitudinally polarized in order to obtain the most conservative UL. This UL is consistent with the *BABAR* result contributed to the *XIIIrd* International Symposium on Lepton and Photon, 2007⁵.

BABAR has recently published the first observation for two new charmless decay modes, $B^+ \rightarrow a_1(1260)^+K^0$ and $B^0 \rightarrow a_1(1260)^-K^+$. The branching fractions are $(34.9 \pm 5.0 \pm 4.4) \times 10^{-6}$ and $(16.3 \pm 2.9 \pm 2.3) \times 10^{-6}$ with a significance of 6.2 and 5.1 standard deviations (σ), respectively⁷. The first uncertainty is statistical and the second systematic. Charge asymmetries are found to be consistent with zero. These new results together with the time-dependent CP parameters extracted from neutral B mesons decays to $a_1(1260)\pi$ ⁸ and with measurements of the CP -averaged decay rates $B \rightarrow K_1(1270)\pi$ and $B \rightarrow K_1(1400)\pi$ will allow to provide a new and independent measurement of the CKM angle α ⁹.

3 $B \rightarrow K\pi\pi$ Dalitz-plot analyses

Two Dalitz-plot analyses of B meson decays to $K\pi\pi$ have been updated by *BABAR*. These $B^0 \rightarrow K^+\pi^-\pi^0$ and $B^\pm \rightarrow K^\pm\pi^\mp\pi^\pm$ decay modes are interesting for several reasons. Two anomalous features measured in $b \rightarrow s$ penguin dominated transitions have attracted substantial interest in recent years. CP asymmetries in these modes show a hint of systematic deviation from standard model predictions and direct CP asymmetries in $K\pi$ are hard to explain¹⁰. Moreover the fact that $K^*\pi$ amplitude can be used to constrain the $(\bar{\rho}, \bar{\eta})$ plane of the UT^{11,12} and also the possibility to study the nature of the scalar $f_x(1300)$ make these modes very attractive. With a sample of 231.8 million $\bar{B}B$ decays the *BABAR* Collaboration has measured the magnitudes and phases of the intermediate resonant and non-resonant amplitudes for B^0 and \bar{B}^0 decays and determined the corresponding CP -averaged branching fractions and charge asymmetries¹³. The inclusive branching fraction and the global CP -violating charge asymmetry are measured to be $\mathcal{B}(B^0 \rightarrow K^+\pi^-\pi^0) = (35.7_{-1.5}^{+2.6} \pm 2.2) \times 10^{-6}$ and $\mathcal{A}_{CP} = -0.030_{-0.051}^{+0.045} \pm 0.055$, respectively. The branching fraction of the decays $B^0 \rightarrow K^{*0}(892)\pi^0$ is observed with a significance of 5.6 σ , the measured branching fraction of the decay $B^0 \rightarrow K^{*+}(892)\pi^-$ is in agreement with previous *BABAR* and Belle measurements^{15,16}. No evidence for direct CP -violation in any resonant sub-decay has been found and the decay rate of $B^0 \rightarrow \rho^-K^+$ is found to be lower by 2 σ with respect to the Belle measurement¹⁶.

The Dalitz-plot analysis of the charged B mesons decays to $K^\pm\pi^\mp\pi^\pm$, using a sample of 383 million $\bar{B}B$ pairs, has found an evidence above 3 σ of direct CP -violation in $B^+ \rightarrow \rho^0K^+$, with a CP violation parameter $\mathcal{A}_{CP} = (+0.44 \pm 0.10 \pm 0.04_{-0.13}^{+0.05})$ where the third uncertainty represent the model dependence¹⁴. The total branching fraction $\mathcal{B}(B^+ \rightarrow K^+\pi^-\pi^+) = (54.4 \pm 1.1 \pm 4.5 \pm 0.7) \times 10^{-6}$ is compatible with the Belle measurement¹⁷. The $K^{*0}(892)\pi^+$, $(K\pi)_0^*\pi^+$ and $K_2^{*0}(892)\pi^+$ charge asymmetries are all consistent with zero, as expected¹⁸. Contributions from $f_2(1270)K^+$ and $f_x(1300)K^+$ are found to be necessary to obtain a good fit to the data with $f_x(1300)$ being a scalar with parameters $m_{f_x} = (1479 \pm 8)\text{MeV}/c^2$ and $\Gamma_{f_x} = (80 \pm 19)\text{MeV}$ where the errors are statistical only.

4 Measurements related to the CKM angle γ

The UT angle γ is related to the complex phase of the CKM matrix element V_{ub} through $V_{ub} = |V_{ub}|e^{-i\gamma}$. Over the years various methods have been proposed to measure the angle γ using either charged and neutral B decays. The GLW method¹⁹ has been implemented to work with intermediate D^0 and \bar{D}^0 mesons decay to a CP eigenstate. In the ADS method²⁰ the D^0 from the favored $b \rightarrow c$ amplitude is reconstructed in the doubly-Cabibbo suppressed final state, while the \bar{D}^0 from the $b \rightarrow u$ suppressed amplitude is reconstructed in the favored final state. In the GGSZ (Dalitz) method²¹ the D^0 and the \bar{D}^0 are reconstructed in the same $K_S^0\pi^+\pi^-$ three body final state. This method is based on the analysis of the $K_S^0\pi^+\pi^-$ Dalitz distribution and can to some extent be considered as a mixture of the two previous methods.

BABAR has recently reported the first analysis extracting γ from neutral B meson decays in $B^0 \rightarrow D^0(\bar{D}^0)K^{*0}$ with $D^0 \rightarrow K_S\pi^+\pi^-$ ²². The data sample consists of 371 million $B\bar{B}$ events. The charge of the kaon from K^* tags the flavor of the neutral B. The $D^0(K_S\pi^+\pi^+)$ mesons are analyzed with a well-known Dalitz model plus a K-Matrix formalism for the S-wave component of the $\pi^+\pi^-$ system. The interference of the signal decay amplitude with other $B^0 \rightarrow (\bar{D})^0(K\pi)_{non-K^*}^*$ amplitudes is taken into account using effective variables. The value of the angle γ as a function of r_S , the magnitude of the ratio between $b \rightarrow u$ and $b \rightarrow c$ amplitudes have been extracted. Combining this result with the available information on r_S , γ was found to be $(162 \pm 56)^\circ$ or $(342 \pm 56)^\circ$ and $r_S < 0.55$ at 95% probability²².

An improved measurement of the CKM CP -violating phase γ through a Dalitz-plot analysis of neutral D meson decays to $K_S^0\pi^+\pi^-$ and $K_S^0K^+K^-$ produced in the processes $B^\mp \rightarrow DK^\mp$, $B^\mp \rightarrow D^*K^\mp$ with $D^* \rightarrow D\pi^0, D\gamma$ and $B^\mp \rightarrow DK^{*\mp}$ with $K^{*\mp} \rightarrow K_S^0\pi^\mp$ has recently been updated by the *BABAR* Collaboration using a sample of 383 million $B\bar{B}$ pairs²³. These B meson decays are theoretically clean and are unlikely to be affected by new physics. The extracted CKM angle γ is $(76 \pm 22 \pm 5 \pm 5)^\circ \pmod{180^\circ}$ where the first error is statistical, the second is the experimental systematic uncertainty and the third reflects the uncertainty on the description of the Dalitz-plot distribution. This result has a significance of direct CP violation ($\gamma \neq 0$) of 3.0 σ .

A measurement of the decay $B^\pm \rightarrow D_{CP} K^\pm$ has recently been updated by the *BABAR* Collaboration using a sample of 382 million $B\bar{B}$ events²⁴. The D_{CP}^0 mesons have been reconstructed in both a non- CP eigenstate and in CP (CP -even and CP -odd) eigenstates. The ratios ($R_{CP\pm}$) of charge-averaged partial rates and of the partial-rate of the charge asymmetries ($A_{CP\pm}$) have been measured, $A_{CP+} = 0.27 \pm 0.09(stat) \pm (0.04)(syst)$, $A_{CP-} = -0.09 \pm 0.09(stat) \pm (0.02)(syst)$, $R_{CP+} = 1.06 \pm 0.10(stat) \pm 0.05(syst)$ and $R_{CP-} = 1.03 \pm 0.10(stat) \pm 0.05(syst)$.

BABAR has reported an improved measurement of the branching fractions of neutral B meson decays to $D_s^{(*)+}\pi^-$ and $D_s^{(*)-}K^+$ and performed the first branching fraction and polarization measurement of $D_s^{(*)+}\rho^-$ and $D_s^{(*)-}K^{*+}$ in a sample of 381 million $B\bar{B}$ pairs²⁵. The measured branching fractions and longitudinal polarization fractions for the new modes are $\mathcal{B}(B^0 \rightarrow D_s^{(*)+}\rho^-) = (4.4_{-1.2}^{+1.3} \pm 0.8) \times 10^{-5}$, $f_L(B^0 \rightarrow D_s^{(*)+}\rho^-) = (0.86_{-0.28}^{+0.26} \pm 0.15)$, $\mathcal{B}(B^0 \rightarrow D_s^-K^{*+}) = (3.6_{-0.9}^{+1.0} \pm 0.4) \times 10^{-5}$, $\mathcal{B}(B^0 \rightarrow D_s^{*-}K^{*+}) = (3.0_{-1.2}^{+1.4} \pm 0.3) \times 10^{-5}$ and $f_L(B^0 \rightarrow D_s^{(*)-}K^{*+}) = (0.96_{-0.31}^{+0.38} \pm 0.08)$.

5 Search for baryonic B decays and $B \rightarrow D^+D^-$ and \bar{D}^0D^0

Several baryonic B decays have been observed over the past years by *BABAR* and Belle. A search for the doubly charmed baryonic decays $\bar{B}^0 \rightarrow \Lambda_c^+\bar{\Lambda}_c^-$ at Belle using a data sample of 520 million $B\bar{B}$ events is presented. No significant signal has been found and an upper limit of $\mathcal{B}(\bar{B}^0 \rightarrow \Lambda_c^+\bar{\Lambda}_c^-) < 6.2 \times 10^{-5}$ at 90% confidence level has been set. The result is significantly below the naïve extrapolation from the branching fraction of the decay $B^- \rightarrow \Xi_c^0\bar{\Lambda}_c^-$ assuming

a simple factor of $|V_{cd}/V_{cs}|^2$ Cabibbo-suppressed²⁶.

Belle has also reported the branching fraction measurements of $B^0 \rightarrow p\bar{p}K^{*0}$ and $B^+ \rightarrow p\bar{p}K^{*+}$ with a sample of 535 million $B\bar{B}$ pairs. The measured branching fractions are $\mathcal{B}(B^0 \rightarrow p\bar{p}K^{*0}) = (1.18_{-0.25}^{+0.29}(stat.) \pm 0.11(syst.)) \times 10^{-6}$ and $\mathcal{B}(B^+ \rightarrow p\bar{p}K^{*+}) = (3.38_{-0.60}^{+0.73}(stat.) \pm 0.39(syst.)) \times 10^{-6}$ with a statistical significance of 7.2σ and 8.8σ , respectively. The polarization of the K^* contains important information on the decay dynamics; the K^{*0} meson is found to be almost 100% in the helicity zero state, compared to $(32 \pm 17 \pm 9)\%$ for the K^{*+} meson²⁷.

An improved measurement of the $B^+ \rightarrow D^+\bar{D}^0$ and $B^0 \rightarrow D^0\bar{D}^0$ decays based on 657 million $B\bar{B}$ events is also reported²⁸. The measured branching fraction is $\mathcal{B}(B^+ \rightarrow D^+\bar{D}^0) = (3.85 \pm 0.31 \pm 0.38) \times 10^{-4}$ with charge asymmetry $A_{CP}(B^+ \rightarrow D^+\bar{D}^0) = 0.00 \pm 0.08 \pm 0.02$. An upper limit for $B^0 \rightarrow D^0\bar{D}^0$ has been set for the decay $\mathcal{B}(B^0 \rightarrow \bar{D}^0D^0) < 0.42 \times 10^{-4}$.

References

1. N. Cabibbo Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973)
2. M. Gronau and D. London, Phys. Rev. Lett. **65**, 3381 (1990); H. J. Lipkin *et al.*, Phys. Rev. D **44**, 1454 (1991); M. Gronau, Phys. Lett. B **265**, 389 (1991).
3. Y. Grossman and H. R. Quinn, Phys. Rev. D **58**, 017504 (1998); J. Charles, Phys. Rev. D **59**, 054007 (1999); M. Gronau *et al.*, Phys. Lett. B **514**, 315 (2001).
4. M. Gronau and J. Zupan, Phys. Rev. D **70**, 074031 (2004).
5. B Aubert *et al.*, [BABAR Collaboration], arXiv:0708.1630[hep-ex]
6. For the $a_1(1260)$ meson we use the short notation a_1 .
7. B Aubert *et al.*, Phys. Rev. Lett. **100**, 051803 (2008);
8. B Aubert *et al.*, Phys. Rev. Lett. **98**, 181803 (2007);
9. M. Gronau and J. Zupan, Phys. Rev. D **73**, 057502 (2006).
10. M. Gronau, Int. J. Mod. Phys. A **22**, 1953 (2007); E. Lunghi and A. Soni, JHEP **0709**, 053 (2007).
11. M. Ciuchini, M. Pierini and L. Silvestrini, Phys. Rev. D **74**, 051301 (2006)
12. M. Gronau, D. Pirjol and A. Soni, arXiv:0712.3751[hep-ph]
13. B Aubert *et al.*, [BABAR Collaboration], arXiv:0711.4417[hep-ex]
14. B Aubert *et al.*, [BABAR Collaboration], arXiv:0803.4451[hep-ex]
15. B Aubert *et al.*, [BABAR Collaboration], Phys. Rev. D **73**, 031101 (2006).
16. P. Chang *et al.*, [Belle Collaboration], Phys. Lett. B **599**, 148 (2004).
17. A. Garmash *et al.*, [Belle Collaboration], Phys. Rev. Lett. **96**, 251803 (2006).
18. M. Beneke and M. Neubert, Nucl. Phys. **B675**, 333 (2003); C.-W. Chiang, M. Gronau, Z. Luo, J. L. Rosner, and D. A. Suprun, Phys. Rev. **D69**, 034001 (2004);
19. M. Gronau and D. London, Phys. Lett. B **253**, 483 (1991); M. Gronau and D. Wyler, Phys. Lett. B **265**, 172 (1991);
20. D. Atwood, I. Dunietz and A. Soni, Phys. Rev. Lett. **78**, 3257 (1997).
21. A. Giri, Yu. Grossman, A. Soffer and J. Zupan, Phys. Rev. D **68**, 054018 (2003).
22. B Aubert *et al.*, [BABAR Collaboration], arXiv:0805.2001[hep-ex]
23. B Aubert *et al.*, [BABAR Collaboration], arXiv:0804.2089[hep-ex]
24. B Aubert *et al.*, [BABAR Collaboration], arXiv:0802.4052[hep-ex]
25. B Aubert *et al.*, [BABAR Collaboration], arXiv:0803.4296[hep-ex]
26. K. Abe *et al.*, [Belle Collaboration], arXiv:0708.1105[hep-ex]
27. J.-H. Chen *et al.*, [Belle Collaboration], arXiv:0802.0336[hep-ex]
28. K. Abe *et al.*, [Belle Collaboration], arXiv:0708.1668[hep-ex]