

Recent results on two-photon physics at BABAR

Vladimir DRUZHININ*

Budker Institute of Nuclear Physics

E-mail: druzhinin@inp.nsk.su

We present measurements of the $\gamma\gamma^* \rightarrow \pi^0$ transition form factor for the momentum transfer range $Q^2=4-40 \text{ GeV}^2$ and the $\gamma\gamma^* \rightarrow \eta_c$ transition form factor for the range $Q^2=2-50 \text{ GeV}^2$. The results of the measurement of the η_c mass, total and two-photon widths are also presented.

*European Physical Society Europhysics Conference on High Energy Physics
July 16-22, 2009
Krakow, Poland*

*Speaker.

1. Introduction

The diagram for the process of the two-photon production of the pseudoscalar meson is shown in Fig. 1(left). The effect of strong interactions in this process is described only one form factor $F(q_1^2, q_2^2)$ depending on the squared momentum transfers to the electrons.

The electrons in such process are scattered predominantly at small angle. Therefore, the two-photon processes are usually studied in so called no-tag mode with both final electrons undetected. In this case the virtual photons are practically real, the momentum transfers squared are close to zero. In no-tag mode the meson-photon transition form factor at zero q^2 's and the two-photon width of the meson are measured. In single tag-mode the one of the final electron is detected. The corresponding virtual photon is highly off-shell. From the measurement of the cross section we extract more rich information: the dependence of the meson form factor on $Q^2 = -q_1^2$.

At large Q^2 perturbative QCD (pQCD) predicts that the transition form factor can be represented as a convolution of a calculable hard scattering amplitude for $\gamma\gamma^* \rightarrow q\bar{q}$ with a nonperturbative meson distribution amplitude (DA), $\phi(x, Q^2)$ [1]. The latter can be interpreted as the amplitude for the transition of the meson with momentum p into two quarks with momenta px and $p(1-x)$. The experimental data on the transition form factor can be used to test different phenomenological models for DA.

The cross section of the process $e^+e^- \rightarrow e^+e^-P$ falls very rapidly with increase of Q^2 (Q^{-6} for π^0). Therefore, a precise measurement of the transition form factor can be performed only at high luminosity e^+e^- machines. We present the results of the measurements of the transition form factors for π^0 and η_c mesons performed by the BABAR detector at the PEP-II e^+e^- collider. The results are based on data with integrated luminosity of about 450 fb^{-1} collected at the center-of-mass energy of 10.6 GeV. The single-tag events are selected with detected and identified electron and with fully reconstructed π^0 or η_c . It is required that the transverse momentum of electron-plus-meson system be low and the missing mass in an event be close to zero.

2. Measurement of the $\gamma^*\gamma \rightarrow \pi^0$ transition form factor [2]

The π^0 meson is detected via its decay into two photons. The two-photon invariant mass spectrum for selected π^0 candidates is shown in Fig. 1(right). The clear π^0 peak is seen. The main non-peaking background process is so called virtual Compton scattering (VCS), the process $e^+e^- \rightarrow e^+e^-\gamma$ with one of the final electrons directed along the beam axis. The VCS photon together with a soft photon, for example from beam background, may give an invariant mass value close to the π^0 mass. The peaking background comes from the process of two-photon production of two π^0 's. This background is estimated from data and is about 10% of signal events. The total number of signal events determined from the fit to the mass spectrum in Fig. 1(right) is about 13000. This number is an order of magnitude large than the statistics of the previous measurement of the form factor by CLEO [3].

We measure the form factor in the Q^2 region from 4 to 40 GeV^2 . The lower Q^2 limit is determined by the detector acceptance for the electron. For $Q^2 > 40 \text{ GeV}^2$ we do not see evidence of a π^0 signal over background. The data were divided into 17 Q^2 intervals. For each Q^2 interval the mass spectrum is fitted by a sum of signal and background distributions. From the measured Q^2




Figure 1: **Left panel:** The Feynman diagram for the process of the pseudoscalar meson two-photon production. **Right panel:** The two-photon invariant mass spectrum for data events with $4 < Q^2 < 40 \text{ GeV}^2$ and fitting curves.

spectrum we determine the differential cross section for $e^+e^- \rightarrow e^+e^-\pi^0$ and the transition form factor. The result for the form factor is shown in Fig.2. The errors shown are combined statistical and Q^2 -dependent systematic. There is also Q^2 -independent error equal to 2.3%. Main sources of the systematic uncertainties are background subtraction, data-MC simulation difference in the detector response, and the model uncertainty due to the unknown q_2^2 dependence of the form factor.

The comparison of our results with previous measurements [4, 3] is shown in Fig. 2(left). In the Q^2 range from 4 to 9 GeV^2 our results are in reasonable agreement with the measurements by the CLEO collaboration [3], but have significantly better precision.

The horizontal dashed line indicates the asymptotic limit for the form factor. The value of the asymptotic limit ($Q^2 F(Q^2) = \sqrt{2} f_\pi \approx 0.185 \text{ GeV}$) is predicted by pQCD. The measured form factor exceeds the asymptotic limit at $Q^2 > 10 \text{ GeV}^2$. This is an unexpected behavior; most models for the pion DA give form factor approaching the limit from below (see, e.g., Ref. [5] and references therein). Our data in the range from 4 to 40 GeV^2 are well described by the function

$$Q^2 |F(Q^2)| = A \left(\frac{Q^2}{10 \text{ GeV}^2} \right)^\beta \quad (2.1)$$

with $A = 0.182 \pm 0.002 \text{ GeV}$ and $\beta = 0.25 \pm 0.02$ (dotted line in Fig. 2(left)). The effective Q^2 dependence of the measured form factor is $\sim 1/Q^{3/2}$.

Fig. 2(right) demonstrates the comparison of our measurement with the result of the NLO QCD calculations performed by Bakulev, Mikhailov, and Stefanis [6] for the three models of the pion DA: asymptotic [7], Chernyak-Zhitnitsky (CZ) [8], and the DA derived from QCD sum rules with non-local condensates (BMS) [9]. There is a large difference between data and theory in Q^2 dependence. We conclude that all these models are inadequate for $Q^2 < 15 \text{ GeV}^2$. For $Q^2 > 20 \text{ GeV}^2$ the theoretical uncertainties are expected to be smaller. In this region our data lie above




Figure 2: The $\gamma\gamma^* \rightarrow \pi^0$ transition form factor multiplied by Q^2 . The dashed line indicates the asymptotic limit for the form factor. The dotted curve at the left panel shows the interpolation given by Eq.(2.1). The solid and dotted lines, and shaded band at the right panel show the predictions for the form factor for the CZ [8], asymptotic (ASY) [7], and BMS [9] models of the pion distribution amplitude, respectively.

asymptotic limit and are consistent with CZ model. It should be noted that the CZ DA is widest of the three DA's discussed.

There are theoretical works which appeared after the publication of our result. Mikhailov and Stefanis [10] argue that the growth of form factor cannot be explained by higher-order pQCD and power corrections. Other works [11, 12, 13, 14] consider flat or very wide pion DA. With such distribution amplitude the Q^2 dependence observed by BABAR is reproduced well.

3. Measurement of the $\gamma^*\gamma \rightarrow \eta_c$ transition form factor

The two-photon η_c production is studied both in no-tag and in single-tag modes. The η_c is reconstructed via its decay to $K_S K^- \pi^+$. The $KK\pi$ mass spectra for no-tag events is shown in Fig. 3(left) The η_c and J/ψ peaks are clearly seen. The J/ψ 's are produced in initial state radiation (ISR) process $e^+e^- \rightarrow J/\psi\gamma$. From the fit to the mass spectrum we determine η_c parameters:

$$m = 2982.2 \pm 0.4 \pm 1.5 \text{ MeV}/c^2,$$

$$\Gamma = 31.7 \pm 1.2 \pm 0.8 \text{ MeV},$$

$$\Gamma(\eta_c \rightarrow \gamma\gamma)B(\eta_c \rightarrow K\bar{K}\pi) = 0.379 \pm 0.009 \pm 0.030 \text{ keV}.$$

These results are preliminary. Main sources of systematic uncertainties on the mass and width are unknown background shape and possible interference between η_c and non-resonant two-photon $KK\pi$ amplitudes. The uncertainty on the detection efficiency dominates in the systematic uncertainty of $\Gamma(\eta_c \rightarrow \gamma\gamma)B(\eta_c \rightarrow K\bar{K}\pi)$. The results for the mass and width are in an agreement with the previous BABAR measurement [15]: $m = 2982.5 \pm 1.1 \pm 0.9 \text{ MeV}/c^2$ and $\Gamma = 34.3 \pm 2.3 \pm 0.9 \text{ MeV}$, obtained using 88 fb^{-1} data. The current PDG values for these parameters




Figure 3: Left panel: The $K_S K^\pm \pi^\mp$ invariant mass distribution and fitted curve for no-tag data events. **Right panel:** The $\gamma\gamma^* \rightarrow \eta_c$ transition form factor normalized to $F(0)$ (points with error bars). The solid curve shows the interpolation given by a monopole form. The dotted curve shows the leading order pQCD prediction from Ref. [20].

are $m = 2980.5 \pm 1.2 \text{ MeV}/c^2$ and $\Gamma = 27.4 \pm 2.9 \text{ MeV}$ [16]. The obtained value of the product $\Gamma(\eta_c \rightarrow \gamma\gamma)B(\eta_c \rightarrow K\bar{K}\pi)$ agrees with the PDG value $0.44 \pm 0.05 \text{ keV}$ [16], and also with the recent CLEO measurement $0.407 \pm 0.022 \pm 0.028 \text{ keV}$ [17].

We select $520 \pm 40 \pm 20$ single-tag η_c events. This number can be compared with 8 ± 5 events selected in the previous single-tag η_c measurement by L3 [18]. The single-tag data were divided into 11 Q^2 intervals. For each interval we fit to the $KK\pi$ mass spectrum and determine number of events with η_c . From the ratio of the measured Q^2 spectrum to the number of the no-tag η_c events we extract the normalized η_c transition form factor shown in Fig. 3(right). The errors shown are combined statistical and Q^2 -dependent systematic. There is also Q^2 -independent error equal to 4.3%. Main source of the systematic error is the systematic uncertainty on detection efficiency.

The form factor data are fitted by the monopole function $|F(Q^2)/F(0)| = 1/(1 + Q^2/\Lambda)$. The result of the fit is shown in Fig. 3(right) by solid line. The pole parameter Λ is found to be $\Lambda = 8.5 \pm 0.6 \pm 0.7 \text{ GeV}^2$. This value is in reasonable agreement with that expected from vector dominance, namely $\Lambda = m_{J/\psi}^2 = 9.6 \text{ GeV}^2$, and with the result of the lattice QCD calculation $\Lambda = 8.4 \pm 0.4 \text{ GeV}^2$ [19]. The dotted curve in Fig. 3(right) shows results of the leading-order pQCD calculation of Ref. [20]. The data lie systematically below this prediction.

4. Summary

The $\gamma^* \gamma \pi^0$ transition form factor has been measured for Q^2 range from 4 to 40 GeV^2 . The unexpected Q^2 -dependence for the form factor is observed for $Q^2 > 10 \text{ GeV}^2$. The data lie above the asymptotic limit. This indicates that the pion distribution amplitude should be wide. The measurement stimulated development of new models for the form-factor calculation, in particular, with flat distribution amplitude [12, 13, 14].

The $\gamma^* \gamma \eta_c$ transition form factor has been measured for Q^2 range from 2 to 50 GeV². The form factor data are well described by the monopole form with pole parameter about 9 GeV². The data are in reasonable agreement with both QCD and VDM predictions.

References

- [1] G. P. Lepage and S. J. Brodsky, Phys. Rev. D **22**, 2157 (1980).
- [2] B. Aubert *et al.* [The BABAR Collaboration], Phys. Rev. D **80**, 052002 (2009).
- [3] J. Gronberg *et al.* [CLEO Collaboration], Phys. Rev. D **57**, 33 (1998).
- [4] H. J. Behrend *et al.* [CELLO Collaboration], Z. Phys. C **49**, 401 (1991).
- [5] N. G. Stefanis, Nucl. Phys. Proc. Suppl. **181-182**, 199 (2008).
- [6] A. P. Bakulev, S. V. Mikhailov, and N. G. Stefanis, Phys. Rev. D **67**, 074012 (2003); Phys. Lett. B **578**, 91 (2004).
- [7] G. P. Lepage and S. J. Brodsky, Phys. Lett. B **87**, 359 (1979).
- [8] V. L. Chernyak and A. R. Zhitnitsky, Nucl. Phys. B **201**, 492 (1982) [Erratum-ibid. B **214**, 547 (1983)].
- [9] A. P. Bakulev, S. V. Mikhailov, and N. G. Stefanis, Phys. Lett. B **508**, 279 (2001) [Erratum-ibid. B **590**, 309 (2004)].
- [10] S. V. Mikhailov and N. G. Stefanis, Nucl. Phys. B **821**, 291 (2009).
- [11] A. E. Dorokhov, arXiv:0905.4577 [hep-ph].
- [12] A. V. Radyushkin, arXiv:0906.0323 [hep-ph].
- [13] M. V. Polyakov, arXiv:0906.0538 [hep-ph].
- [14] H. Li and S. Mishima, arXiv:0907.0166 [hep-ph].
- [15] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **92**, 142002 (2004).
- [16] Particle Data Group, 2009 partial update for the 2010 edition, <http://pdg.lbl.gov>.
- [17] D. M. Asner *et al.* [CLEO Collaboration], Phys. Rev. Lett. **92**, 142001 (2004).
- [18] M. Acciarri *et al.* [L3 Collaboration], Phys. Lett. B **461**, 155 (1999).
- [19] J. J. Dudek and R. G. Edwards, Phys. Rev. Lett. **97**, 172001 (2006).
- [20] T. Feldmann and P. Kroll, Phys. Lett. B **413**, 410 (1997).