

FERMI LAT OBSERVATIONS OF LS I +61°303:
FIRST DETECTION OF AN ORBITAL MODULATION IN GEV GAMMA RAYS

A. A. ABDO^{2,3}, M. ACKERMANN⁴, M. AJELLO⁴, W. B. ATWOOD⁵, M. AXELSSON^{6,7}, L. BALDINI⁸, J. BALLE⁹, G. BARBIELLINI^{10,11},
D. BASTIERI^{12,13}, B. M. BAUGHMAN¹⁴, K. BECHTOL⁴, R. BELLAZZINI⁸, B. BERENJI⁴, R. BLANDFORD⁴, E. D. BLOOM⁴,
E. BONAMENTE^{15,16}, A. W. BORGLAND⁴, J. BREGEON⁸, A. BREZ⁸, M. BRIGIDA^{17,18}, P. BRUEL¹⁹, T. H. BURNETT²⁰,
G. A. CALIANDRO^{17,18}, R. A. CAMERON⁴, P. A. CARAVEO²¹, J. M. CASANDJIAN⁹, E. CAVAZZUTI²², C. CECCHI^{15,16}, Ö. ÇELİK²³,
E. CHARLES⁴, S. CHATY⁹, A. CHEKHTMAN^{3,24}, C. C. CHEUNG²³, J. CHIANG⁴, S. CIPRINI^{15,16}, R. CLAUS⁴, J. COHEN-TANUGI²⁵,
L. R. COMINSKY²⁶, J. CONRAD^{6,27,28,29}, S. CORBEL⁹, R. CORBET^{23,30}, S. CUTINI²², C. D. DERMER³, A. DE ANGELIS³¹,
A. DE LUCA³², F. DE PALMA^{17,18}, S. W. DIGEL⁴, M. DORMODY⁵, E. DO COUTO E SILVA⁴, P. S. DRELL⁴, R. DUBOIS^{1,4}, G. DUBUS³³,
D. DUMORA^{34,35}, C. FARNIER²⁵, C. FAVUZZI^{17,18}, S. J. FEGAN¹⁹, W. B. FOCKE⁴, M. FRAILIS³¹, Y. FUKAZAWA³⁶, S. FUNK⁴,
P. FUSCO^{17,18}, F. GARGANO¹⁸, D. GASPARRINI²², N. GEHRELS^{23,37}, S. GERMANI^{15,16}, B. GIEBELS¹⁹, N. GIGLIETTO^{17,18},
F. GIORDANO^{17,18}, T. GLANZMAN⁴, G. GODFREY⁴, I. A. GRENIER⁹, M.-H. GRONDIN^{34,35}, J. E. GROVE³, L. GUILLEMOT^{34,35},
S. GUIRIEC³⁸, Y. HANABATA³⁶, A. K. HARDING²³, M. HAYASHIDA⁴, E. HAYS²³, A. B. HILL^{1,33}, R. E. HUGHES¹⁴, G. JÓHANNESSEN⁴,
A. S. JOHNSON⁴, R. P. JOHNSON⁵, T. J. JOHNSON^{23,37}, W. N. JOHNSON³, T. KAMAE⁴, H. KATAGIRI³⁶, J. KATAOKA³⁹, N. KAWAI^{40,41},
M. KERR²⁰, J. KNÖDLSER⁴², M. L. KOCIAN⁴, F. KUEHN¹⁴, M. KUSS⁸, J. LANDE⁴, S. LARSSON^{6,28}, L. LATRONICO⁸, F. LONGO^{10,11},
F. LOPARCO^{17,18}, B. LOTT^{34,35}, M. N. LOVELLETTE³, P. LUBRANO^{15,16}, G. M. MADEJSKI⁴, A. MAKEEV^{3,24}, M. MARELLI²¹,
M. N. MAZZIOTTA¹⁸, J. E. MCENERY²³, C. MEURER^{6,28}, P. F. MICHELSON⁴, W. MITTHUMSIRI⁴, T. MIZUNO³⁶, C. MONTE^{17,18},
M. E. MONZANI⁴, A. MORSELLI⁴³, I. V. MOSKALENKO⁴, S. MURGIA⁴, P. L. NOLAN⁴, E. NUSS²⁵, T. OHSUGI³⁶, A. OKUMURA⁴⁴,
N. OMODEI⁸, E. ORLANDO⁴⁵, J. F. ORMES⁴⁶, D. PANEQUE⁴, J. H. PANETTA⁴, D. PARENT^{34,35}, V. PELASSA²⁵, M. PEPE^{15,16},
M. PESCE-ROLLINS⁸, F. PIRON²⁵, T. A. PORTER⁵, S. RAINO^{17,18}, R. RANDO^{12,13}, P. S. RAY³, M. RAZZANO⁸, N. REA^{47,48},
A. REIMER⁴, O. REIMER^{4,49}, T. REPOSEUR^{34,35}, S. RITZ²³, L. S. ROCHESTER⁴, A. Y. RODRIGUEZ⁴⁸, R. W. ROMANI⁴, F. RYDE^{6,27},
H. F.-W. SADROZINSKI⁵, D. SANCHEZ¹⁹, A. SANDER¹⁴, P. M. SAZ PARKINSON⁵, J. D. SCARGLE⁵⁰, C. SGRÒ⁸, M. S. SHAW⁴,
A. SIERPOWSKA-BARTOSIK⁴⁸, E. J. SISKIND⁵¹, D. A. SMITH^{34,35}, P. D. SMITH¹⁴, G. SPANDRE⁸, P. SPINELLI^{17,18}, E. STRIANI^{43,52},
M. S. STRICKMAN³, D. J. SUSON⁵³, H. TAJIMA⁴, H. TAKAHASHI³⁶, T. TAKAHASHI⁵⁴, T. TANAKA⁴, J. B. THAYER⁴, J. G. THAYER⁴,
D. J. THOMPSON²³, L. TIBALDO^{12,13}, D. F. TORRES^{1,48,55}, G. TOSTI^{15,16}, A. TRAMACERE^{4,56}, Y. UCHIYAMA⁴, T. L. USHER⁴,
V. VASILEIOU^{30,57}, N. VILCHEZ⁴², V. VITALE^{43,52}, A. P. WAITE⁴, P. WANG⁴, B. L. WINER¹⁴, K. S. WOOD³, T. YLINEN^{6,27,58},
M. ZIEGLER⁵

Draft version July 24, 2009

ABSTRACT

This *Letter* presents the first results from the observations of LS I +61°303 using Large Area Telescope data from the *Fermi Gamma-Ray Space Telescope* between 2008 August and 2009 March. Our results indicate variability that is consistent with the binary period, with the emission being modulated at 26.6 ± 0.5 days. This constitutes the first detection of orbital periodicity in high-energy gamma rays (20 MeV–100 GeV, HE). The light curve is characterized by a broad peak after periastron, as well as a smaller peak just before apastron. The spectrum is best represented by a power law with an exponential cutoff, yielding an overall flux above 100 MeV of $0.82 \pm 0.03(\text{stat}) \pm 0.07(\text{syst}) 10^{-6}$ ph cm⁻² s⁻¹, with a cutoff at $6.3 \pm 1.1(\text{stat}) \pm 0.4(\text{syst})$ GeV and photon index $\Gamma = 2.21 \pm 0.04(\text{stat}) \pm 0.06(\text{syst})$. There is no significant spectral change with orbital phase. The phase of maximum emission, close to periastron, hints at inverse Compton scattering as the main radiation mechanism. However, previous very high-energy gamma ray (>100 GeV, VHE) observations by MAGIC and VERITAS show peak emission close to apastron. This and the energy cutoff seen with *Fermi* suggest the link between HE and VHE gamma rays is nontrivial.

Subject headings: binaries: close — stars: variables: other — gamma rays: observations — X-rays: binaries — X-rays: individual (LS I +61°303)

¹ Corresponding authors: R. Dubois, richard@slac.stanford.edu; A. B. Hill, adam.hill@obs.ujf-grenoble.fr; D. F. Torres, dtorres@ieec.uab.es.

² National Research Council Research Associate, National Academy of Sciences, Washington, DC 20001

³ Space Science Division, Naval Research Laboratory, Washington, DC 20375

⁴ W. W. Hansen Experimental Physics Laboratory, Kavli Institute for Particle Astrophysics and Cosmology, Department of Physics and SLAC National Accelerator Laboratory, Stanford University, Stanford, CA 94305

⁵ Santa Cruz Institute for Particle Physics, Department of Physics and Department of Astronomy and Astrophysics, University of California at Santa Cruz, Santa Cruz, CA 95064

⁶ The Oskar Klein Centre for Cosmo Particle Physics, AlbaNova, SE-106 91 Stockholm, Sweden

⁷ Department of Astronomy, Stockholm University, SE-106 91 Stockholm, Sweden

⁸ Istituto Nazionale di Fisica Nucleare, Sezione di Pisa, I-56127 Pisa, Italy

⁹ Laboratoire AIM, CEA-IRFU/CNRS/Université Paris Diderot, Service d'Astrophysique, CEA Saclay, 91191 Gif sur Yvette, France

¹⁰ Istituto Nazionale di Fisica Nucleare, Sezione di Trieste, I-34127 Trieste, Italy

¹¹ Dipartimento di Fisica, Università di Trieste, I-34127 Trieste, Italy

¹² Istituto Nazionale di Fisica Nucleare, Sezione di Padova, I-35131 Padova, Italy

¹³ Dipartimento di Fisica “G. Galilei”, Università di Padova, I-35131 Padova, Italy

¹⁴ Department of Physics, Center for Cosmology and Astro-Particle Physics, The Ohio State University, Columbus, OH 43210

¹⁵ Istituto Nazionale di Fisica Nucleare, Sezione di Perugia, I-06123 Perugia, Italy

¹⁶ Dipartimento di Fisica, Università degli Studi di Perugia, I-06123 Perugia, Italy

¹⁷ Dipartimento di Fisica “M. Merlin” dell’Università e del Politecnico di Bari, I-70126 Bari, Italy

1. INTRODUCTION

The high-mass X-ray binary LS I +61°303 (=V615 Cas) has long been plausibly associated with a high-energy (HE, 20 MeV–100 GeV) gamma-ray source, although never before confirmed. The discovery of the *COS B* source 2CG 135+01 (Hermsen et al. 1977) quickly brought attention to this binary system’s Be star localized within its error box, because of its unusual periodic radio emission (Gregory et al. 1979) and its X-ray emission (Bignami et al. 1981). 2CG 135+01 was to

remain one of the brightest sources known in the HE gamma-ray sky, with a flux of $\sim 10^{-6}$ ph s $^{-1}$ cm $^{-2}$ above 100 MeV (Swanenburg et al. 1981). In the 1990s, EGRET detected the source with high confidence at the same average flux level and derived a power-law photon index of $\Gamma = 2.05 \pm 0.06$ (Kniffen et al. 1997). Although there are no other objects of note (radio-loud AGN, or pulsars) coinciding with the 3EG source (Hartman et al. 1999), its positional uncertainty was not small enough to firmly associate the gamma-ray source with the binary. Variability in the EGRET light curve could be neither firmly established nor related to variability at other wavelengths (Tavani et al. 1998; Nolan et al. 2003). Recently, *AGILE* has reported detecting the source at the same flux level (Pittori et al. 2009).

LS I +61°303 is an unusual binary system exhibiting strong variable emission from the radio to X-ray and TeV energies. At radio wavelengths the source has been shown to exhibit radio outbursts that are modulated on an orbital period of 26.4960 ± 0.0028 days (Taylor & Gregory 1982; Gregory 2002). The phase of radio maximum has also been shown by Gregory (2002) to vary with a super-orbital period of 1667 ± 8 days. Observations of orbital modulation in the optical place constraints on the binary system parameters. The binary has an eccentric orbit ($e=0.55-0.72$) and the Be star radial velocity is consistent with a neutron star companion or, if the orbital inclination is $\leq 25^\circ$, with a $\geq 3M_\odot$ black hole (Hutchings & Crampton 1981; Casares et al. 2005). Significant uncertainty still exists in key parameters of the orbital solution of the system (Grundstrom et al. 2007; Aragona et al. 2009).

Behavior in the X-ray band is much more complicated. Orbital modulation has been reported with the peak of emission appearing at phases 0.6–0.7 (Paredes et al. 1997; Esposito et al. 2007). However the modulation is not smooth, with short timescale flares and very strong orbit-to-orbit variability (Smith et al. 2009). Broad-band spectral analysis of *XMM-Newton* and *INTEGRAL* data by Chernyakova, Neronov, & Walter (2006) reveal LS I +61°303 to be well fitted by a simple absorbed power-law with a hard photon index, $\Gamma \simeq 1.5$, in the 0.5–100 keV band.

The MAGIC telescope detected a variable very high-energy (VHE >100 GeV) gamma-ray source coincident with LS I +61°303 (Albert et al. 2006); a result that has been independently confirmed by the VERITAS collaboration (Acciari et al. 2008). More recently, the MAGIC collaboration has further reported that the VHE emission is periodic at the 26.5 day orbital period of the system (Albert et al. 2009). The VHE emission is consistently highest close to apastron, when the compact object is farthest from the Be star, and remains undetected at periastron. Like LS 5039 and PSR B1259–63 (Aharonian et al. 2005a,b), and contrary to Cyg X–1 (Albert et al. 2007), LS I +61°303 is a gamma-ray binary with its spectral energy distribution peak-

- ¹⁸ Istituto Nazionale di Fisica Nucleare, Sezione di Bari, 70126 Bari, Italy
¹⁹ Laboratoire Leprince-Ringuet, École polytechnique, CNRS/IN2P3, Palaiseau, France
²⁰ Department of Physics, University of Washington, Seattle, WA 98195-1560
²¹ INFN-Istituto di Astrofisica Spaziale e Fisica Cosmica, I-20133 Milano, Italy
²² Agenzia Spaziale Italiana (ASI) Science Data Center, I-00044 Frascati (Roma), Italy
²³ NASA Goddard Space Flight Center, Greenbelt, MD 20771
²⁴ George Mason University, Fairfax, VA 22030
²⁵ Laboratoire de Physique Théorique et Astroparticules, Université Montpellier 2, CNRS/IN2P3, Montpellier, France
²⁶ Department of Physics and Astronomy, Sonoma State University, Rohnert Park, CA 94928-3609
²⁷ Department of Physics, Royal Institute of Technology (KTH), AlbaNova, SE-106 91 Stockholm, Sweden
²⁸ Department of Physics, Stockholm University, AlbaNova, SE-106 91 Stockholm, Sweden
²⁹ Royal Swedish Academy of Sciences Research Fellow, funded by a grant from the K. A. Wallenberg Foundation
³⁰ University of Maryland, Baltimore County, Baltimore, MD 21250
³¹ Dipartimento di Fisica, Università di Udine and Istituto Nazionale di Fisica Nucleare, Sezione di Trieste, Gruppo Collegato di Udine, I-33100 Udine, Italy
³² Istituto Universitario di Studi Superiori (IUSS), I-27100 Pavia, Italy
³³ Observatoire de Sciences de l’Univers, Université Joseph Fourier, BP 53, 38041 Grenoble CEDEX 9, France
³⁴ CNRS/IN2P3, Centre d’Études Nucléaires Bordeaux Gradignan, UMR 5797, Gradignan, 33175, France
³⁵ Université de Bordeaux, Centre d’Études Nucléaires Bordeaux Gradignan, UMR 5797, Gradignan, 33175, France
³⁶ Department of Physical Sciences, Hiroshima University, Higashi-Hiroshima, Hiroshima 739-8526, Japan
³⁷ University of Maryland, College Park, MD 20742
³⁸ University of Alabama in Huntsville, Huntsville, AL 35899
³⁹ Waseda University, 1-104 Totsukamachi, Shinjuku-ku, Tokyo, 169-8050, Japan
⁴⁰ Cosmic Radiation Laboratory, Institute of Physical and Chemical Research (RIKEN), Wako, Saitama 351-0198, Japan
⁴¹ Department of Physics, Tokyo Institute of Technology, Meguro City, Tokyo 152-8551, Japan
⁴² Centre d’Étude Spatiale des Rayonnements, CNRS/UPS, BP 44346, F-30128 Toulouse Cedex 4, France
⁴³ Istituto Nazionale di Fisica Nucleare, Sezione di Roma “Tor Vergata”, I-00133 Roma, Italy
⁴⁴ Department of Physics, Graduate School of Science, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan
⁴⁵ Max-Planck Institut für extraterrestrische Physik, 85748 Garching, Germany
⁴⁶ Department of Physics and Astronomy, University of Denver, Denver, CO 80208
⁴⁷ Sterrenkundig Instituut “Anton Pannekoek”, 1098 SJ Amsterdam, Netherlands
⁴⁸ Institut de Ciències de l’Espai (IEEC-CSIC), Campus UAB, 08193 Barcelona, Spain
⁴⁹ Institut für Astro- und Teilchenphysik, Leopold-Franzens-Universität Innsbruck, A-6020 Innsbruck, Austria
⁵⁰ Space Sciences Division, NASA Ames Research Center, Moffett Field, CA 94035-1000
⁵¹ NYCB Real-Time Computing Inc., Lattingtown, NY 11560-1025
⁵² Dipartimento di Fisica, Università di Roma “Tor Vergata”, I-00133 Roma, Italy
⁵³ Department of Chemistry and Physics, Purdue University Calumet,

Hammond, IN 46323-2094

⁵⁴ Institute of Space and Astronautical Science, JAXA, 3-1-1 Yoshinodai, Sagami-hara, Kanagawa 229-8510, Japan

⁵⁵ Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain

⁵⁶ Consorzio Interuniversitario per la Fisica Spaziale (CIFS), I-10133 Torino, Italy

⁵⁷ Center for Research and Exploration in Space Science and Technology (CRESTT), NASA Goddard Space Flight Center, Greenbelt, MD 20771

⁵⁸ School of Pure and Applied Natural Sciences, University of Kalmar, SE-391 82 Kalmar, Sweden

ing in HE gamma rays (for a full SED see Dubus 2006; Chernyakova, Neronov, & Walter 2006).

Calculations of the theoretical expectations of the gamma ray emission from LS I +61°303 go back almost three decades (Maraschi & Treves 1981), and there has been a recent burst of activity following the MAGIC detection. Two scenarios have been put forward involving either the relativistic wind of a young, rotation-powered pulsar (Dubus 2006; Sierpowska-Bartosik & Torres 2008, 2009), or the relativistic jet of an accreting black hole or neutron star (Romero, Christiansen, & Orellana 2005; Bednarek 2006; Gupta & Böttcher 2006; Bosch-Ramon et al. 2006). In light of the orbital modulations seen in radio, X-ray, and VHE gamma rays, a detailed light curve in the HE gamma ray domain (where most of the energy is output) is an essential piece to identify the main radiative process at work and model the source.

2. DATA REDUCTION AND RESULTS

The *Fermi Gamma-ray Space Telescope* was launched on 2008 June 11, from Cape Canaveral, Florida. The Large Area Telescope (LAT) is an electron-positron pair production telescope, featuring solid state silicon trackers and cesium iodide calorimeters, sensitive to photons from ~ 20 MeV to > 300 GeV (Atwood et al. 2009). Relative to earlier gamma-ray missions the LAT has a large ~ 2.4 sr field of view, a large effective area (~ 8000 cm 2 for > 1 GeV on axis) and improved angular resolution or point spread function (PSF, better than 1° for 68% containment at 1 GeV). The *Fermi* survey mode operations began on 2008 August 4, after the conclusion of a flawless commissioning period. In this mode, the observatory is rocked north and south on alternate orbits to provide more uniform coverage so that every part of the sky is observed for ~ 30 minutes every 3 hours. Thus *Fermi* is ideally suited for long term all-sky observations. The dataset for this analysis spanned 2008 Aug 4, through 2009 Mar 24. Thus LS I +61°303 was observed for approximately 9 orbital periods.

The data were reduced and analysed using the *Fermi* Science Tools v9r8 package⁵⁹. The standard onboard filtering, event reconstruction, and classification were applied to the data (Atwood et al. 2009), and for this analysis the high-quality ("diffuse") event class is used. Time periods when the region around LS I +61°303 was observed at a zenith angle greater than 105° were also excluded to avoid contamination from Earth albedo photons. With these cuts, a photon count map of a 10° region around the binary is shown in Fig. 1. The alignment of the LAT pointing direction with the celestial frame was calibrated using a large set of high latitude gamma-ray sources to better than $10''$ (Abdo et al. 2009b). The position of LS I +61°303 was found to be R.A. = $02^{\text{h}}40^{\text{m}}22.^{\text{s}}3$, Dec. = $61^\circ13'30''$ (J2000) with a 95% error of 0.069° ; in agreement with the accepted position (Dhawan, Mioduszewski, & Rupen 2006).

2.1. Spectral Analysis

The `glike` likelihood fitting tool was used to perform the spectral analysis, with "Pass 6 v3" (P6_V3) instrument response functions (IRFs); the P6_V3 IRFs are a post-launch update to address gamma-ray detection inefficiencies that are correlated with trigger rate. The 10 degree region around




FIG. 1.— The counts map for 100 MeV–300 GeV in (RA,Dec) of a 10° region around the LS I +61°303 location. The exposure varies by less than 2.5% across the field at a representative energy of 10 GeV. The source is bright and fairly isolated, sitting on a background of Galactic and extragalactic diffuse emission. A fit to the source yields a significance of more than 70σ . The dashed line indicates the Galactic equator ($b=0$); the crosses indicate the location of LS I +61°303 (the brighter source) and a faint nearby point source.




FIG. 2.— Fitted spectrum of LS I +61°303 to the phase-averaged *Fermi* data. The solid red lines are the $\pm 1\sigma$ limits of the *Fermi* cutoff power law; blue (open circle) data points from MAGIC (high state phases 0.5–0.7); black (filled circle) data points from VERITAS (high state phases 0.5–0.8). Data points in the *Fermi* range are likelihood fits to photons in those energy bins. Note that the data from the different telescopes are not contemporaneous, though they do cover multiple orbital periods.

the source was modeled for Galactic and extragalactic diffuse emission, and included one nearby point source at (R.A., Dec) of ($02^{\text{h}}23^{\text{m}}12^{\text{s}}$, $62^\circ0'0''$), too faint to be found in the 3-month Bright Source List (Abdo et al. 2009a). It is important to include this nearby source in the fitting model because at low energies the PSF is sufficiently wide that despite being $\sim 2.2^\circ$ away the PSF wings extend across the location of LS I +61°303 contributing approximately 13% to the flux at this position. Simultaneous modelling of this source accounts for its contribution to the flux in this region and any uncertainty is folded into the statistical error of the flux of LS I +61°303 found by the likelihood fitting tool. The 10° region was chosen to capture the broad PSF obtained at 100 MeV. An alternate fitting method using energy-dependent re-

⁵⁹ See the FSSC website for details of the Science Tools: <http://fermi.gsfc.nasa.gov/ssc/data/analysis/>




FIG. 3.— The 100 MeV to 20 GeV 1-day binned *Fermi* light curve of LS I +61°303 covering the period 2008 August 4 through 2009 March 24. The vertical lines indicate the zero phase from Gregory (2002).

gions of interest was used, yielding compatible results that were folded into the systematic errors.

The Galactic diffuse emission was modeled using GALPROP, described in Strong, Moskalenko, & Reimer (2004) and Strong (2007), updated to include recent H I and CO surveys, more accurate decomposition into Galactocentric rings, and many other improvements, including some from comparison with LAT data (Abdo et al. 2009b). The GALPROP run designation for our model is 54_59varh7S. The diffuse sources contribute $\sim 95\%$ of the observed photons shown in Fig. 1.

Initially a simple power law, $E^{-\Gamma}$, was fit to the orbital phase-averaged data yielding a photon index of $\Gamma \sim 2.42$. However, as indicated in Fig. 2, the energy spectrum appears to turn over at energies ~ 6 GeV. The possibility of an exponential cutoff was investigated, in the form $E^{-\Gamma} \exp[-(E/E_{\text{cutoff}})]$. The chance probability to incorrectly reject the power law hypothesis was found to be 1.1×10^{-9} . The best fit exponential cutoff returns a test statistic (Mattox et al. 1996) significance value of about 4770, or roughly 70σ . The photon index is $\Gamma = 2.21 \pm 0.04$ (stat) ± 0.06 (syst); the flux above 100 MeV is $(0.82 \pm 0.03$ (stat) ± 0.07 (syst)) $\times 10^{-6}$ ph cm $^{-2}$ s $^{-1}$ and the cutoff energy is 6.3 ± 1.1 (stat) ± 0.4 (syst) GeV (see below for a discussion of systematics). A total of 135,659 photons were found in the 10° region. Evaluating the fit parameters, 6467 ± 80 photons were observed from LS I +61°303 above 100 MeV. Fig. 2 shows the best fit cutoff power law model as well as the fluxes fit per energy bin and archival data from MAGIC (Albert et al. 2009)

and VERITAS (Acciari et al. 2008).

A number of effects are expected to contribute to the systematic errors. Primarily, these are uncertainties in the effective area and energy response of the LAT as well as background contamination. These are currently estimated by using outlier IRFs that bracket our nominal ones in effective area. These are defined by envelopes above and below the P6_V3 IRFs by linearly connecting differences of (10%, 5%, 20%) at $\log(E/\text{MeV})$ of (2, 2.75, 4) respectively. Other potential sources of systematic effects investigated are: fitting technique; cuts applied (zenith angle, minimum and maximum energies); and details of the diffuse modeling. The systematic errors estimated using the bracketing IRFs were found to be greater than these additional effects, hence the bracketing IRF results were quoted for the upper limits on the systematics.

2.2. Timing Analysis

LAT light curves were extracted using aperture photometry. The LAT point spread function is strongly energy dependent and, particularly since LS I +61°303 is located in the Galactic plane, there is also significant contribution to the flux within an aperture from diffuse emission and point sources that depends on the aperture size and the energy range used. The aperture and energy band employed were independently chosen to maximize the signal-to-noise level. The optimum aperture radius was found to be approximately 2.4° in the energy range 100 MeV–20 GeV. The time resolution of the light curve was 11,478 s, equal to twice the *Fermi* orbital period. Exposures were calculated using `gtexposure` and




FIG. 4.— Power spectrum of the light curve. The vertical line indicates the known orbital period from Gregory (2002), coinciding with a strong peak in the spectrum, while the horizontal lines indicate the marked significance levels.

used to determine the count rate in each time bin. In the exposure calculation, the spectral shape is assumed to be a power-law with a photon index of 2.4. The 1-day binned light curve is shown in Fig. 3. Contributions from the nearby source and Galactic and extragalactic diffuse backgrounds were estimated based on the spectral fit and subtracted from the light curve

A search was made for periodic modulation by calculating the periodogram of the light curve (Lomb 1976; Scargle 1982). Since the exposure of the time bins was variable, the contribution of each time bin to the power spectrum was weighted based on its relative exposure. The periodogram of the unbinned, unsmoothed light curve is shown in Fig. 4. The vertical line marks the Gregory (2002) orbital period and a highly significant peak is detected at this period. The significance levels marked are for a “blind” search with 500 independent frequency steps, however, the effects of the tuning of the aperture radius and energy range are not taken into account. The period and its error from the LAT observations were estimated using a Monte-Carlo approach: light curves were simulated using the observed LS I +61°303 light curve and randomly shuffling the data points within their statistical errors, assuming Gaussian statistics. The corresponding periodogram was then calculated and the location of the peak at ~ 26.5 days recorded. From $\sim 250,000$ simulations the distribution of values gives an estimation of the measured orbital period and its associated error of 26.6 ± 0.5 days (1σ).

The binned LAT light curve folded on the Gregory (2002) period with zero phase at MJD 43,366.2749 (Gregory et al. 1979) is shown in Fig. 5. The folded light curve shows a large modulation amplitude with maximum flux occurring slightly after periastron passage. The overall light curve can be fit reasonable well by a simple sine wave, yielding a reduced χ^2_ν of 1.4 for 1682 d.o.f. However, if we use the known orbital period and ephemeris of the system (Gregory 2002) to fit a sine wave to each of the individual 9 orbits observed then we find that the best fit amplitude varies between 6.8 ± 0.9 and $2.2 \pm 0.9 \times 10^{-7}$ ph cm $^{-2}$ s $^{-1}$, which suggests some orbit-to-orbit variability.

2.3. Phase resolved spectral analysis




FIG. 5.— Folded light curve of LS I +61°303 binned in orbital phase, see text for details. The dashed-lines indicate periastron and apastron as given by Aragona et al. (2009)

The possibility of the spectral shape changing across the orbit was explored by running `gtlike` fits for phase-folded bins of 0.1 width. The reduced statistics in each phase bin result in a cutoff not being statistically required to fit the data and so a simple power law model is used. There is no significant dependence of photon index on phase; a fit to a constant value returns a reduced χ^2_ν of 1.4 for 9 d.o.f, consistent with no variation.

3. DISCUSSION AND CONCLUDING REMARKS

The *Fermi* data enable for the first time the detection of a modulation in GeV gamma rays at the orbital period of a binary system. The derived period is in excellent agreement with the radio and optical-based ephemeris Gregory (2002). The COS-B source 2CG 135+01 is now firmly identified as the gamma-ray counterpart to LS I +61°303, resolving a 30-year long suspicion that the two were associated. With the identification originally based on localization only, the detection of orbital-modulated very high-energy emission (> 100 GeV, VHE) from LS I +61°303 by MAGIC and VERITAS (Albert et al. 2006, 2009; Acciari et al. 2008) had already provided very strong support in favour of this association.

LS I +61°303 is detected at a mean flux level above 100 MeV consistent with that seen by EGRET and *AGILE*. Averaged over the orbital modulation, the source persists as one of the brightest high-energy gamma-ray sources in the sky over a timescale of decades (see Abdo et al. 2009a, Bright Source List, in which this source is the 15th brightest). The folded *Fermi* light curve peaks around phase 0.3, which is compatible with periastron passage (when the compact object is closest to the Be star) according to the latest radial velocity studies (Aragona et al. 2009). This contrasts with the behavior at very high energies where peak flux occurs at phases 0.6-0.7 and detections are achieved only at phases ranging from 0.5 to 0.8, before or at apastron. In X-rays, LS I +61°303 also appears to peak at phases 0.6-0.7 (Paredes et al. 1997; Esposito et al. 2007) whereas the radio peak occurs over a wide range of phases depending upon a 4-year super-orbital cycle (Gregory 2002).

The average *Fermi* and EGRET spectra have compatible power law indices and fluxes taking into account systematics, but the *Fermi* spectrum also shows a cutoff at approxi-

mately 6 GeV. There is no evidence for a phase-dependence of the spectral shape and hence, the index or cutoff energy. VERITAS reports upper limits during the only VHE observations that are contemporary with *Fermi*, covering only part of one orbit from phase 0 to 0.75 (up to 2008 November 9, Holder et al. 2009). The later phases have short exposure times. Moreover, the past VHE history of the source shows several non-detections at phases 0.6–0.7 (Acciari et al. 2008; Albert et al. 2009), perhaps due to variability from one orbital cycle to the other. The *Fermi* light curve displays signs of orbit-to-orbit variability superposed on the mean behavior, with the primary peak always around phase 0.3. Such variability could be attributed to changing conditions in the Be star wind, affecting the interaction with the pulsar wind or relativistic jet. Indeed, optical spectra show evidence for changes in wind emission with the orbit (Zamanov et al. 1999).

The obvious radiative process to invoke in the HE and VHE range is inverse Compton scattering of the abundant stellar photons into gamma rays by a population of electrons accelerated in the vicinity of the compact object (e.g. in a relativistic jet or in a pulsar wind). Then, all else being equal, the peak flux phase is determined by where the seed photon density is highest and by geometry; favorable when the high energy electrons are seen behind the star by the observer, e.g. Dubus, Cerutti, & Henri (2008); Khangulyan, Aharonian, & Bosch-Ramon (2008); Sierpowska-Bartosik & Torres (2008). Superior conjunction is close in phase to periastron passage in LS I +61°303 ($\phi_{\text{per}} - \phi_{\text{sup}} = 0.07$ to 0.17 depending on the orbital solution). Hence, having the *Fermi* flux peak close to periastron is consistent with inverse Compton emission from electrons located close to the compact object. The cutoff in the average spectrum could arise due to radiative losses (because of different accelerating conditions for electrons, because of the magnetic field amplitude in the relativistic jet or the pulsar wind along the orbit and/or because of the greater photon density at periastron), or due to a varying maximum energy for accelerated electrons or to pair production on stellar photons for gamma rays above ≈ 50 GeV (Dubus 2006; Sidoli et al. 2006; Cerutti, Dubus, & Henri 2008; Sierpowska-Bartosik & Torres 2009). In the latter case, cascade emission might also be seen in the *Fermi* range. All these effects introduce phase-

dependent spectral changes. Hadronic interactions related to crossings of the Be star's equatorial wind (disk) could also contribute (Chernyakova, Neronov, & Walter 2006). This would provide an independently varying spectral component to explain why the HE and VHE emission peak at different phases and vary with orbital cycle. The expectation is that hadronic interactions would result in two asymmetric peaks in the light curve whose amplitude depends upon the intercepted matter density during the crossings and occurring at phases *a priori* unrelated to periastron passage but on the orientation of the orbit of the compact object relative to the Be star disk.

Continued monitoring by *Fermi* combined with dedicated campaigns by pointed instruments is needed to better constrain spectral variability and establish the multiwavelength connections: how do orbit-to-orbit variations compare in different energy ranges? Are there separate HE and VHE spectral components?

The *Fermi*-LAT Collaboration acknowledges generous ongoing support from a number of agencies and institutes that have supported both the development and the operation of the LAT as well as scientific data analysis. These include the National Aeronautics and Space Administration and the Department of Energy in the United States, the Commissariat à l'Énergie Atomique and the Centre National de la Recherche Scientifique / Institut National de Physique Nucléaire et de Physique des Particules in France, the Agenzia Spaziale Italiana and the Istituto Nazionale di Fisica Nucleare in Italy, the Ministry of Education, Culture, Sports, Science and Technology (MEXT), High Energy Accelerator Research Organization (KEK) and Japan Aerospace Exploration Agency (JAXA) in Japan, and the K. A. Wallenberg Foundation, the Swedish Research Council and the Swedish National Space Board in Sweden.

Additional support for science analysis during the operations phase from the following agencies is also gratefully acknowledged: the Spanish CSIC and MICINN and the Istituto Nazionale di Astrofisica in Italy.

We thank the anonymous referee for useful and constructive comments.

Facility: Fermi

REFERENCES

- Abdo A. A., et al., 2009a, *ApJ*, 700, 597
 Abdo A. A., et al., 2009b, *APh*, *submitted*, arXiv:0904.2226
 Acciari V. A., et al., 2008, *ApJ*, 679, 1427
 Aharonian F., et al., 2005a, *Sci*, 309, 746
 Aharonian F., et al., 2005b, *A&A*, 442, 1
 Albert J., et al., 2009, *ApJ*, 693, 303
 Albert J., et al., 2007, *ApJ*, 665, L51
 Albert J., et al., 2006, *Sci*, 312, 1771
 Aragona C., McSwain M. V., Grundstrom E. D., Marsh A. N., Roettenbacher R. M., Hessler K. M., Boyajian T. S., Ray P. S., 2009, *ApJ*, 698, 514
 Atwood W. B., et al., 2009, *ApJ*, 697, 1071
 Bednarek W., 2006, *MNRAS*, 368, 579
 Biggami G. F., Caraveo P. A., Lamb R. C., Markert T. H., Paul J. A., 1981, *ApJ*, 247, L85
 Bosch-Ramon V., Paredes J. M., Romero G. E., Ribó M., 2006, *A&A*, 459, L25
 Casares J., Ribas I., Paredes J. M., Martí J., Allende Prieto C., 2005, *MNRAS*, 360, 1105
 Cerutti B., Dubus G., Henri G., 2008, *A&A*, 488, 37
 Chernyakova M., Neronov A., Walter R., 2006, *MNRAS*, 372, 1585
 Dhawan V., Mioduszewski A., Rupen M., 2006, in *Proceedings of the VI Microquasar Workshop: Microquasars and Beyond* (Como, Italy), 52
 Dubus G., 2006, *A&A*, 456, 801
 Dubus G., Cerutti B., Henri G., 2008, *A&A*, 477, 691
 Eposito P., Caraveo P. A., Pellizzoni A., de Luca A., Gehrels N., Marelli M. A., 2007, *A&A*, 474, 575
 Gregory P. C., 2002, *ApJ*, 575, 427
 Gregory P. C., et al., 1979, *AJ*, 84, 1030
 Grundstrom E. D., et al., 2007, *ApJ*, 656, 437
 Gupta S., Böttcher M., 2006, *ApJ*, 650, L123
 Hartman R. C., et al., 1999, *ApJS*, 123, 79
 Hermsen W., et al., 1977, *Nature*, 269, 494
 Holder, J. et al., 2009, *proc. "High Energy Phenomena in Massive Stars"*, Jaen, Spain, in prep.
 Hutchings J. B., Crampton D., 1981, *PASP*, 93, 486
 Khangulyan D., Aharonian F., Bosch-Ramon V., 2008, *MNRAS*, 383, 467
 Kniffen D. A., et al., 1997, *ApJ*, 486, 126
 Lomb N. R., 1976, *Ap&SS*, 39, 447
 Maraschi L., Treves A., 1981, *MNRAS*, 194, 1P
 Mattox J. R., et al., 1996, *ApJ*, 461, 396
 Nolan P. L., Tompkins W. F., Grenier I. A., Michelson P. F., 2003, *ApJ*, 597, 615
 Paredes J. M., Martí J., Peracaula M., Ribo M., 1997, *A&A*, 320, L25
 Pittori C., et al., 2009, arXiv, arXiv:0902.2959
 Romero G. E., Christiansen H. R., Orellana M., 2005, *ApJ*, 632, 1093
 Seargle J. D., 1982, *ApJ*, 263, 835

- Sidoli L., Pellizzoni A., Vercellone S., Moroni M., Mereghetti S., Tavani M., 2006, *A&A*, 459, 901
- Sierpowska-Bartosik A., Torres D. F., 2009, *ApJ*, 693, 1462
- Sierpowska-Bartosik A., Torres D. F., 2008, *APh*, 30, 239
- Smith A., Kaaret P., Holder J., Falcone A., Maier G., Pandel D., Stroh M., 2009, *ApJ*, 693, 1621
- Strong A. W., 2007, *Ap&SS*, 309, 35
- Strong A. W., Moskalenko I. V., Reimer O., 2004, *ApJ*, 613, 962
- Swanenburg B. N., et al., 1981, *ApJ*, 243, L69
- Tavani M., Kniffen D., Mattox J. R., Paredes J. M., Foster R., 1998, *ApJ*, 497, L89
- Taylor A. R., Gregory P. C., 1982, *ApJ*, 255, 210
- Zamanov R. K., Martí J., Paredes J. M., Fabregat J., Ribó M., Tarasov A. E., 1999, *A&A*, 351, 543