SPIN-UNITARY SPIN SPLITTING OF SU(8) SUPERMULTIPLETS*

C. A. Nelson Department of Physics State University of New York at Binghamton Binghamton, New York 13901

ABSTRACT

The surprising narrowness of the J or $\psi(3, 1)$ is interpreted as indication of a pure $c \overline{c}$ state, and hence as evidence for the SU(8) \rightarrow SU(6) \times SU(2)_S \times U(1)_Y symmetry breaking chain $(\tilde{S}_c = charmed quark spin generators, Y_c = hypercharm$ generator) instead of an approximate $SU(8) \rightarrow SU(4) \times SU(2)_{S}$ chain $(\overline{S} = quark$ spin generators) which would imply strong mixings. Decompositions under both chains of the s wave $q \bar{q}$ meson states of the 64 = 1 + 63 of SU(8) and of the 3q baryon states of the three-particle symmetric 120 representation are given. The most general mass splitting operators with breaking in the Y and Y directions for these two multiplets are derived which commute with the Casimir operators of the SU(6) \times SU(2)_{S_c} \times U(1)_{Y_c} chain, which contain only one- and two-body operators, and which are invariant under rotations. Two independent mass relations follow for mesons containing charmed quarks; six, for baryons containing charmed quarks. In an appendix, for reference relative to previous SU(6) symmetric quark model mass analyses, the reduced numerical coefficients as determined by the meson 36 of SU(6) are listed.

(Submitted to Phys. Rev.)

*Work supported in part by the U.S. Energy Research and Development Administration and in part by a State University of New York Research Foundation Award.

I. INTRODUCTION

Our purpose is to discuss here two topics from the standpoint of the charmed symmetric quark model:

(i) The ground state 64 = 1 + 63 (meson) and 120 (baryon) representations of SU(8) together with their decompositions under the subgroups SU(6) × SU(2)_S × U(1)_Y and SU(4) × SU(2)_S where S stands for spin. The SU(2)_S subgroup acts on the c quark's spin, and Y_c is the hypercharm operator, see Section 3, with eigenvalues -1/4, -1/4, -1/4, and 3/4 for, respectively, the p, n, λ , and c type quarks.

(ii) The mass operator for these states in the $SU(6) \times SU(2)_{S_c} \times U(1)_{Y_c}$ chain which is derived by extending the one- and two-body force analysis of the SU(6) symmetric quark model¹ which previously gave the successful mass formulas¹⁻³ for the baryons, e.g., the Gürsey-Radicati formula⁴ for the 56 of SU(6) theory.^{4,5} Electromagnetic effects will be ignored.⁶

The motivation, of course, is the recent discovery⁷ of narrow resonances J or $\psi(3.1)$, and $\psi'(3.7)$, which can be interpreted as charmed⁸ quark-antiquark objects, $J^{PC} = 1^{--}$, $I^{G} = 0^{-}$, with N = 0 and 2 harmonic oscillator quanta excited, respectively. The N = 2 state is either a radial or an orbital excitation. It is important to recognize that present difficulties with the charm interpretation (the rise of R to 5.3 ± 0.6 at 7.8 GeV, the absence of narrow peaks in missing mass plots, the absence of increased kaon to pion production ratio, etc.), principally involve phenomena in $e \overline{e} \rightarrow$ hadrons above the transition region at about 3.6 - 4.1 GeV. Hence, these difficulties may, in fact, not exist if the transition region is due to excitation of first the charm, and then at about 3.9GeV of the color degrees of freedom which would be a natural occurrence⁹⁻¹¹ in the Han-Nambu version of the three-quartet model. In this case, the details of the discussion in this article apply to the SU(3) "-color singlet states. On the other hand, this type of mass analysis remains relevant, though not in detail, even if additional heavy quarks $^{12-14}$ are found to be necessary, because this analysis preserves successful SU(6) results and accepts the heavy quark explanation, based on the Zweig rule, of the narrowness of the new particles. A single charmed quark is certainly the simplest of such heavy quark models.

Lastly, we emphasize the basic contrast between (a) the present spectra and mass analyses in which the $c \overline{c}$ purity of the ψ and ψ' is given greatest importance, and (b) various previous analyses¹⁵ in which broken SU(4) is treated in analogy with broken SU(3) so as to derive mass relations and mass mixing angles and to predict specific mass values from existing data, but where $\psi \sim c \overline{c} + \epsilon (p \overline{p} + n \overline{n}) + \delta \lambda \overline{\lambda}$ results with $\epsilon, \delta \neq 0$.

We first discuss the mesons, Sections 2 and 3, and then the baryons, Section 4.

-3-

II. SPIN-UNITARY SPIN SPLITTING OF THE MESON 64 SUPERMULTIPLET OF SU(8)

We will make use of the well-known fact¹⁶ that the breaking of an approximate symmetry group can be simply expressed in terms of a "chain" of successively smaller subgroups which are valid to an increasingly better approximation. The prime example is the chain $SU(6) \rightarrow SU(3) \times SU(2)_{S_{\alpha}}$, S_{q} stands for the spin of the non-charmed quarks, with SU(3) \rightarrow SU(2)₁ × U(1)_V in SU(6) theory. Here, for instance, the hypercharge operator, which breaks the SU(3) symmetry, is conserved at the level of the smaller SU(2) $_{\rm I} \times$ U(1) $_{\rm Y}$ subgroup. The eigenvalues of commuting sets of generators in the chain provide quantum numbers with which to label, in practice uniquely, the states in the irreducible representations of the initial approximate symmetry group. Often two or more chains are relevant physically, and then superposition effects occur such that the physical resonances are eigenstates of neither chain. In the preceding example, there is also the chain SU(6) \rightarrow SU(4)_N × SU(2)_S × U(1)_Y with $SU(4)_{N} \rightarrow SU(2)_{I} \times SU(2)_{S_{N}}$, S_{N} stands for the spin of the n and p type quarks, whose existence is announced by the "mixing" of the I = Y = 0 pairs of s wave meson states, the $\phi - \omega$ and $\eta - \eta'$.

In the SU(8) theory there are two analogous reduction chains, the "SU(6) chain"

$$SU(8) \rightarrow SU(6) \times SU(2)_{S_{C}} \times U(1)_{Y_{C}}$$
 (1)

where the SU(6) subgroup is that discussed above; it acts on the p, n, and λ type quarks. The other chain is

$$SU(8) \rightarrow SU(4) \times SU(2)_{S}$$
 (2)

The SU(4) subgroup here acts on the p, ⁿ, λ and c type quarks; it does not involve spin and is not to be confused with the SU(4)_N subgroup of SU(6) theory. It has the further reduction

SU(4) \rightarrow SU(3) \times U(1)_{Y_c}

where SU(3) is the usual group for the p, n, and λ type quarks. This second chain we will call the "SU(4) chain."

The physical resonances, as we noted, need not be eigenstates of either chain so we will, first, consider the meson eigenstates in each chain separately. The ground state of an s wave, Fermi quark-antiquark pair with negative parity and spin J = S = 0, 1 is the reducible 64 of SU(8)

$$8 \times 8^* = 64 = 1 + 63$$
.

In the "SU(6) chain," the direct sum

$$64 = \begin{bmatrix} 1, 1 \end{bmatrix}_{0}^{0} + \begin{bmatrix} 35, 1 \end{bmatrix}_{0}^{0}$$
$$+ \begin{bmatrix} 6, 2^{*} \end{bmatrix}_{1}^{-1} + \begin{bmatrix} 6^{*}, 2^{1} \\ 1 \end{bmatrix}_{1}^{1} + \begin{bmatrix} 1, 1 \end{bmatrix}_{2}^{0} + \begin{bmatrix} 1, 3 \end{bmatrix}_{2}^{0}$$

with the notation dim SU(6), dim SU(2)_{S_c} $\begin{bmatrix} Y_c \\ y_c \end{bmatrix}_{n_c}^{Y_c}$ where n_c is the total number of

charmed quarks plus charmed antiquarks. Mesons associated with the first two representations contain no charmed quarks and are the familiar ones from the SU(6) theory. Continuing the chain, there next are the further reductions, $SU(6) \rightarrow SU(3) \times SU(2)_{S_q}$ with the notation (dim SU(3), dim SU(2)_{S_q})

$$1 = (1, 1)$$

$$35 = (8, 1) + (1, 3) + (8, 3)$$

$$6 = (3, 2)$$

$$6^* = (3^*, 2^*)$$

and similarly for the other SU(6) chain SU(6) \rightarrow SU(4)_N × SU(2)_{S_λ} × U(1)_Y with

the notation $\left(\dim SU(4)_{N}, SU(2)_{S_{\lambda}}\right)_{n_{\lambda}}^{Y}$

$$1 = (1, 1)_{0}^{0}$$

$$35 = (15, 1)_{0}^{0} + (4^{*}, 2)_{1}^{-1} + (4, 2^{*})_{1}^{1}$$

$$+ (1, 1)_{2}^{0} + (1, 3)_{2}^{0}$$

$$6 = (4, 1)_{0}^{0} + (1, 2)_{1}^{-1}$$

$$6^{*} = (4^{*}, 1)_{0}^{0} + (1, 2^{*})_{1}^{1}$$

The SU(6) \rightarrow SU(3) \times SU(2)_S chain yields, for the n_c \neq states, after recoupling the spins by $\overline{S} = \overline{S}_q + \overline{S}_c$,

$$\begin{bmatrix} 6, 2^* \end{bmatrix}_{1}^{-1} = (3, 2) 1^{-1} + (3, 2) 0^{-1}$$
$$\begin{bmatrix} 6^*, 2 \end{bmatrix}_{1}^{1} = (3^*, 2^*) 1^{-1} + (3^*, 2^*) 0^{-1}$$
$$\begin{bmatrix} 1, 1 \end{bmatrix}_{2}^{0} = (1, 1) 0^{-1}$$
$$\begin{bmatrix} 1, 3 \end{bmatrix}_{2}^{0} = (1, 1) 1^{-1}$$

with the notation (dim SU(3), dim SU(2)_{Sq})^{J^P}, J = S for the 64 representation. The (3*, 2*) 1⁻ consists of the isospin singlet F*+ = $(\overline{\lambda} c)^+$ and a doublet $D^{*+} = (\overline{n} c)^+$ and $D^{*0} = (\overline{p} c)^0$. The (3*, 2*) 0⁻ consists of a singlet F⁺ = $(\overline{\lambda} c)^+$ and a doublet $D^+ = (\overline{n} c)^+$ and $D^0 = (\overline{p} c)^0$. The $[6, 2^*]_1^{-1}$ contains their antiparticles. The $[1,3]_2^0$ is the J^P = 1⁻ isospin singlet $\phi_c^0 = (\overline{c} c)^0$, and the $[1,1]_2^0$ is the 0⁻ singlet $\eta_c^0 = (\overline{c} c)^0$.

On the other hand, for the "SU(4) chain" under SU(8) \rightarrow SU(4) × SU(2)_S these SU(8) representations decompose into

$$1 = \{1, 1\}$$

63 = {15, 1} + {1, 3} + {15, 3}

with the notation {dim SU(4), dim SU(2)_S}. Then, under SU(4) \rightarrow SU(3)×U(1)_Y_c the SU(4) representations decompose into

$$1 = 1_{2}^{0}$$

15 = 1_{0}^{0} + 8_{0}^{0} + 3_{1}^{1} + 3_{1}^{*-1}

with the notation dim SU(3) $_{n_c}^{Y_c}$. The corresponding wave functions can be easily written down; we only note that in this chain the eigenstates are superpositions of $\omega_8 - \phi_1 - \psi_c$, and of $\eta_8 - \eta_1 - \eta_c$.

For several reasons, we will assume that the SU(6) \times SU(2)_{S_c} \times U(1)_{Y_c} subgroup of SU(8) and the chain associated with it are of major importance for the breaking of SU(8) for mesons and baryons. First and foremost, the striking narrowness of the $\psi(3,1)$ and the $\psi'(3,7)$ suggest that they are pure $c \overline{c}$ states due to some new dynamical invariance principle, for example, n_c is exactly conserved in the strong interactions responsible for the mass spectra. In particular, the $\psi(3.1)$ will be identified with the $[1,3]_2^0$ irreducible representation of the SU(6) × SU(2) $S_c \times U(1) Y_c$ subgroup of SU(8). Note that decay modes such as $\psi \rightarrow 5\pi$ can go, for instance, via unitarity corrections rather than from mixings of the quark content of the ψ and ψ '. This¹⁷ has been pointed out in the context of Zweig rule suppressions, e.g., $\phi \rightarrow K\overline{K}$ and $K\overline{K} \rightarrow 3\pi$ both have connected duality diagrams so $\phi \rightarrow 3\pi$, with a hairpin diagram, can go via unitarity corrections which are difficult to distinguish from ϕ being other than an eigenstate of the SU(6) \rightarrow SU(4)_N × SU(2)_S × U(1)_Y chain. However, in the case of the ϕ resonance, the mass spectrum, see Appendix B, indicates that the latter $\phi \sim \lambda \overline{\lambda} + \epsilon \ (p \overline{p} + n \overline{n}), \ \epsilon \neq 0$, indeed occurs. Second, the lowest mesons can

be identified in the 64 of SU(8) with the 1 and 35 of SU(6), and this 1 + 35 can be identified with the $[1 + 35, 1]_0^0$ representation of the SU(6) × SU(2)_{S_c} × U(1)_{Y_c} subgroup of SU(8). Also, as discussed in Section 4, the lowest baryons can be similarly identified in the 120 of SU(8) with the 56 of SU(6), and this 56 can be identified with the $[56, 1]_0^0$ irreducible representation of the SU(6) × SU(2)_{S_c} × U(2)_{S_c} × U(1)_{Y_c} subgroup of SU(8).

Thus, in our derivation of the mass splitting operators for the s wave meson 64 = 1 + 63 and baryon 120 representations of SU(8), we will assume that the operator (i) commutes with the Casimir operators of the SU(6), SU(2) $_{
m S}$, and $U(1)_{Y}$ subgroups of this chain, and is invariant under rotations. We want the derivation to be a direct extension of SU(6) analyses in the symmetric quark model used to rederive¹ the Gürsey-Radicati result, used to study the first excited baryon multiplet, the $(70, 1)_1$ in the notation (dim SU(6), $L^p)_N$ with N the number of orbital or radial quanta excited in harmonic oscillator shells, and used³ to treat uniformly all of the baryon multiplets with N = 0, 1, or 2 harmonic oscillator excitation quanta. Hence, we will assume that the mass splitting operator (ii) contains only one- and two-body operators, 18 and that it (iii) transforms like a linear combination of three types of terms which transform, respectively, as a singlet, as the hypercharge operator under SU(3), and as the hypercharm operator under SU(4). For one-body operators, this transformation assumption is equivalent to mass splitting between the non-strange and strange quarks, and to an independent mass splitting between the non-charmed and charmed guarks.

III. MESON MASS OPERATOR AND INDEPENDENT MASS FORMULAE

We use a formalism in terms of the generators of SU(8) to derive the ground state SU(6) × SU(2)_S × U(1)_Y meson and baryon mass splitting operators. The 63 generators of SU(8), I_{Mr}^{Nr} with M = 1, 2, 3, 4 or p, n, λ , c for SU(4) and r = 1, 2 or \uparrow, \downarrow for SU(2)_S, are constructed from Fermi creation and annihilation operators for s-wave quarks and antiquarks in the charmed symmetric quark model in Appendix A. The SU(8) commutation relations, which can be easily computed from Eq. (A1), are

$$\begin{bmatrix} \mathbf{I}_{\mathrm{Mr}}^{\mathrm{Ns}}, \ \mathbf{I}_{\mathrm{M'r'}}^{\mathrm{N's'}} \end{bmatrix} = \boldsymbol{\delta}_{\mathrm{M'r'}}^{\mathrm{Ns}}, \ \mathbf{I}_{\mathrm{Mr}}^{\mathrm{N's'}} - \boldsymbol{\delta}_{\mathrm{Mr}}^{\mathrm{N's'}} \ \mathbf{I}_{\mathrm{M'r'}}^{\mathrm{Ns}}, \quad .$$
(3)

The charm operator C with eigenvalue 1 (-1) for a c quark (antiquark) and 0 for p, n, λ quarks and their antiquarks is not a linear combination of generators of SU(8) so we introduce the operator

$$Y_{c} \equiv C - \frac{3}{4}B = I_{c\uparrow}^{c\uparrow} + I_{c\downarrow}^{c\downarrow}$$
(4)

with B the baryon number operator. This relation is the analogue of $Y = S + B = \mathscr{I}_{\lambda\uparrow\uparrow}^{\lambda\uparrow} + \mathscr{I}_{\lambda\downarrow}^{\lambda\downarrow}$ which relates the hypercharge SU(3) generator and strangeness operator for the p, n, and λ quarks. The generators of SU(6) and its subgroups will be denoted by script letters to distinguish them from SU(8) generators. Since the c quark has B = 1/3, S = 0, the phenomenological extension to include the c quark is $Y = \mathscr{I}_{\lambda\uparrow\uparrow}^{\lambda\uparrow} + \mathscr{I}_{\lambda\downarrow}^{\lambda\downarrow} = B + S - \frac{1}{3}C$.

In Section 2, we discussed the relevant reduction chains which occur in SU(8). Generators¹⁹ for the subgroups in these chains are tabulated below:

-10-

I

:

Subgroup	Generators
SU(4)	$I_{M}^{N} = I_{Ms}^{Ns}$; M = 1, 2, 3, 4
SU(2) _S	$S_{r}^{s} = I_{Mr}^{Ms} = (S_{c} + \mathscr{P})_{r}^{s}$; r = 1, 2
SU(3)	$\mathscr{I}_{q}^{q'} = I_{q}^{q'} - \frac{1}{3} \delta_{q}^{q'} I_{p}^{p'} = I_{q}^{q'} + \frac{1}{3} \delta_{q}^{q'} Y_{c}; q = 1, 2, 3$
U(1) _Y c	$Y_c = I_c^c = -I_p^p$
SU(6)	$\mathscr{I}_{qr}^{q's} = I_{qr}^{q's} + \frac{1}{6} \delta_{q}^{q'} \delta_{r}^{s} Y_{c}$
SU(2) _{Sc}	$(\mathbf{S}_{c})_{r}^{s} = \mathbf{I}_{cr}^{cs} - \frac{1}{2} \boldsymbol{\delta}_{r}^{s} \mathbf{Y}_{c}$
SU(2) Sq	$(\mathscr{P})_{r}^{s} = I_{qr}^{qs} + \frac{1}{2} \delta_{r}^{s} Y_{c} = (\mathscr{P}_{N} + \mathscr{P}_{\lambda})_{r}^{s}$
SU(2) _I	$(\mathcal{I}_{N})_{m}^{n} = \mathcal{I}_{m}^{n} - \frac{1}{2} \delta_{m}^{n} Y; m = 1, 2$
U(1) _Y	$Y = -\mathcal{J}_3^3 = \mathcal{J}_m^m$
SU(4) _N	$(\mathcal{I}_{N})_{mr}^{ns} = \mathcal{I}_{mr}^{ns} - \frac{1}{4} \delta_{m}^{n} \delta_{r}^{s} Y$
SU(2) _{SN}	$(\mathscr{P}_{N})_{r}^{s} = \mathscr{P}_{mr}^{ms} - \frac{1}{2} \delta_{r}^{s} Y$
SU(2) _S λ	$(\mathscr{P}_{\lambda})_{r}^{s} = \mathscr{I}_{3r}^{3s} + \frac{1}{2} \delta_{r}^{s} Y$

The tensor operators in the mass formula will be expressed in terms of the Casimir operators for the various subgroups. Casimir operators needed are

$$C_{2}^{(8)} = \frac{1}{2} \begin{bmatrix} I_{Mr}^{Ns} , I_{Ns}^{Mr} \end{bmatrix}_{+}^{Ns}, \quad C_{2}^{(4)} = \frac{1}{2} \begin{bmatrix} I_{M}^{N} , I_{N}^{M} \end{bmatrix}_{+}^{Ns}$$

$$C_{2}^{(2)} (S) = \frac{1}{2} \begin{bmatrix} S_{r}^{S} , S_{r}^{r} \end{bmatrix}_{+}^{T} = 2S (S + 1)$$

$$C_{2}^{(6)} = \frac{1}{2} \begin{bmatrix} \mathcal{J}_{qr}^{q's} , \mathcal{J}_{q's}^{qr} \end{bmatrix}_{+}^{T} = \frac{1}{2} \begin{bmatrix} I_{qr}^{q's} , I_{q's}^{qr} \end{bmatrix}_{+}^{T} - \frac{1}{6} Y_{c}^{2}$$

$$C_{2}^{(3)} = \frac{1}{2} \begin{bmatrix} \mathcal{J}_{q}^{q'} , \mathcal{J}_{q'}^{q} \end{bmatrix}_{+}^{T} = \frac{1}{2} \begin{bmatrix} I_{qr}^{q's} , I_{qr}^{qr} \end{bmatrix}_{+}^{T} - \frac{1}{6} Y_{c}^{2}$$

$$C_{2}^{(3)} = \frac{1}{2} \begin{bmatrix} \mathcal{J}_{q}^{q'} , \mathcal{J}_{q'}^{q} \end{bmatrix}_{+}^{T} = \frac{1}{2} \begin{bmatrix} I_{q'}^{q's} , I_{q's}^{qr} \end{bmatrix}_{+}^{T} - \frac{1}{3} Y_{c}^{2}$$

$$C_{2}^{(2)} (I) = \mathcal{J}_{m}^{n} \mathcal{J}_{n}^{m} - \frac{1}{2} Y^{2} = 2I (I + 1)$$

$$C_{2}^{(4)} (N) = \mathcal{J}_{mr}^{ns} \mathcal{J}_{ns}^{mr} - \frac{1}{4} Y^{2}$$

and those for the several SU(2) subgroups describing the spins of particular sets of quarks. All these Casimir operators can be expressed as bilinear terms in the SU(8) generators of the form $[X, Y]_+$.

We can now derive²⁰ the mass splitting operator for the s wave meson 64 = 1 + 63. Our assumptions require the mass operator be a quadratic polynomial in the generators of SU(8), commute with $C_2^{(6)}$, $C_2^{(2)}$ (S_c), and Y_c^2 , be invariant under rotations, and transform like a linear combination of three types of terms which, respectively, transform as a singlet, as Y under SU(3), and as Y_c under SU(4). For mesons, the mass operator must be invariant under charge conjugation. We group the 63 generators of SU(8) into seven types: $I_q^{q'}$, I_c^c , I_q^c and I_c^q , I_{qr}^{qs} , I_{cr}^{cs} , $I_{qr}^{q's}$, I_{qr}^{cs} and I_{cr}^{qs} . Modulo pieces to make them traceless, $I_q^{q'}$ are the generators of SU(3); I_c^c is the generator of U(1)_{Y_c}; I_q^c and its adjoint, I_c^q , are generators of SU(4) not contained in the SU(3) and U(1)_{Y_c} subalgebras, etc.

-11-

The most general term linear in the generators and a scalar under rotations must be a linear combination of the generators of SU(4). Generators $I_q^{q'}$ and I_c^{c} are clearly admissible. We next consider the linear combination of the remaining generators $\mathscr{O} = a_c^q I_q^c + b_q^c I_c^q$. Using the identity

$$\left[\begin{bmatrix} X Y \end{bmatrix}_{+} Z \end{bmatrix}_{-} + \left[\begin{bmatrix} Y Z \end{bmatrix}_{+} X \end{bmatrix}_{-} + \left[\begin{bmatrix} Z X \end{bmatrix}_{+} Y \end{bmatrix}_{-} = 0$$
(5)

and Eq. (1),

$$\frac{1}{2} \left[\left[\mathbf{I}_{c}^{c}, \mathbf{I}_{c}^{c} \right]_{+}^{c}, \mathbf{a}_{c}^{q} \mathbf{I}_{q}^{c} + \mathbf{b}_{q}^{c} \mathbf{I}_{c}^{q} \right]_{-}^{c} \right]$$
$$= -\mathbf{a}_{c}^{q} \left[\mathbf{I}_{c}^{c}, \mathbf{I}_{q}^{c} \right]_{+}^{c} + \mathbf{b}_{q}^{c} \left[\mathbf{I}_{c}^{c}, \mathbf{I}_{c}^{q} \right]_{+}^{c} \right]_{-}^{c}$$

Since symmetrized expressions which have different numbers of different types of generators are linearly independent, the conditions that this commutator vanish are

$$a_{c}^{q} = b_{q}^{c} = 0$$
 (V q). (6)

Thus n_{λ} and n_{c} are the only admissible one-body terms which satisfy the transformation requirement and which are invariant under charge conjugation.

For bilinear terms in the generators of SU(8), invariance under rotations implies that there are two classes of terms that can be considered separately: those constructed from $I_q^{q'}$, I_c^c , I_q^c and I_c^q , and those constructed from I_{qr}^{qs} , I_{cr}^{cs} , $I_{qr}^{q's}$, I_{qr}^{cs} and I_{cr}^{qs} . We write the terms quadratic in the generators in the form $[XY]_+$ so that they are linearly independent of the terms linear in the generators. The most general term in $I_q^{q'}$, I_c^c , and I_q^c and its adjoint is a linear combination of the six expressions of the form $[XY]_+$. Of these, only $\begin{bmatrix} I_q^{q'}, I_p^{c} \end{bmatrix}_+$ when commuted with $\frac{1}{2} \begin{bmatrix} I_c^{c}, I_c^{c} \end{bmatrix}_+$ yields combinations of the form

 $\left[\mathbf{I}_{c}^{c},\left[\mathbf{I}_{q}^{q'},\mathbf{I}_{p}^{c}\right]_{+}\right]_{+}$

so these can be considered separately. The vanishing of

$$\frac{1}{2} \left[\begin{bmatrix} \mathbf{I}_{c}^{c}, \mathbf{I}_{c}^{c} \end{bmatrix}_{+}^{}, \mathbf{a}_{qq'p} \begin{bmatrix} \mathbf{I}_{q}^{q'}, \mathbf{I}_{p}^{c} \end{bmatrix}_{+} \right]$$
$$= -\mathbf{a}_{qq'p} \begin{bmatrix} \mathbf{I}_{c}^{c}, \begin{bmatrix} \mathbf{I}_{q}^{q'}, \mathbf{I}_{p}^{c} \end{bmatrix}_{+} \end{bmatrix}_{+}$$

implies that

$$a_{qq'p} = 0$$
 (Yq, q', p) . (7)

The adjoint $\begin{bmatrix} I_{q'}^{q}, I_{c}^{p} \end{bmatrix}_{+}^{+}$ is also eliminated. The same argument with $I_{q}^{q'} \rightarrow I_{c}^{c}$ eliminates terms of the form $\begin{bmatrix} I_{c}^{c}, I_{q}^{c} \end{bmatrix}_{+}^{c}$ and its adjoint. Similarly the terms $\begin{bmatrix} I_{q}^{c}, I_{q'}^{c} \end{bmatrix}_{+}^{c}$ and their adjoint are excluded. Under commutation with

$$\frac{1}{2} \begin{bmatrix} \mathbf{I}_{\mathrm{cr}}^{\mathrm{cs}}, & \mathbf{I}_{\mathrm{cs}}^{\mathrm{cr}} \end{bmatrix}_{+},$$

only

$$\begin{bmatrix} I_q^c, I_c^{q!} \end{bmatrix}_+$$

leads to combinations of

$$\left[I_{q}^{c}, \left[I_{cs}^{cr}, I_{cr}^{q's}\right]_{+}\right]_{+}$$

and its adjoint; so, this term can be considered separately. The vanishing of

$$\frac{1}{2} \left[\begin{bmatrix} \mathbf{I}_{cr}^{cs}, \mathbf{I}_{cs}^{cr} \end{bmatrix}_{+}^{*}, \mathbf{a}_{qq'} \begin{bmatrix} \mathbf{I}_{q}^{c}, \mathbf{I}_{c}^{q'} \end{bmatrix}_{+}^{*} \right]$$
$$= \mathbf{a}_{qq'} \left\{ \begin{bmatrix} \mathbf{I}_{q}^{c}, \begin{bmatrix} \mathbf{I}_{cs}^{cr}, \mathbf{I}_{cr}^{q's} \end{bmatrix}_{+}^{*} \right]_{+}^{*}$$
$$- \begin{bmatrix} \mathbf{I}_{c}^{q'}, \begin{bmatrix} \mathbf{I}_{cs}^{cs}, \mathbf{I}_{qs}^{cr} \end{bmatrix}_{+}^{*} \end{bmatrix}_{+}^{*}$$

implies that

$$a_{qq'} = 0$$
 ($\forall q, q'$). (8)

This then leaves as admissible terms

$$\begin{bmatrix} I_{q}^{q'}, I_{p}^{p'} \end{bmatrix}_{+}, \begin{bmatrix} I_{c}^{c}, I_{c}^{c} \end{bmatrix}_{+}, \text{ and } \begin{bmatrix} I_{q}^{q'}, I_{c}^{c} \end{bmatrix}_{+}$$

Linear combinations of these lead to

$$\left[\mathcal{J}_{q}^{p}, \mathcal{J}_{p}^{q} \right]_{+}$$

which transforms as a singlet,

$$\mathbf{y}_{\mathbf{q}}^{\mathbf{q}'} \left[\mathscr{I}_{\mathbf{q}'}^{\mathbf{p}}, \mathscr{I}_{\mathbf{p}}^{\mathbf{q}} \right]_{+}$$

which transforms as Y, and

$$c_{M}^{N} \left[I_{N}^{O}, I_{O}^{M} \right]_{+} - \frac{1}{4} \left[I_{N}^{O}, I_{O}^{N} \right]_{+}$$

which transforms as Y_c and a singlet. Here the numerical diagonal matrices $y_q^{q'}$ = diag (1/3, 1/3, -2/3) and c_M^N = diag (-1/4, -1/4, -1/4, 3/4) have been used.

Next, we discuss the second class of bilinear terms: those formed from generators transforming as vectors under rotations. These terms must also commute with the Casimir operators $C_2^{(6)}$, $C_2^{(2)}(S_c)$ and with Y_c^2 so the candidates are the rotation-vector analogues of the surviving first class terms; plus those constructed with $(S_c)_r^s$ and $(\mathscr{S})_r^s$. These analogues are

$$\begin{bmatrix} I_{qr}^{q's}, I_{ps}^{p'r} \end{bmatrix}_{+}$$
, $\begin{bmatrix} I_{cr}^{cs}, I_{cs}^{cr} \end{bmatrix}_{+}$, and $\begin{bmatrix} I_{qr}^{q's}, I_{cs}^{cr} \end{bmatrix}_{+}$

Since the admissible terms linear in the generators are $I_q^{q'}$ and I_c^{c} , the only candidates constructed from $(S_c)_r^s$ and $(\mathscr{G})_r^s$ are

$$\left[\left(\mathbf{S}_{c} \right)_{r}^{s}, \left(\mathbf{S}_{c} \right)_{s}^{r} \right]_{+}, \left[\left(\mathscr{I} \right)_{r}^{s}, \left(\mathscr{I} \right)_{s}^{r} \right]_{+}, \left[\left(\mathbf{S}_{c} \right)_{r}^{s}, \left(\mathbf{I}_{cs}^{r} \right)_{+}^{s} \right]_{+}$$

and

 $\left[\left(\mathcal{G}\right)_{r}^{s}, I_{qs}^{q'r}\right]_{+}.$

Commutation with Y_c^2 , $C_2(S_c)$, and $C_2^{(6)}$ shows that all these terms enter. The additional linear combinations of bilinear terms, satisfying the transformation requirement, are

$$\begin{bmatrix} \mathscr{I}_{qr}^{ps}, \mathscr{I}_{ps}^{qr} \end{bmatrix}_{+}, \quad y_{q}^{q'} \begin{bmatrix} \mathscr{I}_{q'r}^{ps}, \mathscr{I}_{ps}^{qr} \end{bmatrix}_{+}, \quad c_{M}^{N} \begin{bmatrix} I_{Ns}^{Or}, I_{Or}^{Ns} \end{bmatrix}_{+} \\ -\frac{1}{4} \begin{bmatrix} I_{Ns}^{Or}, I_{Or}^{Ns} \end{bmatrix}_{+}, \quad \begin{bmatrix} s_{r}^{s}, s_{s}^{r} \end{bmatrix}_{+}, \\ y_{q}^{p} \begin{bmatrix} \mathscr{I}_{ps}^{qr}, \mathscr{I}_{r}^{s} \end{bmatrix}_{+}, \quad \text{and} \quad c_{M}^{N} \begin{bmatrix} I_{Ns}^{Mr}, s_{r}^{s} \end{bmatrix}_{+}.$$

Finally, we collect the admissible terms and obtain the mass formula for

the s wave meson 64 = 1 + 63. The terms, together with their form in terms of quantum numbers, are tabulated below:

Form in terms of quantum numbers	In form of generators
<u>2 Linear Terms:</u>	
nλ	$Y(q) - Y(\overline{q})$
ⁿ c	$Y_{c}(q) - Y_{c}(\overline{q})$
9 Bilinear Terms:	
C ₂ ⁽⁶⁾	$\frac{1}{2} \left[\mathscr{I}_{\mathrm{qr}}^{\mathrm{ps}}, \mathscr{I}_{\mathrm{ps}}^{\mathrm{qr}} \right]_{\mathrm{+}}$
2 S (S+1)	$\frac{1}{2} \left[s_r^s, s_s^r \right]_+$
C ₂ ⁽³⁾	$\frac{1}{2} \left[\mathscr{I}_{q}^{p}, \mathscr{I}_{p}^{q} \right]_{+}$
I (I+1) - $\frac{1}{4}$ Y ² - $\frac{1}{6}$ C ⁽³⁾ ₂	$\frac{1}{2} y_{q}^{q'} \left[\mathscr{I}_{q'}^{p}, \mathscr{I}_{p}^{q'} \right]_{+}$
$2 S_{\lambda} (S_{\lambda} + 1) - C_{2}^{(4)} (N) + \frac{1}{4}Y^{2} + \frac{1}{3}C_{2}^{(6)}$	$-y_q^{q'} \left[\mathscr{I}_{q'r}^{ps}, \mathscr{I}_{ps}^{qr} \right]_+$
$\mathbf{S_N}(\mathbf{S_N}+1) - \mathbf{S_\lambda}(\mathbf{S_\lambda}+1) - \frac{1}{3}\mathbf{S_q}(\mathbf{S_q}+1)$	$\left[\frac{1}{2}\left[\mathscr{G}_{r}^{s}, y_{q}^{p}\mathscr{G}_{ps}^{qr}\right]_{+}\right]$
$\frac{2}{3} Y_c^2 - C_2^{(3)}$	$c_{M}^{N} \left[I_{N}^{O}, I_{O}^{M} \right]_{+} - \frac{1}{4} \left[I_{N}^{O}, I_{O}^{N} \right]_{+}$
$2 S_{c} (S_{c} + 1) + \frac{1}{3} Y_{c}^{2} - C_{2}^{(6)}$	$\left[c_{M}^{N} \left[I_{Ns}^{Or}, I_{Or}^{Ns} \right]_{+} - \frac{1}{4} \left[I_{Ns}^{Or}, I_{Or}^{Ns} \right]_{+} \right]_{+}$
$S_q (S_q + 1) - S_c (S_c + 1) - \frac{1}{2} S (S + 1)$	$-\frac{1}{2}\left[S_{r}^{s}, c_{M}^{N}I_{Ns}^{Mr}\right]_{+}$

I

...

These give the following twelve parameter mass formula:

$$M = m_{0} + m_{1}n_{\lambda} + m_{2}C_{2}^{(6)} + m_{3} 2S(S+1) + m_{4}C_{2}^{(3)}$$

$$+ m_{5} \left[I(I+1) - \frac{1}{4}Y^{2} \right] + m_{6} \left[2S_{\lambda}(S_{\lambda}+1) - C_{2}^{(4)}(N) + \frac{1}{4}Y^{2} \right]$$

$$+ m_{7} \left[2S_{N}(S_{N}+1) - 2S_{\lambda}(S_{\lambda}+1) - \frac{1}{3} 2S_{q}(S_{q}+1) \right]$$

$$+ m_{8}n_{c} + m_{9}Y_{c}^{2} + m_{10} 2S_{c}(S_{c}+1)$$

$$+ m_{11} 2S_{q}(S_{q}+1)$$
(9)

This mass formula predicts the following independent equalities:

$$F^* - D^* = F - D$$
 (10)

the strange-nonstrange mass differences for the pseudoscalar and vector SU(3) triplets, and antitriplets, are equal; and from the observed mass splittings of the states in the meson 36 of SU(6), the magnitude of this mass difference

F - D = 76 MeV linear mass formula

$$= 0.074 \text{ GeV}^2 \quad \text{quadratic mass formula} \tag{11}$$

The numerical reduced coefficients as determined by the 36 are given in Appendix B. The quadratic formula for a D mass of 2 GeV implies an almost degenerate F mass of 2.02 GeV. Equation (10), as well as Eq. (11), is an SU(8) prediction, since the four states are in the $[6, 2^*]_1^{-1}$ multiplet of SU(6)×SU(2)_{S_c}× U(1)_{Y_c} which involves recoupling the charmed and non-charmed quark spins.

Note that in Eq. (9) the SU(6) breaking terms enter with the same coefficients for each SU(6) multiplet. This is the same as for the coefficients of the SU(3) breaking terms in the Gürsey-Radicati formula and in the SU(8) baryon mass formula obtained below, Eq. (13). Here Eq. (9) mixes the 1 and 63 of SU(8) only as a consequence of the standard 1 and 35 mixing of η_1 and η_8 in the SU(6) theory. Relative to the SU(6) \rightarrow SU(4)_N × SU(2)_{S₁} chain, the $\eta - \eta$ ' mixing is due to the $C_2^{(6)}$ and $C_2^{(3)}$ terms; the breaking of ideal $\phi - \omega$ mixing is due to the $C_2^{(3)}$ term, i.e., it is solely responsible for ϕ not being a pure $\lambda \overline{\lambda}$ state. The choice $M_{\omega} = M_{\rho}$ requires, in addition to $C_2^{(3)}$ being absent, the absence of $\left[I(I + 1) - \frac{1}{4}Y^2\right]$. From the viewpoint of an SU(3) singlet-octet system, n_{λ} mixes the singlet and octet whereas $\left[I(I + 1) - \frac{1}{4}Y^2\right]$ only breaks the octet; however, the m_6 and m_7 terms also mix the states and break the octet. If systematic use of SU(6) is used to classify the terms, as in the SU(6) irreducible tensor approach, the m_5 , m_6 , and m_7 terms arise 16,21 from SU(6) tensors ${}^{35}T^{8,1}_{35}$, ${}^{35}T^{8,1}_{189}$, and ${}^{35}T^{8,1}_{405}$. Irreducible tensor operators are labeled ${}^{m}T_{dim SU(3), dim SU(2)S}^{dim SU(2)}$ where "m" specifies the SU(6) state of q and/or \overline{q} . Experience $2^{2,3}$ with baryon levels and past confusions 2^{22} over inadequate meson mass operators indicate that such operators with the larger SU(6) representations should not be excluded, but should be retained as has been done here. This means that only mixing angles can be predicted for the s wave mesons of the 36 of SU(6); however, these predictions alone are significant for decay tests.

IV. MASS OPERATOR AND INDEPENDENT MASS FORMULAE FOR THE BARYON 120 SUPERMULTIPLET

For completeness, we first discuss the "SU(4) chain" reduction of the totally symmetric three-particle representation of SU(8), the 120, in which we place the baryons. As in the symmetric quark model, to be consistent with the spin and statistics theorem, we assume that there exists an SU(3)" -color degree of freedom and that the states in the 120 are in the totally antisymmetric three-particle representation of SU(3)" -color, the singlet. The direct sum

$$120 = \{20_{\rm s}, 4\} + \{20_{\rm m}, 2\}$$

in the SU(8) \rightarrow SU(4) \times SU(2)_S chain with the notation {dim SU(4), dim SU(2)_S} where a permutation symmetry subscript, s = symmetric and m = mixed, suffices to distinguish the two twenty-tuplet Young diagrams. Under SU(4) \rightarrow SU(3) \times U(1)_{Y₂},

$$20_{m} = 8_{0} + 6_{1} + 3_{1}^{*} + 3_{2}$$
$$20_{s} = 10_{0} + 6_{1} + 3_{2} + 1_{3}$$

with the notation dim $SU(3)_{n_c}$. Note that $n_c = C$, i.e., n_c has the charm eigenvalue, for states containing no antiquarks. While the SU(3) decuplet and octet are obtained by this reduction, their physical relation as submultiplets of the 56 of SU(6) is not made manifest by the SU(4) reduction chain. Hence, we return to the SU(6) chain.

The relevant reductions of the 120 under the SU(8) \rightarrow SU(6) \times SU(2) ${\rm S_{c} \times U(1)}_{\rm Y_{c}}$ chain are

$$120 = [56, 1]_0 + [21, 2]_1 + [6, 3]_2 + [1, 4]_3$$

with the notation, as before, of $\left[\dim SU(6), \dim SU(2)_{S_{c}}\right]_{n_{c}}$ and then under

$$SU(6) \rightarrow SU(3) \times SU(2)_{S_q}$$

 $56 = (10, 4) + (8, 2)$
 $21 = (6, 3) + (3^*, 1)$
 $6 = (3, 2)$
 $1 = (1, 1)$

with the notation $(\dim SU(6), \dim SU(2)_{S_q})$. In order to obtain the physical states with $n_c \neq 0$, the spin of the charmed and non-charmed quarks must be recoupled, i.e., $\overline{S} = \overline{S_q} + \overline{S_c}$. This yields $[21, 2]_1 = (6, 3)\frac{3}{2}^+ + (6, 3)\frac{1}{2}^+ + (3^*, 1)\frac{1}{2}^+$ $[6, 3]_2 = (3, 2)\frac{3}{2}^+ + (3, 2)\frac{1}{2}^+$

 $\begin{bmatrix} 1, 4 \end{bmatrix}_3 = (1, 1) \frac{3^+}{2}$

with the notation $(\dim SU(3), \dim SU(2)_{Sq})^{J^{P}}$, S = J for the 120 representation. It is a straightforward exercise to tabulate the wave functions for the thirteen charmed states, and from their composition in terms of p, n, λ and c type quarks to read off their respective I, B, Y and C quantum numbers. The (6,3) consists of an isotopic spin singlet $(\lambda \lambda c)^{O}$, a doublet $(n \lambda c)_{S}^{O}$ and $(p \lambda c)_{S}^{+}$, and a triplet $(nn c)^{O}$, $(np c)_{S}^{+}$ and $(pp c)^{++}$. The (3*, 1) consists of a doublet $(n \lambda c)_{A}^{O}$ and $(p \lambda c)_{A}^{+}$, and a singlet $(np c)_{A}^{+}$. The (3, 2) consists of a singlet $(\lambda c c)^{+}$ and a doublet $(n c c)^{+}$ and $(p c c)^{++}$. The S and A subscripts denote the permutation symmetry of the two-particle combination of p, n, λ quarks.

For the 120 representation of SU(8) there exists the identify

$$2S_{c}(S_{c} + 1) = C_{c} + \frac{1}{2}C_{c}^{2}$$
(12)

-20-

Thus, by the derivation of Sec. 3, the most general $SU(6) \times SU(2)_{S_c} \times U(1)_{Y_c}$ mass formula for the 120 which satisfies our conditions has eleven parameters and is

$$M = m_{0} + m_{1}Y + m_{2}C_{2}^{(3)} + m_{3}\left[I(I+1) - \frac{1}{4}Y^{2}\right]$$

$$+ m_{4}C_{2}^{(6)} + m_{5}2S(S+1) + m_{6}Y_{c} + m_{7}Y_{c}^{2}$$

$$+ m_{8}\left[2S_{\lambda}(S_{\lambda}+1) - C_{2}^{(4)}(N) + \frac{1}{4}Y^{2}\right]$$

$$+ m_{9}\left[2S_{N}(S_{N}+1) - 2S_{\lambda}(S_{\lambda}+1)\right] + m_{10}2S_{q}(S_{q}+1)$$
(13)

On the 56 of SU(6) only the first four terms, the Gürsey-Radicati formula, are independent. Here, as in SU(6) theory, for baryons the two-body dominance assumption has led to a significant simplification.

For the 56 of SU(6) the Gürsey-Radicati formula yields four independent sum rules. For the thirteen additional states in the 120 of SU(8), Eq. (13) predicts the following six new independent equalities:

$$\left[(ppc) - (p\lambda c)_{S} \right]_{(6,3)} \frac{1}{2}^{+} = \left[(p\lambda c)_{S} - (\lambda\lambda c) \right]_{(6,3)} \frac{1}{2}^{+}$$
(14)

$$\begin{bmatrix} (ppc) - (p\lambda c)_{S} \\ (6,3) \frac{3^{+}}{2} \end{bmatrix} = \begin{bmatrix} (p\lambda c)_{S} - (\lambda\lambda c) \\ (6,3) \frac{3^{+}}{2} \end{bmatrix}$$
(15)

$$\begin{bmatrix} (pcc) - (\lambda cc) \\ (3, 2) \frac{1}{2}^{+} \end{bmatrix}_{(3, 2)}^{+} = \begin{bmatrix} same \\ (3, 2) \frac{3}{2}^{+} \end{bmatrix}_{(3, 2)}^{+} = \begin{bmatrix} (ppc) \\ (6, 3) \frac{3}{2}^{+} \end{bmatrix}_{(3, 2)}^{+} = \begin{bmatrix} (ppc) \\ (6, 3) \frac{1}{2}^{+} \end{bmatrix}_{(6, 3)}^{+} = \begin{bmatrix} (ppc) \\ (6, 3) \frac{1}{2}^{+} \end{bmatrix}_{(6, 3)}^{+} = \begin{bmatrix} (ppc) \\ (6, 3) \frac{1}{2}^{+} \end{bmatrix}_{(6, 3)}^{+} = \begin{bmatrix} \frac{1}{5} \begin{bmatrix} (npc \\ A - (p\lambda c) \\ A \end{bmatrix}_{(3^{*}, 1)}^{+} = \begin{bmatrix} \frac{4}{5} \left\{ N - \frac{1}{4} (\Lambda + 3\Sigma) \right\} \\ = -188 \text{ MeV} \end{bmatrix}_{(19)}^{+}$$

where the subscripts denote $(\dim SU(3), \dim SU(2)_{S_q})^{J^P}$. Equalities (14) and (15) specify equal spacing for both of the two SU(3) sextets, Eq. (16) specifies that this spacing is also common, Eq. (17) specifies a common spacing for the two SU(3) triplets, Eq. (18) specifies the same separation between isotopic spin multiplets in the $J^P = 3/2^+$ and $1/2^+$ levels for the triplets as for the sextets, and Eq. (19) specifies a relation between the splitting of the anti-triplet and those of the other charm levels and the nucleon octet.

Equations (14) and (15) are SU(3) equal spacing statements. Both Eq. (16) and (17) are SU(8) results for recoupling of \overline{S}_q and \overline{S}_c is involved, and clearly Eq. (18) and (19) are SU(8) results.

-22-

V. SUMMARY

We again emphasize, from the point-of-view of future ψ spectroscopy, that the mass relations derived in this article preserve $c \ c \ c$ purity of the $\psi(3, 1)$ resonance. We studied the charmed symmetric quark model for mesons and baryons using approximate SU(6) × SU(2) $S_c \times U(1)_{Y_c}$ symmetry with breaking in the Y and Y_c directions in order to resolve mass degeneracies among resonances in the same sub-multiplets. To reduce the number of possible mass formulas for baryons, we assumed that one- and two-body contributions to the mass splitting operator dominate. For the six meson levels containing charmed quarks, we predicted two new independent mass relations. For the corresponding thirteen new baryon levels, we predicted six new mass relations. In an appendix, for reference relative to previous SU(6) symmetric quark model mass analyses, we gave the reduced numerical coefficients as determined by the meson 36 of SU(6).

ACKNOWLEDGMENTS

The author wishes to thank the Joint Awards Council/University Awards Committee for a State University of New York Research Foundation Award. He also thanks SLAC for its fine hospitality during the completion of this work.

APPENDIX A: SECOND QUANTIZED FORMALISM

We introduce a set of Fermi creation and annihilation operators¹ for the s-wave quarks and antiquarks, $a_{Mr}^{q''\dagger}$, $a_{p''}^{Ns}$, $b_{q''}^{Mr\dagger}$, and $b_{Ns}^{p''}$ where M = 1, 2, 3, 4 for SU(4), r = 1, 2 for SU(2)_S, and q'' = 1, 2, 3 for SU(3)"-color. The spin and SU(4) indices for the antiquark operators can be grouped together since complex conjugate representations in SU(2) are unitarily equivalent to the original ones. The generators of SU(8) constructed in terms of these are

$$I_{Mr}^{Ns} = I(q)_{Mr}^{Ns} + I(\overline{q})_{Mr}^{Ns}$$
(A1)

with

$$I(q)_{Mr}^{Ns} = a_{Mr}^{q''} a_{q''}^{Ns} - \frac{1}{8} \delta_{Mr}^{Ns} a_{Ot}^{q''} a_{q''}^{Ot}$$

$$I(\overline{q})_{Mr}^{Ns} = -\left(b_{q''}^{Ns\dagger} b_{Mr}^{q''} - \frac{1}{8} \delta_{Mr}^{Ns} b_{q''}^{Ot\dagger} b_{Ot}^{q''}\right)$$
(A2)

From these expressions, $Y_c(q) = \frac{1}{4}(-N_q + 3N_c)$ and $\overline{Y}_c(\overline{q}) = -\frac{1}{4}(-N_{\overline{q}} + 3N_{\overline{c}})$. These lead in the s-wave meson mass splitting operator, for a system composed of a fixed number of quarks and antiquarks, to a single charge conjugation in-variant term.

$$\mathbf{n}_{c} = \mathbf{N}_{c} + \mathbf{N}_{\overline{c}} = \left(\mathbf{Y}_{c}(\mathbf{q}) - \overline{\mathbf{Y}}_{c}(\overline{\mathbf{q}})\right) + \frac{1}{4}\left(\mathbf{N} + \overline{\mathbf{N}}\right) . \tag{A3}$$

This is the number operator for the total number of charmed quarks and antiquarks.

From the other linearly independent terms in the $SU(6) \times SU(2)_{S_c} \times U(1)_{Y_c}$ meson and baryon mass splitting operators in the text, this second quantized formalism can also be used to extract specific dynamical parameters characterizing single quarks and the two-body inter-quark forces. Such an explicit interpretation in the three-quartet model of the forces responsible for the observed hadronic mass splittings brings these mass operators in closer contact with more basic quantum field theory approaches to quark dynamics, for example, gauge fields on a lattice and the bag model.

APPENDIX B: NUMERICAL COEFFICIENTS AS DETERMINED BY MESON 36 OF SU(6)

On the s-wave meson 36 multiplet of SU(6) the terms in the meson mass formula derived in the text, Eq. (9), reduce to the first eight terms. For each term a normalization factor $\mathcal{N}_i = \left(T_{\max}^i - T_{\min}^i + 1\right)^{-1}$ is introduced so as to treat them in a comparable manner. It is in order of the terms in Eq. (9), 1, 1/3, 1/13, 1/5, 1/7, 1/5, 1/13, and 1/9. The reduced coefficients, $M_i = m_i^2 / N_i$, for the linear (quadratic) mass formula as determined from the experimental data²³ are 1.000, -0.489, -1.853, 0.594, 1.800, -0.411, 0.593, and 0.530 (0.975, -0.353, -2.364, 0.697, 1.967, -0.710, -0.176, and 0.054) in units of GeV (GeV²). The last term does not contribute significantly to the quadratic mass formula. Otherwise, for a simultaneous treatment of $J^{P} = 0^{-1}$ and 1 states it does not seem possible to reduce the number of terms a priori, for example, by abstracting rules from the nearness of mesons to eigenstates of the SU(6) \rightarrow SU(4) \times SU(2) $_{S_2}$ chain. Note that for both the linear and quadratic formulas, the two terms with largest reduced coefficients are $C_2^{(6)}$ and $C_2^{(3)}$ which are the operators responsible for $\eta - \eta$ and $\phi - \omega$ mixing of the associated eigenstates of the SU(6) \rightarrow SU(4)_N × SU(2)_S chain.

The ideal mixing angle is

$$\theta_{\rm SU(4)_N} = \tan^{-1} (1/\sqrt{2}) = 35^{\circ} 16'$$

to be compared with the empirical mixing angles $\theta_{\rm V} = 37^{\circ} 27^{\circ}$ (linear), $39^{\circ} 59^{\circ}$ (quadratic) and $\theta_{\rm P} = -24^{\circ}$ (linear), $-10^{\circ} 33^{\circ}$ (quadratic) as determined from $\sin^2 \theta_{\rm V} = \left[\phi - \frac{1}{3} (4 \,\mathrm{K}^* - \rho)\right] / \left[\phi - \omega\right]$, etc.

-26-

FOOTNOTES AND REFERENCES

- 1. O. W. Greenberg, U. of Maryland Report 680 (1967) (unpublished).
- O. W. Greenberg and M. Resnikoff, Phys. Rev. <u>163</u>, 1844 (1968);<u>172</u>, 1850 (E) (1968); D. R. Divgi and O. W. Greenberg, Phys. Rev. <u>175</u>, 2024 (1968); D. R. Divgi, U. of Maryland Report 836 (1968) (unpublished).
- R. H. Dalitz, U. of Oxford Report presented at Triangle Meeting on Low Energy Hadron Physics, November 1973, at Smolenice, Czechoslovakia;
 R. Horgan and R. H. Dalitz, Nucl. Phys. <u>B66</u>, 135 (1973); <u>B71</u>, 546 (E) (1974); R. Horgan, Nucl. Phys. <u>B71</u>, 514 (1974).
- 4. F. Gürsey and L. A. Radicati, Phys. Rev. Letters <u>13</u>, 173 (1964);
 B. Sakita, Phys. Rev. <u>136</u>, B1756 (1964).
- See, for example, A. Pais, Rev. Mod. Phys. <u>38</u>, 215 (1966), and
 F. J. Dyson, <u>Symmetry Groups in Nuclear and Particle Physics</u> (W. A. Benjamin, Inc., Advanced Book Program, Reading, Mass., 1966).
- 6. Mass relations based on approximate SU(8) symmetry should be no better than those based on SU(6) because of the larger mass splittings. With the linear Gürsey-Radicati formula, there is a limit to the accuracy with which all the masses in the 56 can be fit; it is to get within 11 MeV of all of them.
- J. J. Aubert <u>et al.</u>, Phys. Rev. Letters <u>33</u>, 1404 (1974); J.-E. Augustin <u>et al.</u>, Phys. Rev. Letters <u>33</u>, 1406 (1974); C. Bacci <u>et al.</u>, Phys. Rev. Letters <u>33</u>, 1408 (1974); G. S. Abrams <u>et al.</u>, Phys. Rev. Letters <u>33</u>, 1453 (1974).
- B. J. Bjorken and S. L. Glashow, Phys. Letters <u>11</u>, 255 (1964); S. L.
 Glashow, J. Iliopoulos, and L. Maiani, Phys. Rev. D2, 1285 (1970).

- C. A. Nelson, "Excitation of Charm, then Color, in ee → Hadrons,"
 State University of New York at Binghamton Technical Report No. 3/34/75.
- 10. J. C. Pati and A. Salam, U. of Maryland Technical Report No. 75-056.
- M. Machacek, Y. Tomozawa, and S. K. Yun, U. of Michigan Report UMHE 75-10.
- 12. R. M. Barnett, Phys. Rev. Letters <u>34</u>, 41 (1975).
- 13. M. Suzuki, Berkeley Report (1975).
- 14. H. Harari, Phys. Letters 57B, 265 (1975); SLAC-PUB-1589 (1975).
- Many authors have carried out such broken SU(4) analyses; see, for example, M. K. Gaillard, B. W. Lee, and J. L. Rosner, Rev. of Mod. Physics <u>47</u>, 277 (1975) and S. Borchardt, V. S. Mathur, and S. Okubo, Phys. Rev. Letters <u>34</u>, 38 (1975); <u>34</u>, 236 (1975).
- M. A. B. Bég and V. Singh, Phys. Rev. Letters <u>13</u>, 418 (1964); <u>13</u>, 509 (1964).
- O. W. Greenberg, Proceedings of Orbis Scientiae II, Coral Gables, Florida (1975), U. of Maryland Technical Report No. 75-064.
- P. Federman, H. R. Rubenstein and I. Talmi, Phys. Letters <u>22</u>, 208 (1966). See also A. W. Hendry, Nuovo Cimento <u>48</u>, 780 (1967).
- 19. Note that for a particular spin group, say SU(2)_S, the familiar spin generators in the spherical basis are given by $S_{\pm}^{c} = (\sigma_{\pm})_{r}^{s} I_{cs}^{cr}$, and $S_{3}^{c} = \frac{1}{2} (\sigma_{3})_{r}^{s} I_{cs}^{cr}$.
- 20. This method of derivation was first used in O. W. Greenberg andC. A. Nelson, Phys. Rev. 179, 1354 (1969).
- 21. L. Licht, U. S. Naval Ordnance Laboratory Report (1968) (unpublished).

- J. Schwinger, Phys. Rev. <u>135</u>, B816 (1964); H. Harari and M. A. Rashid, Phys. Rev. <u>143</u>, 1354 (1966); and R. H. Dalitz, Report presented at the Topical Conference on Meson Spectroscopy, Philadelphia (May 1968).
- 23. Particle Data Group, Physics Letters 50B, No. 1 (1974).