ERRATA TO

A QUARK-BINDING STRING

S.-H.H. Tye

p. 7 equation following Eq. (2.17)

$$\sqrt{-g}\mathscr{P}^{\alpha}_{\mu} = \frac{\delta\mathscr{L}}{\delta\tau^{\mu}_{\alpha}} = \dots$$

p. 8, line 1:

 $\sqrt{-g} \mathscr{P}^{\alpha}_{\mu j} (u^{\beta}) = \dots$

p. 10, 3rd line from bottom:

$$\mathscr{P}^{\beta}_{\mu} = - \mathbf{T}^{\beta\alpha} \boldsymbol{\tau}_{\alpha\mu}$$

p. 51, Ref. 9. See, e.g., L. P. Eisenhart, Riemannian Geometry (Princeton....
p. 55, Fig. 2. ...hypercharges. Quantum mechanically, quark wave functions are spread out.

December 1975

SLAC - PUB-1673 November 1975 (T)

A QUARK-BINDING STRING *

S. - H. H. Tye Stanford Linear Accelerator Center, Stanford University, Stanford, California 94305

ABSTRACT

A relativistic invariant extended model of hadrons is constructed. The finite region where quarks can move freely is a line in a three dimensional space. Physically, the construction resembles that of the MIT bag and the SLAC bubble. In the absence of quarks, the model reduces to the relativistic massless string model. We then solve explicitly the closed string version where massless quarks move only in the clockwise direction. The resulting quantum system does not satisfy Lorentz algebra. However, this negative result indicates that for an open (or closed) string, the quantum system is consistent if there are 22 quark fields. A quark-confining string is also constructed and discussed. This is obtained as an improved version of the quark-binding string. It incoporates many physically desirable features.

(Submitted for publication)

^{*} Work supported by the U.S. Energy Research and Development Administration.

I. Introduction

Numerous phenomenological successes of the dual resonance amplitudes have stimulated the search for the physical picture underlying the model. Nambu and others (1) observed that the spectrum of the Veneziano model (2) is identical to that of a vibrating string. Subsequently the free string formalism has been studied in detail, starting from an action written in terms of the area swept out by the string (3). More recently, the string picture has also emerged in many different investigations (4), and its relevance to hadron physics is generally accepted. Of course, the string picture is, at best, an approximate description of hadron physics. Its significance lies at least partly in the fact that it can be explicitly solved. This allows one to calculate its dynamical and static properties for comparison with experiments, starting from the exact quantum relativistic solution of the string; scattering (5) and other interactions can be introduced as perturbations around the free string solution, maintaining quantum mechanics and relativity at every order of perturbation. This approach towards hadron physics is attractive since it is the only approach we have rest experience, namely, in quantum electrodynamics; there the free photon and electron are first solved completely, and the electromagnetic interaction is then introduced as a perturbation around the exact solutions. Whether this gives an approximate description of hadrons we can tell only after the program has been carried out completely, beyond the free string version.

The present string formalism has a few unsatisfactory features. Among them are (1) the quantized string exists only in 26 dimensional Minkowski space ; (2) lack of incorporation of quarks (i.e. Dirac particles) and (maybe) their confinement. Numerous attempts have been made to overcome on e or both of these shortcomings. This work maybe considered as the first step of yet another attempt to improve the present string formalism. The formulation is motivated purely by the physical

1

picture.

We construct the system of a string upon which physical quarks are confined. (By "physical" we mean the quarks are spinors in the four dimensional Minkowski space but behave as scalars under a conformal transformation of the parameters describing the string.) We then proceed to solve the closed string along which massless quarks move around it in either the clockwise or the anti-clockwise direction. The mathematical formulation of this is summarized in Table I. The resulting quantized system does not satisfy Lorent_z covariance, for the same reason as the Nambu string. However, the results do indicate that the complete system (with quarks moving both clockwise and anticlockwise around a closed string, <u>or</u> for an open string) will have Lorentz covariance if the number of quarks is 22. This speculation will be formulated as a conjecture.

Our string formalism is modeled after that of the M.I.T. bag $^{(6)}$ and the SLAC bubble $^{(7)}$, and the quantization procedure is modeled after that of the free string $^{(3)}$ and of the SLAC bubble in three dimensional Minkowski space $^{(8)}$ (which is equivalent to a quark-binding closed string).

From the investigation of the quark-binding string we find it has a few undesirable features: its ground state is an empty string state; it also has exotic states, i. e., states with any number of quark excitation modes. To remove these bad features, a quark-confining string is constructed. Here the quarks come in color triplets interacting with color SU(3) gluon fields. Because we demand there be no local color electric field pointing out of the string, only color singlet states exist. There are no gluon degrees of freedom; only color electric fields are left. Also there are no pure string modes, so that the ground state of this string is a meson; the quarks and the antiquarksmay have masses, in which case,

 $\mathbf{2}$

they sit at the ends of the string, one at each end, linked by the appropriate electric flux line, which also defines the string.

Section II contains the formalism. In Section III we solve the closed string with only clockwise moving quarks. In Section IV we examine the quantized system, and discover that Lorentz covariance is not satisfied. Section V contains discussions and speculations. Here we speculate what the spectrum for the quarkbinding string would look like. (By "quark-binding" we mean the absence of massless free point-like quarks.) However, a string with any number of quark and antiquark modes can be present. In this section we also discuss how a quarkconfining string can be written down in the action formalism. (By "quarkconfining," we mean only color singlet states exist.) Some of the technical details and supplementary discussions are relegated to the four appendices.

-3-

II. Formalism

The geometry of a relativitic string is that of a two dimensional subspace (V_2) imbedded in a four dimensional flat Minkowski space (S_4) . To describe a quark-binding string we must develop a convenient set of variables in the language of differential geometry (9).

Let X_{μ} and u^{α} be the coordinates in the Minkowski space S_4 (with signature (l, -l, -l, -l)) and the curved space V_2 (with signature (l, -l)) respectively. The shape and position of the string is described by (see figure l.)

$$X_{\mu} = X_{\mu}(u^{\alpha})$$
 $\mu = 0, 1, 2, 3$ (2.1)
 $\alpha = 0, 1$

where the choice of the parameters $\mathcal{U}^{\mathcal{A}}$ is arbitrary. We assume that X_{μ} is differentiable. (Our notation will be such that the indices $\boldsymbol{\triangleleft}, \boldsymbol{\beta}, \boldsymbol{\delta}, \boldsymbol{\delta}$ refer to the v_2 space and $\mu, \nu, \boldsymbol{\lambda}, \boldsymbol{\beta}$ refer to the s_4 space). The quantity

$$\mathcal{T}^{\mu}_{\alpha} \equiv \frac{\partial \chi^{\mu}}{\partial u^{\alpha}} \equiv \chi^{\mu}_{\ \alpha} \equiv \partial_{\alpha} \chi^{\mu} \qquad (2.2)$$

is tangential to the string and is a vector in both $\rm V_2$ and $\rm S_4$. The induced metric for the $\rm V_2$ space is given by

$$g_{\alpha\beta} = \tau^{\mu}_{\alpha} \tau_{\beta\mu} = \tau_{\alpha} \cdot \tau_{\beta}$$
(2.3)

which is a covariant symmetric tensor with signature (1, -1).

At each point on the string, a plane perpendicular to the tangent vectors can be defined:

$$U_{\mu\nu} = \frac{\varepsilon_{\mu\nu\rho\lambda} T_{\chi}^{\rho} T_{\beta}^{\rho}}{\sqrt{-g'}} \qquad \times \pm \beta \qquad (2.4)$$

where $\mathcal{E}_{\mu\nu\rho\lambda}$ is totally ant isymmetric, with $\mathcal{E}_{0123} = 1$, and $g = \det(g_{\alpha\beta}) = g_{00}g_{11} - g_{01}^2$. An alternative way to define this plane is by introducing two space-like outward

normals $N_{\mu}(u^{\alpha})$ and $M_{\mu}(u^{\alpha})$:

$$m \cdot T_{\alpha} = n \cdot T_{\alpha} = 0$$

They are normalized to unity,

$$m_{\mu}m^{\mu} = m^2 = n^2 = -1$$
 (2.6)

(2.5)

and, for convenience, taken to be perpendicular to each other,

$$\gamma \cdot m = 0 \tag{2.7}$$

Now $U\mu\nu$ can be defined including its orientation in terms of n_{μ} , m_{μ}

$$U_{\mu\nu} = N_{\mu}M_{\nu} - M_{\mu}N_{\nu}$$
(2.8)

The Minkowski metric at each point on the string can now be expressed in terms of the curvilinear co-ordinates provided by the set of linearly independent vectors:

$$\eta_{\mu\nu} = \tau^{\alpha}_{\mu}\tau_{\alpha\nu} - \eta_{\mu}\eta_{\nu} - m_{\mu}m_{\nu} \qquad (2.9)$$

where $\tau^{\alpha}_{\mu} = g^{\alpha\beta} \tau_{\beta\mu}$; $g^{\alpha\beta}$ is the inverse of $g_{\alpha\beta}$, and is a contravariant symmetric tensor. The tengential components of any vector ∇_{μ} can be projected out, $\nabla_{\mu\mu} = (\nabla \cdot \tau^{\alpha}) \tau_{\alpha\mu} = \nabla^{\alpha} \tau_{\alpha\mu}$.

Let us introduce some more quantities that are needed later. The derivatives of the normals

$$n_{\mu 1 \chi} = h_{\mu \beta} T_{\mu}^{\beta} + V_{\chi} m_{\mu}$$
(2.10a)

$$m_{\mu i \alpha} = l_{\alpha \beta} T^{\beta}_{\mu} - \nu_{\alpha} n_{\mu}$$
(2.10b)

introduce the torsion $V_{\alpha} = N^{\mu}(m_{\mu 1 \alpha}) = -m^{\mu}(m_{\mu 1 \alpha})$ and the symmetric curvature tensors

$$h\alpha\beta = n_{1\alpha} \cdot T_{\beta} = -n \cdot T_{\alpha 1\beta} = -n \cdot T_{\beta 1\alpha} = -n \cdot T_{\beta 1\alpha}$$

$$(2.11a)$$

$$L\alpha\beta = -m \cdot T_{\alpha 1\beta}$$

$$(2.11b)$$

$$\tau_{\alpha \parallel \beta}^{\mu} = \tau_{\alpha \perp \beta}^{\mu} - \left\{ \chi_{\beta}^{\nu} \right\} \tau_{\gamma}^{\mu} = h_{\alpha \beta} n^{\mu} + l_{\alpha \beta} m^{\mu}$$
(2.12)

The "Christoffel symbol of the second kind is a function of the metric

$$\begin{cases} \chi \\ \alpha \beta \\ = \pm g^{\chi} \left[g_{\alpha 81\beta} + g_{8\beta1\alpha} - g_{\alpha\beta18} \right] \\ = \tau^{\chi} \tau_{\alpha1\beta} \end{cases}$$
(2.13)

Let us now introduce the quark fields. Since they are confined on the string, we can write

$$\Psi_{j} = \Psi_{j} \left(u^{\alpha} \right)$$
(2.14)

j = l, 2, ..., N, where N is the number of quark species. (Throughout, repeated indices are summed unless specified otherwise; for individual quark field, the index is suppressed.) The Dirac matrices, which obey

$$\{\chi^{\mu},\chi^{\nu}\} = \chi^{\mu}\chi^{\nu} + \chi^{\nu}\chi^{\mu} = 2\eta^{\mu},$$

are chosen to be

$$Y_{i} = \begin{bmatrix} 0 & \sigma_{i} \\ -\sigma_{i} & 0 \end{bmatrix}, \quad Y_{0} = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} \quad Y_{5} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$

where τ_{λ} are the Pauli matrices. $A = a_{\mu} \gamma^{\mu}$, for any vector a_{μ} Now we can write down the action of free quarks confined on a string ⁽¹⁰⁾ ($\hbar = c = 1$)

$$S = \int d^2 u^{\chi} \sqrt{-g} \left\{ \frac{i}{2} \overline{\psi_j} \right\}_{ii} \psi_j - \zeta \right\} = \int d^2 u^{\chi} \mathcal{L}$$
(2.16)

$$j_{\mu}n^{\mu} = (\overline{\psi}\gamma_{\mu}\psi)n^{\mu} = j_{\mu}m^{\mu} = 0 \qquad (2.17)$$

$$p_{\mu}^{\chi}n^{\mu} = p_{\mu}^{\chi}m^{\mu} = 0$$

where
$$\mathcal{P}_{\mu}^{\alpha} = \frac{St}{ST_{\alpha}^{\mu}} = \frac{S\sqrt{-g}}{ST_{\alpha}^{\mu}} \frac{1}{\sqrt{-g}} t + \sqrt{-g} \left[\frac{i}{2} \overline{\Psi}_{j} \frac{ST_{\alpha}^{j}}{ST_{\alpha}^{\mu}} \frac{J_{\alpha}^{j}}{\sqrt{j}} \right]$$

$$= T_{\mu}^{\alpha} t - \frac{i}{2} \sqrt{-g} \overline{\Psi}_{j} \left[g^{\alpha\beta} (A n_{\mu} + i\pi m_{\mu}) + T_{\mu}^{\beta} t^{\alpha} \right] \frac{J_{\alpha}^{\beta}}{\sqrt{j}} \Psi_{j}$$

It is clear that the string part of $\mathcal{P}^{\mathscr{A}}_{\mu}$ is already tangential. Because the quarks are all distinguishable from each other (different quantum numbers) we demand that the normal components of $\mathcal{P}^{\mathscr{A}}_{\mu}$ of each quark must vanish:

$$P_{\mu j}^{\chi}(u^{\beta}) = \frac{i}{2} \int -g g^{\alpha \beta} \overline{\Psi}_{j}(Hn_{\mu} + Hm_{\mu}) \partial_{\beta} \Psi_{j} = 0$$

j not summed

$$\overline{\psi}_{H}\overline{\partial}_{x}\psi = \overline{\psi}_{H}\overline{\partial}_{x}\psi = 0$$

or

(2.18)

(2.19)

(egn 2.18);

This simply means the momentum of each quark is tangential and may be considered as the normal components of the quark equation of motion.

Although the quarks are confined along the string, they are physical spinors in the S_4 space; hence continuity of each quark field across the string must also be satisfied:

$$\overline{\psi}\psi = 0$$

Eqn (2.17), (2.18) and (2.19) can be considered as constraints on the spinor fields. Together with the action (2.16), the system is completely determined except for boundary conditions. The latter depends on whether the string is open or closed. For an open string, the second momentum "current" density and the charge current densities must vanish at the ends of the string. For a closed string, all variables must be periodic (anti-periodic for fermions) in the parameter that measures along the length of the string.

The physical picture of the model is clear. The string moves and vibrates while the quarks are moving left and/or right along the string, The distribution and the motion of the quarks affect the shape of the string, which in turn defines the path of the quark motion. Since there is no direct interaction among the quarks, the left moving and the right moving quarks pass each other freely. However, the complete solution of the quark fields is not a superposition of the solutions of the left and right moving quark fields, because of their non-linear influence on each other via the geometry of the string.

8

In this work we shall study the string when there is only left moving or $\operatorname{culy}^{\mathsf{ruly}}$ right moving quarks. We discover that this system can be solved completely. The complete solution of the subsystem allows one to speculate (a rather natural guess, we believe) at the complete solution of the quark-binding string we have just formulated.

When there are only left moving quarks on the string, the boundary condition of the open string at the ends cannot be satisfied, since quarks will flow in and out there. Hence we can only study a closed string; the left moving quarks become quarks moving clockwise around the string. The first thing we have to do is to project out the clockwise moving quarks.

We note that from eqn (2.7),

AHA = - HAA

which provides a signature for the plane $\lim_{\mu \nu} \nu$. This can be used to specify the direction. The projection operators ω_{\pm} must satisfy

$$\psi = (\omega_{+} + \omega_{-})\psi = \psi_{+} + \psi_{-}$$

and

$$(\omega_{\pm})^2 = \omega_{\pm}$$

Let us define

$$\omega_{\pm} = \frac{1 \pm i \sqrt{5} + 4}{2}$$
(2.20a)

and study only the case when one projection of the spinor field is present, say

$$\lambda V_5 + + + \Psi = \Psi_+ \qquad (2.20b)$$

That this projects out the left moving quarks will become clear later. From now on we shall drop the +'' index whenever it is obvious. The quark species index is implied in eqn (2.20b) also.

9

The inclusion of λ_{15}^{\prime} in the choice of ω_{+} in eqn (2.20) automatically implies eqn (2.19) and (2.17)

$$\overline{\Psi}\Psi = \overline{\Psi} (\lambda \delta S H H \Psi)$$

$$= (\lambda \overline{\Psi} S H H) \Psi$$

$$= -\overline{\Psi} \Psi = 0$$

and

ļ

$$j \cdot n = (\overline{\psi} \delta_{\mu} \psi) n^{\mu} = \psi + \psi$$

$$= \overline{\psi} + (\overline{\chi} \delta_{5} + \overline{\psi} + \psi) \qquad (2.21a)$$

$$= (\overline{\chi} \overline{\psi} \delta_{5} + \overline{\psi} + \overline{\psi} + \overline{\psi} + \overline{\psi} + \overline{\psi} = 0$$

(2.2lb)

and similarly for

j.m = 0

It is useful to note that the normal components of the axial vector current is also zero

$$j_{sp}n^{m} = \overline{\psi}H\gamma_{s}\psi = i f_{\mu}m^{m} = 0$$

Hence, using eqn (2.9)

$$j_{5\mu} = (\overline{\psi} \, \overline{\psi}^{*} \gamma_{5} \, \psi) \, \tau_{x\mu} = J_{5}^{x} \, \tau_{x\mu}$$

That is, both the vector and axial-vector currents are tangential.

Next we demand the energy momentum "current" density of the string be tangential also.

$$P^{\beta}_{\mu} = -\int -g T^{\beta \chi} T_{\chi \mu}$$

where

$$\tau^{\alpha\beta} = \mathcal{C}g^{\alpha\beta} + \mathcal{O}^{\alpha\beta} + \frac{\dot{\lambda}}{2}\overline{\psi}_{j$$

This follows from eqn (2.18) and the Dirac equation (2.25) written down below.

which can be rewritten as, using eqn (2.10), (2.11) and (2.21)

$$\overline{\Psi}H\overline{\partial}_{x}\Psi = -\lambda \partial_{x}[\overline{\Psi}HY_{5}\Psi] - \lambda\overline{\Psi}Y_{5}(\partial_{x}H)\Psi$$

$$= \lambda V_{x}\overline{\Psi}Y_{5}H\Psi - \lambda L_{x}\beta \overline{\Psi}Y_{5}\overline{\Psi}\Psi$$

$$= \lambda L_{x}\beta J_{5}^{\beta} = C \qquad (2.23)$$

$$\overline{\Psi}H\overline{\partial}_{x}\Psi = \lambda h_{x}\beta J_{5}^{\beta} = 0$$

Hence we have a closed system when the axial vector current is identically zero.

It is now striaghtforward to derive the equations of motion of the system. The Euler-Lagrange equation for the string co-ordinates is

$$\frac{\partial_{\beta} S \mathcal{L}}{S \partial_{\beta} X^{\mu}} = -(J - g T^{\beta \alpha} T_{\alpha \mu})_{1\beta}$$
$$= J - g (-T^{\beta \alpha} T_{\alpha \mu})_{1\beta} = 0 \qquad (2.24)$$

The Euler-Lagrange equation of the quark field becomes

$$\frac{st}{s\overline{\psi}} = \partial_{x}\frac{st}{s\overline{\psi}}$$

$$L.H.S = i \int -g t^{\alpha} \partial_{\alpha} \psi$$

and

I

$$R.H.S = -\partial_{x}\left[i\sqrt{-g}t^{x}\psi\right] = -\frac{i}{2}\sqrt{-g}\left[t^{x}_{ix}\psi + t^{x}\right]_{x}\psi$$

Introducing the mean curvatures

$$2k_1 = h^{2} \times 2k_2 = \lambda^{2} \times 2k_2$$

the Dirac equation becomes, using eqn (2.12),

$$i t' \partial_x \Psi = -i (\beta_1 A + \beta_2 M) \Psi$$

It follows that $\Theta^{\alpha\beta} = \frac{1}{2}\overline{\psi}_{j} t^{\alpha} \overline{j}^{\beta} \psi_{j}$ since $\overline{\psi} t^{\alpha} \overline{j}_{\alpha} \psi = 0$. Projecting out the normal and tangential components of eqn (2.24) and using eqn (2.12), we get

$$T^{\alpha\beta}_{\mu\alpha} = 0$$

$$= G g^{\alpha\beta} + \frac{\lambda}{2} \overline{\Psi}_{j} \overline{\Psi}^{\alpha} \overline{\partial}^{\beta} \Psi_{j}$$

$$(2.26)$$

and

$$h_{\alpha\beta}T^{\alpha\beta} = l_{\alpha\beta}T^{\alpha\beta} = 0$$
(2.27)

which is just eqn (ii) and (iii) in Table I.

Note that these equations of motion can also be derived using the constraints (2.17) - (2.19). Hence they are also the equations of motion for the complete (i.e. with both left and right moving quarks) system.

It is straightforward to show that eqn (2.26) is redundant: one can derive $T^{\alpha}f_{\mu\alpha} = g^{\beta\gamma}(T^{\alpha}\gamma_{\mu\alpha}) = 0$

using eqn. (2.21), (2.23) and (2.25). This means that not all the dynamical variables, namely the quark fields and the string co-ordinates, are independent. This is a reflection of "gauge invariance" under reparametrization.

To summarize, the system of a closed string, along which massless quarks obeying the projection (2.20) move freely, is completely specified by eqn (2.25) (2.27) together with the periodicity condition on all dynamical variables and the vanishing of the axial vector current

$$js\mu = \overline{\Psi} \gamma_{\mu} \gamma_{s} \Psi = 0$$

(2.28)

Classically, any system involving fermions is plagued with negative energies. In particular the fermion energy density is not positive definite. We shall ignore this difficulty by assuming that the fermion energy density is positive definite, which is the case after proper fermion-antifermion identification and fermi statistics are taken into account. This is discussed in Section IV.

In the absence of quark fields, the quark-binding string action reduces to the Nambu, or the empty string action. This is clear from the original action (2.16) and also from the final spectrum, as we shall see later.

7

III. Solution

The scheme of solving the set of coupled non-linear equations of motion (2.25) and (2.27) is as follows: first we derive some properties of the system, which are essential for the solution. We then solve the geometry of the closed string in terms of the fermion distribution by adopting a particular coordinate choice. Finally we go back to solve the Dirac equations in terms of the geometry. Supple-

mentary discussions and details are relegated to the three appendices A, B and C.

A. Properties of the System

Here we derive some of the properties of the system. In particular, we show that there exists a parameter choice such that the string coordinates obey the free string equation of motion. For convenience we write

$$\mathcal{G} \not \sim \mathcal{G} = \begin{pmatrix} A & V \\ V & B \end{pmatrix}$$
(3.1)

so that $g = AB - V^2 < 0$, since the signature of the metric is (1, -1). Let us also introduce, for any function $W(u^4)$, the following notation

$$\tilde{W} = \frac{\partial W}{\partial u^{\circ}}$$
 $W' = \frac{\partial W}{\partial u^{\circ}}$

Using Eq. (2.4) and (2.8), the projection (2.2) can be expressed in terms of $\tau_{\alpha\mu}$

$$i \sqrt{5} + H = \frac{\sqrt{-7} \sqrt{7}}{\sqrt{-9}}$$
(3.2)

Next we observe that the current is a null vector; in fact (see Appendix A)

$$f_{i} \Psi_{R} = (\Psi_{i} \vee_{\mu} \Psi_{i}) \vee_{\mu} \Psi_{R} = 0$$
i not summed (3.3)

for both i = k and $i \neq k$. Multiplying $\overline{\Psi}_k$ on the left, we get

$$J_{i\mu}J_{R}^{m} = J_{i\alpha}J_{R}^{\alpha} = 0 \qquad (3.4)$$

Also

l

$$J_{x} \theta^{x\beta} = 0$$

where $\theta^{\star\beta}$ is the fermion part of the energy momentum tensor (2.22), or

$$J_{\alpha} T^{\alpha \beta} = \mathcal{C} J^{\beta}$$
(3.5)

Using the Dirac equation (2.25), we also obtain

$$\overline{\mathcal{J}}_{\parallel \times}^{\times} = 0 \tag{3.6}$$

To study the string geometry, we can, for the moment, concentrate only on Eq. (2.26), (2.27), (3.4), (3.5) and (3.6). The quarks appear only in the form of the current and their energy momentum density $e^{i\frac{1}{2}\beta}$.

The original action is independent of the choice of the parameters u_{α} . Since $T^{\alpha\beta}$ is symmetric and has signature (1, -1) (see Appendix B), we choose $T^{\alpha\beta}$ to be off-diagonal:

$$T^{\alpha\beta} = CT^{\alpha\beta} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

That this is possible should be clear: given $T^{\alpha\beta}$ (u_0, u_1) , a local coordinate system can be found: $(u^0, u^1) \rightarrow (\tau, \tau)$ such that $T^{\alpha\beta}$ (τ, τ) is off-diagonal. Using Eq. (3.1) and (3.6), $T^{\alpha\beta}$ becomes

$$T^{\alpha\beta} = \frac{\mathcal{C}}{\mathcal{V}} \begin{pmatrix} 0 & 1\\ 1 & 0 \end{pmatrix}$$
(3.7)

As we shall see, $V = \mathbf{x} \cdot \mathbf{x}'$ must be nonzero. Using Eq. (3.5)

$$J_{\beta}J^{\beta} = J_{x}T^{x\beta}J_{\beta} = 2\underline{C}J_{\alpha}J_{1} = 0$$

We find that either J_0 or J_1 must be zero. We choose (this defines our sign convention)

$$J_1 = 0$$

Then from Eq. (3.4)

$$J_{x}J^{x} \equiv q^{x\beta} J_{x}J_{\beta} = 0$$

$$g_{\alpha\beta} = \begin{pmatrix} A & V \\ V & O \end{pmatrix}$$

and

$$g^{\times\beta} = \frac{1}{V} \begin{pmatrix} 0 & 1 \\ 1 & -A_{/V} \end{pmatrix}$$

so that $-g = V^2 > 0$. This gives

$$\Theta_{\alpha\beta} = \mathcal{C}\left(\begin{array}{c}A & 0\\0 & 0\end{array}\right) = \frac{\lambda}{2}\overline{\Psi}_{j}\overline{\Psi}_{\alpha}\overline{\partial}_{\beta}\Psi_{j}$$

It should be clear that we cannot choose $J_{i1} = 0$ for some quarks while $J_{k0} = 0$ for some other quarks.

The relation between the coordinates and the quark fields is manifest

$$CA = C\dot{x}^{2} = \frac{\lambda}{2} \dot{\Psi}_{j} \not\equiv 0 \dot{\delta} \dot{\Psi}_{j}$$
(3.8)

The Christoffel symbols of the second kind now simplify to

Using these and Eq. (2.26)

I

$$- \frac{1}{1} \frac{1}{1} = 0$$

we obtain A' = 0, so that the Christoffel symbols are further simplified:

Eq. (2.27) immediately gives

$$h_{01} = l_{01} = 0$$

Eq. (2.12) becomes, when written explicitly

$$\ddot{x}_{\mu} = \dot{x}_{\mu}(\ln v)' + hoo M_{\mu} + loo M_{\mu} + (\frac{A}{2v} - \frac{Av}{v^2}) x'_{\mu}$$
(3.10)

$$X''_{\mu} = \dot{X}_{\mu} (\ln V)' + h_{\mu} n_{\mu} + l_{\mu} m_{\mu}$$
 (3.11)

$$\dot{X}'_{\mu} = 0 \tag{3.12}$$

The last of which allows us to write

$$\chi_{\mu}(\tau,\sigma) = S_{\mu}(\sigma) + G_{\mu}(\tau)$$
(3.13)

B. The Geometry

The imbedding of the string in Minkowski space requires the introduction c_{5}^{f} the metric and the curvature tensors. The consistency of the geometry is guaranteed by the integrability conditions, namely, the Gauss-Codazzi equations⁽⁹⁾.

$$R_{x\beta xS} = (h_{xx} h_{\beta S} - h_{x\beta} h_{xS}) + (l_{xx} l_{\beta S} - l_{x\beta} l_{xS})$$

$$h_{x\beta ||x} - h_{xx ||\beta} = v_{x} l_{x\beta} - v_{\beta} l_{xX}$$

$$l_{x\beta ||x} - l_{xx ||\beta} = -v_{x} h_{x\beta} + v_{\beta} h_{xX}$$

where the Riemann curvature tensor is given by

$$R_{\beta\gamma\delta}^{x} = \{x_{\beta}\}_{1\delta} - \{x_{\beta}\}_{1\delta}^{x} + \{y_{\delta}\}_{1\delta}^{y} + \{$$

When the curvature tensors are all constructed from the metric tensor, the Gauss-Codazzi equations are automatically satisifed. However, since we are going to solve the string by making a coordinate choice which apparently breaks Lorentz covariance, it is reassuring to check that the integrability constraints are satisfied.

From its symmetry properties, it is clear that the Riemann tensor has only one independent component. Using Eq. (3.9) the Gauss-Codazzi equations become

$$h_{00} + \frac{1}{V} \left(\frac{\dot{A}}{2} - \frac{AV}{V} \right) h_{11} = V \dot{\lambda}_{00}$$
(3.14a)

$$\hat{\chi}_{00}' + \frac{1}{V} \left(\frac{\dot{A}}{2} - \frac{AV}{V} \right) \hat{\chi}_{11} = -\nu' \hat{h}_{00}$$
(3.14b)

$$\ddot{\mathbf{h}}_{\mathbf{n}} = \ddot{\boldsymbol{\nu}} \, \boldsymbol{\chi}_{\mathbf{n}} \tag{3.15a}$$

$$\dot{\nu}h_{\rm H} = -\dot{k}_{\rm H} \tag{3.15b}$$

We observe that the theory has conformal invariance; that is, Eq. (3.10), (3.11), (3.12), (3.14) and (3.15) are form invariant under the conformal transformation

 $\tau \rightarrow e(\tau)$ $\sigma \rightarrow f(\tau)$

Hence we can choose, from eqn (3.13)

$$\chi_{+} = p(\tau + \sigma) = \chi_{0} + \chi_{3}$$
(3.16)

where p is a constant. Then the constraint among the string coordinates

$$g_{11} = \mathbf{x}'^2 = \mathbf{c}$$

can be written as

ņ

$$\chi'_{-} = \frac{1}{P} \left(\chi'_{1}^{2} + \chi'_{2}^{2} \right)$$
(3.17)

and Eq. (3.8) becomes

$$\dot{\chi}_{-} = \frac{i}{p} \left(\dot{\chi}_{1}^{2} + \dot{\chi}_{2}^{2} + A \right)$$
 (3.18)

Hence x_1 , x_2 and A can be taken to be the independent dynamical variables, with

$$2V = (\tilde{x}_1 - \tilde{x}_2)^2 + (x_1' - x_2')^2 + A$$

Similarly $h_{\alpha\beta}$ and $\ell_{\alpha\beta}$ can be expressed in terms of x_1 , x_2 and A. The Gauss-Codazzi equations can then be explicitly verified (see Appendix C).

C. The Dirac Equation

Equation (3.2) becomes

$$\tau_0 \tau_1 = V(1 - i \sqrt{s} + t + t)$$

 \mathbf{or}

$$\vec{t}_1 \vec{t}_3 = V(1 + i \sqrt{3} + H)$$

Therefore, with $A = \nabla \overline{\nabla} \overline{\nabla} \sigma$

$$\overline{\psi}, \dot{\Psi} = \frac{1}{A} \overline{\psi}_{c}(\overline{\psi}, \overline{\psi},) \Psi = 0$$
(3.19)

Differentiating with respect to τ , we obtain, using Eq. (3.13)

$$\partial_{\sigma}(\vec{\tau},\psi) = \vec{\tau}_{i}\psi = 0$$

Differentiating with respect to \mathcal{T} , we obtain, using Eq. (3.12)

$$\dot{x}\psi' = -(\dot{h}_{11} + + \dot{z} + \dot{z}_{11} + \dot{z})\psi$$

Note that

$$k_{1} = \frac{1}{2} k_{\alpha}^{\alpha} = -\frac{A}{2V^{2}} k_{11}$$

$$k_{2} = -\frac{A}{2V^{2}} k_{11}$$

Substituting the above four equations into the Dirac equation (3.25), we obtain

$$g'' t_0 \psi' = -\frac{A}{2V^2} (f_{11} + L_{11} + L_{11}) \psi$$

Multiplying this on the left by \neq_0 and using Eq. (3.12), the Dirac equation becomes

$$\psi' = - \frac{\cancel{t} \cdot \cancel{t}}{2 \vee} \psi$$

Equation (3.19) puts a constraint among the components of Ψ . Using this to eliminate the dependent components, we can write Ψ in terms of the two unknown functions u, v, Γ u γ

$$\psi = \begin{bmatrix} u \\ x''/p \\ -x'e/p \\ v \end{bmatrix}$$

where $\chi'_{\gamma} = \chi'_{\ell}^{*} = \chi'_{1} + \Lambda \chi'_{2}$

The Dirac equation can now be written in terms of u, v

$$u' = - \frac{(x'_{1} - \dot{x}_{1}) x''_{1}}{2v} u$$

$$v' = - \frac{(x'_{1} - \dot{x}_{1}) x''_{1}}{2v} v$$

Solving this, we obtain

$$\Psi = \begin{bmatrix} 1 \\ x'_{\tau/p} \\ 0 \\ c \end{bmatrix} \perp F(\tau) + \begin{bmatrix} 0 \\ 0 \\ -x'_{\ell/p} \\ 1 \end{bmatrix} R G(\tau)$$
(3.20a)

7

where

$$L = R^{+} = c_{xp} \left[-\frac{1}{2} \int \frac{(x_{x}' - \dot{x}_{y}) x_{y}''}{V} d\sigma_{i} \right]$$
(3.20b)

and $F(\tau)$ and $G(\tau)$ are arbitrary functions of τ . This is an implicit solution of ψ since L is a function of A (via V) which in turn is a function of the quark fields (see Eq. (3.8) and (3.18)).

Since the axial vector current is tangential, the only nontrivial component of $j_{5\mu}$ in Eq. (2.28) is, using Eq. (3.19)

$$\overline{\Psi} \neq_{o} Y_{5} \Psi = \frac{2}{P} \left(G^{\dagger} G - F^{\dagger} F \right) = 0$$

This means F and G differ by a phase only; we choose F = G. It is clear that if $\dot{\Psi}$ is a solution of the system, so is

$$\Psi_{c} = C\Psi^{*} = (\lambda \vartheta_{2})\Psi^{*}$$

where C $\widecheck{\textbf{X}}^0$ is the charge conjugation matrix. Explicitly, we have

$$i S_5 M H 4_c = 4_c$$

and

$$\lambda \neq^{\alpha} J_{\alpha} \Psi_{c} = -\lambda (\lambda_{1} + \lambda_{2} + \lambda_{2}) \Psi_{c}$$

Here

$$\begin{aligned} \dot{\Psi}_{c} &= i \tilde{V}_{2} \begin{bmatrix} LF \\ x_{i/p} LF \\ -x_{i/p} RF \\ RF \end{bmatrix}^{*} = \begin{bmatrix} L \\ x_{i'/p} L \\ -x_{i/p} R \\ R \end{bmatrix} F^{*} \end{aligned}$$

This means Ψ should have both quark and antiquark solutions. Using its antiperiodic property (with the period τ_0 taken to be 1) $F(\tau)$ can be expanded into Fourier components. Hence

$$\psi = \mathcal{N} \begin{bmatrix} L \\ \chi'_{i/p} L \\ -\chi'_{i/p} R \end{bmatrix} \xrightarrow{\infty} b_{m} e^{-2\pi i m \tau}$$
(3.21)

where the prime indicates the sum over half-integers with coefficients b_m , \mathbf{N} is the normalization to be determined. A Dirac field has fermion and antifermion solutions each with two spin polarizations. The projection (2.20) is a geometric one, leaving one solution each for the fermion and the antifermion. The confinement on a two dimensional manifold has deprived the Dirac particles of their conventional spin structures. The reader probably would have guessed this from the fact that the energy-momentum tensor $T^{\alpha}f^{\beta}$ is symmetric without the introduction of any "superpotential" terms. We should also expect the absence of the spin part of the angular momentum tensor density $M^{\alpha\mu\nu}$, which is given by

$$M^{\alpha\mu\nu} = \chi^{\mu} P^{\alpha\nu} - \chi^{\nu} P^{\alpha\mu} + \frac{1}{2} \overline{\Psi}_{j} [\overline{\psi}^{\alpha} \sigma^{\mu\nu} + \sigma^{\mu\nu} \overline{\psi}^{\alpha}] \Psi_{j}^{\alpha} (3.22)$$

where

$$\sigma^{\mu\nu} = \frac{1}{2} \begin{bmatrix} \chi^{\mu}, \chi^{\nu} \end{bmatrix}$$

It is straightforward to verify this is indeed ture, that is,

$$\frac{1}{2}\overline{\psi}\left[\overline{\psi}^{\alpha}\sigma^{\mu\nu}+\sigma^{\mu\nu}\overline{\psi}^{\alpha}\right]\psi=0$$

using Eq. (3.20) and (3.21). Note that

$$LR = LL^{T} = V$$

Using this, we see that **C**A is quite simple despite/the complicated form of Ψ ,

$$CA = \frac{\lambda}{2} \overline{\Psi_{j}} \overline{\psi_{j}} \overline{\psi_{j}} \overline{\psi_{j}}$$

= $\frac{2\lambda}{P} \left(F_{j}^{\dagger} \overline{F_{j}} - \overline{F_{j}}^{\dagger} \overline{F_{j}} \right)$ (3.23)

Poincaré covariance should be satisfied, since the original formulation is Lorentz covariant.

IV. The Quantized System

A. Non-covariant Quantization

We choose our units such that C = 1. We shall restore it only at the end. The energy momentum is given by

$$P_{\mu} = \int d\Sigma_{\alpha} \int -g P_{\mu}^{\alpha}$$
(4.1)

where the integral is to be taken at any fixed time t

$$t = \tau + \sigma \tag{4.2}$$

This means the integration is over any space-like submanifold; where the space-like parameter

$$g = \frac{1}{2} (\tau - \sigma) \tag{4.3}$$

actually measures along the string. To see this, we observe that P_{-} is the generator of infinitesimal translations in the x_{+} direction (see Eq. (3.16)); therefore the Hamiltonian should be given by

$$H = P_+ P_- \tag{4.4}$$

To evaluate H, let us expand x^{μ} as follows, which is permitted by Eq. (3.13)

$$\chi^{\mu}(\tau,\sigma) = \frac{1}{2\sqrt{\pi}}q^{\mu} + \sqrt{\pi} \ a_{o}^{\mu}\tau + \sqrt{\pi} \ c_{o}^{\mu}\sigma + \frac{\lambda}{2\sqrt{\pi}}\sum_{n\neq o} \left(\frac{a_{n}}{n}e^{-2\pi i n\tau} + \frac{c_{n}}{n}e^{-2\pi i n\sigma}\right)$$
(4.5)

where the period of σ is also taken to be one. For fixed t, the periodicity condition of x^{μ} is

$$\mathbf{x}^{\mathbf{M}}(\mathbf{t},\mathbf{p}) = \mathbf{x}^{\mathbf{M}}(\mathbf{t},\mathbf{p}+\mathbf{i})$$

This implies $a_o^{\mu} = C_o^{\mu}$. Reality of x^{μ} implies

 $q_{\mu} = q_{\mu}^{\star}$ $a_{n}^{\dagger} = a_{-n}$ $a_{o} = a_{o}^{\star}$ $c_{n}^{\dagger} = c_{-n}$

Equation (3.16) then gives

I

$$q_{+} = a_{n}^{+} = c_{n}^{+} = 0 \qquad n \neq 0$$

$$p = \sqrt{\pi} a_{0}^{+} \qquad (4.6)$$

Therefore, from Eq. (3.17) and (-4.5) we obtain

$$C_{n} = 2 \frac{\tilde{d}_{n}}{a_{0}^{+}}$$

$$\tilde{d}_{n} = \frac{1}{2} \sum_{l=-\infty}^{\infty} C_{-l} C_{n+l} = \frac{1}{2} \sum_{l=0}^{\infty} \left(c_{-l}^{1} C_{n+l}^{1} + c_{-l}^{2} C_{n+l}^{2} \right)^{(4.7)}$$

To calculate the dependent modes a_n^- , we have to fix the normalization of the fermion fields. Consider the conserved charge Q

$$Q = \int d\Sigma_{x} \int -g \int^{x} (\Psi t^{x} \Psi)$$

= $\int d\Sigma_{x} \sqrt{-g} (\Psi t^{x} \Psi)$
= $\frac{4}{p} \int_{0}^{1} d\tau F^{\dagger} F$
= $N^{2} \frac{4}{p} \sum_{n}^{1} b_{n}^{\dagger} b_{n}$ (4.8)

This suggests the normalization $\mathbf{N}^2 = p/4$ to be convenient one. Now, we can put Eq. (3.18), (3.23), (4.5), (4.6) and (4.8) together to obtain

$$a_{n} = 2 \frac{d_{n}}{d_{0}^{+}}$$

$$d_{n} = \frac{1}{2} \sum_{k} a_{-k} \cdot a_{n+k} + \sum_{m}' (m + \frac{n}{2}) b_{jm}^{+} b_{j,m+n} \qquad (4.9)$$

and

$$P_{-} = \frac{2\sqrt{\pi}}{\alpha \epsilon} \left(\frac{1}{4} + \frac{1}{4} \right)$$

where periodicity requires $C_0^- = G_0^-$, G_0^-

$$\mathcal{I}_{c} = \widetilde{\mathcal{I}}_{c}$$
 (4.10)

The Hamiltonian is now given by

$$H = 4\pi (f_{o} + f_{o})$$

= $4\pi \left[\sum_{n=1}^{\infty} (a_{n}^{\dagger} \cdot a_{n} + c_{n}^{\dagger} \cdot c_{n}) + a_{o}a_{o} + \sum_{n} inb_{jm}^{\dagger} b_{jm} \right]$
(4.11)

where all the independent modes appear in bilinear forms. Furthermore, H is already diagonalized. This suggests that, if we introduce canonical quantization to these modes, any function which can be expressed in these modes automatically satisfies the Hamiltonian equation of motion. We shall not burden the reader with details. The quantized system is specified as follows:

$$d_{m} = b_{-m} \qquad m > c \qquad (4.12)$$

are respectively the annihilation and creation operators for antiquark modes while b_m , b_m^{\dagger} (m > 0) are those for quark modes. They where b_m are the set of the set

$$\{b_{in}, b_{im}^{\dagger}\} = \{d_{in}, d_{im}^{\dagger}\} = S_{m,n} S_{ij}$$

(4.13)

where i and j are quark species indices. We also have

$$[an, a_{\ell}] = n \, \delta n, -\ell \, \delta^{\Lambda j}$$
(4.14a)

$$[c_n, c_n^3] = n \delta_{n,-1} \delta^{n,-1}$$
 (4.14b)

$$\left[\begin{array}{c} q^{\lambda}, a^{\lambda}_{\sigma} \end{array}\right] = \lambda S^{\lambda} \delta \qquad (4.14c)$$

$$[q_{-}, a_{0}^{+}] = -\lambda$$
 (4.14d)

where the upper indices i, j = 1, 2 are the space-time indices. All other commutators (or anti-commutators) among b_m , d_m , c_n , a_n , q^i , q_- , a_0^j and a_0^+ are zero. Ambiguities arising from the ordering of operators are resolved by normal ordering with respect to the ground state $| \circ \rangle$

$$b_{m}|_{07} = d_{m}|_{07}$$
 $m > 0$
 $a_{n}^{i}|_{07} = C_{n}^{i}|_{07}$ $n > 0$ (4.15)

Hence the charge operator becomes

$$G = \sum_{m>0}^{\prime} \left(b_m b_m - d_m^{\dagger} d_m \right) + G_0$$
(4.16)

Also

so
$$\widehat{J}_{c} = \sum_{n=1}^{\infty} C_{n}^{\dagger} \cdot C_{n} + \frac{\widehat{a}_{c}^{2}}{2}$$

 $\widehat{J}_{o} = \sum_{n=1}^{\infty} \widehat{a}_{n}^{\dagger} \cdot \widehat{a}_{n} + \frac{\widehat{a}_{o}^{2}}{2} + \sum_{m \neq o}^{\prime} m \left(\underset{\text{bjm}}{\text{bjm}} \underset{\text{bjm}}{\text{bjm}} + \underset{\text{djm}}{\text{djm}} \right)$
(4.17)

where

$$C_{0}^{-} a_{0}^{+} = f_{0}^{-} - f_{0}^{-}$$

$$a_{0}^{-} a_{0}^{+} = f_{0}^{-} - x_{0}$$

 α_0 , $\overline{\alpha}_0$ and Q_0 are constants arising from normal ordering. Since the charge of the empty string is zero, we shall assume $Q_0 = 0$.

Condition (4.10) can be imposed weakly, i.e. all physical states 127 must satisfy

$$(\widehat{d}_{0} - \chi_{0}) | \Im \rangle = (\widehat{d}_{0} - \overline{\chi}_{0}) | \Im \rangle$$

$$(4.18)$$

Since a_0^{μ} is the momentum of the string, the mass ()/(²) spectrum follows immediately from the Hamiltonian:

$$m^{2} = 4\pi \mathcal{E} \left[\mathbf{J}_{0} + \mathbf{\tilde{J}}_{0} - \mathbf{a}_{0}^{2} - \mathbf{d}_{0} - \mathbf{\tilde{J}}_{0} \right]$$
(4.19)

where we restore the dimensional parameter ζ . The Regge trajectories are linear with slope $\alpha' = 1/8\pi \zeta$. and intercept $(\alpha_0 + \overline{\alpha}_0)$.

Some useful commutators are given in Appendix D.

B. No Lorentz Covariance

Since the quantization is non-covariant, Lorentz covariance has to be examined. To construct the angular momentum operators, the ordering problem arises again. This can be resolved by symmetrizing the classical expression in x_{μ} and $\vartheta^{\alpha}\nu$ to ensure that $M^{\mu\nu}$ is a Hermitian operator

$$M^{\mu\nu} = \int d\Sigma_{x} M^{\alpha\mu\nu} \int \overline{g}$$
$$= \frac{1}{2} \int d\Sigma_{x} \sqrt{-g} \left[x^{\mu} p^{x\nu} - x^{\nu} p^{x\mu} + p^{\alpha\nu} x^{\mu} - p^{\alpha\mu} x^{\nu} \right]$$

since we have shown earlier that the spin part of $M^{\alpha\mu\nu}$ is zero. The angular momentum operators are

$$M^{12} = q^{2} a_{w}^{2} - q^{2} a_{w}^{1} - \lambda \sum_{n=1}^{\infty} \frac{1}{n} \left[a_{-n}^{1} a_{n}^{2} - a_{-n}^{2} a_{n}^{1} + C_{-n}^{1} C_{n}^{2} - C_{-n}^{2} C_{-n}^{1} C_{n}^{1} \right]$$

$$M^{\lambda +} = q^{\lambda} a_{w}^{\lambda}$$

$$M^{\lambda -} = -\frac{1}{2} \left(q^{2} a_{w}^{+} + a_{w}^{+} q^{-} \right) - 4\pi E p$$

$$M^{\lambda -} = M_{+}^{\lambda -} + M_{-}^{\lambda -} - q^{-} a_{w}^{\lambda} - \frac{2\pi}{a_{w}^{+}} \left[a_{w}^{\lambda} E + E a_{w}^{\lambda} \right] p$$
where
$$M_{+}^{\lambda -} = \frac{1}{2} \left[q^{\lambda} \frac{1}{a_{w}^{2}} (d_{w} - d_{w}) + \frac{1}{a_{w}^{2}} (d_{w} - d_{w}) q^{\lambda} \right]$$

$$-\frac{\lambda}{a_{w}^{2}} \sum_{n=1}^{\infty} \frac{a_{-n}^{\lambda} d_{n} - d_{-n} a_{n}^{\lambda}}{n}$$

$$M_{-}^{\lambda -} = \frac{1}{2} \left[q^{\lambda} \frac{1}{a_{w}^{2}} (\tilde{L}_{w} - \tilde{L}_{w}) + \frac{1}{a_{w}^{2}} (\tilde{L}_{w} - d_{w}) q^{\lambda} \right]$$

$$E = (f_{w} - d_{w}) - (\tilde{J}_{w} - \tilde{J}_{w})$$

The presence of an explicit dependence on φ does not violate the conference of angular momentum since we have $\Xi \mid \Im \rangle = 0$ and E commutes with all $M^{\mu\nu}$ and the Hamiltonian. It is straightforward to verify that all

commutation relations of $M^{\mu\nu}$ are satisfied except $[M^{1-}, M^{2-}]$, which must be zero for the Lorentz algebra to close. (Appendix) contains some useful commutators.) Define

$$d'_n = d_n - \alpha \cdot \alpha_n$$
$$\vec{d}'_n = \vec{d}_n - \alpha \cdot \alpha_n$$

The troublesome part of M^{i-} can be subdivided into the left moving part

$$M_{L}^{\lambda-} = -\frac{i}{a_{t}^{\star}} \sum_{n=1}^{\infty} \frac{a_{n}^{\star} dn - d_{n} a_{n}^{\star}}{n}$$
(4.20)

and the right moving part

I

$$M_{R}^{\lambda-} = -\frac{\lambda}{a_{v}} \sum_{n=1}^{\infty} \frac{C_{-n} \tilde{d}_{n} - \tilde{d}_{-n} C_{n}}{n}$$
(4.21)

It is straightforward (though tedious) to verify the following

$$a_{\perp}^{+2} \left[M_{\perp}^{\hat{\lambda}-}, M_{\perp}^{\hat{\lambda}-} \right] = 2 \left(\frac{1}{2c} - \frac{N+2}{24} \right) \sum_{n=1}^{\infty} \frac{a_{n}^{\hat{\lambda}} a_{n}^{\hat{\lambda}} - a_{-n}^{\hat{\lambda}} a_{n}^{\hat{\lambda}}}{n} + \left(1 - \frac{N+2}{24} \right) \sum_{n=1}^{\infty} n \left(a_{-n}^{\hat{\lambda}} a_{n}^{\hat{\lambda}} - a_{-n}^{\hat{\lambda}} a_{n}^{\hat{\lambda}} \right) (4.22)$$

$$a_{e}^{\dagger 2} \left[M_{L}, M_{R}^{\star -} \right] = 0$$

and

$$d_{0}^{+2} \left[M_{R}^{\lambda^{-}}, M_{R}^{j^{-}} \right] = 2\left(\frac{2}{L_{0}} - \frac{N'+2}{24} \right) \sum_{n=1}^{\infty} \frac{c_{n}^{\lambda} c_{n}^{j} - c_{-n}^{j} c_{n}^{\lambda}}{n} + \left(1 - \frac{N'+2}{24} \right) \sum_{n=1}^{\infty} n \left(c_{-n}^{\lambda} c_{n}^{j} - c_{-n}^{j} c_{n}^{\lambda} \right)$$
(4.23)

where N and N' are the number of quarks moving left and right respectively. For the subsystem we have been considering so far, N' = 0. Hence the/term in Eq. (4.23) does not vanish in the commutator $[M^{1-}, M^{2-}]$ and the Lorentz algebra is not closed. However, it is still interesting to note that the commutator of the left moving part of $[M^{1-}, M^{2-}]$ vanishes if N = 22 and $\alpha_0 = 1$ (where Eq. (4.22) is used). If, at the beginning, we choose

$$\lambda \delta_5 + \mu + \psi_j = -\psi_j$$
 $j = 1, 2, \cdots N$

£

instead of the projection (2.2), then the left moving part of $[M^{1-}, M^{2-}]$ will not vanish while the right moving part vanishes if N' = 22 and $\overline{\alpha}_0 = 1$. The inclusion of N_s number of massless scalar fields moving in the same direction as the fermion fields (say, left moving) changes the condition on the number of fermions to⁽¹¹⁾

$$N \leq N + N_S = 22$$

That this non-linear representation of the Lorentz algebra does not close is due to the normal ordering of the operators. This is purely a quantum mechanical effect.

V. Discussions and Speculations

A. The quark-Binding String

We have shown that the well defined, covariant subsystem of quarks moving clockwise around a closed string do not have a relativistic quantum mechanical solution. Mathematically it is due to the non-linear dependence among the dynamical variables. We are very much aware that there may exist singular modes in the solution which we have discarded by fiat ; their presence may restore the much wanted Lorentz covariance. Also it is likely that our quantization procedure is incorrect; to resolve the difficulty, we just have to find the proper quantization. In either case, it may require slight modifications in the original

formulation of the string. Barring these possibilities, we believe that quantum mechanics and relativity are putting very strong constraints on the non-linear system. To interpret boldly the negative results we are facing, we shall go into a wide spree of speculations. No apology is offered to the cautious readers, since the unsupported statements to be made are so precise that we are confident they can be proven or disproven in the very near future.

We shall formulate our speculations in the form of a conjecture. First we make a few observations, keeping in mind also the Nambu string in 3 and in 26 dimensions and the quark binding string in three dimensions.

(1) If we include some massless scalar particles moving clockwise around the closed string, then the clockwise part of the Lorentz algebra will close if the total number of Dirac and scalar quarks is 22. The anti-clockwise part of the Lorentz algebra remains not satisfied. (2) If we consider the other projection of the quark field, $\lambda \delta_5 + \Psi \Psi = -\Psi$, the quarks will be moving anti-clockwise, i.e. $\Psi \propto F(\sigma)$.

(3) The spectrum of the subsystem contains the spectrum of the empty string (i. e. Nambu string). We expect the spectrum of the complete system to contain the spectrum of the closed string with quarks moving clockwise only or with quarks moving anti-clockwise only.

(4) The string has only one dimensional parameter ζ ; it is not surprising that the Regge trajectories are straight lines.

(5) For the open string to exist, we need a linear combination of the left and right moving quark waves such that the string is a closed system: that is, the energy-momentum and current densities vanish at the ends of the string.

(6) Despite the complicated looking form of the implicit solution of the quark fields, Eq. (3.21), the quark part of the Hamiltonian (4.10) is very simple.

Now we state the speculation:

Conjecture1: Consider the following mass spectrum

$$\mathcal{D}_{1}^{2} = 4 \pi G \left[\sum_{n \neq 0}^{\infty} a_{n}^{\dagger} \cdot a_{n} + \sum_{j=1}^{N} \sum_{m \neq 0}^{\prime} m \left(b_{jm}^{\dagger} b_{jm} + d_{jm}^{\dagger} d_{jm} \right) \right] \\ + \sum_{k=N+1}^{22} S_{nk}^{\dagger} S_{nk} - \alpha_{0} \right]$$

where a_{p} , s_{n} obey the commutator (4.14a); b_{m} and d_{m} obey Eq. (4.13). Its Lorentz algebra can be defined consistently to be identical to that given in Section IV with the following modifications

 $\begin{aligned} & \mathcal{C}_{n} = 0 \quad , \quad \overline{\mathcal{A}}_{o} = 0 \\ & \overline{\mathcal{A}}_{n} = 0 \quad E = 0 \\ & \overline{\mathcal{A}}_{n} = \frac{1}{2} \sum_{k} : \left(\alpha_{-1} \cdot \alpha_{n+1} + S_{-1} \cdot S_{n+k} \right) : + : \sum_{k} \left(m + \frac{n}{2} \right) b_{jm}^{\dagger} b_{j,m+n} : \end{aligned}$

$$Sn | 07 = 0 \qquad n > 0$$

where

$$d_{m} = b - m$$
$$d_{m} = b^{\dagger} - m$$
$$s - n = s^{\dagger} n$$

We interpret b_m and d_m to be quark and antiquark oscillators; s_n to be massless scalar field oscillators. ⁽¹²⁾ Closure of the Lorentz algebra requires $\alpha_0 = 1$. Hence the lowest "mesonic" state $b_{\frac{1}{2}}^{\dagger} d_{\frac{1}{2}}^{\dagger} |0\rangle$ has zero mass. $\alpha_0 = 1$ also prevents condensation: the mass of a state with M pairs of quark-antiquark is heavier than M "mesonic" states each with one of the M pairs of quark-antiquark. We speculate that this mass spectrum (and its Lorentz algebra) belongs to the (open) quark-binding string with the addition of (22-N) scalar fields

The Regge slope is $\alpha' = 1/4\pi \zeta$ for the open string. We also speculate that the closed quark-binding string has no fermionic states in its spectrum.

B. The Quark-Confining String

The quark-binding string model formulated in Section II has an arbitrary constant & introduced as a tension to prevent the quark fields from expanding without bound. It is the analogue of the MIT bag constant (as a pressure) and bubble the SLAC constant (as a surface tension). It may be the effective binding force of some quark-gluon interactions, so that the quark-binding string can be considered as a phenomenological description of some quark-gluon interacting field theory. However, the present formalism is at best preliminary: there is no quark confinement; that is, there are exotic states in the spectrum. Also the presence of pure vibrational modes imply the existence of hadronic states which have no quark modes. Experimentally there is no evidence of such states.

To improve the string model, we present our second conjecture.

 $\frac{\text{Conjecture 2:}}{S = \int d^{2}u^{\alpha} \sqrt{-g} \left(\overline{\psi_{j}} \overline{\mathcal{T}}^{\alpha} \left(\frac{\lambda}{2} \overline{\partial_{x}} - e A^{\alpha}_{\alpha} \overline{\mathcal{T}}^{\alpha} \right) \psi_{j} - \frac{1}{4} F^{\alpha}_{\alpha\beta} F^{\alpha \times \beta} \right)}$ $= \int d^{2}u^{\alpha} \mathcal{L}$ $F^{\alpha}_{\alpha\beta} = F^{\alpha}_{\mu\nu} \overline{\mathcal{T}}^{\mu}_{\alpha} \overline{\mathcal{T}}^{\nu}_{\beta} = \partial_{\alpha} A^{\alpha}_{\beta} - \partial_{\beta} A^{\alpha}_{\alpha} + e f^{abc} A_{bx} A_{c\beta}$

and $F^{a}_{\alpha\beta} = 0$ at the ends of the string.

 $A^{a}_{\alpha} = \tau^{\mu}_{\alpha} A^{a}_{\mu}$ are non-abelian gauge fields in the color SU(3) adjoint representation. Quarks come in color triplets. They must obey the same constraints in Table I as before to ensure the string is a closed system. Fermion masses $(M\bar{\Psi}\Psi)$ can of course be added if needed. f^{abc} is the group structure constant :

$$[T^a, T^b] = f^{abc} T^c$$

where T^a is an SU(3) matrix, a = 1, 2, ..., 8. This model we refer to as

the quark-confining string, since only color singlet states exist. If necessary, we can also introduce any number of color singlet scalar fields into the Lagrangian

 $t \rightarrow t' + \int -g \left[\frac{1}{2} \partial^2 \phi_i \partial_x \phi_i - \frac{1}{2} \mu_i^2 \phi_i^2 \right]$

This may be necessary to saturate the Lorentz algebra. Notice that the string constant ζ is not included. Intuitively we see that there is no color vector gluons, since $F^{a}_{\alpha\beta}$ has only one component, namely $F^{a}_{C_{1}} = E^{a}_{;}$ A^{a}_{1} can be removed by an appropriate gauge choice. This leaves only the "Coulomb potential," which provides an attractive force along the string, very much like a tension in the presence of quarks. Hence the spectrum contains color singlet states with quark and vibrational excitations. In the absence of quarks, the tension (i.e. Coulomb disappears, so that the scalar fields are not bound. Hence one force) also this can hope that the relativistic invariant quark-confining string model has only mesons and baryons among its low lying states; there are no states which have only vibrational, color and/or scalar modes. At high energies, there also exist color singlet states with 3N quarks plus M quark-antiquark pairs. If the Regge intercept is positive, which is probable, these states would have very large widths (to decay to normal baryons and mesons) and therefore very difficult to observe.

We believe this quark-confining string, where exact color symmetry and quark-binding (along a line element) are put in by hand, offers enough computational simplifications to make it a worthwhile approach for studying quark confinement.

We may also consider the Han-Nambu color. The color gauge field involved is abelian. One may hope to solve this model exactly. In practice, it is like putting the Schwinger model in a dynamical curved space, which is imbedded in Minkowski space. It is straightforward to write down the action :

$$S = \int d^{2}u^{\alpha} \int -g \left[\overline{\Psi}_{j} \neq^{\alpha} \left(\frac{\lambda}{2} \overline{\partial}_{\alpha} - e_{j} A_{\alpha} \right) \Psi_{j} - \frac{F^{2}}{4} - G - M_{j} \overline{\Psi}_{j} \Psi_{j} \right]$$

where, for the moment, we include the constant $\,\zeta\,$. The equations of motion are (see fig. 2a)

and Ψ obeys the usual constraints. Neglecting the quark term in $\tau^{\alpha\beta}$, we see that, in the coordinate $\int_{\alpha} \phi = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$

$$T^{00} = G + \frac{E^2}{2} = G + \left(\frac{F^{01}}{2}\right)^2 > 0$$

Since the electric field (E^2) term is positive, we can take $C \rightarrow O$. The boundary conditions at the ends of the string must be imposed to ensure the system is closed:

$$E = 0$$
 at ends
 $P^{P}_{\mu} = 0$

This means each end of the string must end with a quark (or an antiquark) as

in fig. 2b. The baryon string has a configuration shown in fig. 2c. For the non-abelian case, fig. 2(d, e) may also have to be considered. The coupling constant \mathcal{C} is the only dimensional parameter in the model in the absence of the constant (\mathcal{L}) and masses M_j ($e_j \ll e$). In the presence of quark masses, the Regge trajectories will not be linear for the low mass states.

For the case where $\left(\frac{M}{2}\right)^2 \ll j$, the system should be solved in the absence of masses and treat the mass terms perturbatively. For the case $\left(\frac{M}{2}\right)^2 \gg j$, it is straightforward to show, (and is left as an exercise for the reader) that, to the lowest order approximation, the potential between a quark and an antiquark rises linearly $V(\gamma) \propto \gamma$.

41

Acknowledgments

The author would like to thank his colleagues at SLAC, especially R. Giles, for helpful discussions.

Appendix A

The best way to show $\dot{x}_i \dot{\Psi}_k = 0$ is probably the straightforward way. Let

$$\mathbf{q} = \begin{bmatrix} \mathbf{a} \\ \mathbf{v} \\ \mathbf{a} \\ \mathbf{z} \\ \mathbf{z} \end{bmatrix}$$

Equation (2.20) puts constraints among the four components. Using the Dirac representation (2.15) the constraints become

$$v = gu = \frac{1 + \lambda (m_{\ell} n_{\ell} - m_{\ell} n_{-})}{\lambda (m_{\ell} n_{\ell} - m_{\ell} n_{\ell})} u$$

$$j = f w = \frac{1 + i (m_{-}n_{+} - m_{2}n_{2})}{i (n_{-}m_{2} - m_{-}n_{2})} u^{3}$$

where

Ł

$$M_+ = M_0 \pm M_3$$

$$m_1 = m_1 + i m_2 = m_1^2$$

and similarly for n_{\pm} and $n_{\ell,~\gamma}$. Using Eq. (2.6) and (2.7), we obtain

$$f^{\dagger}g = fg^{\dagger} = -1$$

Now

$$\dot{j} = \begin{bmatrix} c & -j_{c} - j_{c} \vec{\sigma} \\ -j_{c} + \bar{j}_{c} \vec{\sigma} & c \end{bmatrix} = (\bar{\psi} \chi_{\mu} \psi) \chi^{\mu}$$

Hence

$$\dot{q} \Psi_{h} = \begin{bmatrix} -\dot{j}_{+} W_{h} - j_{k} \bar{j}_{h} \\ -\dot{j}_{+} W_{h} - j_{-} \bar{j}_{h} \\ -\dot{j}_{-} W_{h} - j_{-} \bar{j}_{h} \\ \bar{j}_{+} W_{h} - \bar{j}_{+} \bar{v}_{h} \end{bmatrix} \propto 1 + f^{\dagger}g = 0$$

where

7

ş

$$j_{+} = 2u^{+}u + 2j^{+}j$$

 $j_{-} = 2v^{+}v + 2w^{+}w$
 $j_{l} = 2v^{+}u - 2j^{+}w$
 $j_{r} = 2u^{+}v - 2w^{+}j$

where Ψ (in \mathbf{j}_{μ}) and $\Psi_{\mathbf{k}}$ can either be the same spinor or different spinors.

Appendix B

 $T^{\alpha\beta}$ can be chosen to be off-diagonal and symmetric if it is symmetric and has signature (1, -1) to begin with. The proof is almost identical to that given for Ref. (8). Since $g^{\alpha\beta}$ is already symmetric, we need to show $Q^{\alpha\beta}$ is symmetric. $\Theta^{\alpha\beta}$ is a tensor; hence it is sufficient to show that it is symmetric at any given point in some coordinate system. At any given point, we can choose locally geodesic coordinates

$$g_{x\beta} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$g_{x\beta} = 0$$

Equation (3.2) becomes

$$t't^\circ = -\lambda\delta s H H = -t^\circ t'$$

It is then straightforward to show that $e^{01} = e^{10}$ in this coordinate system using the Dirac Equation (2.25) and Eq. (2.21). To show that $T^{\alpha,\beta}$ has signature (1,-1) all we need is to show that det $(T^{\alpha/\beta}) < 0$, where

$$dit(T^{\alpha,3}) = \frac{\mathcal{E}^2}{\mathcal{G}} + \frac{\mathcal{E}}{\mathcal{G}}\mathcal{E}^{\alpha} + det(\mathcal{E}^{\alpha,3})$$

Dirac equation (2.25) implies

$$\frac{1}{2}\overline{\psi}t^{*}\overline{\partial}_{x}\psi=0$$

and the light-like currents (Appendix A) implies the vanishing of the last term. Since $g^{\alpha,3}$ also has signature (1,-1) then

$$det(T^{x,3}) = \frac{e^2}{g} < 0$$

That $T^{\alpha,3}$ can be chosen to be off-diagonal is, of course, due to the arbitrariness in the choice of the parametrization. This is also related to the fact that our phase space is constrained. To see this, let us consider the canonical momentum $\mathcal{P}^{\circ}_{\mu} = \frac{SL}{S \times P}$

₹-Э ,

and the shifted coordinate

$$y_{i\mu} = \tau_{i\mu} - \frac{\hbar}{2C} q_j Y_{\mu} \partial_i q_j$$

Naively one may expect them to be independent. However, they are constrained by the following relations:

$$P^{\circ}_{\mu} Y_{1} = 0$$

$$P^{\circ}_{\mu} P^{\circ \mu} + e^{2} Y_{1\mu} Y^{1\mu} = 0$$

Appendix C

The Gauss-Codazzi equations involve the torsion ν_{α} which depends on the choice of the normals n_{μ} and m_{μ} . To verify the Gauss-Codazzi equations in the coordinates chosen, we first make an explicit choice of n_{μ} and m_{μ} ; we then calculate all the curvature tensors and the torsion and sbustitute them directly into the equations.

The normal plane

$$\mathcal{U}_{\mu\nu} = n_{\mu}m_{\nu} - m_{\mu}n_{\nu}$$

is invariant under the rotation

$$M_{\mu} \rightarrow M_{\mu} \cos \varepsilon S + M_{\mu} \sin \Theta$$

$$M_{\mu} \rightarrow -n_{\mu} \sin \varepsilon + m_{\mu} \cos \Theta$$

Let us choose n_{μ} such that $n_{+} = n_{0} + n_{3} = 0$. (This is a valid choice since n_{μ} is space-like.) To do so, it is convenient to introduce some notations. Explicitly Eq. (2.4) gives

$$a = VU_{+-} = 2(\dot{x}_{2}\dot{x}_{1} - \dot{x}_{1}\dot{x}_{2}')$$

$$b = VU_{+1} = \dot{x}_{2}\dot{x}_{1}' - \dot{x}_{1}\dot{x}_{2}' = p(\dot{x}_{2} - \dot{x}_{2}') = pA$$

$$c = VU_{+2} = \dot{x}_{+}\dot{x}_{1}' - \dot{x}_{1}\dot{x}_{+}' = p(\dot{x}_{1}' - \dot{x}_{1}) = -pt$$

$$d = VU_{12} = \frac{1}{2}(\dot{x}_{+}\dot{x}_{-}' - \dot{x}_{-}\dot{x}_{+}') = \frac{1}{2}(\dot{x}_{-}' - \dot{x}_{-})$$

$$d = VU_{-1} = \dot{x}_{-}\dot{x}_{2}' - \dot{x}_{2}\dot{x}_{-}'$$

$$f = VU_{-2} = \dot{x}_{1}\dot{x}_{-}' - \dot{x}_{-}\dot{x}_{1}'$$

where

I

$$V = \int -g$$

Note that bf = ad + CE

and

$$V = \frac{t^2 + s^2 + A}{2} = d^2 - \frac{a^2}{4} - be - cf$$

Now it is straightforward to construct $\mathbf{n}_{\mu},\ \mathbf{m}_{\mu}$ in terms of $\dot{\mathbf{x}}_{\mu},\ \mathbf{x}_{\mu}'$:

$$\begin{split} H_{\mu} &= (H_{+}, H_{-}, H_{1}, H_{2}) \\ &= \frac{1}{\sqrt{b^{2} + c^{2}}} (0, a, b, c) \\ H_{\mu} &= \frac{1}{2\sqrt{b^{2} + c^{2}}} (2b^{2} + 2c^{2}, a^{2} + 2be + 2cf, ba + 2cd, ac - 2bd) \end{split}$$

Using Eq. (3.11), we now obtain

$$t_{11} = \frac{t x_1'' + s x_2''}{\sqrt{s^2 + t^2}}$$

$$f_{11} = \frac{s x_1'' - t x_2''}{\sqrt{s^2 + t^2}}$$

$$v_e = -m_{\mu} it^{\mu} = \frac{t \dot{x}_2 - s \dot{x}_1}{s^2 + t^2}$$

$$v_f = \left(\frac{1}{s^2 + t^2} - \frac{1}{V}\right) \left(s x_1'' - t x_2''\right)$$

It is clear that the second Gauss-Codazzi Equation (3.15) is satisfied.

The first Gauss-Codazzi equation (3.14) can also be verified easily once we obtain the other components of the curvature tensors. They are, using Eq. (3.10),

$$h_{cc} = \frac{s\ddot{\imath}_{1} - t\ddot{\imath}_{2}}{\sqrt{s^{2} + t^{2}}}$$

$$l_{cc} = \frac{1}{\sqrt{\sqrt{s^{2} + t^{2}}}} \left(A\dot{V} - \frac{\dot{A}V}{2} - \dot{V}V \right)$$

where

$$\dot{V} = \frac{\dot{A}}{2} + t\ddot{x}_1 + s\ddot{x}_2$$

For the record, we note that the "mean curvature" is given by

$$k = \sqrt{k_1^2 + k_2^2} = -\frac{A}{2V^2} \left(\chi_1^{1/2} + \chi_2^{1/2}\right)^{1/2}$$

Appendix D

Given the commutators among the independent operators a_n, c_n, b_m, d_m, q_i, q_j, a_0^i and a_0^+ , all commutators can be calculated. We give some of the useful ones. [Qj, him] = Sij bim [Gig , dim] = - Sig dim $[dn, a_{2}] = -l a_{2+n}$ $[L_{n}, b_{m}] = -(m+\frac{n}{2})b_{m+n}$ $\begin{bmatrix} 2 \\ m \\ bm \end{bmatrix} = \begin{pmatrix} m - \frac{m}{2} \end{pmatrix} \vec{b} m - n$ $\left[\widehat{f}_{n}, C_{l}\right] = -lC_{l+n}$ $[q_{-}, \frac{1}{a_{+}^{+}}] = \frac{\lambda}{(a_{+}^{+})^{2}}$ $[dn, dm] = (n-m)dn+m + Sn, -m(n^3-n)(N+1)$ $\left[\widetilde{\mathcal{A}}_{n},\widetilde{\mathcal{A}}_{m}\right] = (n-m)\widetilde{\mathcal{A}}_{n+m} + S_{n,-m}(\underline{n^{3}-n})$

where N is the number of quark fields. The constant terms in the last two commutators play the same role as c-number Schwinger terms.

References

- Y. Nambu, in <u>Symmetries and Quark Models</u>, edited by R. Chand (Gordon and Breach, New York, 1970); H. Nielsen, in <u>High Energy Physics</u>, <u>Pro-</u> <u>ceedings of the Fifteenth International Conference on High Energy Physics</u>, <u>Kiev, 1970</u>, edited by V. Shelest (Naukova Dumka, Kiev, USSR, 1972); L. Susskind, Nuovo Cimento <u>69A</u>, 457 (1970); T. Takabayashi, Prog. Theor. Phys. 43, 1117 (1970).
- 2. G. Veneziano, Nuovo Cimento 57A, 190 (1968) and numerous papers thereafter.
- Y. Nambu, lectures at Copenhagen Summer Symposium, 1970 (unpublished);
 P. Goddard, J. Goldstone, C. Rebbi and C. B. Thorn, Nucl. Phys. <u>B56</u>, 109 (1973). See also L. N. Chang and F. Mansouri, Phys. Rev. <u>D5</u>, 2535 (1972).
- 4. See, e.g., H. B. Nielsen and P. Olesen, Nucl. Phys. <u>B61</u>, 45 (1973);
- K. Wilson, Phys. Rev. <u>D10</u>, 2445 (1974);
 A. M. Polychov, Landau Institute Preprint (1974)
 Y. Nambu, Phys. Rev. <u>D10</u>, 4262 (1974); J. Kogut and L. Susskind,
 Phys. Rev. <u>D11</u>, 395 (1975); S. Mandelstam, Phys. Lett. <u>53B</u>, 476 (1975);
 L. D. Faddeev, Max-Planck-Institut (Munchen) preprint MPI-PAE/Pth 16 (1974).
 Z. F. Ezawa and H. C. Tze, DAMTP Preprint (1975)
 S. Mandelstam, Nucl. Phys. <u>B64</u>, 205 (1973).
- A. Chodos, R. L. Jaffe, K. Johnson, C. B. Thorn and V. F. Weisskopf, Phys. Rev. <u>D9</u>, 3471 (1974).
- W. A. Bardeen, M. S. Chanowitz, S. D. Drell, M. Weinstein and T. M. Yan, Phys. Rev. <u>D11</u>, 1094 (1975); R. C. Giles, SLAC preprint SLAC-PUB-(1975).
- 8. R. C. Giles and S. H. H. Tye, SLAC preprint, SLAC-PUB- (1975).
- See e. g. L. P. Eisenhart, Differential Geometry (Princeton University Press, Princeton, 1949). See. R. C. Giles, Ref. (7), for further references and also a lucid discussion on the imbedding of hypersurfaces.
- 10. After the completion of this work we were informed by L.-N. Chang of the

work by L.-N. Chang and F. Mansouri in Workshop on Current Problems in High Energy Physics, John Hopkins University, January 1974, where a similar model has been considered.

- 11. R. Staffin (unpublished)
- 12. Compare with K. Bardakci and M. B. Halpern, Phys. Rev. D3, 2493 (1971).
- 13. The constraint $\overline{\Psi}\Psi = 0$ may be relaxed in the presence of quark mass. The derivation of the equations of motion does not use it. Quark masses may also require further relaxation of the constraints on the fermion fields.

| TABLE 1 |
|---------|
|---------|

| The Quark-Binding String | |
|---|--------------|
| $\lambda \partial_{\mu} \Psi = -\frac{\lambda}{2} \left(h^{\alpha} + l^{\alpha} + l^{\alpha$ | (i) |
| $-(ch'_{x} = \frac{\lambda}{2}\overline{\Psi}_{j}(J_{x}H)\vec{J}^{x}\Psi_{j}$ $-(ch'_{x} = \frac{\lambda}{2}\overline{\Psi}_{j}(J_{x}H)\vec{J}^{x}\Psi_{j}$ | (ii) (ii) |
| and $\overline{\psi}\psi = j \cdot n = j \cdot m = p_k^{\alpha} \cdot n = p_k^{\alpha} \cdot m = 0$ | |
| The Subsystem | |
| Equations (i), (ii), and (iii), and, for each quark $\lambda J_5 H_4 \Psi = \Psi$ $j_{5\mu} = \Psi J_{\mu} J_5 \Psi = 0$ | |

Proper boundary conditions have to be included for an open or a closed string. The constraints need not be independent. The subsystem describes massless quarks moving clockwise around the closed

string.























Quark-confining strings. In (C), 2/3 and 1/3 are the Abelian color hypercharges.