SLAC-PUB-1691 ITP-519 FNL-PUB-75-92-THY COO-3075-131 December 1975 (T)

A STUDY OF THE LONGITUDINAL KINK MODES OF THE STRING

W. A. Bardeen^{*}

Fermi National Accelerator Laboratory Batavia, Illinois 60510

Itzhak Bars[‡]

Department of Physics, Yale University New Haven, Connecticut 06520

Andrew J. Hanson*

Stanford Linear Accelerator Center Stanford University, Stanford, California 94305

and

R. D. Peccei[‡]

Institute of Theoretical Physics, Department of Physics Stanford University, Stanford, California 94305

ABSTRACT

We examine the massless limit of a model for the massive relativistic Nambu string. The system possesses longitudinal kink modes excluded from the standard lightlike gauge treatment. We demonstrate the equivalence of these modes to those proposed by Patrascioiu. The classical nonlinear field theory of the twodimensional string is shown to be a completely integrable Hamiltonian system. The Hamiltonian is expressed in terms of normal mode action variables alone; the mass-squared spectrum is linear in the Bohr-Sommerfeld approximation. The difficulties of canonical quantization are exposed using a particular timelike gauge which admits commuting center-of-mass coordinates.

^{*}Research supported in part by the U.S. Energy Research and Development Administration.

^{*}Research (Yale Report COO-3075-131) supported in part by the U.S. Energy Research and Development Administration, under Contract No. AT(11-1) 3075.

 $^{^{\}ddagger}$ Research supported in part by the National Science Foundation, grant PHY 75-18444.

INTRODUCTION

The massless relativistic string model derived from the Nambu action functional [1] is widely believed to be connected in an essential way with the dual models for strong interactions. It is therefore puzzling that despite the desirable features of the dual models, the free string quantization procedure of Goddard, Goldstone, Rebbi and Thorne [2] (GGRT) and the interacting string quantization of Mandelstam [3] succeed only in 26 dimensional spacetime. Patrascioiu [4] has pointed out that this phenomenon may occur due to the omission of longitudinal modes of oscillation which appear when the massless string action principle is suitably generalized. Patrascioiu's modes belong to a class of solutions with discontinuous derivatives which appear to be excluded from the GGRT solutions due to a singularity in their choice of coordinates.

Our purpose here is to reexamine the string with particular attention to the longitudinal modes in two spacetime dimensions. We argue that a physically sound procedure for constructing the Hamiltonian is to define the massless relativistic string as the smooth massless limit of a <u>massive</u> relativistic string [5]. Then the longitudinal modes remain in the theory as subtle minima of the action principle: In two dimensions, these motions appear as massless limits of the solutions to the massive Euler equations, but cannot be derived from the standard Euler equations if the masses are set to zero before the variation. We note that the string is an essentially nonlinear system if the constraint equations are taken into account; the longitudinal modes of the string behave precisely like kink solutions of a nonlinear field theory, which enormously complicates any attempt at canonical quantization of the independent modes.

Our eventual aim is to write down the full quantum Hamiltonian and Poincaré group generators for the longitudinal-plus-transverse string oscillations in D dimensions, and then investigate positivity and Lorentz covariance for D = 4.

This program remains for the moment incomplete, so this work will concentrate on limited sectors of the full theory.

We begin by deriving the Hamiltonian dynamics of the massive relativistic string system. This yields a system of particles interacting via a relativistic potential which becomes linear in two dimensions. Then we take the massless limit of the massive theory, obtaining longitudinal modes of oscillation. We thus find a more physical understanding of the motions proposed by Patrascioiu, who used different methods. One section is devoted to showing the equivalence of our simplest mode to Patrascioiu's orthonormal gauge solution. Working mostly in two spacetime dimensions, where no transverse solutions exist, we approximate the quantum mechanics of the longitudinal oscillations by using a semiclassical Bohr-Sommerfeld approach. For the D = 2 string with an arbitrary mass distribution, we find in the massless limit a simple form for the Hamiltonian in terms of action variables alone. The theory of the kink solutions of the D = 2 string is thus a completely integrable Hamiltonian system, comparable in spirit to the classical system found by Faddeev and Takhtajan [6] for the sine-Gordon equation. Finally, using a gauge proposed by Rohrlich [7] to separate the Newton-Wigner center-of-mass coordinates of the string, we examine the Dirac bracket algebra of the fully constrained system and expose the difficulties of canonical quantization. An Appendix gives the Dirac bracket analysis and an alternative gauge-invariant approach to the oscillators of the constrained system.

1. MASSLESS LIMIT OF MASSIVE RELATIVISTIC SYSTEMS

Classical relativistic theories with points moving at the speed of light require extra care in the definition of the system. Here we review the theory of a massless relativistic scalar particle, so that we may later apply the resulting intuition to the string model.

We take as our Lagrangian

$$L = -\mu \left[-x_{\tau}^{2}\right]^{1/2}$$
(1.1)

where

$$x_{\tau}^{2} \equiv x_{\tau}^{\mu} g_{\mu\nu} x_{\tau}^{\nu} \equiv \frac{d\vec{x}}{d\tau} \cdot \frac{d\vec{x}}{d\tau} - \left(\frac{dx^{0}}{d\tau}\right)^{2} . \qquad (1.2)$$

The canonical momenta

$$p^{\alpha} = \frac{\partial L}{\partial x_{\alpha,\tau}} = \frac{\mu x_{\tau}^{\alpha}}{[-x_{\tau}^2]^{1/2}}$$
(1.3)

obey the constraint

$$p^2 + \mu^2 \approx 0 \tag{1.4}$$

and have canonical Poisson brackets (PB) given by

$$\{p^{\mu}(\tau), x^{\nu}(\tau)\} = -g^{\mu\nu} . \qquad (1.5)$$

In the timelike gauge

$$x^{0}(\tau) \approx \tau$$
, (1.6)

the Hamiltonian becomes

$$H = p^{0} = \left[\vec{p}^{2} + \mu^{2}\right]^{1/2} .$$
 (1.7)

If we now take the limit $\mu \not \rightarrow 0$ with p^{α} finite, we find a finite Hamiltonian

$$H = \left| \overrightarrow{p} \right| , \qquad (1.8)$$

even though the action functional <u>vanishes</u>. We also recover from Eq. (1.3) the usual result that for p^{α} to remain finite as the particle becomes massless, the particle must move at the speed of light,

$$x_{\tau}^2 = 0$$
 (1.9)

Another approach to the massless classical relativistic particle might be to consider the action functional

$$I[x] = -\int_{\tau_1}^{\tau_2} d\tau \ [-x_{\tau}^2]^{1/2} , \qquad (1.10)$$

where $x^{\mu}(\tau_1)$ and $x^{\mu}(\tau_2)$ have a lightlike separation. If we prohibit paths connecting $x^{\mu}(\tau_1)$ and $x^{\mu}(\tau_2)$ which make $[-x_{\tau}^2] \leq 0$ at any point, then there are no legal variations of the action functional; only the single lightlike path connecting the two points is permitted, and $I[x] \equiv 0$.

We interpret these observations as an indication that to find the correct Hamiltonian for a massless classical system, it may be necessary to begin with a massive system and take the limit as the mass goes to zero.

2. THE MASSIVE STRING

The Nambu action [1] describes a relativistic massless string. Because points on the string can move at the speed of light in this theory, the Hamiltonian may contain terms of the type found in the preceding section, which do not follow from the standard variational methods.

We are thus motivated to expand the techniques of the previous section and consider the theory of a massive relativistic string. When the string is massive, all points move with velocities less than the speed of light, so the standard methods may be applied without ambiguity to find the Hamiltonian. The Hamiltonian of the massless relativistic string is then found to be the smooth limit of the massive Hamiltonian as the mass goes to zero, as was the case for the point particle in Section 1.

The action for a massive relativistic string is

$$S[x] = \int_{\tau_1}^{\tau_2} d\tau \int_{0}^{\pi} d\sigma \left\{ -\mu \left[-x_{\tau}^2 \right]^{1/2} - \gamma \left[-G \right]^{1/2} \right\}, \qquad (2.1)$$

which has been briefly considered by Chodos and Thorne [5]. $x^{\mu}(\tau,\sigma)$ is a D-component field on (τ,σ) space and

$$G = x_{\tau}^{2} x_{\sigma}^{2} - (x_{\tau} \cdot x_{\sigma})^{2} . \qquad (2.2)$$

We define

$$p^{\alpha}(\tau,\sigma) = \mu x_{\tau}^{\alpha} [-x_{\tau}^{2}]^{-1/2}$$

$$K^{\alpha}(\tau,\sigma) = \{x_{\tau}^{\alpha} x_{\sigma}^{2} - x_{\sigma}^{\alpha} (x_{\tau} \cdot x_{\sigma})\} [-G]^{-1/2}$$

$$N^{\alpha}(\tau,\sigma) = \{x_{\sigma}^{\alpha} x_{\tau}^{2} - x_{\tau}^{\alpha} (x_{\tau} \cdot x_{\sigma})\} [-G]^{-1/2},$$
(2.3)

so that we may write the canonical momenta as

$$\theta^{\alpha}(\tau,\sigma) = \frac{\delta S}{\delta x_{\alpha,\tau}} = p^{\alpha}(\tau,\sigma) + \gamma K^{\alpha}(\tau,\sigma) , \qquad (2.4)$$

where neither p^{α} nor \textbf{K}^{α} are true canonical variables. The nonvanishing canonical Poisson brackets are

$$\{ \ell^{\alpha}(\tau,\sigma), x^{\beta}(\tau,\sigma') \} = -g^{\alpha\beta} \delta(\sigma-\sigma')$$
(2.5)

and $ensuremath{\mathcal{P}}^{\alpha}$ obeys the constraint

$$\mathcal{O}^{2} + 2\gamma [(\mathcal{O} \cdot x_{\sigma})^{2} + \mu^{2} x_{\sigma}^{2}]^{1/2} + \gamma^{2} x_{\sigma}^{2} + \mu^{2} \approx 0$$
(2.6)

due to $\tau\text{-reparametrization}$ invariance. If we choose a timelike gauge such as $x^0\approx\tau,$ we find the Hamiltonian

$$H[\mu] = \int_{0}^{\pi} d\sigma \, \ell^{0}(\tau, \sigma) = \int_{0}^{\pi} d\sigma \, \{\vec{\ell}^{2} + 2\gamma [(\vec{\ell} \cdot \vec{x}_{\sigma})^{2} + \mu^{2} \vec{x}_{\sigma}^{2}]^{1/2} + \gamma^{2} \vec{x}_{\sigma}^{2} + \mu^{2}\}^{1/2} \quad (2.7)$$

At this point, we may take the mass parameter $\mu \rightarrow 0$ with $\mathcal{O}^{\mu}(\tau,\sigma)$ fixed, yielding the Hamiltonian

$$H = \int_{0}^{\pi} d\sigma \left\{ (\vec{p} + \gamma \vec{K})^{2} + 2\gamma | \vec{p} \cdot \vec{x}_{\sigma} \right| + \gamma^{2} \vec{x}_{\sigma}^{2} \right\}^{1/2}$$
(2.8)

for the massless relativistic string. This differs from the usual massless string Hamiltonian

$$H_{0} = \gamma \int_{0}^{\pi} d\sigma (\vec{k}^{2} + \vec{x}_{\sigma}^{2})^{1/2} , \qquad (2.9)$$

even though the action functionals appear to be the same in this limit. To understand the difference between these two formulations of the Hamiltonian, we recall that the canonical momenta in Eq. (2.8) are given by Eq. (2.4), where $\vec{k} \cdot \vec{x}_{\sigma}$ is formally zero. When $\vec{p} = 0$, the two Hamiltonians are identical. However, as $\mu \neq 0$, $\vec{p}(\tau, \sigma)$ need not be zero in our formulation in any region of σ for which the string moves at the velocity of light $(x_{\tau}^2 = 0)$, as is evident from Eq. (2.3). Our approach thus makes it apparent that the string may have more general motions than those considered in the standard treatment of GGRT. The Nambu action is invariant under σ -reparametrizations as well as τ -reparametrizations. Placing mass on the string as in Eq. (2.1) breaks the σ -reparametrization invariance. However, the theory described by the Hamiltonian (2.8) regains the invariance. This may be seen by examining the generator of σ -reparametrizations

$$\Sigma[\mathbf{f}] = \int_{0}^{\pi} d\sigma \ \mathbf{f}(\sigma) \, \boldsymbol{\theta} \cdot \mathbf{x}_{\sigma} \quad . \tag{2.10}$$

The function $f(\sigma)$ is arbitrary save for the constraints

$$f(0) = f(\pi) = 0$$
 (2.11)

which follow from the requirement that the end points of the string map into themselves. $x^{\mu}(\tau,\sigma)$ and $\mathscr{O}^{\mu}(\tau,\sigma)$ transform under the action of $\Sigma[f]$ as

$$\{x^{\mu}(\sigma), \Sigma[\mathbf{f}]\} = \mathbf{f}(\sigma) \ x_{\sigma}^{\mu}(\sigma)$$
(2.12)

$$\mathscr{P}^{\mu}(\sigma), \Sigma[\mathbf{f}] = \partial_{\sigma} [\mathbf{f}(\sigma) \, \mathscr{P}^{\mu}(\sigma)] .$$
 (2.13)

One can now compute the bracket of H in Eq. (2.8) with $\Sigma[f]$. We find

$$\{H, \Sigma[f]\} = \int_{0}^{\pi} d\sigma \left[\frac{\partial P^{0}}{\partial \dot{x}_{\sigma}} \cdot \frac{\partial \left(f(\sigma)\dot{x}_{\sigma}\right)}{\partial \sigma} + \frac{\partial P^{0}}{\partial \vec{\sigma}} \cdot \frac{\partial \left(f(\sigma)\vec{P}\right)}{\partial \sigma}\right]$$

$$= \int_{0}^{\pi} d\sigma \left[f(\sigma)\frac{\partial P^{0}}{\partial \sigma} + \frac{\partial f(\sigma)}{\partial \sigma}\left\{\frac{\partial P^{0}}{\partial \dot{x}_{\sigma}} \cdot \dot{\vec{x}}_{\sigma} + \frac{\partial P^{0}}{\partial \vec{P}} \cdot \vec{P}\right\} \right]$$

$$= \int_{0}^{\pi} d\sigma \left[f(\sigma)\vec{P}^{0}\right] / \partial \sigma = 0 .$$

$$(2.14)$$

Equation (2.14) implies that $\Sigma[f]$ is a constant of motion. Hence we will be able to use σ -reparametrizations to further simplify the theory.

3. MASSLESS LIMIT OF A STRING WITH MASSIVE ENDS

The general Hamiltonian for the string system proposed in the previous section incorporates the possibility of many complex motions. To understand more clearly the properties of such motions, we devote this section to a detailed analysis of the simplest sector of the theory with $\vec{p}(\tau,\sigma) \neq 0$: We restrict ourselves to purely longitudinal motion and allow only the end-points to move at the speed of light in the massless limit. To illustrate the desired motion it is sufficient to consider the simplified action

$$S[x] = \int_{\tau_1}^{\tau_2} d\tau \{-\mu[-x_{\tau}^2(0)]^{1/2} - \mu[-x_{\tau}^2(\pi)]^{1/2} - \gamma \int_{0}^{\pi} d\sigma [-G]^{1/2} \}$$
(3.1)

describing a pair of point masses joined by a string. Even for $\mu \neq 0$, this action is invariant under the usual invariance group of σ -reparametrizations obeying $\tilde{\sigma}$ $(\tau, 0) = 0$, $\tilde{\sigma}$ $(\tau, \pi) = \pi$. In the massless limit, this theory corresponds to treating the Hamiltonian (2.8) in a specific gauge. This connection will be made clearer when we study gauge transformations between equivalent systems in Section 4. For $0 < \sigma < \pi$, we find the equations of motion

$$K_{\tau}^{\alpha} + N_{\sigma}^{\alpha} = 0 .$$
 (3.2)

At the endpoints, we have

$$p_{\tau}^{\ \alpha}(0) + \gamma N^{\alpha}(0) = 0$$

$$p_{\tau}^{\ \alpha}(\pi) - \gamma N^{\alpha}(\pi) = 0 . \qquad (3.3)$$

The variables $p^{\alpha},\ K^{\alpha}$ and N^{α} are defined as in Eq. (2.3). The conserved total momentum is

$$P^{\alpha} = \int_{0}^{\pi} d\sigma \, \mathscr{C}^{\alpha} = p^{\alpha}(0) + p^{\alpha}(\pi) + \gamma \int_{0}^{\pi} d\sigma \, K^{\alpha} , \qquad (3.4)$$

and the conserved Lorentz-transformation generator is

$$M^{\alpha\beta} = \int_{0}^{\pi} d\sigma (x^{\alpha} \, \ell^{\beta} - x^{\beta} \, \ell^{\alpha}) \\ = x^{\alpha}(0) p^{\beta}(0) - x^{\beta}(0) p^{\alpha}(0) + x^{\alpha}(\pi) p^{\beta}(\pi) - x^{\beta}(\pi) p^{\alpha}(\pi) + \gamma \int_{0}^{\pi} d\sigma (x^{\alpha} K^{\beta} - x^{\beta} K^{\alpha}) \quad (3.5)$$

A. Massive Classical Solutions

In order to solve the equations of motion, we must fix the gauge to eliminate arbitrary functions. Choosing the timelike gauge,

$$x^0 \approx \tau$$
 (3.6)

to fix the scale of $\boldsymbol{\tau},$ we find the Hamiltonian

$$H = \left[\vec{p}^{2}(0) + \mu^{2}\right]^{1/2} + \left[\vec{p}^{2}(\pi) + \mu^{2}\right]^{1/2} + \gamma \int_{0}^{\pi} d\sigma \left[\vec{K}^{2}(\sigma) + \vec{x}_{\sigma}^{2}(\sigma)\right]^{1/2} .$$
(3.7)

If we consider only longitudinal motions, so that we are effectively in two spacetime dimensions, then $\vec{k} = 0$ and $\vec{x}_{\sigma} \equiv x_{\sigma}$ giving

$$H = [p^{2}(0) + \mu^{2}]^{1/2} + [p^{2}(\pi) + \mu^{2}]^{1/2} + \gamma \int_{0}^{\pi} d\sigma |x_{\sigma}| .$$
 (3.8)

The Euler equations (3.2), (3.3) which we must solve are [5]

$$\frac{\partial}{\partial \sigma} \left\{ \frac{x_{\sigma}}{[x_{\sigma}^2]^{1/2}} \right\} = 0 , \quad 0 < \sigma < \pi$$
(3.9a)

$$\frac{\partial}{\partial \tau} \left\{ \frac{\mu x_{\tau}}{[1 - x_{\tau}^{2}]^{1/2}} \right\} \neq \frac{\gamma x_{\sigma}}{[x_{\sigma}^{2}]^{1/2}} = 0, \quad \{\sigma = 0, \sigma = \pi \}.$$
(3.9b)

Using Eq. (3.9a) and σ -reparametrization invariance, we see that we can make $x_{\sigma}^{(\tau,\sigma)}$ independent of σ for $0 < \sigma < \pi$, so that [8]

$$\int_{0}^{\pi} d\sigma |x_{\sigma}| = |x(\tau,\pi) - x(\tau,0)|. \qquad (3.10)$$

The Hamiltonian (3.8) then becomes

$$H = [p^{2}(0) + \mu^{2}]^{1/2} + [p^{2}(\pi) + \mu^{2}]^{1/2} + \gamma |x(\pi) - x(0)| . \qquad (3.11)$$

We can check classical Poincaré invariance of the two spacetime dimensional system by using Eq. (3.5) to derive the boost generator

$$B \equiv M^{01} = \tau \left(p(0) + p(\pi) \right) - x(0) \left[p^{2}(0) + \mu^{2} \right]^{1/2} - x(\pi) \left[p^{2}(\pi) + \mu^{2} \right]^{1/2} - \frac{1}{2} \gamma \left(x(0) + x(\pi) \right) \left| x(\pi) - x(0) \right| .$$
(3.12)

Taking the total momentum $P = p(0) + p(\pi)$ from Eq. (3.4), we verify the Poincaré group Poisson bracket algebra

 $\{P,H\} = 0$ $\{B,P\} = -H$

 $\{B,H\} = -P$.

The Hamiltonian (3.11) therefore describes a classically Poincaré-invariant system.

The solutions of the Euler equations are now of the form

$$x(\tau,\sigma) = \pm \left(\frac{2\sigma}{\pi} - 1\right) \left\{ -\left[\left(\tau - \tau_0\right)^2 + \mu^2/\gamma^2\right]^{1/2} + \left[\pi^2/4 + \mu^2/\gamma^2\right]^{1/2} \right\}$$
(3.13)

in the rest frame, where $P = p(0) + p(\pi) = 0$. For $-\frac{\pi}{2} < \tau < \frac{\pi}{2}$, one chooses the (+) sign and sets $\tau_0 = 0$; this solution joins continuously onto the next τ region as shown in Fig. 3.1 provided that for $\frac{\pi}{2} < \tau < \frac{3\pi}{2}$, one chooses $\tau_0 = \pi$ and the (-) sign, etc. We observe that at $\sigma = 0$ and $\sigma = \pi$,

$$[1 - x_{\tau}^{2}]^{1/2} = (\mu/\gamma) [(\tau - \tau_{0})^{2} + \mu^{2}/\gamma^{2}]^{-1/2} , \qquad (3.14)$$

so that the first term in the $\sigma = 0$, π boundary condition (3.9b) remains finite as $\mu \neq 0$. For any μ , the momentum of the end points is thus

$$p(\sigma = 0, \pi) = \pm \gamma(\tau - \tau_0)$$
 (3.15)

In Fig. 3.2, we plot p(0), $p(\pi)$ and $x_{\sigma}(\sigma)$ as a function of τ .

B. Massless Limit

In the massless limit, our Hamiltonian (3.8) with the gauge choice (3.10) becomes

$$H = |p_0| + |p_{\pi}| + \gamma |x_{\pi} - x_0| , \qquad (3.16)$$

where $x_0 = x(\tau, \sigma = 0)$, etc. It is amusing to observe that the potential energy in this Hamiltonian has precisely the form of a one-space-dimension Coulomb potential, so this system is closely related to the two-spacetime dimensional relativistic "hydrogen atom". Examining the $\mu \neq 0$ limit of the classical rest frame solution (3.13), we find

$$x(\tau,\sigma) = \pm (1 - \frac{2\sigma}{\pi}) \{ |\tau - \tau_0| - \frac{\pi}{2} \},$$
 (3.17)

while Eq. (3.15) for the momentum continues to be valid. The resulting motion is pictured in Fig. 3.3. This motion will be shown in Section 4 to be identical to the simplest longitudinal string mode proposed by Patrascioiu [4].

C. Action-Angle Variables and Bohr-Sommerfeld Quantization

We now exhibit the center-of-mass momentum P of the system by changing to the variables

$$P = p_{0} + p_{\pi}$$

$$R = \frac{1}{2} (x_{0} + x_{\pi})$$

$$k = \frac{1}{2} (p_{\pi} - p_{0})$$

$$r = x_{\pi} - x_{0} .$$
(3.18)

Then the Hamiltonian (3.14) becomes

$$H = \left| \frac{1}{2} P - k \right| + \left| \frac{1}{2} P + k \right| + \gamma |r| .$$
 (3.19)

Hamilton's equations now tell us that

$$\dot{\mathbf{r}} = \frac{\partial H}{\partial k} = \varepsilon \left(k - \frac{1}{2} P \right) + \varepsilon \left(k + \frac{1}{2} P \right)$$

$$\dot{\mathbf{k}} = -\frac{\partial H}{\partial \mathbf{r}} = -\gamma \varepsilon \left(\mathbf{r} \right) , \qquad (3.20)$$

where $\varepsilon(z) = (algebraic sign of z) = z/|z|$. When we plot the motion in τ for arbitrary initial energy H = E and momentum P, Eq. (3.19) generates the closed phase space trajectory of Fig. 3.4. In the rest frame the total momentum P vanishes, and the vertical lines at constant r in Fig. 3.4 disappear.

We now define the action variable

$$J = \oint k dr \tag{3.21}$$

where the integral is around the path of Fig. 3.4 and k(E) is the solution of

$$E = \left| \frac{1}{2} P - k \right| + \left| \frac{1}{2} P + k \right| + \gamma |\mathbf{r}| .$$
 (3.22)

The result is

$$J = 4 \int_{0}^{\gamma^{-1}(E-P)} dr \left[\frac{1}{2}(E-\gamma r)\right] = \frac{1}{\gamma}(E^{2}-P^{2}) . \qquad (3.23)$$

The invariant mass-squared is thus

$$M^2 \equiv E^2 - P^2 = \gamma J \tag{3.24}$$

and J is manifestly Lorentz-invariant.

The Bohr-Sommerfeld approximation for the quantum mechanical energy levels of the center-of-mass system is simply

$$M^{2} = \gamma J = 2\pi \hbar \gamma \ (n + \text{const.}), \quad n = 0, 1, 2... \tag{3.25}$$

The mass-squared spectrum of this sector of the longitudinal massless string thus rises linearly with n in this semiclassical approximation.

The exact quantum-mechanical mass spectrum of this system is presumably given by setting P = 0 and examining the integral equation

$$(2|k_{op}| + \gamma|\mathbf{r}|)\psi(\mathbf{r},t) = M\psi(\mathbf{r},t) . \qquad (3.26)$$

Here we may represent $|k_{op}|$ as

$$\left|k_{op}\right| \psi(\mathbf{r},t) = \int_{-\infty}^{\infty} d\mathbf{r}' \ G(\mathbf{r},\mathbf{r}') \ \psi(\mathbf{r}',t)$$
(3.27)

where

$$G(\mathbf{r},\mathbf{r'}) = \lim_{\varepsilon \to 0} \frac{1}{\pi} \int_{0}^{\infty} d\mathbf{k} \operatorname{cosk}(\mathbf{r}-\mathbf{r'}) e^{-\varepsilon \mathbf{k}}$$

= $P\left\{\frac{1}{\pi(\mathbf{r}-\mathbf{r'})^{2}}\right\}$ (3.28)

and P denotes the principal value.

D. Lightlike Gauge

Throughout the preceding treatment, we have employed the timelike gauge $x^0 \approx \tau$. Since the usual GGRT analysis is conducted in the lightlike gauge, one might ask what happens when we use a lightlike gauge to study longitudinal modes. Defining $x^{\pm} = (x^0 \pm x)/\sqrt{2}$, let us begin by choosing the gauge

$$x^{\dagger}(\tau,\sigma) \approx \tau.$$
 (3.29)

In the x_{σ}^{-} -uniform gauge, the action (3.1) then yields the Hamiltonian

$$H \equiv P^{-} = \frac{\mu^{2}}{2P_{0}} + \frac{\mu^{2}}{2P_{\pi}} + \gamma |x_{\pi} - x_{0}|$$
(3.30)

where we have written

$$p_{0} = p^{+}(\tau, \sigma = 0) \ge 0 , p_{\pi} = p^{+}(\tau, \sigma = \pi) \ge 0 ,$$

$$x_{0} = x^{-}(\tau, \sigma = 0) , \qquad x_{\pi} = x^{-}(\tau, \sigma = \pi) . \qquad (3.31)$$

Now we define the total (+) momentum

$$P \equiv P^{+} = P_{0} + P_{\pi}$$
 (3.32)

and make the canonical transformation (3.18) with null-plane metric variables replacing the timelike variables. The Hamiltonian then becomes

$$H = \frac{1}{2}\mu^{2}P(\frac{1}{4}P^{2} - k^{2})^{-1} + \gamma |r| . \qquad (3.33)$$

The invariant mass-squared is simply [9]

$$M^{2} = 2PH = \frac{\mu^{2}}{\frac{1}{4} - \kappa^{2}} + 2\gamma |\rho| , \quad |\kappa| \leq \frac{1}{2} , \qquad (3.34)$$

where the transformation

$$\kappa = k/P \qquad \rho = rP \qquad (3.35)$$

has eliminated all reference to P and made Lorentz invariance manifest.

It is clear that the $\mu \neq 0$ limit of the lightlike gauge description of this system is quite pathological, in contrast to the timelike gauge. However, the semiclassical spectrum of M^2 agrees with that found in the timelike gauge. We can see this explicitly by calculating the action-angle variables for $\mu \neq 0$ using the standard techniques. We find the result

$$J = 2P \int_{0}^{E\gamma^{-1}(1-2\mu^{2}/EP)} dr \left[1 - \frac{2\mu^{2}}{P(E-\gamma r)}\right]^{1/2}$$

= $\frac{1}{\gamma} \left(2PE\alpha - 2\mu^{2} \ln \left[\frac{1+\alpha}{1-\alpha}\right]\right)$
= $\frac{1}{\gamma} \left\{M(M^{2} - 4\mu^{2})^{1/2} - 4\mu^{2} \ln \left[\frac{M}{2\mu} + \left(\frac{M^{2}}{4\mu^{2}} - 1\right)^{1/2}\right]\right\}$ (3.36)

where

$$\alpha = \left[1 - 2\mu^2 / EP\right]^{1/2} . \tag{3.37}$$

The variable J is well-behaved as μ^2 \rightarrow 0, giving

$$M^2 = 2PE = \gamma J$$
, (3.38)

so the Bohr-Sommerfeld quantum spectrum is the same as in the timelike case.

4. COMPARISON TO ORTHONORMAL GAUGE RESULTS

The motion shown in Fig. 3.3 strongly resembles a class of solutions to the massless string studied by Patrascioiu [4] in a timelike orthonormal gauge. In this Section, we will review Patrascioiu's orthonormal gauge analysis and show that his solutions can be mapped identically into ours by an appropriate gauge transformation.

A. Orthonormal Gauges

We begin by reviewing the properties of the orthonormal gauge. We recall that the orthonormality conditions

$$x_{\tau}^2 + x_{\sigma}^2 = 0$$

$$\mathbf{x}_{\mathsf{T}} \cdot \mathbf{x}_{\mathsf{T}} = \mathbf{0} \tag{4.1}$$

allow the massless string Euler equations to be written as

$$x_{TT}^{\mu} - x_{GG}^{\mu} = 0 , \qquad (4.2)$$

$$x_{\sigma}^{\mu} = 0, \text{ at } \sigma = 0, \pi.$$
 (4.3)

These equations can then be solved in the form

$$x^{\mu}(\tau,\sigma) = q^{\mu} + \frac{p^{\mu}\tau}{\pi\gamma} + \frac{i}{\pi\gamma} \sum_{n \neq 0} \frac{1}{n} \alpha_{n}^{\mu} \cos n\sigma e^{-in\tau}$$
(4.4)

subject to the constraints

$$L_{0} = \frac{1}{2} \sum_{m \neq 0}^{n} \alpha_{m} \cdot \alpha_{-m} + \frac{1}{2} P^{2} = 0$$

$$L_{n} = \frac{1}{2} \sum_{m \neq 0, n}^{n} \alpha_{m} \cdot \alpha_{n-m} + P \cdot \alpha_{n} = 0.$$
(4.5)

In the orthonormal gauge the canonical momentum is simply

$$\mathscr{C}^{\mu}(\tau,\sigma) = \gamma x_{\tau}^{\mu}(\tau,\sigma) \quad . \tag{4.6}$$

It we take the nonvanishing canonical PB of the Fourier components of x^{μ} to be

$$\{P^{\mu}, q^{\nu}\} = -g^{\mu\nu}$$

$$\{\alpha_{m}^{\mu}, \alpha_{n}^{\nu}\} = -i\pi\gamma \ g^{\mu\nu} \ m\delta_{m, -n}$$
(4.7)

then the equal-t PB of ${\ensuremath{\it P}}^\mu$ and x^ν are

$$\{ \mathcal{C}^{\mu}(\tau,\sigma), x^{\nu}(\tau,\sigma') \} = -g^{\mu\nu} \Delta(\sigma,\sigma') . \qquad (4.8)$$

Here

$$\Delta(\sigma,\sigma') = \sum_{n=-\infty}^{\infty} \left[\delta(\sigma - \sigma' + 2n\pi) + \delta(\sigma + \sigma' + 2n\pi) \right]$$
(4.9)

is just that modification of the periodic delta function necessary for compatibility with the boundary condition (4.3).

B. Simple Longitudinal Motions

Patrascioiu has observed that a general solution to Eqs. (4.1)-(4.3) is

$$\mathbf{x}^{0} = \tau \mathbf{E}/\pi\gamma$$

$$\vec{\mathbf{x}}(\tau,\sigma) = \vec{\mathbf{a}} + \frac{\mathbf{E}}{2\pi\gamma} \left[\vec{\mathbf{f}}(\tau+\sigma) + \vec{\mathbf{f}}(\tau-\sigma) \right]$$
(4.10)

provided

$$\vec{f}'(\tau + 2\pi) = \vec{f}'(\tau)$$

$$[\vec{f}'(\tau + \sigma)]^2 = [\vec{f}'(\tau - \sigma)]^2 = 1$$
(4.11)

where $\vec{f'}(z) = \partial \vec{f} / \partial z$. We note that

$$\vec{\mathcal{P}}(\tau,\sigma) = \vec{\gamma x_{\tau}} = \frac{E}{2\pi} \left[\vec{f}'(\tau+\sigma) + \vec{f}(\tau-\sigma) \right]$$

$$\vec{x}_{\sigma}(\tau,\sigma) = \frac{E}{2\pi\gamma} \left[\vec{f}'(\tau+\sigma) - \vec{f}'(\tau-\sigma) \right].$$
(4.12)

The conserved total momentum may thus be written as

$$\vec{P} = \int_{0}^{\pi} d\sigma \ \vec{C}(\tau,\sigma) = \frac{E}{2\pi} \left[\vec{f}(\tau+\pi) - \vec{f}(\tau-\pi) \right]$$
(4.13)

for arbitrary $\boldsymbol{\tau}.$

Suppose we now choose the simplest rest frame solution

$$a \equiv a^{(1)} = -E/2\gamma$$

$$f(z) \equiv f^{(1)}(z) = |z|_{periodic} = \sum_{n=-\infty}^{\infty} c_n e^{inz}$$

$$a^{(i)} = f^{(i)} = 0, i \neq 1,$$
(4.14)

where

$$c_{n} = \frac{1}{\pi n^{2}} [(-1)^{n} - 1]$$

$$c_{0} = \frac{\pi}{2} .$$
(4.15)

We now show that this motion is precisely that shown in Fig. 3.3 and examined in Section 3 as the massless limit of the string with massive ends. Working from the plot of f(z) in Fig. 4.1, we may construct $\mathscr{P}(\tau,\sigma)$, $x(\tau,\sigma)$, $x_{\sigma}(\tau,\sigma)$ for appropriate values of τ to find the motions presented in Fig. 4.2. We can thus see directly how this solution contrives to have $\mathscr{P} x_{\sigma} = 0$ without forcing either $\mathscr{P} \equiv 0$ or $x_{\sigma} \equiv 0$. If we follow the motion of the end points on Fig. 4.2, we find that the endpoints of our solution, shown in Fig. 3.3, follow the same paths. The motion of the endpoints is physical and cannot be gauged away, so we conclude that the physical meanings of the two pictures are very likely the same.

The motions of the interior points in Figs. 3.3 and 4.2 differ because this motion depends on the σ -gauge chosen. We can complete the identification of the two systems by making a σ -reparametrization which maps the orthonormal gauge into the uniform-x_{σ} gauge. The appropriate transformation is found by requiring

$$\mathbf{x}(\tau,\sigma) = \tilde{\mathbf{x}}(\tau,\tilde{\sigma} \ (\tau,\sigma)) \tag{4.16}$$

and taking the σ -derivative,

$$\mathbf{x}_{\sigma} = \frac{\partial \tilde{\sigma}}{\partial \sigma} \, \tilde{\mathbf{x}}_{\tilde{\sigma}} \quad . \tag{4.17}$$

Now suppose x_{σ} is in the orthonormal gauge (uniform \mathcal{P}^0) but \tilde{x}_{σ} is <u>independent</u> of $\tilde{\sigma}$ (uniform- x_{σ} gauge). Then we can do the integral immediately, finding

$$\tilde{\sigma}(\tau,\sigma) = F(\tau) \int_{0}^{\sigma} d\rho x_{\rho}(\tau,\rho) , \qquad (4.18)$$

where $F(\tau) = [\tilde{x}_{\tilde{\sigma}}]^{-1}$. The requirement that the endpoints stay fixed,

$$\tilde{\sigma}(\tau, 0) = 0$$

 $\tilde{\sigma}(\tau, \pi) = \pi$, (4.19)

then fixes the function $F(\tau)$ uniquely, giving

$$\tilde{\sigma}(\tau,\sigma) = \pi \frac{\int_0^0 d\rho \ x_\rho(\tau,\rho)}{\int_0^{\pi} d\rho \ x_\rho(\tau,\rho)} .$$
(4.20)

Taking $x_{\sigma}^{}(\tau,\sigma)$ from Fig. 4.2, we plot $\tilde{\sigma}(\tau,\sigma)$ in Fig. 4.3.

The effect of this transformation is clearly to map the finite-width plateaus in \mathscr{C} and the zero-regions of x_{σ} in Fig. 4.2 into the endpoints $\tilde{\sigma} = 0$, $\tilde{\sigma} = \pi$, of the new system. For example, when $0 < \tau < \frac{\pi}{2}$, $\tilde{\sigma}$ vanishes for the entire interval $0 < \sigma < \tau$, where $x_{\sigma} = 0$ and $\mathscr{C} \neq 0$.

If we define the momentum accumulating near an endpoint as

$$p_0(\tau) = \int_0^{\pi/2} d\sigma \ \ell(\sigma)$$
 (4.21)

in the orthonormal gauge, we see from Fig. 4.2 that p_0 increases linearly with τ for $0 < \tau < \frac{\pi}{2}$. In the x_{σ} -uniform gauge, $\tilde{\ell}(\tilde{\sigma})$ vanishes except at $\tilde{\sigma} = 0$, π , so $\tilde{\ell}(\tilde{\sigma})$ necessarily becomes proportional to a delta function at the endpoints. $p_0(\tau)$ is thus precisely identifiable with our endpoint momentum (3.13), which does indeed change linearly with time.

In Fig. 4.4, we plot the new functions $\tilde{\ell}(\tilde{\sigma})$, $\tilde{x}(\tilde{\sigma})$, $\tilde{x}_{\tilde{\sigma}}(\tilde{\sigma})$ versus $\tilde{\sigma}$ and τ . Comparison to the properties of our solution (3.15) for the $\mu \rightarrow 0$ limit of the massive-end string shows that the motions are identical. Having demonstrated the equivalence of the orthonormal gauge and x_{σ} -uniform gauge treatment, we remark that the latter approach has a more obvious physical interpretation; the system consists solely of massless point particles interacting via linear potentials.

C. Longitudinal Fourier Components

Because of Eq. (4.11), the longitudinal solutions are not linearly superimposable and behave like <u>kink solutions</u> of a nonlinear field theory. In fact, if we identify the coefficients in Eq. (4.4) with those in Eqs. (4.15) and (4.16), we find

$$q = P = 0$$

$$i\alpha_n = Enc_n = \frac{E}{\pi n} [(-1)^n - 1],$$
 (4.22)

so that the extraction of quantizable amplitudes from the Fourier coefficients of the classical solution is nontrivial.

It is instructive to note that the solution (4.22) satisfies the constraints (4.5) in an unusual nonlinear manner. For L_0 , we find

$$\left(\frac{2\pi^{2}}{E^{2}}\right)L_{0} = \frac{\pi^{2}}{E^{2}}\sum_{m\neq 0}^{\infty} \alpha_{m} \alpha_{-m} - \pi^{2}$$
$$= 4\sum_{m=1}^{\infty} \frac{1}{m^{2}} \left[1 - (-1)^{m}\right] - \pi^{2}$$
$$= 6 \zeta(2) - \pi^{2} \equiv 0 , \qquad (4.23)$$

where $\zeta(n)$ is the Riemann zeta function. Examining L_n , $n \neq 0$, we see that

$$\left(\frac{2\pi^{2}}{E^{2}}\right)L_{n} = \frac{\pi^{2}}{E^{2}}\sum_{m\neq 0,n} \alpha_{m} \alpha_{n-m} = \left[1 + (-1)^{n}\right]\sum_{m\neq 0,n} \frac{(-1)^{m}-1}{m(n-m)}, \qquad (4.24)$$

which vanishes trivially for odd n. For even n, one may use

$$\frac{1}{m(n-m)} = \frac{1}{n} \left(\frac{1}{m} + \frac{1}{n-m} \right)$$
(4.25)

and the fact that $(-1)^{m+n} = (-1)^m$ to prove that there is a cancellation of terms in the sum which makes $L_n \equiv 0$.

D. N Fold Longitudinal Solutions

We saw in the beginning of this section that the simplest longitudinal solution (4.14) of the constraint Eq. (4.11) corresponded to our solution (3.15) with only the endpoints of the string moving at the speed of light. More general solutions for f(z) give longitudinal motions with arbitrary numbers of interior points moving at the speed of light.

The only restrictions on f'(z) are that it be periodic and have unit magnitude. If we confine ourselves to the rest frame $\vec{P} = 0$, then Eq. (4.13) implies that f(z) itself has period 2π . Wherever there is a discontinuity in the slope of f(z), there is a point moving at the speed of light; we will hereafter refer to these points as "folds".

In Fig. 4.5, we depict f(z) for a mode which has N folds; for simplicity, we have chosen f(z) = f(-z) so that the initial σ -positions

 $c_n, n = 1, ..., N$

of each discontinuity are identifiable with the initial position of each fold. For this solution, the initial momentum density $\mathscr{C}(0,\sigma)$ vanishes for all σ . More general choices for f(z), with $f(z) \neq f(-z)$, have 2N free parameters giving nontrivial initial momentum configurations.

5. HAMILTONIAN APPROACH TO N FOLD MODES

In the previous two sections, we examined the Hamiltonian for longitudinal oscillations of the string with endpoints only moving at the speed of light and related that motion to a particular solution of the orthonormal gauge string equations. Motivated by the existence of N fold solutions in the orthonormal gauge, we now turn our attention to the Hamiltonian description of longitudinal string oscillations with N interior points moving at the velocity of light.

For pure longitudinal modes, the Hamiltonian (2.8) can be written

$$H = \int_{0}^{\pi} d\sigma \left\{ \left| p(\tau, \sigma) \right| + \gamma \left| x_{\sigma}(\tau, \sigma) \right| \right\}$$
(5.1)

where a σ -gauge remains to be chosen. We argued in Sec. 2 that $p(\tau,\sigma)$ remains nonzero in the massless limit only in those σ -regions where the string moves at the velocity of light. By making a suitable choice of σ -gauge, we may write H in the form

$$H = \sum_{n=0}^{N+1} |p_n| + \gamma \sum_{n=0}^{N} |x_{n+1} - x_n|$$
(5.2)

where $x_0 = x(\sigma = 0)$, $x_{N+1} = x_{\pi} = x(\sigma = \pi)$ and similarly for p_0 and p_{N+1} . Here $x_n(\tau)$ labels the n-th point moving at the velocity of light along the string, and $p_n(\tau)$ its conjugate momentum.

In this section, we will first analyze the motion and action angle variables for the one-fold mode. Then we give the form of the general N fold Hamiltonian expressed in terms of action variables alone.

A. One Fold Mode

The string with one interior point moving at the velocity of light is described by the Hamiltonian

$$H = |p_0| + |p_1| + |p_{\pi}| + \gamma |x_1 - x_0| + \gamma |x_{\pi} - x_1|$$
(5.3)

It is convenient to rewrite the above in terms of a set of relative coordinates and momenta. Defining the new coordinates and momenta

$$r_{1} = x_{1} - x_{0}$$

$$r_{2} = x_{\pi} - x_{1}$$

$$R = \frac{1}{3}(x_{0} + x_{1} + x_{\pi})$$
(5.4)

and

$$k_{1} = \frac{1}{3} (-2p_{0} + p_{1} + p_{\pi})$$

$$k_{2} = \frac{1}{3} (-p_{0} - p_{1} + 2p_{\pi})$$

$$P = p_{0} + p_{1} + p_{\pi} , \qquad (5.5)$$

we find that the Hamiltonian takes the form

$$H = \left|\frac{1}{3}P - k_{1}\right| + \left|\frac{1}{3}P + k_{1} - k_{2}\right| + \left|\frac{1}{3}P + k_{2}\right| + \gamma |r_{1}| + \gamma |r_{2}| .$$
(5.6)

Hence in the P = 0 frame the Hamiltonian reduces to

$$H(P = 0) = M = |k_1| + |k_2 - k_1| + |k_2| + \gamma |r_1| + \gamma |r_2| .$$
(5.7)

Yet another set of interesting variables is the choice

$$k_{\pm} = k_1 \pm k_2$$

 $r_{\pm} = \frac{1}{2} (r_1 \pm r_2)^2$. (5.8)

In this case

$$M = \frac{1}{2} |k_{+} + k_{-}| + \frac{1}{2} |k_{+} - k_{-}| + |k_{-}| + \gamma |x_{+} + x_{-}| + \gamma |x_{+} - x_{-}| .$$
(5.9)

To proceed with the Bohr-Sommerfeld quantization, we must analyze the most general periodic motions in phase space. We begin by choosing the initial conditions

$$k_{1} = k_{2} = 0$$

$$r_{1} = -\frac{mc}{\pi\gamma}$$

$$r_{2} = (\pi - c) \frac{m}{\pi\gamma}$$
(5.10)

where m is the rest-mass and c corresponds to another constant of the motion. This particular choice of initial configuration is suggested by the type of one-fold orthonormal gauge solution discussed in Section 4.D, for which all momenta initially vanish. Hamilton's equations of motion determine the subsequent periodic motion of the system, which we plot in phase space in Fig. 5.1. If we calculate the action-angle variables in $(x_1,k_1;x_2,k_2)$ -space, we find that J_1 and J_2 are independent of c:

$$J_{1} = \oint k_{1} dr_{1} = m^{2}/2\gamma$$

$$J_{2} = \oint k_{2} dr_{2} = m^{2}/2\gamma .$$
(5.11)

This is expected because of the exact symmetry of the Hamiltonian (5.7) under $1 \Leftrightarrow 2$ interchange.

A typical phase space diagram in (k_{\pm}, r_{\pm}) -space is given in Fig. 5.2. The exhibited τ -sequence of numbers corresponds to the initial conditions (5.10). The action variables J_{\pm} and J_{\pm} computed by integrating over these phase-space orbits depend explicitly on c; moreover, the regions $0 < c < \frac{\pi}{3}$, $\frac{\pi}{3} < c < \frac{2\pi}{3}$, $\frac{2\pi}{3} < c < \pi$, must be treated separately. For arbitrary c, one finds the result

$$J_{\pm} = \oint k_{\pm} dr_{\pm} = \frac{m^2}{\gamma} \quad [\frac{1}{2} \pm g(c)] , \qquad (5.12)$$

where

$$g(c) = \frac{7}{2} \frac{c}{\pi} \left(\frac{c}{\pi} - 1\right) + \frac{3}{2} \left\{ \left|\frac{c}{\pi} - \frac{2}{3}\right| + \left|\frac{c}{\pi} - \frac{1}{3}\right| \right\} - \frac{9}{2} \left|\frac{c}{\pi} - \frac{2}{3}\right| \left|\frac{c}{\pi} - \frac{1}{3}\right| .$$
(5.13)

We plot J_{\pm} vs. c in Fig. 5.3. Now we may eliminate the constant of motion c to determine M^2 uniquely in terms of J_{\pm} , with the result

24

$$M^2/\gamma = m^2/\gamma = J_+ + J_- = J_+ + J_2$$
 (5.14)

 J_{\perp} and J_{\perp} are clearly interpretable as normal modes. We see that when

$$c = \pi/2$$
 (5.15a)

then

$$J_{+} = 0$$
, $J_{-} = m^2 / \gamma$. (5.15b)

The motion consists of the center point oscillating in opposition to the two ends, which remain coincident throughout the motion. This is a completely folded string. When

$$c = 0 \text{ or } c = \pi$$
 (5.16a)

then

$$J_{+} = m^{2}/\gamma$$
, $J_{-} = 0$. (5.16b)

In this case, the endpoints move exactly as in the no-fold problem of Section 3; the "center" point attaches itself alternately to one endpoint or the other, always remaining on the right side of the string (or always on the left side). In this case there are no folds in the string. We plot these normal mode motions in Fig. 5.4.

B. N Fold Modes

We have shown in our explicit analysis of the no-fold mode that

$$H(P = 0) = M = [\gamma J]^{1/2}$$
(5.17)

while for the one-fold mode

$$H(P = 0) = M = [\gamma(J_{+} + J_{-})]^{1/2}$$

= [\gamma(J_{1} + J_{2})]^{1/2}. (5.18)

 J_+ was identifiable with a normal mode indistinguishable from the no-fold motion described by J; J corresponded to a double cycle of the completely symmetric

folded mode of the one-fold motion, as can be seen from Fig. 5.4. It is clearly of interest to know whether similar expressions hold for arbitrary P and for an arbitrary number of folds. We will now show in fact that for N folds, the Hamiltonian can be written in terms of P^2 and a sum of appropriate action variables alone.

We begin by defining the local action variable

$$J(\sigma) = \oint \mathcal{O}(\tau, \sigma) \, dx(\tau, \sigma)$$

one
period

$$= \int_{-\pi}^{\pi} d\tau \, \mathcal{O}(\tau, \sigma) \, x_{\tau}(\tau, \sigma) \, . \qquad (5.19)$$

In the timelike orthonormal gauge of Section 4, $\mathscr{C} = \gamma x_{\tau}$ and $x(\tau, \sigma)$ is given by Eq. (4.10). Then we may consider the direct sum of all the action variables, which we write as

$$\int_{0}^{\pi} d\sigma J(\sigma) = \frac{1}{2\gamma} \int_{-\pi}^{\pi} d\tau \int_{-\pi}^{\pi} d\sigma \, \theta^{2}(\tau, \sigma) \\ = \frac{E^{2}}{4\pi^{2}\gamma} \int_{-\pi}^{\pi} d\tau \int_{-\pi}^{\pi} d\sigma \left[1 + f'(\tau + \sigma)f'(\tau - \sigma)\right] .$$
(5.20)

In the last line, we have used Eqs. (4.11) and (4.12). Changing variables from (τ,σ) to $(\tau+\sigma,\tau-\sigma)$ and using Eq. (4.13), we find

$$\int_{0}^{\pi} d\sigma J(\sigma) = \frac{E^2 - P^2}{\gamma} \equiv \frac{M^2}{\gamma} . \qquad (5.21)$$

This argument extends trivially to D dimensions, with the result

$$M^{2} = E^{2} - \vec{P}^{2} = \gamma \sum_{i=1}^{D-1} \int_{0}^{\pi} d\sigma J_{i}(\sigma) .$$
 (5.22)

In two spacetime dimensions, we may revert to the $x_{\rm g}^{}$ -uniform gauge and write Eq. (5.21) as

$$M^{2} = \gamma \sum_{n=1}^{\infty} J_{n}$$
(5.23)

where

$$J_{n} = \oint_{\substack{\text{complete}\\\text{cycle}}} k_{n} \frac{dr_{n}}{n}$$
(5.24)

and (P,Q; k_n, r_n) comprise a complete set of canonically conjugate pairs of variables.

We saw in the one fold problem that a particular choice of (k_n, r_n) makes one of the J_n 's (J_+) a pure no-fold variable, the other (J_-) a pure single-fold variable. In addition, the no-fold variables (k_+, r_+) went through one cycle while the single-fold variables (k_-, r_-) went through two complete cycles, as seen in Figs. 5.2(a) and (b). It is easy to convince oneself that the completely symmetric N fold string, with orthonormal-gauge initial σ -conditions

$$c_n = \frac{n\pi}{N+1}$$
, $n = 1, 2, ..., N$, (5.25)

undergoes N + 1 identical cycles while the no-fold variables finish one complete cycle. The appropriate Bohr-Sommerfeld quantization applies to the variables

$$\widetilde{J}_{n} = \frac{1}{n} J_{n} = \oint_{\substack{\text{single} \\ \text{cycle}}} k_{n} dr_{n}, \qquad (5.26)$$

where the integral in Eq. (5.26) is over one cycle in the (k_n, r_n) subspace instead of one cycle in the full phase space. We thus take

$$M^{2} = \gamma \sum_{n=1}^{\infty} n \widetilde{J}_{n}$$
(5.27)

where \tilde{J}_1 represents a pure no-fold string, \tilde{J}_2 is associated with a string having one fold in the middle, \tilde{J}_3 with two folds, etc. The Bohr-Sommerfeld quantization rule is

$$\tilde{J}_{n} = 2\pi\hbar[\ell_{n} + \text{const}]$$
, $\ell_{n} = 0, 1, 2, ...$ (5.28)

The Hamiltonian describing the motion for any number of folds, given appropriate initial conditions, is then

$$H = [P^{2} + \gamma \sum_{n=1}^{\infty} n \tilde{J}_{n}]^{1/2} .$$
 (5.29)

Defining $\theta_n(\tau)$ to be the canonical conjugate to $\tilde{J}_n(\tau)$, we find from Hamilton's equations

$$\omega_{n} \equiv \stackrel{\bullet}{\theta}_{n} = \frac{\partial H}{\partial J_{n}} = \frac{n\gamma}{H} = \text{constant} . \qquad (5.30)$$

Thus

$$\theta_{n}(\tau) = \theta_{n}(0) + \frac{n\gamma\tau}{H} , \qquad (5.31)$$

where $\theta_n(0)$ is a constant of motion. Since H is independent of $\theta_n(\tau)$, each of the \tilde{J}_n 's is also a constant of motion. Thus we have been able to express the N fold string problem as a completely integrable Hamiltonian system. Equation (5.4) for the two-dimensional string parallels exactly the Faddeev-Takhtajan Hamiltonian [6] for the classical sine-Gordon equation; both give the complete solution to their respective nonlinear classical systems.

6. TOWARDS CANONICAL QUANTIZATION

Although we have completely solved the classical longitudinal string, the quantum theory does not follow trivially from the diagonalized Hamiltonian (5.29). Even the no-fold Schrödinger equation (3.24) could not be solved for the exact quantum spectrum. Motivated by the desire to have local commutation relations among $\mathscr{C}^{\mu}(\tau,\sigma)$ and $x^{\nu}(\tau,\sigma)$, we devote this section to a timelike orthonormal gauge treatment of the string constraints. The constrained oscillators in two spacetime dimensions should then determine the canonical properties of the longitudinal string motions.

The first step is to separate the overall center-of-mass variables of the system from those describing the intrinsic motion. In the lightlike orthonormal gauge,

$$x^{+} = P^{+} \tau / \pi \gamma$$

 $\theta^{-} = P^{+} / \pi$, (6.1)

the standard overall coordinate variables

$$Q^{\mu} = \frac{M^{\mu+}}{P^{+}}$$
(6.2)

happily commuted with one another. Unfortunately, as we saw at the end of Section 3, treating longitudinal oscillations in the lightlike gauge is exceedingly difficult. In the timelike gauge

$$x^{0} = P^{0}\tau/\pi\gamma$$

$$\theta^{0} = P^{0}/\pi$$
(6.3)

the coordinates are

$$Q^{\mu} = \frac{M^{\mu 0}}{P^{0}} \quad . \tag{6.4}$$

These \vec{Q} 's do not commute (except in two dimensions), and hence are unsuitable center-of-mass coordinates; nevertheless, this system can be mapped directly into the lightlike-gauge variables using DDF variables [10], with the result that the transverse sectors of the string in the two gauges are equivalent [11].

Here we investigate a timelike gauge condition used by Rohrlich [7] which produces appropriate commuting (Newton-Wigner) coordinates for the string's center-of-mass. By going to two spacetime dimensions, we are able to clearly exhibit the complicated nature of the longitudinal modes appearing in the classical system.

A. Newton-Wigner Coordinates

In order to illuminate the canonical properties of the orthonormal gauge Fourier coefficients (4.4), we wish to separate out the Newton-Wigner coordinates Q^{μ} of the center-of-mass. A general expression for Q^{μ} is

$$Q^{\mu} = \frac{M^{\mu 0}}{P^{0} + P} - \frac{M^{\mu \nu} P_{\nu}}{P(P^{0} + P)} + \frac{P^{\mu} M^{0 \nu} P_{\nu}}{PP^{0}(P^{0} + P)}$$
(6.5)

where

$$P = [-P^{2}]^{1/2} = [(P^{0})^{2} - \vec{P}^{2}]^{1/2}$$
(6.6)

and $Q^0 \equiv 0$ defines Q^{μ} as a timelike-gauge variable. Using the canonical Poisson bracket algebra of the Poincaré group,

$$\{M^{\mu\nu}, P^{\alpha}\} = g^{\mu\alpha} P^{\nu} - g^{\nu\alpha} P^{\mu}$$
(6.7)

$$\{M^{\mu\nu}, M^{\alpha\beta}\} = g^{\mu\alpha}M^{\nu\beta} - g^{\nu\alpha}M^{\mu\beta} + g^{\mu\beta}M^{\alpha\nu} - g^{\nu\beta}M^{\alpha\mu} , \qquad (6.8)$$

we find

$$\{Q^{\mu}, Q^{\nu}\} = 0$$

$$\{Q^{\mu}, P^{\alpha}\} = g^{\mu\alpha} - g^{0\alpha} \frac{P^{\mu}}{P^{0}} .$$
 (6.9)

 P^0 is seen to act as the Hamiltonian, since

$$\{\vec{Q}, P^0\} = \vec{P}/P^0$$
 (6.10)

is the velocity of the center-of-mass. If $M^{\mu\nu}$ has the form

$$M^{\mu\nu} = Q^{\mu}P^{\nu} - Q^{\nu}P^{\mu} + S^{\mu\nu} , \qquad (6.11)$$

then the definition (6.5) of Q^{μ} is an identity if

$$S^{\mu\nu}P_{\nu} - P S^{\mu0} = 0 . (6.12)$$

Taking $J^{i} = \frac{1}{2} \varepsilon^{ijk} S^{jk}$ as the independent components of $S^{\mu\nu}$, we see that the constraint (6.12) implies the usual Wigner representation of the classical boost:

$$M^{0i} = (\vec{N})_{i} = \left(-P^{0\dot{Q}} + \frac{\vec{J} \times \vec{P}}{P^{0} + P}\right)_{i} , \qquad (6.13)$$

For massless states, this formalism is modified to replace \vec{Q} and \vec{J} by appropriate massless-particle variables [12].

In any orthonormal gauge the spin matrix $S^{\mu\nu}$ for the string model takes the form

$$S^{\mu\nu} = \frac{i}{\pi\gamma} \sum_{n\neq 0} \frac{1}{n} \alpha_n^{\mu} \alpha_{-n}^{\nu} .$$
(6.14)

Thus Eq. (6.12) implies that the constraints on the Fourier components (4.4) necessary to give Newton-Wigner \dot{q} 's are

$$g_n \equiv \alpha_n^0 (P^0 + P) - \dot{\alpha}_n \cdot \vec{P} = 0 .$$
 (6.15a)

along with

$$q^0 = 0$$
 . (6.15b)

The analogs of the gauge conditions (6.1) and (6.3) on the canonical fields are

$$x^{0}(\tau,\sigma)(P^{0}+P) - \vec{x}(\tau,\sigma)\cdot\vec{P} - \frac{M^{0\mu}P}{P^{0}} = \frac{\tau}{\pi\gamma}P(P^{0}+P)$$

$$\theta^{0}(\tau,\sigma)(P^{0}+P) - \vec{P}\cdot\vec{\theta}(\tau,\sigma) = \frac{1}{\pi}P(P^{0}+P) . \qquad (6.16)$$

B. Dirac Brackets

The canonical Poisson brackets (4.7) are not compatible with the constraints (4.5) and (6.15). One may, however, define modified brackets — the Dirac brackets [13] — which are manifestly consistent with the constraints. In the Appendix we define and calculate the Dirac brackets for the Fourier components of the string in the Newton-Wigner gauge; we also present an alternative technique, using gauge-invariant variables, which yields equivalent results.

The computations in the Appendix give the following formulas for the Dirac brackets, distinguished hereafter by an asterisk:

$$\{q^{\mu}, q^{\nu}\}^{*} = 0$$

$$\{q^{\mu}, p^{\alpha}\}^{*} = g^{\mu\alpha} - g^{0\alpha} \frac{p^{\mu}}{p^{0}}$$

$$\{\alpha_{m}^{i}, \alpha_{n}^{j}\}^{*} = -i\pi\gamma \, \delta^{ij} m \delta_{m, -n} + m (P^{j} \alpha_{n+m}^{i} T_{n} - P^{i} \alpha_{n+m}^{j} T_{m}) (\pi\gamma)^{2}$$

$$+ m\alpha_{n}^{j} \{P^{i} (U_{m} - V_{m}) - \sum_{\substack{k \neq 0, m}} V_{k} \alpha_{m-k}^{i}\} (\pi\gamma)^{2}$$

$$- m\alpha_{m}^{i} \{P^{j} (U_{n} - V_{n}) - \sum_{\substack{k \neq 0, m}} V_{k} \alpha_{n-k}^{j}\} (\pi\gamma)^{2}$$

$$- 2mn(\pi\gamma)^{2} \sum_{\substack{k \neq 0, m-n}} T_{k} \alpha_{m-k}^{i} \alpha_{n+k}^{j}$$

$$\{q^{i}, \alpha_{n}^{j}\}^{*} = -\frac{n\alpha_{n}^{j}}{M^{2}(H+M)} \sum_{\substack{k \neq 0}} \frac{1}{k} \alpha_{k}^{i} \alpha_{-k}^{0}$$

$$+ \frac{n}{M(H+M)} \sum_{\substack{k \neq 0, -n}} \frac{1}{k} \alpha_{-k}^{0} \alpha_{n+k}^{j} .$$

$$(6.19)$$

Here T_n , U_n , V_n are defined in the Appendix. Since the constraints (4.5) are now strongly valid, we are free to define

$$M^{2} = -P^{2} = \sum_{m \neq 0} \alpha_{m} \cdot \alpha_{-m}$$
(6.20)

even though α_n^0 contains implicit M^2 dependence; similarly, we may now take $H \equiv P^0 = \left[\vec{P}^2 + M^2\right]^{1/2}$ (6.21)

as a dependent variable. Equations (6.17) indicate that canonical quantization is straightforward for \vec{q} and \vec{P} . Unfortunately the Dirac brackets (6.18) and (6.19) are so complicated that canonical quantization is decidedly nontrivial.

C. Two-Dimensional System

In two spacetime dimensions, the only simple solution of the constraint equations (4.5) is

$$\alpha_{n}^{\mu} = 0$$

 $P^{\mu}P_{\mu} = 0$
(6.22)

so x^{μ} has only translational degrees of freedom,

$$x^{0} = \frac{|P|\tau}{\pi\gamma}$$

$$x^{1} = q + \frac{P\tau}{\pi\gamma} \quad . \tag{6.23}$$

The Newton-Wigner coordinate (6.5) becomes simply

$$Q = \frac{M^{10}}{P^0}$$
 (6.24)

However, the Dirac brackets (6.18) evaluated in two dimensions indicate that a highly nontrivial Hamiltonian system still exists if $\alpha_n^{\mu} = 0$ is excluded. These brackets presumably give all the available information about the classical Hamiltonian dynamics of the longitudinal modes. We exhibit the brackets of $\alpha_n^{\ 1} \equiv \alpha_n$ in the rest frame P = 0, H = M, for simplicity:

$$\{P,q\}^{*} = -1$$

$$\{q,H\}^{*} = P/H = 0$$

$$\{H,\alpha_{n}\}^{*} = \frac{in\alpha_{n}}{M}$$

$$\{q,\alpha_{n}\}^{*} = -\frac{n}{2M^{2}} \sum_{\substack{\ell \neq 0 \ n}} \frac{1}{k} \alpha_{\substack{\ell}} \alpha_{n-k}$$

$$\{P,\alpha_{n}\}^{*} = 0$$

$$\{\alpha_{m},\alpha_{n}\}^{*} = -i(\pi\gamma m\delta_{m,-n} + \frac{mn}{M^{2}} \sum_{\substack{\ell \neq 0 \ m,-n}} \frac{1}{k} \alpha_{m-k} \alpha_{n+k}) .$$
(6.25)

Even in two dimensions, we see that canonical quantization of the longitudinal modes will be difficult.

7. CONCLUSION

We have investigated the massless limit of a model for the massive Nambu string and have found that we recover longitudinal modes of oscillation excluded from the GGRT treatment. These oscillations are kink-like, not linearly superimposable, and are identifiable with those proposed by Patrasioiu.

The model was examined in detail in two spacetime dimensions and the Hamiltonian expressed in terms of normal-mode action variables alone. The resulting theory was a completely integrable classical Hamiltonian system. The Bohr-Sommerfeld approximation to the quantum mechanics gave a linear, integer-spaced, mass-squared spectrum.

Finally, we separated the simultaneously measurable Newton-Wigner coordinates of the string center-of-mass and examined the Dirac brackets of the oscillators occurring in the fully constrained, orthonormal-gauge system. The brackets exhibit complex structure for the longitudinal mode oscillators even when we go to the rest frame in two spacetime dimensions. Our attempts to find a canonical quantization procedure have so far failed. We are thus still far from our goal of showing that the longitudinal modes permit the string to be quantized for dimensions other than twenty-six.

ACKNOWLEDGMENTS

Three of us (W.A.B., A.J.H., R.D.P.) are grateful to the Aspen Center for Physics for its hospitality during a portion of this work. We thank J. Goldstone and C. Thorne for sharing with us some of their insights on this problem.

APPENDIX

1. Dirac Brackets

The canonical Poisson brackets of a constrained system can be replaced by Dirac brackets [13] which are compatible with all constraints. Let us denote $\langle by \{\phi_{\alpha}\}, \phi_{\alpha} \approx 0$, the entire collection of constraints, and write

$$C_{\alpha\beta} = \{\phi_{\alpha}, \phi_{\beta}\} \quad . \tag{A.1}$$

The Dirac bracket of two canonical variables A and B is then defined as

$$\{A,B\}^* = \{A,B\} - \{A,\phi_\alpha\} C_{\alpha\beta}^{-1} \{\phi_\beta,B\} .$$
(A.2)

It is clear that the Dirac bracket of a canonical variable with any one of the constraints vanishes identically.

For the case at hand, the set of ϕ_{α} 's is given by Eqs. (4.5) and (6.15),

$$\{\phi_{\alpha}\} = \{L_{n \geq 0}, L_{0}, L_{n \leq 0}; g_{n \geq 0}, q^{0}, g_{n \leq 0}\} \approx 0$$
 (A.3)

We exhibit below the matrix $C_{\alpha\beta}$ defined by (A.1) with ϕ_{α} given by Eq. (A.3), and its inverse $C_{\alpha\beta}^{-1}$. Defining

$$H \equiv p^{0}$$

$$M \equiv [-p^{2}]^{1/2}$$

$$R = \frac{H+M}{M}$$

$$T_{n} = \frac{i}{n\pi\gamma (H+M)M}$$

$$U_{n} = -\frac{\alpha_{n}^{0}}{H} T_{n}$$

$$V_{n} = \frac{H-M}{HM} \alpha_{n}^{0} T_{n}$$

(A.4)

we find

Using $C_{\alpha\beta}^{-1}$ and Eq. (A.2), we obtain for the Dirac brackets the formula

$$\{A,B\}^{*} = \{A,B\} - \{A,L_{n}\} V_{-n}\{L_{0},B\} + \{A,L_{0}\} V_{n}\{L_{-n},B\} - \{A,L_{-n}\} 2T_{n}\{L_{n},B\}$$
$$- \{A,L_{-n}\} T_{n}\{g_{n},B\} - \{A,L_{0}\} \frac{1}{H} \{q^{0},B\} + \{A,L_{0}\} U_{n}\{g_{-n},B\}$$
$$+ \{A,q^{0}\} \frac{1}{H} \{L_{0},B\} - \{A,g_{-n}\} U_{n}\{L_{0},B\} - \{A,g_{-n}\} T_{n}\{L_{n},B\} .$$
(A.7)

Here the implicit sums over n exclude n = 0. Equations (6.17), (6.18) and (6.19) in the main text follow directly from Eq. (A.7).

2. Gauge-Invariant Oscillators

A simple extension of the concepts introduced in Ref.11 allow us to construct gauge-invariant classical oscillators obeying the Newton-Wigner gauge constraints (6.15). The Newton-Wigner coordinates Q^{μ} given by Eq. (6.5) are already gauge-invariant, and satisfy $Q^{0} = 0$. The analogs of the DDF variables [10] are given by

$$A_{n}^{\mu} = \frac{\gamma}{2} \int_{-\pi}^{\pi} d\theta \, \frac{dx^{\mu}(\theta)}{d\theta} \, \exp \, in\pi\gamma \left\{ \frac{x^{0}(\theta) \, (H+M) - \vec{x}(\theta) \cdot \vec{P}}{M(H+M)} \right\}$$
(A.8)

where in any orthonormal gauge $X^{\mu}(\theta)$ may be expressed in the form

$$X^{\mu}(\theta) = q^{\mu} + \frac{p^{\mu}\theta}{\pi\gamma} + \frac{i}{\pi\gamma} \sum_{n \neq 0} \frac{1}{n} \alpha_{n}^{\mu} e^{-in\theta} .$$
 (A.9)

This expression and Eq. (4.7) define the canonical Poisson brackets of $\chi^{\mu}(\theta)$, and hence the canonical properties of A_n^{μ} are determined. Note also that

$$\frac{\gamma}{2} \left[X^{\mu}(\pi) - X^{\mu}(-\pi) \right] = P^{\mu} . \tag{A.10}$$

All of the arguments in Ref. 11 may now be imitated to analyze the properties of A_n^{μ} . The first result is that A_n^{μ} is invariant under the action of the Virasoro constraints (4.5), which generate the gauge transformations. In addition,

 $A_n^{\ \ \mu}$ itself obeys the Newton-Wigner constraints (6.15a),

$$A_n^0(H+M) - \dot{A}_n \cdot \dot{P} = \delta_{n,0} M(H+M) , \qquad (A.11)$$

where A_n^{μ} with n = 0 is just the momentum:

$$A_0^{\mu} = P^{\mu}$$
 (A.12)

The canonical brackets of $A_n^{\ \mu}$ and Q^{μ} among themselves are identical in form to the Dirac brackets (6.18)-(6.19) of $\alpha_n^{\ \mu}$, q^{μ} and P^{μ} . Unfortunately no explicit solution of the Virasoro constraints obeyed by $A_n^{\ \mu}$ is known, so that the quantum analysis of the closure of the Lorentz algebra cannot be carried out.

REFERENCES

- [1] Y. Nambu, Lectures at Copenhagen Summer Symposium, 1970, unpublished.
- [2] P. Goddard, J. Goldstone, C. Rebbi and C. B. Thorne, Nucl. Phys. <u>B56</u>, 109 (1973).
- [3] S. Mandelstam, Nucl. Phys. B64, 205 (1973).
- [4] A. Patrascioiu, Nucl. Phys. B81, 525 (1974).
- [5] A. Chodos and C. B. Thorne, Nucl. Phys. B72, 509 (1974).
- [6] L. D. Faddeev and L. A. Takhtajan, Theor. Math. Phys. 21, 160 (1974).
- [7] F. Rohrlich, Phys. Rev. Lett. 34, 842 (1975).
- [8] Other choices of gauge lead also to Eq. (3.10), e.g. $x(\tau,\sigma) = \frac{1}{2}\cos\sigma(x(\tau,0) x(\tau,\pi)) + \frac{1}{2}(x(\tau,0) + x(\tau,\pi))$. However, the x_{σ} -uniform gauge is particularly convenient.
- [9] This equation has been derived in a different context by G. t'Hooft, Nucl. Phys. <u>B75</u>, 461 (1975). We would like to thank C. Rebbi for informing us of t'Hooft's work.
- [10] E. Del Giudice, P. DiVecchia and S. Fubini, Ann. Phys. 70, 378 (1972).
- [11] P. Goddard, A. J. Hanson and G. Ponzano, Nucl. Phys. B89, 76 (1975).
- [12] See, e.g., J. Schwinger, <u>Particles, Sources, and Fields</u>, Vol. I (Addison-Wesley, 1970).
- [13] P. A. M. Dirac, Proc. Roy. Soc. <u>A246</u>, 333 (1958);
 A. J. Hanson, T. Regge and C. Teitelboim, "Constrained Hamiltonian Systems", to be published by Accademia Nazionale dei Lincei (Rome, 1975).

FIGURE CAPTIONS

- Fig. 3.1 Solution of the two-dimensional string equations with masses at the ends only.
- Fig. 3.2 τ -dependence of p(0), p(π) and $x_{\sigma}(\sigma)$ for string with massive ends.
- Fig. 3.3 Massless limit solution of the two-dimensional massive-end string.
- Fig. 3.4 Phase-space orbit of the massless string with Hamiltonian (3.17). The arrows give the direction of increasing time.
- Fig. 4.1 Plot of periodic absolute value function, Eq. (4.14).
- Fig. 4.2 Plot of P, x, x_o versus τ for simplest rest frame solution of orthonormal gauge string equations.
- Fig. 4.3 Mapping from orthonormal-gauge parameter σ to the x_{σ}-uniform-gauge parameter $\tilde{\sigma}(\tau,\sigma)$ as a function of τ .
- Fig. 4.4 Plots of $\tilde{\theta}$, \tilde{x} , $\tilde{x}_{\tilde{\sigma}}$ in x_{σ} -uniform gauge as a function of τ . Heights of $\tilde{\theta}(0)$, $\tilde{\theta}(\pi)$ give the coefficient of the appropriate delta function.
- Fig. 4.5 A more general solution f(z) to the constraints (4.11), giving N interior points moving at the speed of light.
- Fig. 5.1 (a) Phase-space orbit of 1-fold system projected on the (k_1, r_1) plane; (b) projection on (k_2, r_2) plane. Numbers label sequential configurations in τ .
- Fig. 5.2 (a) Phase-space orbit of 1-fold system projected on the (k_{+},r_{+}) plane; (b) projection on (k_{-},r_{-}) plane — coincident paths are displaced for clarity. Numbers label sequential configurations in τ .
- Fig. 5.3 (a) The action variable J_{+} as a function of initial condition c; (b) the action variable J .
- Fig. 5.4 (a) x-space motion for $c = \pi/2$, a pure J_ normal mode; (b) x-space motion for $c = \pi$, a pure J₊ normal mode. Note that in case (a), the system returns to its initial configuration in half the time required by case (b).







Figure 3.2 τ -dependence of p(0), p(π) and $x_{\sigma}(\sigma)$ for string with massive ends.



Figure 3.3 Massless limit solution of the two-dimensional massive-end string.



Figure 3.4 Phase-space orbit of the massless string with Hamiltonian (3.17). The arrows give the direction of increasing time.





THIS PAGE LEFT BLANK

.





,





coefficient of the appropriate delta function.



Figure 4.5 A more general solution f(z) to the constraints (4.11), giving N interior points moving at the speed of light.

THIS PAGE LEFT BLANK

٠







Figure 5.1 (b) projection on (k_2, r_2) plane. Numbers label sequential configurations in τ .







Figure 5.2 (b) projection on (k_r) plane — coincident paths are displaced for clarity. Numbers label sequential configurations in τ .







•

Figure 5.3 (b) the action variable $J_{_}.$



Figure 5.4 (a) x-space motion for $c = \pi/2$, a pure J_ normal mode;

(b) x-space motion for $c = \pi$, a pure J_+ normal mode. Note that in case (a), the system returns to its initial configuration in half the time required by case (b).