

A NEW THEORETICAL PREDICTION OF THE GROUND STATE
HYPERFINE SPLITTING IN MUONIUM*

William E. Caswell
Department of Physics, Brown University
Providence, Rhode Island 02912

and

G. Peter Lepage**
Stanford Linear Accelerator Center
Stanford University, Stanford, California 94305

ABSTRACT

We have computed a new contribution to the ground state hyperfine splitting in muonium: $-2 \left(\frac{\alpha}{\pi}\right)^2 \frac{m}{M} \left(\ln \frac{M}{m}\right)^2 E_F = -6.6$ KHz. The new theoretical estimate, $\nu_{th} = 4463\ 297.9$ (7.0) KHz is in good agreement with experiment: $\nu_{exp} = 4463\ 302.35$ (0.52) KHz. The bulk of the uncertainty in the theory is due to the measured value of μ_{μ}/μ_p . Uncalculated theoretical contributions are expected to be no larger than a few KHz.

(Submitted to Physical Review Letters)

*Work supported by Department of Energy.

**Address after September 1, 1978: Laboratory of Nuclear Studies,
Cornell University, Ithaca, New York 14853.

High precision measurements of the ground state hyperfine splitting (hfs) in muonium ($\mu^+ e^-$) allow a detailed test of our understanding of two-body bound states in quantum field theory and particularly in QED. The current experimental value is:¹

$$\nu_{\text{exp}} = 4463\ 302.35\ (0.52)\ \text{KHz}$$

Until recently, theoretical predictions were known only to within 10 - 15 KHz.² The bulk of this uncertainty was due to three sources:

- (1) Uncalculated terms of $O\left(\alpha^2 \frac{m}{M} \ln \frac{M}{m} E_F \sim 6\ \text{KHz}\right)$ coming from two-loop ladder and cross-ladder diagrams ($E_F \sim \frac{8}{3} \alpha^4 m^2/M$ is the lowest order hfs);
- (2) Uncalculated terms of $O\left(\left(\frac{\alpha}{\pi}\right)^2 \frac{m}{M} \left(\ln \frac{M}{m}\right)^2 E_F \sim 3\ \text{KHz}\right)$;
- (3) Uncertainty in the measured value of μ_μ/μ_p leading to possible errors of $\pm 5\ \text{KHz}$.¹

In a new paper, Bodwin, Yennie and Gregorio have demonstrated that terms of the first sort cancel completely.³ In this letter, we describe a calculation of all terms of the second type--the largest remaining contributions to the muonium hfs. These arise from radiative corrections to the electron and photon lines in the one-loop ladder diagrams. We find a total contribution of

$$\Delta E = -2\left(\frac{\alpha}{\pi}\right)^2 \frac{m}{M} \left(\ln \frac{M}{m}\right)^2 E_F = -6.6\ \text{KHz} \quad . \quad (1)$$

Consequently the current theoretical prediction is

$$\nu_{\text{th}} = 4463\ 297.9\ (7.0)\ \text{KHz} \quad .$$

The agreement with experiment is excellent. The major source of error is now in (μ_μ/μ_p) , and improved values for this constant will be available

in the near future. There remains a theoretical uncertainty of a few KHz due to uncalculated terms of the form⁴

$$\left(\frac{\alpha}{\pi}\right)^2 \frac{m}{M} \ln \frac{M}{m} E_F \sim .6 \text{ KHz}$$

$$\alpha^2 \frac{m}{M} E_F \sim 1.1 \text{ KHz} \qquad \frac{\alpha^3}{\pi} E_F \sim .6 \text{ KHz}$$

In what follows, we first review the calculation of the $O\left(\alpha \frac{m}{M} \ln \frac{M}{m} E_F\right)$ terms in the hfs.⁵ Building upon this analysis, we then compute all $O\left(\alpha^2 \frac{m}{M} \left(\ln \frac{M}{m}\right)^2 E_F\right)$ contributions. Finally we present a very simple argument supporting the conclusions of Bodwin et al.³

The terms of $O\left(\alpha \frac{m}{M} \ln \frac{M}{m} E_F\right)$ come from the one-loop ladder graphs (Fig. 1). Factors of $\ln \frac{M}{m}$ can only arise from the integration region $m \ll |k| \ll M$ as only there are the integrals sensitive to both mass scales m and M . Thus we can restrict the integration so that $|k| \geq m$. This is useful for three reasons:

- (1) It prevents double counting of lower order (in α) contributions coming from the nonrelativistic region.
- (2) The relative momenta in the wave functions (i.e., p, q), being nonrelativistic ($\sim \alpha m$), can be neglected in the kernel, and the integration over wave functions trivially performed:

$$\left| \int d^3 p \phi(\vec{p}) \right|^2 = \left| \phi(\vec{x} = 0) \right|^2 \simeq \frac{\alpha^3 m^3}{\pi} \quad (2)$$

- (3) The effects due to binding are negligible ($O(\alpha^2)$ corrections) and the external legs can be put on mass-shell. Consequently the contribution from this region of momentum space in the kernels of Fig. 1 is gauge invariant.⁶

Thus the $O\left(\alpha \frac{m}{M} \ln \frac{M}{m} E_F\right)$ hfs due to the uncrossed ladder (Fig. 1a) is (in Feynman gauge):

$$\frac{|\phi(0)|^2}{3} \frac{ie^4}{(2\pi)^4} \int \frac{d^4k}{k^6} \frac{\text{Tr}^{(e)}\left[\gamma^i \gamma_5 \gamma^\mu k^\nu \frac{1+\gamma_0}{2}\right] \text{Tr}^{(\mu)}\left[\gamma^i \gamma_5 \gamma_\mu (-k^\sigma + M(\gamma^0 + 1)) \gamma_\nu \frac{1+\gamma_0}{2}\right]}{k^2 - 2Mk^0}$$

where we have projected out the hyperfine interaction (spin-spin) using

$$\frac{1}{3} \text{Tr}\left[\gamma^i \gamma_5 \mathcal{M}^{(e)} \frac{1+\gamma_0}{2}\right] \text{Tr}\left[\gamma^i \gamma_5 \mathcal{M}^{(\mu)} \frac{1+\gamma_0}{2}\right] \quad (3)$$

It is readily demonstrated that the cross-ladder graph (Fig. 1b) gives an identical contribution, and therefore the sum of the two is

$$\Delta E_{L+CL} = \frac{16}{3} |\phi(0)|^2 \frac{ie^4}{(2\pi)^4} \int \frac{d^4k}{k^6} \frac{k^{\rightarrow 2} + 3k^2}{k^2 - 2Mk^0}$$

Performing a Wick rotation ($k^0 = i k \cos\phi$, $|\vec{k}| = k \sin\phi$) and symmetrizing in k^0 , we finally obtain

$$\begin{aligned} \Delta E_{L+CL} &= \frac{\alpha}{\pi} \frac{m}{M} E_F \frac{2}{\pi} \int_m^{\sim M} \frac{dk}{k} \int_0^\pi d\phi \sin^2\phi \left[3 + 2 \tan^2\phi\right] \theta(|k \cos\phi| - m) \\ &\approx -3 \frac{\alpha}{\pi} \frac{m}{M} E_F \int_m^{\sim M} \frac{dk}{k} = -3 \frac{\alpha}{\pi} \frac{m}{M} \ln \frac{M}{m} E_F \end{aligned} \quad (4)$$

where only logarithmic terms have been retained.

Terms of $O\left(\alpha^2 \frac{m}{M} \left(\ln \frac{M}{m}\right)^2 E_F\right)$ are due to first order radiative corrections on the electron and photon lines of the graphs just discussed (Fig. 2). When $|k| \gg m$ (and Euclidean), these radiative corrections

modify the bare graphs only by an overall factor of the form

$$K \frac{\alpha}{\pi} \ln \left(\frac{k^2}{m^2} \right) \quad k^2 \gg m^2$$

Introducing such a factor into the integrand of (4) results in a splitting of

$$\Delta E = -3 K \left(\frac{\alpha}{\pi} \right)^2 \frac{m}{M} \left(\ln \frac{M}{m} \right)^2 E_F + O \left(\left(\frac{\alpha}{\pi} \right)^2 \frac{m}{M} \ln \frac{M}{m} E_F \right) \quad (5)$$

As is well known, the constants K for the vacuum polarization, electron propagation and vertex corrections (Fig. 2a, b, c) are 1/3, 1/4 and -1/4, respectively.⁷ Furthermore, it is readily demonstrated that K = 1/4 for the radiative correction in Fig. 2d:

$$\begin{aligned} T_{\gamma e \rightarrow \gamma e} &\approx \frac{-ie^2}{(2\pi)^4} \int \frac{d^4 q}{q} \bar{u}(0) \gamma_\mu \not{q} \not{k} \frac{1}{\not{q} + \not{k}} \not{k} \gamma^\mu u(0) \\ &\approx \frac{-ie^2}{(2\pi)^4} \int \frac{d^4 q}{q} \bar{u}(0) \not{k} \frac{1}{\not{k}} \not{k} u(0) \\ &\approx \frac{\alpha}{4\pi} \ln \left(\frac{k^2}{m^2} \right) T_{\text{Born}} \quad k^2 \gg m^2 \end{aligned}$$

Thus the leading contribution from the kernels in Fig. 2 is just that quoted above (Eq. (1)):

$$\Delta E = -3 \left[2 \left(\frac{1}{3} \right) + \frac{1}{4} + 2 \left(-\frac{1}{4} \right) + \frac{1}{4} \right] \left(\frac{\alpha}{\pi} \right)^2 \left(\frac{m}{M} \right) \left(\ln \frac{M}{m} \right)^2 E_F$$

These diagrams are the only source of $O \left(\alpha^2 \left(\ln \frac{M}{m} \right)^2 E_F \right)$ splittings.

Momenta in first order radiative corrections to the muon line are scaled

by M and thus cannot contribute $\left(\ln \frac{M}{m}\right)^2$ terms. Second order radiative corrections to the one-photon exchange interaction result in no $\left(\ln \frac{M}{m}\right)^2$. The only remaining graphs are the two-loop ladder and cross-ladder graphs considered by Bodwin et al.³ Two of these are illustrated in Fig. 3. Separately these graphs contain terms of $O\left(\alpha^2 \frac{m}{M} \ln \frac{M}{m} E_F\right)$. However, one can demonstrate that the graphs cancel in pairs to this order (of course $O\left(\alpha^2 \frac{m}{M} E_F\right)$ terms may remain) if the following intuitively reasonable assumptions are valid:

- the external momenta can be set to zero since logarithmic contributions come only from the region of relativistic momenta (as above);
- the electron mass can be neglected because logarithmic contributions arise only from the region of momenta $m \ll k, q \ll M$ (in both loops).

Bodwin et al. have verified these by direct computation. The logarithmic contribution of any diagram is then cancelled by that from the diagram obtained by reversing the electron line while leaving all other propagators unchanged (e.g., Fig. 3). This is because the traces (as in Eq. (3)) associated with the electron lines in each diagram are equal but opposite in sign. For example, the electron traces for the diagrams in Fig. 3 are (the γ_0 in $\frac{1 + \gamma_0}{2}$ of Eq. (3) obviously does not contribute here):

$$\begin{aligned}
 \text{(a)} \quad \text{Tr}^{(e)} & \left[\gamma^i \gamma_5 \gamma_\alpha \frac{1}{\not{q}} \gamma_\beta \frac{1}{\not{k}} \gamma_\delta \frac{1}{2} \right] \\
 \text{(b)} \quad \text{Tr}^{(e)} & \left[\gamma^i \gamma_5 \gamma_\delta \frac{-1}{\not{k}} \gamma_\beta \frac{-1}{\not{q}} \gamma_\alpha \frac{1}{2} \right] \\
 & = -\text{Tr}^{(e)} \left[\gamma^i \gamma_5 \gamma_\alpha \frac{1}{\not{q}} \gamma_\beta \frac{1}{\not{k}} \gamma_\delta \frac{1}{2} \right]
 \end{aligned}$$

Thus the logarithmic contribution (and indeed all contributions of $O(\alpha^2 E_F)$ from the region $m \ll k, q \ll M$) due to all two-loop ladder graphs cancel completely.

Needless to say, many of the approximations used in this paper are valid only when computing leading logarithmic terms. Non-leading terms must be analyzed within the context of an exact bound state perturbation theory (see, for example, Ref. 2 and 4). However, it appears likely that the uncertainty due to our ignorance of these terms will for the present be no larger than experimental uncertainties in the relevant physical constants (i.e., $\mu_\mu/\mu_p, \alpha$).

ACKNOWLEDGMENTS

The immediate stimulus for this work was the analysis by Bodwin, Yennie and Gregorio described in Ref. 3. We are indebted to these people for several conversations. We also thank V. Hughes and R. Horgan for fruitful discussions.

This work was supported by the Department of Energy.

REFERENCES

1. D. E. Casperson, T. W. Crane, A. B. Denison, P. O. Egan, V. W. Hughes, F. G. Mariam, H. Orth, H. W. Reist, P. A. Souder, R. D. Stambaugh, P. A. Thompson, and G. zu Putlitz, Phys. Rev. Letters 38, 956 (1977).
2. G. P. Lepage, Phys. Rev. A16, 863 (1977); G. T. Bodwin and D. R. Yennie, Cornell preprint CLNS-383 (Dec. 1977), to be published in Phys. Rep.
3. G. T. Bodwin, D. R. Yennie, and M. A. Gregorio, Cornell preprint (July 1978).
4. A systematic procedure for computing terms of the first two types is described in: W. E. Caswell and G. P. Lepage, SLAC-PUB-2080 (Feb. 1978), to be published in Phys. Rev. A. The methods of Ref. 2 can also be applied. Work on $O(\alpha^3 E_F)$ hfs is now in progress.
5. W. A. Newcomb and E. E. Salpeter, Phys. Rev. 97, 1146 (1955); H. Grotch and D. R. Yennie, Rev. Mod. Phys. 41, 350 (1969).
6. This is true even though the momentum k flowing in the photon lines is cut off at m since the cut-off is symmetric when $k \rightarrow -k$.
7. J. D. Bjorken and S. D. Drell, Relativistic Quantum Mechanics (McGraw-Hill, New York, 1964). These constants are easily computed in much the same manner as the result for Fig. 2d is determined in this letter.

FIGURE CAPTIONS

1. Kernels contributing to $O\left(\alpha \frac{m}{M} \ln \frac{M}{m} E_F\right)$ hfs.
2. Kernels contributing to $O\left(\alpha^2 \frac{m}{M} \left(\ln \frac{M}{m}\right)^2 E_F\right)$ hfs.
3. Two graphs which cancel to $O\left(\alpha^2 \frac{m}{M} \ln \frac{M}{m} E_F\right)$.




Fig. 1






Fig. 2



(a)

7-78



(b)

3438A3

Fig. 3