

TESTING QUANTUM CHROMODYNAMICS*

Stanley J. Brodsky
Stanford Linear Accelerator Center
Stanford University, Stanford, California 94305

ABSTRACT

The difficulties in isolating specific QCD mechanisms which control hadronic phenomena, and the complications in obtaining quantitative tests of QCD are discussed. A number of novel QCD effects are reviewed, including heavy quark and higher twist phenomena, initial and final state interactions, direct processes, multi-particle collisions, color filtering, and nuclear target effects. The importance of understanding hadron production at the amplitude level is stressed.

1. introduction

From the standpoint of hadronic physics of a decade ago it seems extraordinary that we now have a fundamental local Lagrangian field theory of the strong interactions. There is now a huge array of experimental observations which support the premise that the basic degrees of freedom of hadrons and their interactions are the confined quark and gluon fields of

* Work supported by the Department of Energy, contract DE-AC03-76SF00515.

(Closing lecture presented at the XIIIth International Symposium on Multiparticle Dynamics, Volendam, The Netherlands, June 6-11, 1982.)

quantum chromodynamics.¹⁾ The empirical evidence ranges from hadronic spectroscopy (including the heavy quark bound state spectrum, and the emerging evidence for gluonic bound states), the basic phenomena of deep inelastic lepton scattering and massive lepton pair production (consistent with point-like spin 1/2 quarks carrying the electromagnetic and weak currents in hadrons and the QCD-predicted pattern of scale violation), the scaling of $\sigma(e^+e^- \rightarrow X)$ (consistent with SU(3) color, asymptotic freedom), and large momentum transfer exclusive processes such as hadronic form factors (consistent with QCD dimensional counting rules and scale-invariant quark-quark interactions at short distances).

The multiparticle physics results reported at this meeting appear to be consistent with the general patterns of jet development and quantum number flow expected in QCD, including evidence for gluon jets in $e^+e^- \rightarrow q\bar{q}g$ events, quark fragmentation in $\ell p \rightarrow \ell'HX$, and the dominance of hard scattering QCD mechanics in high p_{\perp} direct photon and single hadron production in high energy hadron-hadron collisions. The recent results from PETRA²⁾ on $\gamma\gamma \rightarrow$ jets and the photon structure functions are especially important since they give an immediate and striking verification of the QCD-predicted pointlike coupling of real photons to the quark current at high momentum transfer.

Certainly, at the qualitative level, QCD does provide a viable framework for understanding present hadronic phenomena. The paradox is that despite these successes we are not certain that we are actually testing QCD predictions for hadron dynamics in a truly quantitative way, particularly since many results follow from simple parton ideas or more general principles, independent of the theory.

There are several reasons why quantitative tests of QCD have been so difficult.

i) Even for the simplest processes, many different QCD mechanisms contribute, including initial state corrections (including color correlations)

and "higher twist" terms (non-leading terms in $1/Q^2$), particularly multiparticle coherent effects and heavy quark phenomena. We will discuss some of these complications and the tests needed to isolate specific mechanisms in this review.

ii) It is difficult to gauge the reliability of perturbative QCD predictions in the absence of systematic high order calculations and the uncertainties due to the possible influence of non-perturbative dynamics.

iii) Predictions for inclusive hadron production processes are at present based on probabilistic models for quark and gluon jet hadronization, which by necessity contain ad hoc assumptions. It is not clear whether such predictions test QCD or the jet model Monte Carlo.

A prime example of the difficulty in testing QCD quantitatively is the fact that the coupling strength $\alpha_s(Q^2)$, has not been reliably determined to within 50% accuracy at any momentum scale; there is certainly no direct evidence that $\alpha_s(Q^2)$ decreases logarithmically with momentum transfer. In particular, the analysis of CELLO group at PETRA indicates that the value of $\alpha_s(Q^2)$ derived from $e^+e^- \rightarrow 3$ jet events is strongly sensitive to the particular model used to simulate quark and gluon jet fragmentation-- the Lund model gives values of α_s from 20% to more than 50% larger than that determined using the conventional Hoyer et al., Feynman-Field type models.³⁾ Similarly, as shown in the report to this meeting by S. Ellis, predictions for energy flow correlations and asymmetries at present accessible energies, $Q^2 \sim 1000 \text{ GeV}^2$ are strongly dependent on the model used for the non-perturbative jet hadronization.

It is clearly important to test QCD in a systematic way and that the important underlying dynamics not be misidentified. One can be optimistic that as we reach the higher energies of the new $p\bar{p}$, e^+e^- , ep colliders many of the complications of higher twist terms, threshold effects, and jet fragmentation dependence will be alleviated.⁴⁾ On the theoretical side there is now

concerted efforts to understand the higher order and background effects at a basic level. For example, certain higher twist terms can be related via the hadronic wavefunction to quantities measurable in exclusive processes such as form factors, Compton scattering, $\gamma\gamma$ production of hadron pairs, and heavy quarkonia decays.⁵⁾⁻¹⁴⁾ It has also been recently shown that the scale Q^2 of the leading order term in $\alpha_s(Q^2)$ in the perturbative expansion for many QCD high momentum transfer reactions is not ambiguous but is automatically fixed.¹⁵⁾

In this talk I will give a brief review of some of the known effects which negate straightforward tests of QCD at present energies. QCD can also be tested by verifying novel effects unique to the theory such as color filtering and transparency phenomena, and the "direct" production of $q\bar{q}$ jets in $\pi\pi$ collisions in events unaccompanied by forward hadron production (see Section 5). We emphasize in Section 3 the importance of understanding the Fock state wavefunction of hadrons at equal time on the light cone in order to relate and calculate many different exclusive, semi-inclusive and inclusive processes. The most interesting hadronic phenomena are those which were not predicted (see Section 5)--such as the features of high multiplicity high transverse total energy events and heavy quark production in hadron collisions and the copious baryon production in e^+e^- annihilation. It is of course also important to keep open the possibility that QCD could either fail or be modified at some level -- for example by the breaking of color symmetry or by merging with a more comprehensive theory such as supersymmetry.¹⁶⁾

2. Complications in Testing QCD

It is in the nature of inclusive hadronic processes that virtually any QCD mechanism which can be drawn as a Feynman graph will contribute to the cross section at some level. Perturbative QCD, the operator product expansion, and the factorization ansatz are important guides to the dominant contributions for large momentum transfer reactions. However, the secondary effects are

often not under good theoretical control because of the absence of rigorous bounds or because of parametrization uncertainties. On the other hand, many of these complicating processes constitute novel QCD effects and can be important tests of the theory.

There are a number of reasons why the precise determination of $\alpha_s(Q^2)$ from $e^+e^- \rightarrow q\bar{q}g$ jet events is intrinsically difficult. The primary problem is that the separation between 3-jet $q\bar{q}g$ and two-jet qq events requires detailed, certain knowledge of the transverse momentum k_\perp and longitudinal light-cone fraction $z = (k^0 + k^3)/(p_q^0 + p_{\bar{q}}^3)$ dependence of the quark fragmentation distributions $D(z, k_\perp) = dN/dz/d^2k_\perp$. The only theoretical input from perturbative QCD is at large $z \sim 1$ and/or large k_\perp where the hadron wavefunctions are probed in the far off shell regime. One can show that (modulo logarithmic factors) (see figure 1)

$$D_{M/q}(z) = \frac{dN}{dz} (M/q) \underset{z \rightarrow 1}{\sim} A(1-z)^2 + \frac{C_M}{Q^2} \quad (1)$$

where the C_M/Q^2 term is associated with the longitudinal current.¹⁷⁾ The high twist contribution to jet fragmentation from Fig. 1(b) is^{12,13}

$$\frac{dN}{dk_\perp^2} (M/q) \sim \frac{dN}{dk_\perp^2} (g/q) F_M(k_\perp^2) \propto \frac{\alpha_s}{\pi k_\perp^2} F_M(k_\perp^2) \quad (2)$$

for meson production at large k_\perp . The scale constant C_M can be computed in terms of the meson form factor [see Eq. (3)].¹⁸⁾ (Calculations also show that the distribution in k_\perp and z does not factorize.) In contrast to the QCD forms, standard jet hadronization parametrizations usually assume that the transverse momentum distribution falls as $\exp(-bk_\perp^2)$, and that $D_{M/q}$ is non-vanishing at $z \rightarrow 1$ at large Q^2 . The QCD form (2) for $q \rightarrow q+M$ predicts hadron production at relatively large transverse momentum, reducing the number of events which should be identified as $q\bar{q}g$; i.e., the $F_M(k_\perp^2)/k_\perp^2$ terms (summed




Fig. 1. QCD contributions to the quark fragmentation functions at large k_{\perp} . The direct meson contribution (b) gives a $1/k_{\perp}^4$ power-law tail normalized to the meson form factor.

over all meson states) can be a significant background to the $\alpha_s(k_{\perp}^2)/k_{\perp}^2$ gluon jet signal. The problem is compounded by heavy quark fragmentation uncertainties (e.g. $e^+e^- \rightarrow \bar{c}c \rightarrow \bar{c}\Lambda_c X \rightarrow \bar{c}pX'$), and the production of gluonium states. There are also questions of a more fundamental nature which require an understanding of confinement and non-perturbative effects. The analysis of Gupta and Quinn¹⁹⁾

shows that the standard picture of jet hadronization would certainly break down if all quark masses were large compared to the QCD scale Λ . Thus there must be a hidden analytic dependence

on the quark mass which controls jet fragmentation, and corrections to the jet cross section beyond that indicated by the perturbative expansion.

Many of the results reported at this meeting highlight the importance of isolating heavy quark fragmentation effects. The important prediction of Bjorken²⁰⁾ and Suzuki²¹⁾ that charmed hadrons are produced dominantly at large z in the c -quark fragmentation region appears to be confirmed by e^+e^- annihilation and deep inelastic lepton scattering data.²²⁾ Charm and beauty quark fragmentation thus could account for a substantial fraction of meson and baryon production at large k_{\perp} and z , competing with hard gluon bremsstrahlung processes in e^+e^- annihilation.²³⁾

In the case of deep inelastic lepton scattering the central test of QCD is the evolution of the structure functions. The background effects include:

- i) Higher twist contributions (terms suppressed by powers of $1/Q^2$ at fixed x). Such terms arise from mass insertions, k_{\perp} effects, and coherent lepton-multi-quark scattering contributions. Calculations of the latter

contributions require knowledge of the multi-quark distribution amplitudes of the target. However, absolutely normalized predictions can be obtained for the meson structure functions at $x \rightarrow 1$ since the required valence wave function matrix elements are already determined by the meson electromagnetic form factor. For example, the leading contribution at $x \rightarrow 1$ to the meson longitudinal structure function takes the form ($C_F = 4/3$)¹⁶⁾

$$F_L^M(x, Q^2) \cong \frac{2C_F}{\pi} \frac{x^2}{Q^2} \int_{m^2}^{Q^2} dk^2 \alpha_s(k^2) F_M(k^2) \left[1 + O\left(\frac{k^2(1-x)}{Q^2}\right) \right] \quad (3)$$

Numerically this gives $F_L^M \sim .2x^2/Q^2$ in GeV^2 units. Notice that for $x \rightarrow 1$ and fixed Q^2 this contribution will dominate the transverse current meson structure function which is predicted to decrease as $(1-x)^2$ at $x \rightarrow 1$, modulo logarithmic factors. The higher twist F_L contribution comes from the direct interaction of the pion with the current; its QCD evolution is analogous to that of the photon structure function.

In the case of the nucleon structure functions, the leading twist contribution to F_2 at $x \rightarrow 1$ is predicted to vanish as $(1-x)^3$ or $(1-x)^5$ for quarks with helicity parallel or antiparallel respectively to the nucleon helicity.¹⁷⁾ A recent model calculation of the higher twist $[m^2/Q^2(1-x)^2]^{1,2}$ contributions to the nucleon function yields even larger effects than in the meson case. The analyses of Barnett et al.²⁴⁾ show that present deep inelastic nucleon target data cannot unambiguously separate scale-violating QCD evolution effects in leading twist from the higher twist contributions allowed in QCD. The (remote) possibility that all the observed scale violation is due to higher twist effects or that Λ_{QCD} is very small is not ruled out.

ii) Heavy quark thresholds. As $W^2 = (q+p)^2$ is increased beyond the production threshold for new quark flavors, the structure functions increase at fixed x in a direction opposite to QCD evolution, with an attendant change in the characteristics of the final state. Strong scale-breaking effects associated with the charm threshold are seen directly in the EMC data²⁵⁾ for the

$c(x, Q^2) + \bar{c}(x, Q^2)$ distribution in the nucleon. As emphasized by D. P. Roy,²⁶⁾ this effect can simulate an apparent cancellation of QCD evolution effects in deep inelastic structure functions. It is thus essential to accurately parametrize the quark mass effects. One can distinguish two components to the heavy quark distribution functions in the nucleon: (a) the "extrinsic" interaction dependent component generated by standard QCD evolution in association with the deep inelastic scattering, and (b) the "intrinsic" (i.e., initial condition) component generated by the QCD binding potential and equations of state for the proton.²⁷⁾ The intrinsic component is maximal when all the quarks of the bound state have similar velocities, thus favoring large momentum fractions for the intrinsic c and \bar{c} quarks. This leads to a valence-like distribution for the intrinsic charm quark distribution $c(x) \sim u(x)$ and a possible explanation of the charmed hadron distributions seen at the ISR in pp collisions. The present status of the intrinsic charm contribution is discussed in Section 5 and in Carsten Peterson's contribution to this conference.

iii) Final state QCD interactions.²⁸⁾ These effects do not affect the structure functions measured in deep inelastic scattering, but they do lead to changes in the particle distribution in the final state, e.g., k_{\perp} smearing of the current quark distribution,²⁹⁾ the production of associated hadrons in the central region, and the attendant degradation of the fast hadron momentum distribution. It is particularly interesting to study these effects in nuclear targets, using the nucleus to perturb the evolution of quark and gluon jets in hadronic matter. A general principle, the "formation zone",^{30), 31)} (which can be derived in perturbative QCD) leads to the prediction that radiation collinear with the current quark cannot be induced during passage through the nucleus at high energies.³²⁾ On the other hand, induced central region hadron production proportional to $A^{1/3}$ is allowed by QCD [see Fig. 2(a)]. (We note that such radiation, if verified, violates the usual assumption made in

analyzing hadron-nucleus collisions that the induced central region radiation is always correlated with the number of "wounded" nucleons in the target, and it predicts cascading effects in the central region). [see Fig. 2(b).]




Fig. 2. (a) Production of central rapidity region multiplicity in association with final state interactions in deep inelastic lepton scattering on a nuclear target. (b) Production of central region multiplicity in hadron-nucleon collisions. The cascading interactions lead to a ramp shaped multiplicity distribution rather than a flat plateau in the central rapidity region (see Ref. 33).

iv) Non-additive contributions to the nuclear structure functions.

Among the most important effects are shadowing (and possibly antishadowing) at low x and binding and kinematical corrections at $x \sim 1$. An argument that the shadowing due to traditional Glauber processes is suppressed at Q^2 large compared to a scale proportional to $1/x$ is given in Ref. 28. This is in contrast to standard fixed $W^2 = (q+p)^2$ duality arguments³⁵⁾ which connects shadowing at low x with shadowing of the real photon photoabsorption cross section. In addition, QCD evolution itself may be non-additive in the nuclear number.³⁶⁾ Non-additivity of the nuclear structure functions can also occur because of meson exchange currents (leading to the anomalous A dependence of the sea quark distributions), because of perturbations of the nucleon Fock states due to nuclear binding, or possibly because of "hidden color" components in the nuclear state.⁶⁾

Each of the above QCD complications can lead to significant effects in the cross section for virtually any inclusive process. In the case of Drell-Yan process, initial state interactions²⁸⁾ of the active quarks with spectators lead to (a) increased smearing of the Q_{\perp} distribution of the lepton pair beyond that contained in QCD radiative corrections or the hadronic wavefunctions, (b) target-dependent induced radiation in the central region and the associated

degradation of the quark longitudinal momentum distributions, and (c) a modification of the overall normalization of the pair product cross section $d\sigma/dQ^2 dx$ due to induced color correlations, (at least at subasymptotic Q^2). The color correlation can have anomalous dependence on the nuclear number A relative to that measured at the corresponding kinematic range in deep inelastic scattering.

The dominance of the longitudinal structure functions in the fixed W limit for mesons [as in Eq. (3)] is an essential prediction of perturbative QCD. Perhaps the most dramatic consequence is in the Drell-Yan process $\pi p \rightarrow \ell^+ \ell^- X$; one predicts that for fixed pair mass Q , the angular distribution of the ℓ^+ (in the pair rest frame) will change from the conventional $(1 + \cos^2\theta_+)$ distribution to $\sin^2(\theta_+)$ for pairs produced at large $x_L \rightarrow 1$. The results of the Chicago-Illinois-Princeton experiment³⁷⁾ at FNAL appears to confirm the QCD higher twist contribution with about the expected normalization. Striking evidence for this effect has also been seen in a Gargamelle analysis³⁸⁾ of the quark fragmentation functions in $\nu p \rightarrow \pi^+ \mu^- X$. The results yield a quark fragmentation distribution into positive charged hadrons which is consistent with the predicted form: $dN^+/dz dy \sim B(1-z)^2 + (C/Q^2)(1-y)$ where the $(1-y)$ behavior corresponds to a longitudinal structure function. It is also crucial to check that the $e^+e^- \rightarrow MX$ cross section³⁹⁾ becomes purely longitudinal ($\sin^2\theta$) at large z at moderate Q^2 , and that the observed effects are not kinematical in origin, or due to backgrounds from baryon production.

All of the above QCD complications are compounded in reactions involving large transverse momentum particle or jet production in hadron-hadron collisions. Conventional calculations based on the leading p_\perp^{-4} ($2 \rightarrow 2$) subprocesses suffer severe model-dependent ambiguities due to uncertainties in how to include k_\perp smearing effects, the size of scale breaking effects, and the strong parameter-dependence on the quark and gluon fragmentation processes. In these analyses it is usual to assume on-shell kinematics for the basic parton-

parton cross section. However, when k_{\perp} smearing from the hadron wavefunctions is introduced, this leads to a divergence at zero momentum transfer in the gluon propagator and thus an anomalous sensitivity to an arbitrary low momentum cutoff. The correct use of off-shell kinematics at high k_{\perp} removes this divergence⁴⁰⁾ and also much of the scale violations associated with k_{\perp} smearing. In addition, in standard approaches, the $q \rightarrow M+q$ fragmentation function is forced to be non-vanishing at $z \rightarrow 1$ (e.g., $D_{M/q}(z) \sim (1-z)^2 + C$ with $C \neq 0$); otherwise, one predicts more hadronic momentum collinear with the high p_T trigger hadron than that observed by experiment. In fact, as we have emphasized, QCD predicts that the only non-vanishing contributions to the fragmentation function at $z \rightarrow 1$ are due to higher twist subprocesses: specifically, direct subprocesses such as $gq \rightarrow Mq$ where the high p_T hadron is created in the short distance reaction instead of by fragmentation. In addition, the effects of initial and final state interactions (k_{\perp} smearing, multiple scattering, color correlation effects, associated central region multiplicities) can severely complicate the model calculations.⁴¹⁾

There is another serious difficulty with standard QCD phenomenology. If leading twist $2 \rightarrow 2$ subprocesses dominate direct photon and hadron production at large transverse momentum then one predicts a ratio $R_{\gamma/\pi} \sim f(x_{\perp}, \theta_{cm})$, independent of p_{\perp} at fixed $x_{\perp} = 2p_{\perp}/\sqrt{s}$ and θ_{cm} . The ISR data reported to this meeting, however, is consistent with $R_{\gamma/\pi} \sim p_{\perp}^2$ at fixed x_{\perp} and θ_{cm} at large p_{\perp} . The simplest interpretation of this result is simply that higher twist p_{\perp}^{-6} -scaling "direct" subprocesses such as $qg \rightarrow Mq$ and $q\bar{q} \rightarrow Mg$ dominate meson production the ISR kinematic regime whereas direct photons are produced by standard p_{\perp}^{-4} $q\bar{q} \rightarrow \gamma g$ and $qg \rightarrow qg$ subprocesses. The cross section for the direct meson processes can be precisely normalized^{9), 42)} in terms of the meson form factor since the same moments of the hadron wavefunction (distribution amplitude) appear. The direct processes can dominate the leading twist $qq \rightarrow qq$ and $qg \rightarrow qg$ subprocesses in the ISR regime because quark

fragmentation into fast hadrons is not required; the meson M is made directly in the subprocesses. As in the case of direct photon production, the direct meson is unaffected by final state interactions²⁸⁾ since a point-like component of the meson valence Fock state is involved (see Section 5). The direct processes also lead to significant quantum number correlations with the away-side jet since fermion exchange (rather than gluon exchange) plays an important role in the direct processes. Evidence for such strong correlations has been reported by the BFS and SFM collaborations.³¹⁾

It is clear that a complete formulation of large transverse momentum hadron production which takes into account all the relevant QCD effects, including initial state interactions, k_{\perp} smearing with off-shell kinematics, realistic fragmentation functions, as well as the higher twist direct subprocesses must be given before further progress can be made. In particular a careful separation of the leading twist and direct hadron production processes is required. It is certainly incorrect to force parametrizations such that all high p_T hadron production are attributed to the simplest quark or gluon scattering process.

An important clue to identifying underlying QCD subprocesses is the pattern of hadronic radiation produced in association with each inclusive reaction. The real difficulty in analysing multiparticle reactions quantitatively in QCD is that we do not understand color confinement or even how an individual hadron is formed! It is clear that to understand these problems at a basic level we will have to go beyond a statistical treatment of quark fragmentation and actually analyze hadron production at the amplitude level where coherent effects can be identified. It is clear even from electrodynamics that coherence is crucial for soft photon production; e.g., consider the case where two charged sources are nearly collinear.³⁴⁾ It is also important to look in detail at exclusive QCD processes such as form factors, high momentum transfer Compton scattering and two photon reactions $\gamma\gamma \rightarrow M\bar{M}$ at

fixed θ_{cm} which give the simplest and most direct information on the production of individual hadrons.⁴⁵⁾ It is also conceivable that the basic hadron wavefunction knowledge required for understanding quark and gluon fragmentation processes can be obtained from non-perturbative QCD calculations, such as lattice gauge theory, or the QCD equation of state on the light-cone. We will discuss this in more detail in Section 3.

3. Hadronic Wave Functions in QCD^{6), 46)-48)}

Even though quark and gluon perturbative subprocesses are simple in QCD, the complete description of a physical hadronic process requires the consideration of many different coherent and incoherent amplitudes, as well as the effects of non-perturbative phenomena associated with the hadronic wavefunctions and color confinement. Despite this complexity, it is possible to obtain predictions for many exclusive and inclusive reactions at large momentum transfer provided we make the standard ansatz that the effect of non-perturbative dynamics is negligible in the short-distance and far-off-shell domain. (This assumption appears reasonable since a linear confining potential $V \propto r$ is negligible compared to perturbative $1/r$ contributions.) For many large momentum transfer processes, such as deep inelastic lepton-hadron scattering reactions and meson form factors, one can then isolate the long-distance confinement dynamics from the short-distance quark and gluon dynamics -- at least to leading order in $1/Q^2$. The essential QCD dynamics can thus be computed from (irreducible) quark and gluon subprocesses amplitudes as a perturbative expansion in an asymptotically small coupling constant $\alpha_s(Q^2)$.

For example, the pion form factor at large Q^2 takes the form [see Fig. 3(b)]⁴⁶⁾

$$F_{\pi}(Q^2) = \int_0^1 dx \int_0^1 dy \phi^*(x, Q) T_H(x, y; Q) \phi(y, Q) \quad (4)$$

where

$$\phi_{\pi}(x, Q) = \int^Q \frac{d^2 k_{\perp}}{16\pi^3} \psi_{q\bar{q}/\pi^Q}(x, \vec{k}_{\perp}) \quad (5)$$

is the amplitude for finding the q and \bar{q} in the valence state of the pion collinear up to scale Q with light cone longitudinal momentum fractions x and $1-x$, and

$$T_H = \frac{16\pi C_F \alpha_s [Q^2(1-x)(1-y)]}{(1-x)(1-y)Q^2} \left[1 + \frac{O[\alpha_s(Q^2)]}{\pi} \right] \quad (6)$$

is the probability amplitude for scattering collinear constituents from the initial to the final direction. (The superscript Q in $\psi_{q\bar{q}}^Q$ indicates that all internal loop in $\psi_{q\bar{q}}$ are to be cutoff at $k_{\perp}^2 < Q^2$.) The $\log Q^2$ dependence of the distribution amplitude $\phi(x, Q)$ is determined by the operator product expansion on the light-cone or an evolution equation; its specification at subasymptotic momentum requires the solution to the pion bound state problem. The general form of $F_{\pi}(Q^2)$ is

$$F_{\pi}(Q^2) = \left| \sum_{n=0}^{\infty} a_n \log \frac{-\gamma_n Q^2}{\Lambda^2} \right|^2 C_F \frac{\alpha_s(Q^2)}{Q^2} \times \left[1 + O\left(\frac{\alpha_s(Q^2)}{\pi}\right) + O\left(\frac{m^2}{Q^2}\right) \right] \quad (7)$$

where the γ_n are computable anomalous dimensions. Similar calculations determine the baryon form factors, decay amplitudes⁴⁸⁾ such as $\Upsilon \rightarrow B\bar{B}$ and fixed angle scattering processes (see Figs. 3 and 4) such as Compton scattering, photo-production, and hadron-hadron scattering, although the latter calculations are complicated by the presence (and suppression) of pinch singularities.^{6), 36)} It is interesting to note that $\phi_{\pi}(x, Q^2)$ can be measured directly from the angular θ_{cm} dependence of the $\gamma\gamma \rightarrow \pi^+\pi^-$ and $\gamma\gamma \rightarrow \pi^0\pi^0$ cross sections at large s .⁴⁵⁾




Fig. 3. QCD subprocesses and factorization for high momentum transfer exclusive processes. (a) The $\gamma \rightarrow \pi^0$ transition form factor, measurable in $ee \rightarrow ee\pi^0$ reactions. (b) The meson form factor in factorized form. (c) Contributions to the $\gamma\gamma \rightarrow M\bar{M}$ amplitude. A complete calculation of these processes to leading order in $\alpha_s(Q^2)$ is given in Ref.45.




Fig. 4. Constraints on the baryon wavefunction in QCD. (a) Factorization of the baryon form factor (see Ref. 6). (b) Contribution to quarkonium decay into baryon pairs (see Ref. 48). (c) Calculation of the deep inelastic scattering structure functions from light-cone wavefunctions.

In addition, independent of the form of the mesons wavefunction we can obtain α_s from the ratio⁵⁾

$$\alpha_s(Q^2) = \frac{F_\pi(Q^2)}{4\pi Q^2 |F_{\pi\gamma}(Q^2)|^2} \left[1 + O\left(\frac{\alpha_s(Q^2)}{\pi}\right) \right] \quad (8)$$

where the transition form factor $F_{\pi\gamma}(Q^2)$ can be measured in the two photon reaction $\gamma^*\gamma \rightarrow \pi^0$ via $ee \rightarrow \pi^0 ee$. Equation (8) is in principle one of the cleanest ways to measure α_s . The higher order corrections in α_s are discussed in Ref. 49.

Thus an essential part of the QCD predictions is the hadronic wavefunctions which determine the probability amplitudes and distributions of the quark and gluons which enter the short distance subprocesses. The hadronic wavefunctions provide the link between the long distance non-perturbative and short-distance perturbative physics. Eventually, one can hope to compute the wavefunctions from the theory, e.g., from lattice or bag models, or directly from the QCD equations of motion, as we shall outline below. Knowledge of hadronic wavefunction will also allow the normalization and specification of the power law (higher twist) corrections to the leading impulse approximation results.

The wavefunction $\psi_{q\bar{q}}^\pi(x, k_\perp)$ which appears in Eq. (5) is related to the Bethe-Salpeter amplitude at equal "time" $\tau = t + z$ on the light-cone in $A^+ = 0$ gauge. The quark has transverse momentum k_\perp relative to the pion direction and fractional "light-cone" momentum $x = (k^0 + k^3)/(p^0 + p^3) = k^+/p^+$. The state is off the light cone $k^- = k^0 - k^3$ energy shell. In general a hadron state can be expanded in terms of a complete set of Fock state at equal τ :

$$|\pi\rangle = |q\bar{q}\rangle \psi_{q\bar{q}} + |q\bar{q}g\rangle \psi_{q\bar{q}g} + \dots \quad (9)$$

with

$$\sum_n \int [d^2k_\perp] [dx] |\psi_n(x_i, \vec{k}_{\perp i})|^2 = 1 .$$

(We suppress helicity labels.) At large Q^2 only the valence state contributes to an exclusive process, since by dimensional counting an amplitude is suppressed by a power of $1/Q^2$ for each constituent required to absorb large momentum transfer. The amplitudes ψ_n are infrared finite for color-singlet bound states. The meson decay amplitude (e.g. $\pi^+ \rightarrow \mu^+\nu$) implies a sum rule

$$\frac{a_0}{6} = \frac{1}{2\sqrt{n_c}} f_\pi = \int_0^1 dx \phi_\pi(x, Q) . \quad (10)$$

This result, combined with the constraint on the wavefunction from $\pi^0 \rightarrow \gamma\gamma$ requires that the probability that the pion is in its valence state is $\leq 1/4$. Given the $\{\psi_n\}$ for a hadron, virtually any hadronic properties can be computed, including anomalous moments, form factors (at any Q^2), etc.⁴⁸⁾

The $\{\psi_n\}$ also determine the structure functions appearing in deep inelastic scattering at large Q^2 ($a=q, \bar{q}, g$)

$$G_{a/p}(x, Q) = \sum_n \int_0^Q [d^2k_\perp] [dx] |\psi_n^a(x_i, k_{\perp i})|^2 \delta(x-x_i) \quad (11)$$

where one must sum over all Fock states containing the constituent a and integrate over all transverse momentum d^2k_\perp and the light-cone momentum fractions $x_i \neq x_a$ of the $(n-2)$ spectators. The valence state dominates $G_{q/p}(x, Q)$ only at the edge of phase space, $x \rightarrow 1$. All of the multiparticle x and k_\perp momentum distributions needed for multiquark scattering processes can be defined in a similar manner. The evolution equations for the $G_a(x, Q^2)$ can be easily obtained from the high k_\perp dependence of the perturbative contributions to ψ .

There are many advantages obtained by quantizing a renormalizable local $\tau = t+z$. These include the existence of an orthonormal relativistic wavefunction expansion, a convenient τ -ordered perturbative theory, and diagonal (number-conserving) charge and current operators. The central reason why one can construct a sensible relativistic wavefunction Fock state expansion on the light cone is the fact that the perturbative vacuum is also an eigenstate of the full Hamiltonian. The equation of state for the $\{\psi_n(x_i, k_{\perp i})\}$ takes the form

$$H_{LC} \Psi = M^2 \Psi \quad (12)$$

where

$$H_{LC} = \sum_{i=1}^n \left[\frac{k_{\perp i}^2 + m^2}{x} \right]_i + V_{LC}$$

is derived^{5), 15)} from the QCD Hamiltonian in $A^+ = 0$ gauge quantized at equal τ , and Ψ is a column matrix of the Fock state wavefunctions. Ultraviolet regularization and invariance under renormalization is discussed in Refs. 5,6,15.

A comparison of the properties of exclusive and inclusive cross sections in QCD is given in Table I.⁶⁾ Given the $\{\psi_n\}$ we can also calculate decay amplitudes, e.g. $\bar{T} \rightarrow p\bar{p}$ which can be used to normalize the proton distribution amplitudes [see Fig. 4(b)]. The constraints on hadronic wavefunctions which result from present experiments are given in Refs. 6,15,48. An approximate connection between the valence wavefunctions defined at equal τ with the rest frame wavefunction is also given in Ref. 48, so that one can make predictions from non-perturbative analyses such as bag models, lattice gauge theory, chromostatic approximations, potential models, etc. Other constraints from QCD-sum rules are discussed in Ref. 50.

It is interesting to note that the higher twist amplitudes such as $\gamma q \rightarrow Mq$, $gq \rightarrow Mq$, $q\bar{q} \rightarrow MM$, $q\bar{q} \rightarrow Bq$ which are important for inclusive hadron production reactions at high x_{\perp} can be absolutely normalized in terms of the distribution amplitudes $\phi_M(x,Q)$, $\phi_B(x_i,Q)$, by using the same analysis as used for the analysis of form factors.⁵⁾ In fact "direct" amplitudes such as $\gamma q \rightarrow Mq$, $q\bar{q} \rightarrow Mg$ and $gq \rightarrow Mq$ where the meson acts directly in the subprocess are rigorously related to the meson form factor since the same moment of the distribution amplitude appears in each case.

The light cone Fock state expansion also gives insight into the nature of forward hadron production in soft hadronic collisions. It should be emphasized that the properties of the strong interaction itself may distort the properties of the Fock state wavefunction in such a way that the quark distribution observed in soft collisions and fragmentation processes may differ significantly from that observed in deep inelastic scattering. For example, in the case of the DTU model,⁵¹⁾ at least one valence quark must be at low x in order to initiate the Pomeron "cylinder" interaction. One can also understand

Table I. Comparison of exclusive and inclusive cross section.

Exclusive Amplitudes	Inclusive Cross Sections
$\mathcal{M} \sim \Pi \phi(x_i, Q) \otimes T_H(x_i, Q)$	$d\sigma \sim \Pi G(x_a, Q) \otimes d\hat{\sigma}(x_a, Q)$
$\phi(x, Q) = \int^Q [d^2k_\perp] \psi_{val}^Q(x, k_\perp)$	$G(x, Q) = \sum_n \int^Q [d^2k_\perp] [dx] \psi_n^Q(x, k_\perp) ^2$
Measure ϕ in $\gamma\gamma \rightarrow M\bar{M}$	Measure G in $\ell p \rightarrow \ell X$
$\sum_{i \in H} \lambda_i = \lambda_H$	$\sum_{i \in H} \lambda_i \neq \lambda_H$

EVOLUTION

$$\frac{\partial \phi(x, Q)}{\partial \log Q^2} = \alpha_s \int [dy] V(x, y) \phi(y)$$

$$\frac{\partial G(x, Q)}{\partial \log Q^2} = \alpha_s \int dy P(x/y) G(y)$$

$$\lim_{Q \rightarrow \infty} \phi(x, Q) = \prod_i x_i \cdot C_{\text{flavor}}$$

$$\lim_{Q \rightarrow \infty} G(x, Q) = \delta(x) C$$

POWER LAW BEHAVIOR

$$\frac{d\sigma}{dx} (A+B \rightarrow C+D) \cong \frac{1}{s^{n-2}} f(\theta_{CM})$$

$$\frac{d\sigma}{d^2p/E} (AB \rightarrow CX) \cong \sum \frac{(1-x_T)^{2n-1}}{(Q^2)^{n_{act}-2}} f(\theta_{CM})$$

$$n = n_A + n_B + n_C + n_D$$

$$n_{act} = n_a + n_b + n_c + n_d$$

T_H : expansion in $\alpha_s(Q^2)$

$d\hat{\sigma}$: expansion in $\alpha_s(Q^2)$

COMPLICATIONS

End point singularities
Pinch singularities
High Fock states

Multiple scales
Phase-space limits on evolution
Heavy quark thresholds
Heavy twist multiparticle processes
Initial and final state interactions

this "held back" feature if one assumes that only these Fock states which are very peripheral (possessing a large impact parameter, low x constituent) can interact strongly. The non-interacting constituents thus have more of the beam momentum on the average than they would have in undistorted wavefunctions. For example, counting rules for renormalizable theories predict^{52),53)}

$$G_{e/A}(x) \underset{x \rightarrow 1}{\sim} (1-x)^{2n_s-1} \quad \text{where } n_s \text{ is the minimum number of spectators require to}$$

be stopped. If the hadronic wavefunction is undistorted in hadronic collisions, then one predicts, for example, $dN(pp \rightarrow K^+X)/dx \sim (1-x_L)^5$ (corresponding to 3 spectators). The power-behavior indicated by data however is $\sim (1-x_L)^3$. On the other hand if one quark in the incident hadron is required to be at $x \sim 0$ in order to have a large hadronic cross section, then there is one less spectator to stop. More generally, the held back mechanism reduces⁵³⁾ the power-law fall off by $(1-x_L)^2$ and systematically gives results in good agreement with data. The above considerations also indicate that the impact space distribution of the Fock state wavefunction could be strongly correlated with the x distribution; for example, low x sea quarks may be predominantly peripheral in impact space. Such a correlation would be natural if the sea quarks are pictured as constituents of virtual mesons in the nucleon cloud. Other possible QCD mechanisms without the held back mechanism are discussed in Refs. 52,54.

4. The Perturbative Expansion in QCD

The essential predictive power of perturbative QCD is due to the smallness of the effective coupling $\alpha_s(Q^2)$ and the existence of perturbative expansions for subprocess amplitudes which control larger momentum transfer reactions. There has been significant progress in extending the QCD predictions beyond leading order for a number of inclusive and exclusive reactions.

However, a major ambiguity in the interpretation of these perturbation expansions and assessing the convergence of the series is in the choice of expansion parameter. Given a specific renormalization scheme, one must still specify the argument Q^2 of α_s in order to make a definite prediction from an expansion to finite order. As shown in Ref. 15, the scale of momentum Q^* transfer which appears in the leading order term in α_s (in a given renormalization scheme) in the perturbative expansions is not arbitrary, but is in fact, automatically set by the theory. For example, in QED, the running coupling constant $\alpha(Q)$ is defined to take into account the effects of lepton pair vacuum polarization in each photon propagator at $Q^2 \gg 4m_l^2$, and the approximate scale Q^* as the QED expansion in $\alpha(Q^*)$ is readily identified. In the case of QCD in the leading non-trivial order, $\alpha_s(Q)$ is only a function of $\beta_0 = 11 - 2/3 n_f$. Thus we can use the analytic n_f dependence of the higher order corrections coming from light quark loop contributions to the gluon propagator to identify the correct scale of the α_s expansion in leading order, for almost every QCD processes. The net result is that for almost every reaction except $\Upsilon \rightarrow 3g$, the perturbative expansion appears to have good convergence with reasonably small coefficients of α_s/π . An important example is

$$R_{e^+e^-} = 3 \sum_q e_q^2 \left[1 + \frac{\alpha_s^{\overline{MS}}(Q^*)}{\pi} + 0.0825(\alpha_s/\pi)^2 \right] \quad (13)$$

where $Q^* = 0.71 Q$. We can then use this result to define or measure $\alpha_s^R(Q) \equiv \alpha_s^{\overline{MS}}(0.71Q)$, and then write the perturbative expansion for other observables in terms of α_s^R . The QCD predictions for the deep inelastic moments and decay rates for the 0^- and 1^- quarkonium decays in terms of the first two nontrivial orders in α_s^R is given in Ref. 15.

5. Novel Effects and Unexpected Effects in QCD

One can also test QCD by verifying novel effects which are essentially unique features of the theory. In this section we will list some examples of experimental tests which are specific to gauge theories of the strong interactions:

i) Hadron helicity conservation.

To leading order in $1/Q^2$ an exclusive process at large momentum transfer is dominated by amplitudes which conserve total hadron helicity; independent of photon (or weak boson) polarization. This is a consequence of the vector nature of gluon interactions and the fact that the total quark helicity equals the hadron helicity in the distribution amplitude ($L_z=0$). Many tests and predictions based on this rule are given in Ref. 55. An important prediction is that $\Upsilon \rightarrow pp$ should have a $(1 + \cos^2\theta_{cm})$ angular distribution.

ii) Direct processes.⁵⁶⁾

A surprising feature of QCD is the existence of inclusive processes such as $\pi pp \rightarrow q\bar{q}X$ where the pion's energy and momentum are completely consumed in the large p_\perp $q\bar{q}$ production; i.e., there is no associated hadron production in the forward fragmentation region [see Fig. 5(c)]. The jet cross section based on the $\pi g \rightarrow qq$ (and $\pi q \rightarrow gq$) subprocess is absolutely normalized in terms of the pion form factor (see Ref. 56); compared to the leading $qq \rightarrow qq$ subprocess one has

$$d\sigma(\pi \rightarrow q\bar{q}) \sim F_\pi(p_\perp^2) d\sigma(qq \rightarrow qq) \quad (14)$$

independent of α_s and the pion wavefunction. This process is also identifiable by conservation of the $p^+ = p^0 + p^3$ components between the pion and jet fragments, and the close transverse momentum balance of the q and \bar{q} jets. There is no bremsstrahlung or initial state interactions of the incident pion to leading order in $1/p_\perp^2$ since only the valence qq component of the pion wavefunction at $b_\perp \sim 0(1/p_\perp)$ contributes to this process -- such a state has




Fig. 5. Direct QCD subprocesses in inclusive high momentum transfer reactions. (a) The $gq \rightarrow Mq(p_{\perp}^{-6})$ contribution for producing high p_{\perp} mesons. (b) The $qM \rightarrow \ell\bar{\ell}q(Q^{-2})$ contribution to massive lepton pair production. (c) The $gM \rightarrow q\bar{q}(p_{\perp}^{-6})$ contribution to high p_{\perp} jet production. No hadrons are produced in the meson beam fragmentation region. (d) The $q\bar{p} \rightarrow q\bar{q}(p_{\perp}^{-8})$ contribution to the $p\bar{p} \rightarrow q\bar{q}X$ jet cross section. No hadrons are produced in the \bar{p} fragmentation region.

negligible color (hadronic) interactions. Similarly, in the case of nuclear targets $\pi_D A \rightarrow q\bar{q}X$, the valence pion state at $b_{\perp} \sim 1/p_{\perp}$ penetrates throughout the nuclear volume without any nuclear interactions. The corresponding direct baryon induced reaction based on $\bar{p}q \rightarrow \bar{q}\bar{q}$ is shown in Fig. 5(d). We also note that the amplitude to find three quarks in the proton at small separation with one accompanying spectator meson can be measured by an analogous process $\bar{p}pp \rightarrow \bar{q}\bar{q}\pi X$. Such amplitudes are of interest for predictions of baryon decay in grand unified theories.

In the case of hadron production at large p_{\perp} , the direct subprocess $gq \rightarrow \pi_D q$ produces pions unaccompanied by other hadrons on the trigger side [see Fig. 5(a)]. These p_{\perp}^{-6} QCD processes are absolutely normalized in terms of the meson form factor and can dominate jet fragmentation processes at large x_{\perp} . More generally, an entire set of hadrons and resonances can be produced by

direct subprocesses (e.g., $qq \rightarrow Bq$ and $qB \rightarrow qB$ for baryon production). Such contributions provide a serious background to any measurement where there is a high p_{\perp} single particle trigger. Again, the direct process hadrons have no final state interactions or accompanying collinear radiation to leading order in $1/p_T^2$.

iii) Quasi-elastic reactions in nuclei and color transparency.

As we have noted in Section 3, large momentum transfer exclusive reactions are dominated (to leading order in $1/p_{\perp}^2$) by valence Fock states with small constituent separation. Since such states have negligible hadronic interactions, a large momentum transfer quasi exclusive reaction can take place inside of a nuclear target [e.g. $\pi A \rightarrow \pi p(A'-1)$] without any elastic or inelastic initial or final state hadronic interaction.⁵⁷⁾ The rate for such "clean" reactions is normalized to A times the nucleon target rate.

iv) Diffractive dissociation and color filtering.

The existence of the light cone Fock state expansion for a meson implies a finite probability for the hadron to exist as a valence state at small relative impact parameters. This component of the state will interact only weakly in nuclear matter, whereas the majority part of the Fock state structure will interact strongly. The nucleus thus acts as a "color filter" absorbing all but the weakly interacting Fock components. The consequence is a computable cross section for the diffractive dissociation by a nucleus of pion into relatively high k_{\perp} $q\bar{q}$ jets.⁵⁸⁾ The rate is normalized to the pion decay constant and $\sigma_{\pi A}^{e\bar{e}}$. The θ_{cm} dependence of the jet in the $q\bar{q}$ rest frame is related to the pion distribution amplitude. Predictions for the k_{\perp} dependence of the jets are given in Ref. 58. All of the effects (2)-(4) test a basic feature of QCD: the Fock state structure of the color singlet hadronic wavefunctions and the special features of the valence Fock state.

v) Initial state interactions and color correlations.²⁸⁾

A detailed discussion of the expected effects of initial and final state interactions in a non-Abelian theory is given in a separate contribution to this volume.⁵⁹⁾ The main novel effects are subasymptotic color correlations (in principle A -dependent), and associated production in the central region. It is also possible that the soft particles produced in an initial state interaction could subsequently interact with other valence quark in the beam hadron to produce low mass lepton pairs ($Q^2 < m\sqrt{s}$), low p_{\perp} jets, etc. (see Fig. 6).




Fig. 6. Intermediate mass lepton pairs produced by the annihilation of a valence quark with a central region \bar{q} produced in association with an initial state collision. Such contributions have anomalous nuclear number and energy dependence.

vi) Hadron multiplicity in e^+e^- collisions.

In a remarkable calculation, Basetto, Ciafaloni, Marchesini, and Mueller⁶⁰⁾ have shown from an all orders perturbative analysis that QCD predicts a dip for $dn/dy_{cm} \sim 0$ for particle production in e^+e^- annihilation. This result needs careful experimental confirmation. Although the QCD prediction seems special to gauge theory, it should be noted that the statistical model of Ochs⁶¹⁾ based on a simple branching process also has this feature.

vii) The perturbative coupling strength of gluons to a gluon jet is $9/4$ larger than the coupling to a quark jet. The deviation from a factor of 2 is due to color coherence. Perturbative QCD thus evidently predicts that at some level gluon jets are broader and have a higher multiplicity plateau than quark jets. The fact that gluon jets can be screened by gluonium production may lead to further differences between quark and gluon jets. Polarized gluon jets may also have distinctive features such as oblateness.⁶²⁾

Perhaps the most interesting and important experimental observations are the phenomena not readily explainable or predicted by conventional QCD analyses. We will briefly discuss four examples here:

i) High transverse energy, high multiplicity events.

It is possible that this phenomena, first observed by the NA5 collaboration at the SPS, is a collective or fireball effect, or even a signal of a quark-gluon phase at high temperature. Field, Fox and Kelly⁶³⁾ have attempted to identify these events with the conventional processes, e.g., quark-quark scattering, etc. subprocesses accompanied by multiple hard gluon radiation and subsequent neutralization. However, realistic calculation taking into account energy momentum correlations and interference effects (color correlations, formation zone) between multiple strings seems very difficult. An alternative explanation is that multiple quark/gluon scattering (Glauber) processes are involved at high transverse energies, since at $x_T = E_T/E_{max} \sim 1$ the scattering of all the beam momentum to the transverse direction is required. Using counting rules, the $pp \rightarrow$ jet production cross section receives contributions of nominal order (R is the hadron transverse size)⁶⁴⁾

$$E \frac{d\sigma}{d^3p} \sim \frac{\alpha_s^2}{p_T^4} (1-x_T)^7, \quad \frac{\alpha_s^4}{R^2 p_T^6} (1-x_T)^3, \quad \frac{\alpha_s^6}{R^4 p_T^8} (1-x_T)^{-1}, \quad (15)$$

corresponding to the 2 particle, 4 particle, and 6 particle scattering processes shown in Fig. 7. A realistic calculation requires consideration of all the q and g multiparticle subprocesses, account of scale breaking, etc. The multiscattering terms can clearly dominate at x_T large; they are also characterized by high multiplicity, long-range correlations in rapidity and absence of coplanarity.

ii) Heavy flavor production in hadron collisions.

Standard QCD calculations based on $gg \rightarrow Q\bar{Q}$ subprocesses do not account for the x_L dependence, diffractive character, and magnitude of forward charm




Fig. 7. QCD contributions to high p_T processes. The multipartile 2+2 and 3+3 reactions produce high transverse energy events with a large fraction of the available energy. More complicated reactions with gluons and sea quarks are not shown.

hadron production observed at the ISR. It is also difficult to reconcile the large cross section ($\sigma_{\text{charm}} \sim 1 \text{ mb}$) observed at the ISR with the much smaller cross sections observed at Fermilab energies.

One possible mechanism⁶⁵⁾ for fast charm production in forward pp collisions is that the high momenta of the ud spectators in a nucleon, combined with a centrally produced charmed quark will yield a high momentum $\Lambda_c = |udc\rangle$. This is however contrary to the Bjorken-Suzuki effect, which indicates that the Λ_c should have roughly the same momentum as the charmed quark. This explanation also implies a long-range rapidity separation between the c at $y_{\text{cm}} \sim 0$ and a charmed hadron in the fragmentation region; it also does not account for the observed $(1-x_L)^3$ distribution of the $D^+ = |c\bar{d}\rangle$ meson (unless such mesons always arise from charm baryon decay). Measurements of the x_L distribution of the $\bar{\Lambda}_c$ would be decisive.

The simplest explanation of the observed ISR phenomena is that the proton contains a Fock state $|uudc\bar{c}\rangle$ with the $c\bar{c}$ bound inside the hadron over a relatively long time scale $O(m_c^{-1})$.²⁷⁾ Such a state (analogous to a $\Lambda_c\bar{D}$ component of the nucleon Fock state) could be diffractively dissociated into charmed hadrons in the forward fragmentation region as well as producing

valence-like c and \bar{c} distributions in deep inelastic lepton scattering (see Section 2). If the model is correct, b and t quarks can also be produced at large x_L with substantial cross sections, scaling as $1/M_Q^2$. Such states should contain relatively large transverse momentum--which together with the large quark mass, implies high k_\perp leptons and baryon production; this would also yield a clear signal for t -quark production.

The origin of the intrinsic heavy quark state in the nucleon wavefunction is the heavy quark loop vacuum polarization (and light-by-light scattering) insertions proportional to $1/M_Q^2$ in the QCD potential. As in the case of the potential which appears in the evolution equation for the distribution amplitude,⁵⁾ such contributions do not have a dx/x singularity at small x . The peak of the light-one distribution is at $x_i \sim m_i / \sum_j m_j$ where the state is minimally off-shell. This corresponds to the fact that in the rest system of the proton, the virtual $Q\bar{Q}$ pair is produced dominantly at threshold, at low velocities. This even to lowest order in $1/M_Q^2$ the heavy quarks are produced with a distribution which peaks at large x_Q . There is thus no question that Fock states containing heavy quarks exist at some level in ordinary hadrons -- the real question is the absolute normalization. The vacuum polarization contributions give $P_{Q\bar{Q}} \sim O\alpha_s^2(\lambda^2)\lambda^2/M_Q^2$ where λ^2 is a typical hadronic scale. The bag model calculation of Donoghue and Golowich⁶⁶⁾ gives $P_{c\bar{c}} \sim O(1\%)$. One also could hope to relate the normalization of $P_{Q\bar{Q}}$ to the spin-spin splitting in the baryon system.

The origin and normalization of the heavy quark Fock state components and their systematic dependence on the hadron or nuclear state can yield important clues to the nature of hadron wavefunctions in QCD. Similarly, the production of heavy quark states by diffractive excitation of the intrinsic heavy quark states or a similar mechanism, should also give important insight into the nature of hadronic processes in forward high energy collisions.

iii) The observation of copious baryon production in e^+e^- annihilation is somewhat of a surprise from the standpoint of perturbation QCD ideas. For $z \rightarrow 1$ baryon production should be suppressed by at least a power of $(1-z)$ relative to meson production whereas the data indicates a flat baryon/meson ratio out to $z = 0.5$. In the Lund model⁶⁷⁾ copious baryon production is accounted for by effective diquark production in a non-perturbative tunneling model. Another possibility, suggested by T. DeGrand,²³⁾ is that the fast charmed and beauty baryons produced at large z by the Bjorken-Suzuki mechanism leads by decay to a significant fraction of the large x_L baryons.

6. Conclusions

Despite the formal simplicity of its underlying Lagrangian, QCD has turned out to be an extraordinarily complex theory. Definitive tests have turned out to be very difficult, but there is at present no reason to doubt that QCD is at the basis of all hadron and nuclear dynamics accessible at present energies. Fortunately, as experiment has become more definitive, theoretical analyses has also made definite progress. The goal is to make precise predictions, with systematic control of background effects (such as high twist contribution, threshold effects, initial and final state interactions), as well as to attain a deeper understanding of jet hadronization and the QCD perturbation expansion.

Much of the present uncertainty in QCD predictions is due to the absence of detailed information on hadronic matrix elements. It seems likely that direct calculations of hadronic amplitudes will be possible using lattice gauge theory, or as discussed in Section 3, by solving the QCD equation of state for the Fock states of hadrons at equal $\tau = t+z$. This formalism gives a consistent relativistic wavefunction basis and calculational framework for QCD. One can also use this formalism to obtain many new predictions which are in principle exact, e.g., large momentum transfer exclusive reactions [such as the

meson form factors and $\gamma\gamma \rightarrow M\bar{M}$, new QCD constraints such as hadron helicity conservation, constraints on wavefunctions from decay amplitudes, new methods to determine α_s , and methods to calculate the higher twist and direct subprocesses.

We have also seen that QCD diverges in many ways from the expectations of the parton model. Many of the novel phenomena discussed here such as color correlations in initial state interactions, color filtering and transparency, intrinsic heavy quark Fock states, associated production in hard collisions, color coherence, direct processes, the dominant longitudinal component to the meson structure function, and many nuclear effects⁶⁸⁾ (reduced form factors, anomalous A dependence) were not anticipated within the parton model framework. In addition, the basic hard scattering mechanism for form factors and the QCD breakdown of the exclusive-inclusive connection is contrary to the mechanisms assumed to be dominant in the parton model framework.

We have also emphasized here the importance of understanding the physics of initial and final state interactions in QCD, particularly the breakdown of factorization at large target length, the physics of the formation zone in QCD, and the interesting effects of color correlations. An important clue toward understanding these phenomena as well as the propagation of quark and gluon jets through hadronic matter will be the careful study of nuclear target effects.

ACKNOWLEDGMENT

I wish to thank Professor W. Kittel and his colleagues on the organizing committee for arranging an excellent Symposium at Volendam. I also wish to thank E. Berger, J. Bjorken, G. Bodwin, J. Gunion, T. Huang, R. Hwa, G. P. Lepage, P. Mackenzie, A. Mueller, W. Ochs, C. Peterson, and P. Zerwas for helpful conversations.

REFERENCES

- 1) Reviews of QCD are given in: A.J. Buras, Rev. Mod. Phys. 52, 199 (1980); A.H. Mueller, Phys. Rep. 73C, 237 (1981); E. Reya, Phys. Rep. 69, 195 (1981); G. Altarelli, University of Rome preprint 701 (1978); W. Marciano and H. Pagels, Phys. Rep. 36C, 137 (1978); S.J. Brodsky and G.P. Lepage, Proceedings of the SLAC Summer Institute on particle Physics, 1979. See also "Perturbative Quantum Chromodynamics" (Tallahassee, 1981). AIP Proceedings No. 74, New York, 1981. For recent reviews of the phenomenology of QCD see A.J. Buras, FERMILAB-CONF-81/69-THY and A.H. Mueller, CU-TP-219 (1981), published in the Proceedings of the 1981 International Symposium on Lepton and Photon Interactions at High Energies, Bonn, August 1981.
- 2) J.B. Dainton, this conference. See also, D. Cords, to be published in the Proceedings of the SLAC Summer Institute on Particle Physics (1982).
- 3) P. Renton and J. von Krogh, this conference. See also S.L. Wu, to be published in the Proceedings of the SLAC Summer Institute on Particle Physics (1982), and the 1981 Proceedings of the IVth International Colloquium on Photon-Photon Interactions, Paris (ed. G. London).
- 4) Clear signals for jet production in $\bar{p}p$ collisions at the SPS have been reported. See the reports of T. Himel and D. DiBitonto, to be published in the Proceedings of the SLAC Summer Institute on Particle Physics (1982).
- 5) G.P. Lepage and S.J. Brodsky, Phys. Rev. D22, 2157 (1980).
- 6) S.J. Brodsky, T. Huang and G.P. Lepage, SLAC-PUB-2868, published in the Proceedings of the KFK Summer School on Quarks and Nuclear Forces (1981), "Quarks and Nuclear Forces," edited by D. Fries and B. Zeitnitz, Springfield (1982).
- 7) E.L. Berger and S.J. Brodsky, Phys. Rev. Lett. 42, 940 (1979); E.L. Berger, Phys. Lett. 89B, 241 (1980).
- 8) G.R. Farrar and G.C. Fox, Nucl. Phys. B167, 205 (1980).
- 9) E.L. Berger, T. Gottschalk and D. Sivers, Phys. Rev. D23, 99 (1981).
- 10) J. Gunion, P. Nason and R. Blankenbecler, SLAC-PUB-2937 (1982).
- 11) N.S. Craigie and P. Ratcliffe, Trieste preprint IC/82/47 (1982).

- 12) S.L. Grayson and M.P. Tuite, DAMTP 81/27 (1982).
- 13) E. Laermann, T. Walsh, I. Schmidt and P. Zerwas, SLAC-PUB-2901 (1982).
- 14) T.A. DeGrand, Y.J. Ng, and S.H.H. Tye, Phys. Rev. D16, 3251 (1977). For further discussions of higher twist terms and references, see R.K. Ellis, W. Furmanski and R. Petronzio, CERN-TH-3301 (1982); and R.L. Jaffe and M. Soldate, Phys. Lett. 105B, 467 (1981).
- 15) S.J. Brodsky, G.P. Lepage and P.M. Mackenzie, submitted to the XXith Conference on High Energy Physics, Paris (1982). See also G.P. Lepage, S.J. Brodsky, T. Huang and P.B. Mackenzie, CLNS 82/522 (1982) and the Proceedings of the Banff Summer Institute on Particles and Fields, edited by A. Capri and A.N. Kamal (Plenum Press, 1982) and Ref. 6.
- 16) See, e.g., C.H. Llewellyn Smith, Oxford University preprint 44/82 (1982).
- 17) G.R. Farrar and D.R. Jackson, Phys. Rev. Lett. 35, 1416 (1975); A.I. Vainshtain and V.I. Zakharov, Phys. Lett. 72B, 368 (1978). See also E.G. Drukaev, E.M. Levin and V.A. Rozengauz, Leningrad Nuclear Physics Institute No. 764 (1982).
- 18) E.L. Berger, S.J. Brodsky and G.P. Lepage, in preparation.
- 19) S. Gupta and H.R. Quinn, Phys. Rev. D25, 838 (1982).
- 20) J.D. Bjorken, Phys. Rev. D17, 171 (1978).
- 21) M. Suzuki, Phys. Lett. 71B, 139 (1977).
- 22) C. Peterson, D. Schlatter, I. Schmidt and P. Zerwas, SLAC-PUB-2912 (1982). D. Schlatter, to be published in the Proceedings of the 1982 SLAC Summer Institute on Particles Physics (1982).
- 23) T. DeGrand, Colorado preprint HEP-29a (1982).
- 24) R.M. Barnett and D. Schlatter, Phys. Lett. 112B, 475 (1982); R.M. Barnett, SLAC-PUB-2953 (1982) and Phys. Rev. Lett. 48, 1657 (1982); R.M. Barnett, D. Schlatter and L. Trentadue, Phys. Rev. Lett. 46, 1659 (1981).
- 25) J.J. Aubert et al., Phys. Lett. 110B, 73 (1982).

- 26) D.P. Roy, Phys. Rev. Lett. 47, 213 (1981); R.V. Gavai and D.P. Roy, Z. Phys. C10, 333 (1981). See also A. Dhar and R.M. Godpole, Tata preprint TIFR/TH/82-24.
- 27) S.J. Brodsky, P. Hoyer, C. Peterson and N. Sakai, Phys. Lett. 93B, 451 (1980); S.J. Brodsky, C. Peterson and N. Sakai, Phys. Rev. D23, 11 (1981); S.J. Brodsky and C. Peterson, SLAC-PUB-2888 (1982), and Proceedings of the Topical Conference on Forward Collider Physics, Madison, Wisconsin (1981); C. Peterson, this conference.
- 28) G. Bodwin, S.J. Brodsky and G.P. Lepage, Phys. Rev. Lett. 47, 1799 (1981); in the Proceedings of the 1981 Banff Summer School on Particles and Fields, edited by A.N. Kamal and A. Capri (Plenum Press, 1982); and SLAC-PUB-2927 (1982).
- 29) Possible nuclear effects on transverse momentum have also been considered by C. Michael and G. Wilk, Z. Phys. C10, 169 (1981). See also T. Jaroszewicz and M. Jezabek, Z. Phys. C4, 277 (1980); A. Białas and E. Białas, Phys. Rev. D21, 675 (1980).
- 30) L. Landau and I. Pomeranchuk, Dok. Akademii Nauk SSSR 92, 535 (1953), and 92, 735 (1953); L. Stodolsky, MPI-PAE/pTh 23/75 (1975).
- 31) For an interesting application of the formation zone to gluon jet fragmentation, see I.M. Dremin, Lebedev preprint 250 (1981).
- 32) The preliminary results of the EMC Collaboration [J. Nassalski et al.] are consistent with no nuclear absorption of fast hadrons in the current fragmentation region. There is very little data on associated production in the central region. See L. Hand et al., Acta Phys. Pol. B9, 1987 (1978), Crakow preprint INP-1019/PH (1978).
- 33) See, e.g., S.J. Brodsky, J.F. Gunion and J. Kuhn, Phys. Rev. Lett. 39, 1120 (1977); S.J. Brodsky, SLAC-PUB-2395, published in the Proceedings of the First Workshop on Ultra-Relativistic Nuclear Collisions, LBL (1979).
- 34) G. Bodwin, S.J. Brodsky and G.P. Lepage, in preparation.
- 35) J.D. Bjorken, Acta Phys. Pol. B2, 5 (1975); H. Harari, Proceedings of the 1971 International Symposium on Electron and Photon Interactions at High Energies, Cornell (1971).
- 36) A. Mueller, Columbia preprint (1982), to be published in the Proceedings of the Moriond Conference (1982).

- 37) K.J. Anderson et al., Phys. Rev. Lett. 43, 1219 (1979).
- 38) M. Haguenaer et al., Phys. Lett. 100B, 185 (1981).
- 39) The available SPEAR data does not show any evidence for this effect. (J. Weiss, private communication.)
- 40) W. Caswell, S.J. Brodsky and R. Horgan, Phys. Rev. D18, 2415 (1978).
- 41) Despite the theoretical complications, the great bulk of the data is well represented by models based on leading order QCD. For a comprehensive discussion, see R.D. Field, University of Florida preprint UFTP-82-17 (1982).
- 42) For applications to photoproduction see J. Bagger and J. Gunion, Phys. Rev. D25, 2287 (1982).
- 43) See the ISR reports to this meeting and M.G. Albrow et al., Nucl. Phys. B135, 461 (1978) and references therein. The observed charged correlations for K^- and \bar{p} triggers appear to be much larger than that predicted by leading order QCD processes.
- 44) A color radiation model which includes coherent effects and the $9/4$ factor for gluon jet multiplicity is given in S.J. Brodsky and J.F. Gunion, Phys. Rev. Lett. 37, 402 (1976); other specific non-Abelian effects are discussed in J.F. Gunion and G. Bertsch, Phys. Rev. D25, 746 (1982). Discussion of coherent effects in the Lund model are given by G. Gustafson, Lund preprint LU-TP-82-2 (1982) and B. Anderson, this conference.
- 45) S.J. Brodsky and G.P. Lepage, Phys. Rev. D24, 1808 (1981); P. Damgaard, Cornell preprint CLNS 81/519 (1981).
- 46) S.J. Brodsky and G.P. Lepage, SLAC-PUB-2294, published in "Quantum Chromodynamics," edited by Wm. Frazer and F. Henyey (AIP, 1979), Phys. Lett. 87B, 359 (1979); S.J. Brodsky, Y. Frishman, G.P. Lepage and C. Sachrajda, Phys. Lett. 91B, 239 (1980). See also A.V. Efremov and A.V. Radyushkin, Rev. Nuovo Cimento 3, 1 (1980), Phys. Lett. 94B, 245 (1980); A. Duncan and A. Mueller, Phys. Rev. D21, 1636 (1980), Phys. Lett. 90B, 159 (1980); G.R. Farrar and D.R. Jackson, Phys. Rev. Lett. 43, 246 (1979); V.L. Chernyak and A.R. Whitnishii, JETP Lett. 25, 11 (1977); G. Parisi, Phys. Lett. 43, 246 (1979); M.K. Chase, Nucl. Phys. B167, 125 (1980). V.N. Baier and A.G. Grozin, Nucl. Phys. B192, 476 (1981).
- 47) S.J. Brodsky and G.P. Lepage, Phys. Rev. Lett. 43, 545, 1625(E) (1979); S.J. Brodsky, G.P. Lepage and S.A.A. Zaidi, Phys. Rev. D23, 1152 (1981).

- 48) S.J. Brodsky, T. Huang and G.P. Lepage, SLAC-PUB-2540 (1980); T. Huang, SLAC-PUB-2580 (1980), published in the Proceedings of the XXth International Conference on High Energy Physics, Madison, Wisconsin (1980), and in preparation; S.J. Brodsky and S.D. Drell, Phys. Rev. D22, 2236 (1981).
- 49) E. Braaten, University of Florida preprint UFTP-82-11 (1982); F. del Aguila and M.K. Chase, Nucl. Phys. B193, 517 (1981). Higher order corrections to the distribution amplitude and its evolution equation will be discussed by S.J. Brodsky, Y. Frishman and G.P. Lepage, in preparation.
- 50) B.L. Ioffe and A.V. Smilga, Phys. Lett. 114B, 353 (1982) and references therein.
- 51) See, e.g., A. Capella and J. Tran Thanh Van, Orsay preprint LPTHE 82/2 (1982), and U. P. Sukhatme, review talk presented to this conference.
- 52) R. Blankenbecler and S.J. Brodsky, Phys. Rev. D10, 2973 (1974).
- 53) S.J. Brodsky and J.F. Gunion, Phys. Rev. D17, 848 (1978).
- 54) L. Von Hove, Proceedings of the 12th International Symposium on Multiparticle Dynamics, Notre Dame (1981); J.F. Gunion, Phys. Lett. 88B, 150 (1979), and Moriond Conference, Vol. 1:151 (1980); R.C. Hwa, Oregon preprint OITS-194 (1982), and this conference.
- 55) S.J. Brodsky and G.P. Lepage, Phys. Rev. D24, 2848 (1981).
- 56) E.L. Berger and S.J. Brodsky, Phys. Rev. D24, 2428 (1981).
- 57) This point has also been emphasized by A. Mueller, to be published in the Proceedings of the 1982 Moriond Conference.
- 58) G. Bertsch, S.J. Brodsky, A.S. Goldhaber and J.F. Gunion, Phys. Rev. Lett. 47, 297 (1981).
- 59) S.J. Brodsky, G. Bodwin and G.P. Lepage, SLAC-PUB-2966 (1982), and these Proceedings: A Form of "Weak Factorization" for QCD has been proposed by J.C. Collins, D.E. Soper and G. Sterman, Phys. Lett. 109B, 388 (1982).
- 60) A. Bassetto, M. Ciafaloni, G. Marchesini and A. Mueller, SNS 2/1982 (1982).

- 61) W. Ochs, Max-Planck-Institute (Munich) preprint MPI-PAE/Exp E1-97 (1981), and this conference.
- 62) S.J. Brodsky, T.A. DeGrand and R. Schwitters, Phys. Lett. 79B, 255 (1978). See also I. Dremin, Ref. 31.
- 63) R.D. Field, G.C. Fox and R.L. Kelly, University of Florida preprint UFTP-26-12; and R.D. Field, this conference.
- 64) N. Paver and D. Treleani, Nuovo Cimento A70, 215 (1982); H.D. Politzer, Nucl. Phys. B172, 349 (1980); S.J. Brodsky and J.F. Gunion (unpublished).
- 65) R. Horgan and M. Jacob, Phys. Lett. 107B, 395 (1981).
- 66) J.F. Donoghue and E. Golowich, Phys. Rev. D15, 3421 (1977).
- 67) B. Anderson and G. Gustafson, Ref. 44.
- 68) A discussion of QCD effects in nuclear physics is given in Ref. 6 and S.J. Brodsky, IAS preprint (1982), to be published in the Proceedings of the Conference "New Horizons in Electromagnetic Physics," University of Va (1982).