

# Measurements of $J/\psi$ Decays Into a Vector and a Pseudoscalar Meson<sup>†</sup>

D. Coffman, G.P. Dubois, G. Eigen, J. Hauser, D.G. Hitlin, C.G. Matthews, J.D. Richman,  
W.J. Wisniewski, Y. Zhu

*California Institute of Technology, Pasadena, CA 91125*

M. Burchell, D.E. Dorfan, J. Drinkard, C. Gatto, R.P. Hamilton,<sup>‡</sup> C.A. Heusch, L. Köpke,  
W.S. Lockman, R. Partridge, J. Perrier, H.F.W. Sadrozinski, M. Scarletella,  
T.L. Schalk, A. Seiden, A.J. Weinstein, R. Xu

*University of California at Santa Cruz, Santa Cruz CA 95064*

J.J. Becker, G.T. Blaylock, B.I. Eisenstein, T. Freese, G. Gladding, J. M. Izen, S.A. Plaetzer,  
C. Simopoulos, A.L. Spadafora, I.E. Stockdale, J.J. Thaler, B. Tripsas

*University of Illinois at Urbana-Champaign, Urbana, IL 61801*

U. Mallik

*University of Iowa, Iowa City, IA 52242*

J. Adler, T. Bolton, J. C. Brient, K.O. Bunnell, R.E. Cassell, D.H. Coward, K.F. Einsweiler,  
C. Grab, R.F. Mozley, A. Odian, D. Pitman, R.H. Schindler, W. Stockhausen,  
W. Toki, F. Villa, S. Wasserbaech, N. Wermes, D. Wisinski

*Stanford Linear Accelerator Center, Stanford, CA 94309*

J.S. Brown, T.H. Burnett, V. Cook, A.L. Duncan, A. Li, R. Mir, P.M. Mockett, B. Nemati,  
L. Parrish, H.J. Willutzki

*University of Washington, Seattle, WA 98195*

MARK III Collaboration

---

<sup>†</sup> Work supported in part by the Department of Energy, under contracts DE-AC03-76SF00515, DE-AC02-76ER01195, DE-AC03-81ER40050, DE-AM03-76SF00034, and by the National Science Foundation.

<sup>‡</sup> Deceased.

## ABSTRACT

We present measurements of the two-body decays of the  $J/\psi$  into a vector and a pseudoscalar meson. The data, taken with the Mark III detector at the SLAC  $e^+e^-$  storage ring SPEAR, consist of  $5.8 \times 10^6$  produced  $J/\psi$ 's. The branching ratios for the  $J/\psi$  decays into  $\rho\pi, \rho\eta, \rho\eta', \omega\pi^0, \omega\eta, \omega\eta', \phi\eta, \phi\eta'$ , and  $K^*K$  are measured; an upper limit on  $J/\psi \rightarrow \phi\pi^0$  is obtained. Using the measured branching ratios we obtain parameters of a phenomenological model of  $J/\psi$  decays, indicating that the  $\eta$  and  $\eta'$  are consistent with being composed only of light and strange quarks. The model is used to obtain the mixing angle in the pseudoscalar nonet. The  $\omega\pi^0$  electromagnetic form factor is determined. The upper limit on  $J/\psi \rightarrow \phi\pi^0$  is used to study the contributions from electromagnetic doubly-OZI suppressed decays of the  $J/\psi$ .

## I. INTRODUCTION

Decays of the  $J/\psi$  into vector-pseudoscalar ( $VP$ ) meson pairs are expected to occur mainly by two mechanisms: strong decays via three gluon annihilation (Fig. 1a), and electromagnetic decays through one virtual photon (Fig. 1b). Electromagnetic contributions to  $J/\psi$  decays are important since the strong decays are Okubo-Zweig-Iizuka (OZI) suppressed.<sup>[1]</sup> We can therefore measure the isospin violating electromagnetic decays such as  $J/\psi \rightarrow \omega\pi^0, \rho\eta$  and  $\rho\eta'$  with a sensitivity comparable to the strong decays. Diagrams with more disconnected quark lines can also contribute to the decay rates. An example is the doubly-OZI-suppressed (DOZI) diagram (Fig. 1c). Our recent analysis of  $J/\psi \rightarrow \gamma\omega\phi$  decays<sup>[2]</sup> shows that DOZI processes contribute to  $J/\psi$  decay rates. The lack of flavor correlation between the final state mesons in the isospin violating decay  $J/\psi \rightarrow \phi\pi^0$  requires that the decay proceed via the diagram shown in Fig. 1d. We can thus estimate the importance of this electromagnetic DOZI-suppressed diagram.

A full set of measurements of  $J/\psi \rightarrow VP$  decays enables us to test the

processes involved. We can study the contributions of the different amplitudes to the decay rates, the mixing angle in the pseudoscalar meson nonet and the quark content of its members. The pseudoscalar mixing angle and the quark content of the  $\eta$  and  $\eta'$  mesons reflect any additional contributions (e.g. gluonium or radial excitations) to the meson wavefunctions.<sup>[3]</sup>

Measurements of purely electromagnetic  $J/\psi$  decays can be used to calculate the electromagnetic form factors involved. These form factors are used to test QCD inspired models of mesonic wave functions.<sup>[4]</sup> We have previously calculated electromagnetic form factors in the decays of the  $J/\psi$  to a pair of pseudoscalar mesons,<sup>[5]</sup> and qualitative predictions have been made for their magnitude relative to the electromagnetic form factors in  $J/\psi \rightarrow VP$  decays.<sup>[6]</sup> Assuming that single photon annihilation is the dominant process in the  $J/\psi \rightarrow \omega\pi^0$  decay, its electromagnetic form factor may be calculated by comparing the rate for this decay with that for the crossed channel process  $\omega \rightarrow \gamma\pi^0$ .

We report herein measurements of the branching ratios for  $J/\psi$  decays to all vector ( $\rho, \omega, \phi$  and  $K^*$ ) and pseudoscalar ( $\pi, K, \eta$  and  $\eta'$ ) meson pairs. The data set was taken with the Mark III detector at the SLAC  $e^+e^-$  storage ring SPEAR, at a center of mass energy equal to  $m_{J/\psi}$ . The data, consisting of  $5.8 \times 10^6$  produced  $J/\psi$ 's, were acquired in sets of  $0.9 \times 10^6$ ,  $1.8 \times 10^6$  and  $3.1 \times 10^6$   $J/\psi$ 's. The number of  $J/\psi$ 's in the second data set was found by studying events with 0,1,2,3 and  $> 3$  charged tracks, where the trigger efficiency was determined from  $J/\psi$  decays in events of the type  $\psi' \rightarrow \pi^+\pi^-\psi$ .<sup>[5]</sup> The  $J/\psi$  content of the other two data sets was found by measuring the relative rate of the  $J/\psi \rightarrow \mu^+\mu^-$  decay in each period. The combined sample represents a two-fold increase over the data ( $2.7 \times 10^6$ ) used in our earlier analysis of  $J/\psi \rightarrow VP$  decays.<sup>[7]</sup>

In the next section we discuss the general features of event selection. The analysis features specific to each decay are presented in Section III. The measured branching ratios are given in Table I. In Section IV we investigate the physical implications of our measurements. The set of measured branching ratios is used

to fit the parameters in a model of  $J/\psi$  decays which includes strong, electromagnetic and DOZI-suppressed amplitudes. This model<sup>[8]</sup> also allows for violation of SU(3) invariance and includes a parametrization of the  $\eta$  and  $\eta'$  wavefunctions in terms of light and strange quarks. The mixing angle in the pseudoscalar nonet and the quark contents of the  $\eta$  and  $\eta'$  are calculated. The strength of the electromagnetic DOZI-suppressed processes is estimated. A calculation of the  $\omega\pi^0$  electromagnetic form factor is performed, details of which are given in the Appendix.

## II. EVENT SELECTION

The Mark III detector has been described elsewhere.<sup>[9]</sup> A brief survey follows: Photons are detected as showers in a finely segmented electromagnetic calorimeter which covers 95% of the full  $4\pi$  solid angle, and is fully efficient for photon energies above 0.1 GeV. The angular resolution is 10 mrad, and the energy resolution is  $\sigma_E/E = 0.17/\sqrt{E}$  ( $E$  in GeV). Charged tracks are detected in the drift chamber which covers 85% of the solid angle, and has a momentum resolution of  $\sigma_p/p = 0.015\sqrt{1+p^2}$  ( $p$  in GeV/ $c$ ). The time-of-flight (TOF) system covers 80% of  $4\pi$ , and has a mean resolution of 200 ps, providing  $3\sigma$   $K/\pi$  discrimination up to 0.8 GeV/ $c$ .

The analyses of the individual channels have common features. Vector mesons are detected in the modes  $\rho \rightarrow \pi^+\pi^-$ ,  $\omega \rightarrow \pi^+\pi^-\pi^0$ ,  $\phi \rightarrow K^+K^-$ , and  $K^* \rightarrow K\pi$ . Pseudoscalar mesons are detected in the modes  $\pi^0 \rightarrow \gamma\gamma$ ,  $\eta \rightarrow \gamma\gamma$ ,  $\eta \rightarrow \pi^+\pi^-\gamma$ ,  $\eta \rightarrow \pi^+\pi^-\pi^0$ ,  $\eta' \rightarrow \gamma\rho$ , and  $\eta' \rightarrow \eta\pi^+\pi^-$ . Observing the  $\eta$  and  $\eta'$  decays in more than one mode allows us a cross-check of our measurements. Selection criteria on the invariant mass of a  $\gamma\gamma$  pair forming a  $\pi^0$  or an  $\eta$  differ in various decay modes. The criteria are more stringent when large backgrounds need to be suppressed. The effect on the branching ratios is included in the systematic error.

Each analysis requires events to have the correct number of charged tracks with the sum of their charges equal to zero, and at least the required number

of showers. Spurious photon candidates, arising from the interactions of charged tracks in the shower counters, are minimized by rejecting showers which fall near a charged track and by imposing a minimum-energy requirement.

The selected events are kinematically fitted by imposing energy and momentum constraints. The kinematic fit adjusts the track energies and momenta within the measured errors and detector resolution, so as to satisfy the given event hypothesis. This improves the photon energy measurement and suppresses events with the wrong charged particle assignment. When the number of showers in an event exceeds the minimum, the showers are ordered in decreasing energy, all combinations with up to three additional showers are fitted, and the combination with the minimum  $\chi^2$  is retained.

Scatter plots of the invariant mass of the vector meson candidate versus the invariant mass of the pseudoscalar meson candidate are formed. Invariant mass combinations which fall into the range of the  $V(P)$  meson are selected. The recoiling  $P(V)$  mass spectrum is fitted using the sum of a polynomial background and a Breit-Wigner (BW) curve (width obtained from Ref. 10) convolved with a Gaussian resolution function. The number of  $P(V)$  mesons recoiling against  $V(P)$  mass sidebands is obtained by a similar fit, and is used for background subtraction.

The detection efficiency and mass resolution for each channel are found by a Monte Carlo simulation. This includes the matrix element for the  $J/\psi \rightarrow VP$  decays which completely specifies the final state angular distributions.<sup>[11]</sup> The interactions of photons in the detector are simulated using both detailed shower simulation programs (EGS),<sup>[12]</sup> and real photons from fully reconstructed  $J/\psi$  events.

The fractional systematic error for each branching ratio is obtained by adding contributions from the following terms in quadrature: 8.5% from the  $J/\psi$  flux, 1% from the reconstruction of each charged track, 2% from the reconstruction of each photon, 5% from the kinematic fit, 5% to 10% from the selection criteria,

and 5% to 10% resulting from the uncertainty on the background subtraction.

### III. ANALYSIS

A discussion of the analysis of each final state follows. The resulting branching ratios are given in Table I. Unless specified, the  $\pi^0$  and  $\eta$  in the final state are identified in the  $\gamma\gamma$  mode. On each result the first error is statistical and the second systematic. The average is a weighted mean of the different decay modes. The upper limit is given at the 90% confidence level.

#### A. $J/\psi \rightarrow \rho\pi$

The decay  $J/\psi \rightarrow \rho\pi$  is observed in the  $\pi^+\pi^-\gamma\gamma$  final state. The  $\gamma\gamma$  invariant mass is required to be consistent with a  $\pi^0$  (0.07 to 0.20 GeV/ $c^2$ ). This removes virtually all QED backgrounds, as well as  $\pi^+\pi^-\gamma$  events with an additional spurious shower. A Dalitz plot of one third of the selected events is shown in Fig. 2. The  $\rho\pi$  bands are dominant.<sup>[18]</sup> A study of the population in the Dalitz plot away from the  $\rho\pi$  bands shows no evidence for a population uniform in phase space, or for contributions from other resonances. Possible background from another source (i.e.  $J/\psi \rightarrow K^*K \rightarrow K^+K^-\pi^0$ , with the kaons misidentified as pions) has been estimated by Monte Carlo simulation, and its expected contribution of  $(3.2 \pm 0.4)\%$  has been subtracted from the data. We find the ratio  $\rho^0\pi^0/\rho\pi$  to be  $0.328 \pm 0.005 \pm 0.027$ , in agreement with the value of one third expected from isospin symmetry (see Section IV A).

#### B. $J/\psi \rightarrow \rho\eta$

The decay  $J/\psi \rightarrow \rho\eta$  is observed in two decay modes of the  $\eta$ . For the  $\eta \rightarrow \gamma\gamma$  mode, events in the final state  $\pi^+\pi^-\gamma\gamma$  are kinematically fitted to the hypothesis  $J/\psi \rightarrow \eta\pi^+\pi^-$ , where the requirement of the  $\eta$  mass adds an additional constraint to the fit. The  $\pi^+\pi^-$  mass distribution recoiling against the  $\eta$  is shown in Fig. 3a.

For the  $\eta \rightarrow \pi^+\pi^-\pi^0$  mode, the final state  $\pi^+\pi^-\pi^+\pi^-\gamma\gamma$  is selected. The  $\gamma\gamma$  invariant mass is required to be consistent with a  $\pi^0$  (0.11 to 0.16 GeV/ $c^2$ ).

All  $\pi^+\pi^-\pi^0$  mass combinations consistent with an  $\eta$  (0.53 to 0.57 GeV/ $c^2$ ) are considered. Only one combination per event is found in the  $\eta$  mass region. The  $\pi^+\pi^-$  mass distribution recoiling against the  $\eta$  is shown in Fig. 3b.

To obtain the number of  $J/\psi \rightarrow \rho\eta$  events, the interference from the decay  $J/\psi \rightarrow \omega\eta$ ,  $\omega \rightarrow \pi^+\pi^-$  is taken into account. Although the  $\omega$  decays to  $\pi^+\pi^-$  with a branching ratio of only 1.7%,<sup>[10]</sup> the branching ratio for  $J/\psi \rightarrow \omega\eta$  is one order of magnitude larger than that for  $J/\psi \rightarrow \rho\eta$  (Table I). Interference effects are therefore expected to be significant. The data in Figs. 3a and 3b are fitted with the function  $N(M_{\pi^+\pi^-}) = L(M_{\pi^+\pi^-}) + |A_\rho(M_{\pi^+\pi^-}) + A_\omega(M_{\pi^+\pi^-}) \cdot e^{i\phi}|^2$ , where  $L$  is a polynomial background term and  $A_\rho$  is the contribution of the  $\rho\eta$  channel to the data. The amplitude  $A_\rho = (N_\rho)^{\frac{1}{2}} \cdot F_\rho^{BW}(M_{\pi^+\pi^-})$ , where  $F_\rho^{BW}$  is the appropriate Breit-Wigner term.<sup>[13]</sup> The amplitude  $A_\omega$  is similarly defined. In the fitting procedure,  $N_\rho$ ,  $N_\omega$  and  $\phi$  (the relative phase) are all left free. For the  $\eta \rightarrow \gamma\gamma$  mode we find  $\phi = (46 \pm 15)^\circ$ , the branching ratio for  $J/\psi \rightarrow \rho\eta$  given in Table I, and a branching ratio for  $J/\psi \rightarrow \omega\eta$  of  $(1.35 \pm 0.14 \pm 0.34) \times 10^{-3}$ , in reasonable agreement with that measured in the  $\omega \rightarrow \pi^+\pi^-\pi^0$  mode (Table I). The fit results for the  $\eta \rightarrow \pi^+\pi^-\pi^0$  mode agree within errors.

### C. $J/\psi \rightarrow \rho\eta'$

The decay  $J/\psi \rightarrow \rho\eta'$  is observed in two decay modes of the  $\eta'$ . For the  $\eta' \rightarrow \gamma\rho$  mode, the final state  $\pi^+\pi^-\pi^+\pi^-\gamma$  is selected. Background coming from  $J/\psi \rightarrow \pi^+\pi^-\pi^+\pi^-\pi^0$  is suppressed by rejecting events consistent with the five-pion hypothesis, as determined using a 5C fit. Figure 4a shows the invariant mass of the  $\gamma\rho$  ( $0.50 < M_{\pi^+\pi^-} < 0.85$  GeV/ $c^2$ )<sup>[14]</sup> recoiling against the other  $\rho$  ( $0.60 < M_{\pi^+\pi^-} < 0.90$  GeV/ $c^2$ ) in the event. All possible  $\pi^+\pi^-$  combinations consistent with two  $\rho$ 's are considered. Up to four entries per event are allowed. A clear  $\eta'$  signal is apparent over a large background. The background comes from wrong  $\pi^+\pi^-$  combinations, from the radiative decay  $J/\psi \rightarrow \gamma\rho\rho$ ,<sup>[15]</sup> and from residual  $J/\psi \rightarrow \pi^+\pi^-\pi^+\pi^-\pi^0$  events. The  $\gamma\rho$  events recoiling against  $\rho$  mass sidebands ( $0.2 < M_{\pi^+\pi^-} < 0.4$  and  $1.2 < M_{\pi^+\pi^-} < 1.5$  GeV/ $c^2$ ) show no

significant  $\eta'$  signal.

For the  $\eta' \rightarrow \eta\pi^+\pi^-$  mode, the final state  $\pi^+\pi^-\pi^+\pi^-\gamma\gamma$  is selected. The  $\gamma\gamma$  mass is required to be consistent with an  $\eta$  (0.53 to 0.57 GeV/ $c^2$ ). All  $\pi^+\pi^-$  mass combinations consistent with a  $\rho$  (0.60 to 1.00 GeV/ $c^2$ ) are considered. Figure 4b shows the invariant mass of the  $\eta\pi^+\pi^-$  events recoiling against the  $\rho$ . Only one combination per event is found in the  $\eta'$  mass region. A clear  $\eta'$  signal is apparent, well separated from background. The  $\rho$  mass sidebands show no significant  $\eta'$  signal.

The possible contribution to  $\rho\eta'$  from  $\omega\eta'$ ,  $\omega \rightarrow \pi^+\pi^-$  is estimated to be  $(5 \pm 3)\%$  using our measured rate for  $J/\psi \rightarrow \omega\eta'$  given in Table I. The data is corrected for this effect.

#### D. $J/\psi \rightarrow \omega\pi^0$ and $J/\psi \rightarrow \omega\eta$

The decays  $J/\psi \rightarrow \omega\pi^0$  and  $J/\psi \rightarrow \omega\eta$  are observed in the  $\pi^+\pi^-\gamma\gamma\gamma\gamma$  final state. If at least one  $\pi^0$  is found from any  $\gamma\gamma$  pair ( $0.10 < M_{\gamma\gamma} < 0.17$  GeV/ $c^2$ ), the  $\pi^+\pi^-\pi^0$  invariant mass (Fig. 5a) is formed. Events with  $0.73 < M_{\pi^+\pi^-\pi^0} < 0.83$  GeV/ $c^2$  are selected as  $\omega$  candidates. This procedure is the same in all  $J/\psi \rightarrow \omega P$  channels. The  $\gamma\gamma$  invariant mass recoiling against the  $\omega$  is shown in Fig. 5b. Clear  $\pi^0$  and  $\eta$  signals are apparent. The  $\gamma\gamma$  combinations recoiling against  $\omega$  mass sidebands ( $0.66 < M_{\pi^+\pi^-\pi^0} < 0.72$  and  $0.85 < M_{\pi^+\pi^-\pi^0} < 0.90$  GeV/ $c^2$ ) are used for background subtraction.

The decay  $J/\psi \rightarrow \omega\eta$  is also observed in the  $\pi^+\pi^-\pi^+\pi^-\gamma\gamma\gamma$  final state. The  $\gamma\pi^+\pi^-$  invariant mass recoiling against the  $\omega$  is shown in Fig. 6a. A clear  $\eta$  signal is seen. Curve I in Fig. 6a shows the Monte Carlo simulation of this signal. The branching ratio is determined by taking the total number of events in the  $\eta$  mass region ( $0.50 < M_{\gamma\pi^+\pi^-} < 0.60$  GeV/ $c^2$ ) and subtracting both an estimate of the phase space background, and the contribution from  $\eta \rightarrow \pi^+\pi^-\pi^0$ , where one of the photons from the  $\pi^0$  decay is not detected. This contribution, which populates the mass region from 0.35 to 0.50 GeV/ $c^2$ , is well reproduced

by Monte Carlo simulation (curve II in Fig. 6a). The  $\gamma\pi^+\pi^-$  events recoiling against  $\omega$  mass sidebands are used for background subtraction.

#### E. $J/\psi \rightarrow \omega\eta'$

The decay  $J/\psi \rightarrow \omega\eta'$  is observed in two decay modes of the  $\eta'$ . For the  $\eta' \rightarrow \gamma\rho$  mode the final state  $\pi^+\pi^-\pi^+\pi^-\gamma\gamma\gamma$  is selected. The  $\gamma\pi^+\pi^-$  invariant mass recoiling against the  $\omega$  is shown in Fig. 6b, where the  $\pi^+\pi^-$  invariant mass is required to be consistent with a  $\rho$  ( $0.50 < M_{\pi^+\pi^-} < 0.90 \text{ GeV}/c^2$ ). The  $\omega$  mass sidebands show no significant  $\eta'$  signal.

For the  $\eta' \rightarrow \eta\pi^+\pi^-$  decays, two decay modes of the  $\eta$  are observed. In the  $\eta \rightarrow \gamma\gamma$  mode the final state  $\pi^+\pi^-\pi^+\pi^-\gamma\gamma\gamma\gamma$  is selected. Following the  $\omega$  selection, the other  $\gamma\gamma$  invariant mass is required to be consistent with an  $\eta$  ( $0.52$  to  $0.57 \text{ GeV}/c^2$ ). The  $\eta\pi^+\pi^-$  invariant mass recoiling against the  $\omega$  is shown in Fig. 6c. The  $\omega$  mass sidebands are used for background subtraction. In the  $\eta \rightarrow \pi^+\pi^-\pi^0$  mode the final state  $\pi^+\pi^-\pi^+\pi^-\pi^+\pi^-\gamma\gamma\gamma\gamma$  is selected. Following the  $\omega$  selection, the other  $\gamma\gamma$  invariant mass is required to be consistent with a  $\pi^0$  ( $0.08$  to  $0.20 \text{ GeV}/c^2$ ), and the  $\pi^+\pi^-\pi^0$  invariant mass is required to be consistent with an  $\eta$  ( $0.53$  to  $0.57 \text{ GeV}/c^2$ ). The  $\eta\pi^+\pi^-$  invariant mass recoiling against the  $\omega$  is shown in Fig. 6d. The number of  $\omega\eta'$  events is obtained by subtracting an estimated phase space background. The  $\omega$  mass sidebands are used for background subtraction.

#### F. $J/\psi \rightarrow \phi\pi^0$ and $J/\psi \rightarrow \phi\eta$

The decays  $J/\psi \rightarrow \phi\pi^0$  and  $J/\psi \rightarrow \phi\eta$  are searched for in the  $K^+K^-\gamma\gamma$  final state. The kaons have relatively low momentum and are easily separated from pions by the TOF system. The invariant mass of the two kaons recoiling against the  $\eta$  is shown in Fig. 7a. Events with  $1.005 < M_{K^+K^-} < 1.035 \text{ GeV}/c^2$  are selected to be  $\phi$  candidates. This procedure is used in all  $J/\psi \rightarrow \phi P$  channels. The  $\gamma\gamma$  invariant mass recoiling against the  $\phi$  is shown in Fig. 7b. A clear  $\eta$  signal is seen. The  $\gamma\gamma$  events recoiling against  $\phi$  mass sidebands ( $0.98 < M_{K^+K^-} < 1.00$  and  $1.05 < M_{K^+K^-} < 1.09 \text{ GeV}/c^2$ ) are used for background subtraction.

No  $\pi^0$  signal is visible. An analysis of the population distribution in a scatter plot of  $M_{K^+K^-}$  vs.  $M_{\gamma\gamma}$  predicts  $3.5 \pm 0.5$  background events in the region of the  $\pi^0$  mass. Two events are observed in this region. A 90% confidence level upper limit on the  $B(J/\psi \rightarrow \phi\pi^0)$  is given in Table I.<sup>[16]</sup>

The decay  $J/\psi \rightarrow \phi\eta$  is also seen in the  $K^+K^-\pi^+\pi^-\gamma\gamma$  final state. Following the  $\phi$  and  $\pi^0$  ( $0.08 < M_{\gamma\gamma} < 0.20$  GeV/ $c^2$ ) selection, the  $\pi^+\pi^-\pi^0$  invariant mass is shown in Fig. 8a. The  $\phi$  mass sidebands are used for background subtraction.

#### G. $J/\psi \rightarrow \phi\eta'$

The decay  $J/\psi \rightarrow \phi\eta'$  is observed in two decay modes of the  $\eta'$ . For the  $\eta' \rightarrow \gamma\rho$  mode the final state  $K^+K^-\pi^+\pi^-\gamma$  is selected. Following the  $\phi$  and  $\rho$  ( $0.50 < M_{\pi^+\pi^-} < 0.90$  GeV/ $c^2$ ) selection, the  $\gamma\rho$  invariant mass recoiling against the  $\phi$  is shown in Fig. 8b. The  $\phi$  mass sidebands show no  $\eta'$  signal.

For the  $\eta' \rightarrow \eta\pi^+\pi^-$  mode the final state  $K^+K^-\pi^+\pi^-\gamma\gamma$  is selected. Following the  $\phi$  and  $\eta$  selection ( $0.50 < M_{\gamma\gamma} < 0.60$  GeV/ $c^2$ ), the  $\eta\pi^+\pi^-$  invariant mass recoiling against the  $\phi$  is shown in Fig. 8c. The number of  $\eta'$  events is taken as the number of entries in Fig. 8c, there being no background below 1.1 GeV/ $c^2$ . The  $\phi$  mass sidebands are used for background subtraction.

#### H. $J/\psi \rightarrow K^*K$

The decay  $J/\psi \rightarrow K^*K$  is observed in the channels  $K^{*\pm}K^\mp$  ( $K^{*\pm} \rightarrow K^\pm\pi^0$ ,  $K^{*\pm} \rightarrow K_S^0\pi^\pm$ ) and  $K^{*0}\bar{K}_S^0$  ( $K^{*0} \rightarrow K^\pm\pi^\mp$ ). The  $K_S^0$  is seen in the  $\pi^+\pi^-$  decay mode. Events are required to have at least one kaon identified by the TOF system. In the  $K^+K^-\pi^0$  mode events are rejected if they have a charged track unambiguously identified as a pion, or if the  $K^+K^-$  invariant mass is consistent with a  $\phi$ . The latter criteria rejects only events with  $M_{K^\pm\pi^0} > 1.4$  GeV/ $c^2$ , leaving the  $K^*K$  signal unaffected. All the final states are shown in the Dalitz plots of Fig. 9. To obtain the number of  $K^*K$  events, projections onto the mass axes are taken, and a BW curve with a quadratic background is fitted to the data.

Even though the 4C kinematic fit to  $J/\psi \rightarrow \gamma\gamma K^+K^-$  rejects  $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$  events, a sizeable contamination from  $J/\psi \rightarrow \rho\pi$  remains in the data. Such events are indistinguishable from  $J/\psi \rightarrow K^*K \rightarrow K^+K^-\pi^0$ , and are therefore simulated by Monte Carlo. The predicted contribution which amounts to  $(28\pm 4)\%$  of the events in the signal region is subtracted. The resulting branching ratio for  $J/\psi \rightarrow K^{*\pm}K^\mp$  in the  $K^+K^-\pi^0$  final state, given in Table I, is in good agreement with that measured in the  $K^\pm K_S^0\pi^\mp$  final state. As a further check on the accuracy of the  $\rho\pi$  subtraction the analysis of the  $K^+K^-\pi^0$  final state is repeated with more stringent TOF selection criteria. These criteria require both charged tracks in an event to be identified as kaons, with the momentum of the kaon originating from the  $K^*$  decay to be  $< 0.8 \text{ GeV}/c$ . These criteria reduce the detection efficiency from 20% to 4.8%. The resulting branching ratio for  $J/\psi \rightarrow K^{*\pm}K^\mp$  is  $(5.57 \pm 0.25 \pm 0.78) \times 10^{-3}$ . This is in good agreement with the result in Table I, albeit with a larger statistical error.

#### IV. DISCUSSION OF RESULTS

The branching ratios  $J/\psi \rightarrow VP$  are given in Table I. The results show a greater statistical accuracy than the previous Mark III measurements,<sup>[7]</sup> reflecting the increase in the number of  $J/\psi$  decays studied. The previous limit on  $J/\psi \rightarrow \rho\eta'$ ,  $\eta' \rightarrow \eta\pi^+\pi^-$  is superceded by a measurement using two decay modes of the  $\eta'$ . Due to the limited statistics upon which the previous result was based (three events in the  $\eta'$  mass region) the background estimation was difficult. The current luminosity and improvements in the analysis method (raising the detection efficiency from 9% to 21%) provide us with a clear signal over a well estimated background. The rate for  $J/\psi \rightarrow \omega\eta'$ , previously measured in two modes (one with only four events) is now measured in three modes which yield consistent branching ratios, although the value for the  $\eta' \rightarrow \gamma\rho$  mode is smaller than the previous measurement by 3.7 times the previous systematic error. The branching ratio for  $J/\psi \rightarrow \omega\pi^0$  measured here is lower than the old result by 1.7 times the previous systematic error, due to a better estimate of the

background obtained by fitting the recoiling mass distribution rather than the recoiling momentum spectrum.

Except for the above mentioned changes, the set of new branching ratios given in Table I agree with the previous measurements within  $1\sigma$  of the previous statistical error.

The decays of the  $J/\psi$  (being a  $c\bar{c}$  quark state with well defined initial quantum numbers) are an excellent laboratory for the study of hadron dynamics. The electromagnetic decays play an important role here due to the OZI suppression of hadronic decays via three gluons. This allows us to study  $J/\psi \rightarrow VP$  decay rates in terms of the diagrams shown in Fig. 1a and 1b. The influence of diagrams with more disconnected quark lines, such as the DOZI processes shown in Fig. 1c and 1d, is also investigated. Further, given that the vector meson quark assignments are well established, our measurements are used to study the quark content of the  $\eta$  and  $\eta'$  pseudoscalar mesons.

#### A. The mixing angle of the $\eta$ and $\eta'$ mesons

The widths of  $J/\psi$  into  $VP$  meson pairs can be written as

$$\Gamma(J/\psi \rightarrow VP)_i = |A_i|^2 \times p_{V_i}^3,$$

where  $p_{V_i}$  is the momentum of the vector meson in the rest frame of the  $J/\psi$ .

We analyze our data using a recent phenomenological model,<sup>[8]</sup> which differs from that used to analyze our earlier measurements by including a term representing DOZI processes and that no gluonium contribution is required. Within this model the amplitudes  $A_i$  are coupled with strength  $g$  for the strong decays,  $rg$  for DOZI processes,  $sg$  for possible SU(3) violations, and  $e$  for electromagnetic processes. These constants are relatively real, and the phase of  $e$  relative to  $g$  is defined to be  $\theta_e$ .

Within the pseudoscalar nonet the  $\eta$  and  $\eta'$  mesons can be parametrized by their normalized wavefunctions in terms of only light and strange quarks:<sup>[8]</sup>

$$|\eta\rangle = X_\eta \frac{1}{\sqrt{2}} |u\bar{u} + d\bar{d}\rangle + Y_\eta |s\bar{s}\rangle, \quad |\eta'\rangle = X_{\eta'} \frac{1}{\sqrt{2}} |u\bar{u} + d\bar{d}\rangle + Y_{\eta'} |s\bar{s}\rangle.$$

The pseudoscalar mixing angle  $\theta_p$  can be written in terms of the quark content of the  $\eta$  and  $\eta'$  mesons:<sup>[17]</sup>

$$\tan \theta_p = -\frac{\sqrt{2}X_\eta + Y_\eta}{X_\eta - \sqrt{2}Y_\eta} = \frac{X_{\eta'} - \sqrt{2}Y_{\eta'}}{\sqrt{2}X_{\eta'} + Y_{\eta'}}.$$

The expected contributions of the various processes to the  $J/\psi \rightarrow VP$  amplitudes in the model are given in Table II. The measured reduced branching ratios  $\tilde{B} \equiv B/p_V^3$  are also given in Table II, where common systematic errors due to the normalization have been removed, and the statistical and systematic errors are added in quadrature. The predicted decay amplitudes for  $J/\psi$  decays to the variously charged  $\rho\pi$  states are equal, reflecting the similar  $u\bar{u}, d\bar{d}$  quark nature of these states. Our results (Section III A) are in good agreement with this prediction. By contrast, the amplitudes for  $J/\psi$  decays to neutral and charged  $K^*K$ 's differ due to the different quark content of the neutral ( $d\bar{s}$ ) and charged ( $u\bar{s}$ ) states. This leads to differing electromagnetic contributions to the amplitudes. The necessity for such an isospin violating effect has been noted by several authors.<sup>[18]</sup> Our model predicts that the ratio of neutral over charged  $K^*K$ 's from  $J/\psi \rightarrow K^*K$  should be less than 1. From our results in Table I we find this ratio to be  $0.82 \pm 0.05 \pm 0.09$ , in qualitative agreement with the prediction.

We fit the phenomenological model to the ten reduced branching ratios with the constraints  $X_i^2 + Y_i^2 = 1$  and  $X_{\eta'} = -Y_\eta$ . The fit yields (with a  $\chi^2$  of 10.1 for 4 degrees of freedom)

$$X_\eta = 0.814 \pm 0.014; \quad g = 1.10 \pm 0.04; \quad s = 0.12 \pm 0.03;$$

$$\theta_e = (72 \pm 7)^\circ; \quad e = 0.12 \pm 0.01; \quad r = -0.15 \pm 0.01.$$

This gives  $Y_\eta = -0.581$  and  $\theta_p = (-19.2 \pm 1.4)^\circ$ . The result for  $\theta_p$  is in good agreement with recent two-photon results.<sup>[19]</sup>

A fit to the data, without the above constraints on  $X_i$  and  $Y_i$  yields (with a  $\chi^2$  of 0.02 for one degree of freedom)

$$X_\eta^2 + Y_\eta^2 = 1.00 \pm 0.16 \quad \text{and} \quad X_{\eta'}^2 + Y_{\eta'}^2 = 1.44 \pm 0.26.$$

The values of all the components describing the amplitudes are within the errors of the previous fit results. Varying the magnitudes of the reduced branching ratios to account for possible correlated systematic effects in photon reconstruction efficiencies changes the results for  $X_i^2 + Y_i^2$  within the above errors.

The results of the unconstrained fit indicate that the model where the  $\eta$  and  $\eta'$  consist of only  $u$ ,  $d$  and  $s$  quarks, with no contribution from gluonium, is consistent with the data. Furthermore the results of the constrained fit are consistent with mixing between the ground state octet and singlet states only.

The constrained fit yields a DOZI contribution of 15% of the OZI amplitude. Using the same model with no DOZI contribution ( $r = 0$ ), an unconstrained fit to the data gives a  $\chi^2$  of 25.1 for two degrees of freedom. A constrained fit to the data gives a  $\chi^2$  of 221.0 for five degrees of freedom. Setting  $r = 0$  one expects  $\tilde{B}(J/\psi \rightarrow \omega\eta)/\tilde{B}(J/\psi \rightarrow \omega\eta')$  to be equal to  $\tilde{B}(J/\psi \rightarrow \rho\eta)/\tilde{B}(J/\psi \rightarrow \rho\eta')$ . The values for these ratios from Table II are 8.0 and 1.3 respectively. This illustrates the need for a DOZI contribution in this model.

### B. Contributions From Electromagnetic DOZI Processes

In the model used in the previous section the nature of the DOZI processes was assumed to be hadronic, and any electromagnetic DOZI processes were assumed to be suppressed by at least  $\alpha$  (the fine structure constant). This hypothesis can be tested by considering the decay  $J/\psi \rightarrow \phi\pi^0$ . Unlike the other

$J/\psi \rightarrow VP$  decays, it cannot proceed via any of the processes shown in Fig. 1. It could proceed via an electromagnetic DOZI decay, the lowest order diagram being shown in Fig. 1d. The upper limit on  $\tilde{B}(J/\psi \rightarrow \phi\pi^0)$  (Table II) is one to two orders of magnitude smaller than the reduced branching ratios for all other  $J/\psi \rightarrow VP$  decays. This indicates that the electromagnetic DOZI process shown in Fig. 1d plays a negligible role in  $J/\psi \rightarrow VP$  decays.

### C. The $\omega\pi^0$ Electromagnetic Form Factor

The electromagnetic form factors for the decay of the  $J/\psi$  to two pseudoscalar mesons (such as  $\pi\pi$  or  $KK$ ) have been measured previously.<sup>[6]</sup> For  $J/\psi \rightarrow VP$ , the electromagnetic form factor  $f(q^2)$  of the  $\omega\pi^0$  channel is calculated here. This decay is chosen because, unlike the  $\rho\eta$  and  $\rho\eta'$  decays, the amplitude depends only on  $\epsilon$ , and a good measurement of the crossed channel amplitude  $\omega \rightarrow \gamma\pi^0$  exists. The ratio of electromagnetic form factors at  $q^2 = m_{J/\psi}^2$  and at  $q^2=0$  is given by (see Appendix)

$$\frac{|f(m_{J/\psi}^2)|^2}{|f(0)|^2} = \frac{\alpha}{3} \left( \frac{p_\gamma}{p_\omega} \right)^3 \frac{m_{J/\psi} \Gamma(J/\psi \rightarrow \omega\pi^0)}{\Gamma(\omega \rightarrow \gamma\pi^0) \Gamma(J/\psi \rightarrow \mu^+\mu^-)}.$$

Using the branching ratio for  $J/\psi \rightarrow \omega\pi^0$  (Table I) and taking all other quantities from Ref. 10, we have

$$\frac{|f(m_{J/\psi}^2)|}{|f(0)|} = 0.0335 \pm 0.0059,$$

in agreement with our previous determination.<sup>[7]</sup> This value is three times smaller than that for the  $\pi\pi$  electromagnetic form factor ratio in  $J/\psi$  decay.<sup>[6]</sup> This suppression is in quantitative agreement with that predicted by hadronic helicity conservation in QCD.<sup>[20]</sup>

## V. SUMMARY

New measurements for two-body decays of the  $J/\psi$  into  $\rho\pi$ ,  $\rho\eta$ ,  $\rho\eta'$ ,  $\omega\pi^0$ ,  $\omega\eta$ ,  $\omega\eta'$ ,  $\phi\eta$ ,  $\phi\eta'$  and  $K^*K$ , as well as a new upper limit on  $\phi\pi^0$  are presented. A fit to the data using a phenomenological model requires a hadronic DOZI contribution. The magnitude of the DOZI contribution is  $15\pm 1\%$  of the SOZI amplitude. In this model the  $\eta$  and  $\eta'$  are found to be composed of only light and strange quarks. This rules out any gluonium contribution to the  $\eta'$  wavefunction. Using the results of the model the pseudoscalar mixing angle is found to be  $\theta_p = (-19.2 \pm 1.4)^\circ$ . The decay  $J/\psi \rightarrow \phi\pi^0$  is suppressed relative to the other decays, suggesting the absence of electromagnetic DOZI contributions. The  $\omega\pi^0$  electromagnetic form factor ratio is calculated to be  $\frac{|f(m_{J/\psi}^2)|}{|f(0)|} = 0.0335 \pm 0.0059$ .

We gratefully acknowledge the dedicated effort of SPEAR and the Linear Accelerator operating staff. This work was supported in part by the U.S. National Science Foundation and the U.S. Department of Energy under Contracts No. DE-AC03-76SF00515, No. DE-AC02-76ER01195, No. DE-AC03-81ER40050, and No. DE-AM03-76SF00034.

## APPENDIX

The matrix element for the decay  $J/\psi \rightarrow \omega\pi^0$  is written as the product of two terms,  $\langle \omega\pi^0 | J_\mu | 0 \rangle$  and  $\langle 0 | J_\mu | J/\psi \rangle$ . We calculate

$$\langle \omega\pi^0 | J_\mu | 0 \rangle = f(q^2) \epsilon_{\mu\nu\sigma\rho} e_\omega^\nu p_\omega^\sigma p_\pi^\rho \quad \text{and} \quad \langle 0 | J_\mu | J/\psi \rangle = \sqrt{\alpha} g e_{J/\psi}^\mu,$$

where  $q^2 = m_{J/\psi}^2$ ,  $\alpha$  is the fine structure constant,  $g$  is the coupling constant of the  $J/\psi$  to the virtual photon,  $e_i(p_i)$  is the polarization (momentum) four-vector of particle  $i$ , and  $f(q^2)$  is the electromagnetic form factor. In terms of the matrix element and two-body phase space, the decay width for  $J/\psi \rightarrow \omega\pi^0$  is

$$\Gamma(J/\psi \rightarrow \omega\pi^0) = \frac{4}{3} \pi \alpha^2 g^2 p_\omega^3 |f(m_{J/\psi}^2)|^2,$$

where  $p_\omega$  is the momentum of the  $\omega$  in the rest frame of the  $J/\psi$ .

The decay  $J/\psi \rightarrow \mu^+\mu^-$  is used to obtain  $g$ . In the approximation  $m_\mu^2 \ll m_{J/\psi}^2$ , we have<sup>[21]</sup>

$$\Gamma(J/\psi \rightarrow \mu^+\mu^-) = \frac{4}{3} \pi \alpha^2 g^2 m_{J/\psi}.$$

To normalize the electromagnetic form factor to that of  $q^2=0$ , we use the crossed-process  $\omega \rightarrow \gamma\pi^0$ . The matrix element for this is

$$\langle \pi^0 | J_\mu | \omega \rangle = f(0) \epsilon_{\mu\nu\sigma\rho} e_\gamma^\mu e_\omega^\nu p_\omega^\sigma p_\pi^\rho,$$

which gives

$$\Gamma(\omega \rightarrow \gamma\pi^0) = \frac{\alpha}{3} p_\gamma^3 |f(0)|^2,$$

where  $p_\gamma$  is the momentum of the  $\gamma$  in the rest frame of the  $\omega$ . We thus obtain

$$\frac{|f(m_{J/\psi}^2)|^2}{|f(0)|^2} = \frac{\alpha}{3} \left( \frac{p_\gamma}{p_\omega} \right)^3 \frac{m_{J/\psi} \Gamma(J/\psi \rightarrow \omega\pi^0)}{\Gamma(\omega \rightarrow \gamma\pi^0) \Gamma(J/\psi \rightarrow \mu^+\mu^-)}.$$

## REFERENCES

1. S.Okubo, Phys. Lett. **5**, 165 (1963); G.Zweig, CERN Preprints TH 401, 412 (1964), (unpublished); J.Iizuka, Prog. Theor. Phys. Suppl. **37-38**, 21 (1966).
2. G. Eigen, *Proceedings of the XVII Conference on Multiparticle Dynamics*, ed. by B. Aubert and L. Montanet (Editions Frontieres, Gif-Sur-Yvette, 1986).
3. See, for example, H. J. Lipkin, Phys. Lett. **67B**, 65 (1977); J.L. Rosner, Phys. Rev. D **27**, 1101 (1983), and Refs. 4-15 quoted therein.
4. G.B.Lepage et al, in Particles and Fields 2, Proceedings of the Banff Summer Institute, Banff, Canada, 1981 (ed. A.Z.Capri and A.N.Kamal, (Plenum, New York, 1983) p83 and references therein; S.J.Brodsky et al., *ibid*, p143 and references therein.
5. R.M. Baltrusaitis *et al.*, Phys. Rev. D **32**, 566 (1985).
6. S.J. Brodsky and G.P. Lepage, Phys. Rev. D **24**, 2848 (1981).
7. R.M. Baltrusaitis *et al.*, Phys. Rev. D **32**, 2883 (1985).
8. A. Seiden, H. Sadrozinski and H. Haber, Contributed Paper submitted to the International Europhysics Conference on High Energy Physics, Uppsala, Sweden, June 1987; submitted to Phys. Rev. D.
9. D.Bernstein *et al.*, Nucl. Instr. Meth. **226**, 301 (1984).
10. M. Aguilar-Benitez *et al.*, Phys. Lett. **170B**, 1 (1986).
11. The angular distribution  $J/\psi \rightarrow VP$  is given by  $\frac{d^3\sigma}{d\cos\theta_V d\cos\theta_1 d\phi_1} \propto \sin^2\theta_1[1 + \cos^2\theta_V + \sin^2\theta_V \cos(2\phi_1)]$  where  $\theta_V$  is the angle between the vector meson and the positron direction, and  $\theta_1$  and  $\phi_1$  describe the decay products of the vector meson in its helicity frame. For example, in the three body decays of the  $\omega$ ,  $\theta_1$  and  $\phi_1$  are the angles between the normal to the decay plane and the helicity direction. To fully describe

an event, an isotropic distribution from the decay of the pseudoscalar meson is assumed.

12. R.L. Ford and W.R. Nelson, SLAC-210 (1978) (unpublished).
13. The shape of the  $\rho$  is described by a relativistic Breit-Wigner, cut off above  $1.4 \text{ GeV}/c^2$ , with a width  $\Gamma = \Gamma_0 \frac{m_0}{m} \frac{1+r^2 p_0^2}{1+r^2 p^2} \left(\frac{p}{p_0}\right)^3$ , where  $m$  and  $p$  are the mass and momentum of the  $\pi^+\pi^-$  system.  $\Gamma_0, m_0, p_0$  are taken from Ref. 10. A value of  $r = 3 \text{ (GeV}/c)^{-1}$  best describes the data.
14. The mass of the  $\rho$  coming from the  $\eta'$  decay is shifted downward, as expected for a magnetic dipole transition.
15. R.M. Baltrusaitis *et al.*, Phys. Rev. D **33**, 1222 (1986).
16. The probability of obtaining the experimental result  $N$  if the signal is between  $n$  and  $n + dn$  and the expected mean background is  $b$ , is  $L(n) = N \frac{e^{-(b+n)} (b+n)^N}{N!}$ . The upper limit at a given confidence level is obtained as described in L. Lyons, *Probability and Statistics for Nuclear and Particle Physicists* (Cambridge Univ. Press, 1986) pp. 78-80.
17. R. Kauffman and F.J. Gilman, Phys. Rev. D **36**, 2761 (1987).
18. S. Rudaz, Phys. Rev. D **14**, 298 (1976); H. Kowalski and T. F. Walsh, Phys. Rev. D **14**, 852 (1976).
19. J. Olsson, Invited talk at the *Lepton-Photon Conference*, Hamburg, W. Germany, August 1987; A. Seiden, *Proceedings of the VII International Workshop on Photon-Photon Collisions*, ed. by A. Courau and P. Kessler (Singapore, World Scientific, 1986); H. Aihara, *et al.*, Phys. Rev. Lett. **57**, 2500 (1986).
20. The coupling of the  $\pi^0$  and  $\omega$  via a single form factor gives the  $\omega$  helicity to be  $\lambda_\omega = \pm 1$ . However, momentum conservation and the QCD prediction of hadronic helicity conservation predicts  $\lambda_{\pi^0} = \lambda_\omega = 0$  in  $e^+e^-$  collisions. See Ref. 6.

21. H.Pilkuhn, *Relativistic Particle Physics* (Springer Verlag, New York, 1979), pp. 169-176.

Table I.  $J/\psi \rightarrow VP$  branching ratios.

Channel	Final state	Branching ratio ( $10^{-3}$ )
$\rho\pi$	$\pi^+\pi^-\pi^0$	$14.2 \pm 0.1 \pm 1.9$
$\rho\eta$	$\pi^+\pi^-\eta$	$0.191 \pm 0.013 \pm 0.034$
	$\pi^+\pi^-\pi^+\pi^-\pi^0$	$0.196 \pm 0.029 \pm 0.040$
	<i>average</i>	$0.193 \pm 0.013 \pm 0.029$
$\rho\eta'$	$\pi^+\pi^-\gamma\rho^0$	$0.133 \pm 0.018 \pm 0.024$
	$\pi^+\pi^-\eta\pi^+\pi^-$	$0.097 \pm 0.022 \pm 0.017$
	<i>average</i>	$0.114 \pm 0.014 \pm 0.016$
$\omega\pi^0$	$\pi^+\pi^-\pi^0\pi^0$	$0.482 \pm 0.019 \pm 0.064$
$\omega\eta$	$\pi^+\pi^-\pi^0\eta$	$1.86 \pm 0.06 \pm 0.25$
	$\pi^+\pi^-\pi^0\gamma\pi^+\pi^-$	$1.49 \pm 0.17 \pm 0.22$
	<i>average</i>	$1.71 \pm 0.08 \pm 0.20$
$\omega\eta'$	$\pi^+\pi^-\pi^0\eta\pi^+\pi^-$	$0.169 \pm 0.027 \pm 0.025$
	$\pi^+\pi^-\pi^0\pi^+\pi^-\pi^+\pi^-\pi^0$	$0.154 \pm 0.051 \pm 0.026$
	$\pi^+\pi^-\pi^0\gamma\rho^0$	$0.167 \pm 0.025 \pm 0.024$
	<i>average</i>	$0.166 \pm 0.017 \pm 0.019$
$\phi\pi^0$	$K^+K^-\pi^0$	$<0.0068$
$\phi\eta$	$K^+K^-\eta$	$0.651 \pm 0.049 \pm 0.092$
	$K^+K^-\pi^+\pi^-\pi^0$	$0.677 \pm 0.088 \pm 0.096$
	<i>average</i>	$0.661 \pm 0.045 \pm 0.078$
$\phi\eta'$	$K^+K^-\eta\pi^+\pi^-$	$0.290 \pm 0.050 \pm 0.039$
	$K^+K^-\gamma\rho^0$	$0.327 \pm 0.046 \pm 0.049$
	<i>average</i>	$0.308 \pm 0.034 \pm 0.036$
$K^{*+}K^- + \text{c.c.}$	$K^\pm\pi^\mp K_S^0$	$5.16 \pm 0.16 \pm 0.54$
	$K^+K^-\pi^0$	$5.66 \pm 0.12 \pm 0.83$
	<i>average</i>	$5.26 \pm 0.13 \pm 0.53$
$K^{*0}\bar{K}^0 + \text{c.c.}$	$K^\pm\pi^\mp K_S^0$	$4.33 \pm 0.12 \pm 0.45$

Table II. Amplitudes and reduced branching ratios for  $J/\psi \rightarrow VP$ .

Process $J/\psi \rightarrow$	Amplitude	$\tilde{B}(10^{-3})$
$\rho^+\pi^-, \rho^0\pi^0, \rho^-\pi^+$	$g + e$	$1.556 \pm 0.161$
$K^{*+}K^-, K^{*-}K^+$	$g(1 - s) + e$	$1.017 \pm 0.061$
$K^{*0}\bar{K}^0, \bar{K}^{*0}K^0$	$g(1 - s) - 2e$	$0.836 \pm 0.055$
$\omega\eta$	$(g + e)X_\eta + \sqrt{2}rg(\sqrt{2}X_\eta + Y_\eta)$	$0.632 \pm 0.058$
$\omega\eta'$	$(g + e)X_{\eta'} + \sqrt{2}rg(\sqrt{2}X_{\eta'} + Y_{\eta'})$	$0.079 \pm 0.010$
$\phi\eta$	$(g(1 - 2s) - 2e)Y_\eta + rg(\sqrt{2}X_\eta + Y_\eta)$	$0.287 \pm 0.031$
$\phi\eta'$	$(g(1 - 2s) - 2e)Y_{\eta'} + rg(\sqrt{2}X_{\eta'} + Y_{\eta'})$	$0.182 \pm 0.025$
$\rho^0\eta$	$3eX_\eta$	$0.071 \pm 0.010$
$\rho^0\eta'$	$3eX_{\eta'}$	$0.054 \pm 0.009$
$\omega\pi^0$	$3e$	$0.159 \pm 0.017$
$\phi\pi^0$	$0$	$< 0.0026$

## FIGURE CAPTIONS

1. Diagrams contributing to  $J/\psi$  decays: (a) three gluon annihilation; (b) electromagnetic decay proceeding via  $c\bar{c}$  annihilation into one photon; (c) a purely hadronic DOZI decay; (d) an example of an electromagnetic DOZI decay.
2. The Dalitz plot for  $J/\psi \rightarrow \pi^+\pi^-\pi^0$ . (To emphasize the structure only one third of the data is displayed.)
3. The  $\pi^+\pi^-$  invariant mass for events of the type  $J/\psi \rightarrow \pi^+\pi^-\eta$  in the modes: (a)  $\eta \rightarrow \gamma\gamma$ ; (b)  $\eta \rightarrow \pi^+\pi^-\pi^0$ . The curves show the fit results for: the background (I); background +  $\omega$  (II); background +  $\rho$  (III); and background +  $\omega + \rho$  + interference term (IV). The background is quadratic for (a), linear for (b).
4. (a) The  $\gamma\rho$  invariant mass for events of the type  $J/\psi \rightarrow \rho\rho\gamma$ . (b) The  $\eta\pi^+\pi^-$  invariant mass for events of the type  $J/\psi \rightarrow \rho\eta\pi^+\pi^-$ . The curves show the quadratic background (I) and background +  $\eta'$  (II).
5. (a) The  $\pi^+\pi^-\pi^0$  invariant mass for events of the type  $J/\psi \rightarrow \pi^+\pi^-\pi^0\gamma\gamma$ . (b) The  $\gamma\gamma$  invariant mass for events of the type  $J/\psi \rightarrow \omega\gamma\gamma$ ; the curves show the quadratic background (I), background +  $\pi^0$  (II) and background +  $\eta$  (III).
6. (a) The  $\gamma\pi^+\pi^-$  invariant mass in events of the type  $J/\psi \rightarrow \omega\pi^+\pi^-\gamma$ . The curves show the estimated phase space background (I), background + Monte Carlo simulation of the decay  $J/\psi \rightarrow \omega\eta, \eta \rightarrow \pi^+\pi^-\pi^0$  (II) and background + Monte Carlo simulation of the decay  $J/\psi \rightarrow \omega\eta, \eta \rightarrow \pi^+\pi^-\gamma$  (III). (b) The  $\gamma\rho$  invariant mass in events of the type  $J/\psi \rightarrow \omega\rho\gamma$ . The curves show the quadratic background (I) and the background +  $\eta'$  (II). (c) The  $\eta\pi^+\pi^-$  invariant mass in events of the type  $J/\psi \rightarrow \omega\eta\pi^+\pi^-, \eta \rightarrow \gamma\gamma$ . (d) The  $\eta\pi^+\pi^-$  invariant mass in events of the type  $J/\psi \rightarrow \omega\eta\pi^+\pi^-, \eta \rightarrow \pi^+\pi^-\pi^0$ .

7. (a) The  $K^+K^-$  invariant mass in events of the type  $J/\psi \rightarrow K^+K^-\eta$ .  
 (b) The  $\gamma\gamma$  invariant mass in events of the type  $J/\psi \rightarrow \phi\gamma\gamma$ ; the curves show the quadratic background (I) and the background +  $\eta$  (II).
8. (a) The  $\pi^+\pi^-\pi^0$  invariant mass in events of the type  $J/\psi \rightarrow \phi\pi^+\pi^-\pi^0$ .  
 (b) The  $\gamma\rho$  invariant mass in events of the type  $J/\psi \rightarrow \phi\rho\gamma$ . The curves show the quadratic background (I) and the background +  $\eta'$  (II). (c) The  $\eta\pi^+\pi^-$  invariant mass in events of the type  $J/\psi \rightarrow \phi\eta\pi^+\pi^-, \eta \rightarrow \gamma\gamma$ .
9. (a) The Dalitz plot for  $J/\psi \rightarrow K^\pm K^\mp \pi^0$ . (b) The Dalitz plot for  $J/\psi \rightarrow K^\pm \pi^+ \pi^- \pi^\mp$  events.





Fig. 1



9-87

5840A2

Fig. 2




Fig. 3






Fig. 4



8-87

5480A5

Fig. 5




Fig. 6





Fig. 7







Fig. 8



5840B9

9-87

Fig. 9



9-87  
5840A10

Fig. 10