An Accurate Measurement of the Landau-Pomeranchuk-Migdal Effect

P.L. Anthony,¹ R. Becker-Szendy,¹ P.E. Bosted,² M. Cavalli-Sforza,^{3,*} L.P. Keller,¹

L.A. Kelley,³ S.R. Klein,^{3,4} G. Niemi,¹ M.L. Perl,¹ L.S. Rochester,¹ J.L. White²

¹Stanford Linear Accelerator Center, Stanford, CA 94309 ²The American University, Washington, D.C. 20016

³Santa Cruz Institute for Particle Physics, University of California, Santa Cruz, CA 95064 ⁴Lawrence Berkeley Laboratory, Berkeley, CA 94720

Abstract

In the 1950's, Landau, Pomeranchuk and Migdal predicted that the cross section for bremsstrahlung from highly relativistic particles in dense media would be suppressed due to interference caused by multiple scattering. We have measured the production rates of 5 to 500 MeV photons from 8 and 25 GeV electrons traversing thin gold and carbon targets. We observe, to within 5% accuracy, suppression at the level predicted by Migdal. For extremely thin targets, we observe 'edge effects' where the LPM suppression is reduced near the edges of the targets. We also point out the imprecise foundation of the LPM theory.

Submitted to Physical Review Letters

Work supported by Department of Energy contracts DE-AC03-76SF00515 (SLAC), DE-AC03-76SF00098 (LBL), W-4705-ENG-48 (LLNL) and National Science Foundation grants NSF-PHY-9113428 (UCSC) and NSF-PHY-9114958 (American U.).

In the 1950's Landau, Pomeranchuk and Migdal (LPM) [1,2] predicted that the cross section for bremsstrahlung from highly relativistic particles in dense media is suppressed due to interference caused by multiple scattering. The suppression has its roots in the uncertainty principle. The kinematics of bremsstrahlung requires that the longitudinal momentum transfer between the nucleus and the electron must be small; the uncertainty principle then requires that the transfer must occur over a large longitudinal distance scale (formation zone). If the electron multiple scatters while traversing this zone, the bremsstrahlung amplitude from before and after the scattering can interfere, reducing the amplitude for bremsstrahlung photon emission.

We present here the first quantitative measurement of bremsstrahlung suppression due to the LPM effect. In the past there have been several qualitative tests of the LPM effect using cosmic rays and one qualitative test at an accelerator [3]. The effect is relevant in many areas, but particularly in high energy cosmic ray air showers [4]. Our experiment, SLAC E-146, [5,6] carried out at the Stanford Linear Accelerator Center (SLAC), studied the bremsstrahlung production of 5 to 500 MeV photons by 8 and 25 GeV electrons. We found that the suppression of bremsstrahlung predicted by the LPM theory is correct to within 5%.

Unfortunately, no concise derivations of the LPM effect exist. Qualitative discussions have been given by Galitsky and Gurevich [8] and by Perl [9]. Here we give a brief qualitative discussion and the basic equations. The differential cross section for the bremsstrahlung production of a photon of energy k by an ultrarelativistic electron of energy E and mass m, where $k \ll E$ (complete screening) [10] is given by

$$\frac{d\sigma_{\rm BH}}{dk} = \frac{4\alpha r_e^2}{3k} \Big[\{y^2 + 2[1 + (1-y)^2]\} (Z^2 F_{el} + Z F_{inel}) + (1-y) \frac{(Z^2 + Z)}{3} \Big]$$
(1)

Here $\bar{y} = k/E$, α is the fine structure constant, r_e is the classical radius of the electron, and Z is the nuclear charge. $F_{el} \approx \ln(184/Z^{1/3})$ and $F_{inel} \approx \ln(1194/Z^{2/3})$ are the elastic and inelastic atomic from factors [10]. Eq.(1) has the well known Bethe-Heitler form $(d\sigma_{\rm BH}/dk)_{y\ll 1} \sim 1/k$. In this kinetic regime, $k \ll E$, the angle between the incident electron and the produced photon is small, $\theta_k \approx mc^2/E$, and the average angle between the scattered electron and the incident electron is smaller still. Neglecting these angles, the longitudinal momentum transfer to the atom is $q_{\parallel} \approx k/(2\gamma^2 c)$ where $\gamma = E/mc^2$ [9]. The uncertainty principle then requires that the spatial position of the bremsstrahlung process has a longitudinal uncertainty of $L_{\parallel} = \hbar/q_{\parallel} \approx 2\hbar c\gamma^2/k$. In alternate language, the electron and photon slowly split apart over the distance L_{\parallel} . In a sufficiently dense medium the electron mean free path is much less than L_{\parallel} , so an electron will interact while traversing L_{\parallel} ; in the LPM effect the relevant interaction is multiple scattering. Roughly, bremsstrahlung is suppressed [8] when the mean square multiple scattering angle over the distance L_{\parallel}

$$\theta_s^2 = \left(\frac{E_s}{E}\right)^2 \frac{L_{\parallel}}{X_0} \tag{2}$$

is greater than or equal to θ_k^2 . Here $E_s = mc^2 (4\pi/\alpha)^{1/2} \approx 21$ MeV and X_0 is the radiation length. Therefore $d\sigma/dk$ is suppressed when

$$k < k_{\rm LPM} = \frac{E^2}{E_{\rm LPM}} \tag{3}$$

where $E_{\text{LPM}} = m^2 X_0 \alpha / 8\pi \hbar c = 3.8 \times 10^{12} eV \times X_0(cm)$.

Table I shows that, as Z increases E_{LPM} decreases and k_{LPM} , the electron energy dependent upper limit for the LPM effect, increases. Using semi-classical arguments [1], it can be shown that for $y \ll 1$, $d\sigma_{\text{LPM}}/dk \sim 1/\sqrt{k}$ in contrast to $d\sigma_{\text{BH}}/dk$ which is proportional to 1/k. A more accurate analysis becomes very involved; Migdal based his calculations on scattering theory.

In his formulation [2], Migdal defines a dimensionless variable

- Target	Z	X_0 (cm)	$E_{ m LPM}$ (TeV)	$k_{ m LPM}~({ m MeV})$
С	6 _	- 18.8	71	8.8
Au	79	0.33	1.25	500

TABLE I. Z, X_0, L_{LPM} , and k_{LPM} , the latter for 25 GeV electrons.



FIG. 1. Comparison of $kd\sigma_{\rm LPM}/dk$ and $kd\sigma_{\rm BH}/dk$ for 25 GeV electrons incident on a uranium target. The cross sections are not corrected for the edge effect.

$$s = \frac{1}{2} \left(\frac{y}{1-y}\right)^{1/2} \left(\frac{mc}{\hbar} \frac{mc^2}{E} \frac{\alpha X_0}{8\pi\xi(s)}\right)^{1/2} \tag{4}$$

where $1 \leq \xi(s) \leq 2$. When $y \ll 1$, $s \sim (L_{\text{LPM}}/L_{\parallel})^{1/2}$. When $s \gg 1$, $d\sigma/dk$ is unaffected; when $s \ll 1$, there is strong suppression. Migdal found that

$$\frac{d\sigma_{\rm LPM}}{dk} = \frac{4\alpha r_e^2 \xi(s)}{3k} \{ y^2 G(s) + 2[1 + (1 - y)^2]\phi(s) \} Z^2 \ln\left(\frac{184}{Z^{1/3}}\right)$$
(5)

where G(s) and $\phi(s)$ may be approximated by sums[4]. In the no suppression limit $s \to \infty$, $G(s) \to 1$, and $\phi(s) \to 1$. Comparing $d\sigma_{\text{LPM}}/dk$ [Eq. (6)] with $d\sigma_{\text{BH}}/dk$ [Eq. (1)] in this limit, $d\sigma_{\text{LPM}}/dk$ yields the Z^2F_{el} term but not the ZF_{inel} or the $(Z^2 + Z)$ terms. Our calculations include the ZF_{inel} term by normalizing both our $d\sigma_{\text{LPM}}/dk$ and $d\sigma_{\text{BH}}/dk$ to the radiation length as defined by Tsai [10]. The $(Z^2 + Z)$ term is omitted from both our cross sections and the definition of the radiation length[10]; this is roughly a 2% correction.

Fig. 1 shows the suppression of the LPM cross section as k decreases. Right above the region where the suppression disappears, $d\sigma_{\rm LPM}/dk$ rises slightly above $d\sigma_{\rm BH}/dk$; this occurs because $\xi(s)\phi(s)$ rises slightly above 1.

The experimental layout is shown in Fig. 2. Electron beams with energies of 8 or 25 GeV entered the SLAC End Station A, and interacted in targets of thicknesses 0.1%-6% X_0 . The



FIG. 2. Schematic picture of the experiment.

electrons were then magnetically bent downward by 39 mrad (at the full beam energy) into a wire chamber spectrometer and thence into lead glass blocks which accurately count the electrons. Photons produced in the target were detected 50 meters downstream in a BGO calorimeter array. To minimize backgrounds, the electron path upstream of the calorimeter and the photon flight path were kept in vacuum. The beam ran at 120 pulses/sec, with an intensity adjusted to average 1 electron/pulse, except for the 0.1% gold target, where higher rates were sometimes used. The beam was generated parasitically during SLC collider operation [7]. To avoid possible trigger bias, data were recorded every pulse. This Letter will discuss data taken with gold and carbon targets, as detailed in Table II.

The BGO calorimeter comprises 45 crystals (a 7 by 7 array, with the corners missing). Each crystal is 2 cm square by 20 cm (18 X_0) deep. The calorimeter was read out by PMTs which detected about 1 photoelectron per 30 keV of energy deposition. The PMT

Target	6% C	0.1% Au	0.7% Au	6% Au	
$t \ ({ m mm})$	11.732	0.0039	0.0231	0.1991	
$t~({ m gm/cm^2})$	2.5717	0.0074	0.0445	0.384	
$t(X_0)$	0.0602-	0.0011	0.0068	0.0591	

TABLE II. Target thickness t in mm, gm/cm^2 , and X_0 .

signals were digitized by LeCroy 2282, 12 bit ADCs, with the phototube gain adjusted so that 1 count was about 100 keV. The FWHM calorimeter resolution was 6% for 400 MeV electrons, rising to 8% FWHM for 40 MeV electrons. Three methods were used for calorimeter calibration: electron beams with energies of 400 and 500 MeV; cosmic ray muons; and event reconstruction, requiring that the photon energy in the BGO and the electron energy measured in the wire chamber sum to the beam energy. The BGO temperature was monitored during data taking, and the BGO data was corrected to compensate for temperature variations, using the measured temperature response. Further details of the calibration are provided in Refs. [5] and [6].

Our analysis selected bremsstrahlung events containing containing a photon in the calorimeter plus a single electron in the lead glass blocks. The photon energies were then histogrammed. Figure 3 shows a sampling of our data, together with Monte Carlo predictions, for both the Bethe-Heitler and LPM $dN/d(\log k)$. The plots show $dN/d(\log k)$, with 25 bins per decade of energy, so that the bin width, $\Delta E \approx 0.09E$. Logarithmic bins are used so that the Bethe-Heitler 1/k cross section appears roughly flat.

Figs. 3a and 3b show the number of observed photons per 25 GeV electron traversing the 6% X_0 gold and carbon targets. The gold target shows a large suppression compared to the BH prediction. Carbon, which is much less dense, shows little suppression. Fig. 3c shows the corresponding plots for 8 GeV electrons incident on the gold target; here the LPM effect is much reduced. In all three cases, excellent agreement is seen with the Monte Carlo LPM curves (dashed lines) above k = 30 MeV; below 30 MeV the data is higher than the LPM Monte Carlo.

One shortcoming in Migdal's formulae is that they apply only for infinitely thick targets. In a target of finite thickness, electrons may interact near an edge. Then, the formation zone may extend out of the target so the electrons undergo less multiple scattering, and therefore less suppression. Unfortunately, we have found no satisfactory theoretical treatment of this phenomenon. This can explain the discrepancy between the LPM prediction and data below about 30 MeV in Fig. 3a. This effect is proportionately much larger in thinner targets, as

6



FIG. 3. Measurements with statistical errors only of $dN/d(\log k)$ compared with Monte Carlo calculated theoretical curves The cross sections are given in terms of $dN/d(\log k)/X_0$ where N is the number of events per photon energy bin per incident electron. The photon energy scale is logarithmic with 25 bins per decade, so each bin has a width $\Delta k \sim 0.0964k$. The solid curve is the Bethe Heitler Monte Carlo prediction and the dashed curve gives the LPM Monte Carlo prediction. The figures are: (a) 25 GeV electrons incident on 6% X_0 gold; (b) 25 GeV electrons incident on 6% X_0 carbon; (c) 8 GeV electrons incident on 6% X_0 gold; (d) 25 GeV electrons incident on 0.7% X_0 gold; (e) data and Monte Carlo curves from 25 GeV electrons incident on 0.7% X_0 gold subtracted from 25 GeV electrons incident on 6% X_0 gold, leaving the 'middle' 5.3% X_0 of gold; (f) 25 GeV electrons incident on 0.1% X_0 gold.

seen in Fig. 3d for 25 GeV electrons incident on a 0.7% X_0 gold target. To remove edge effects from our data, we subtract the data from the 0.7% X_0 gold target from the 6% X_0 gold data to get the middle 5.3%, without edge effects. The result is shown in Fig. 3e. Repeating this procedure with Monte Carlo results accounts for multiple photon emission; the agreement is excellent over the entire range 5 < k < 500 MeV. This procedure introduces a small (<2%) systematic error due to additional multiphoton emission associated with the edge effect.

For extremely thin targets, thinner than the formation zone length, LPM suppression should completely disappear because there is too little total multiple scattering to cause suppression. This effect is verified by the data shown in Fig. 3f from 25 GeV electrons passing through a 0.1% X_0 target.

The errors shown on the plots are statistical only. The point-to-point systematic errors vary slowly with k and correspond to a 4.5% uncertainty across the measured range of k. The major sources are photon cluster finding (2%), calorimeter energy calibration (1.5%), calorimeter nonlinearity(3%), backgrounds (1%), uncertainty in the target density (2%), and multiphoton pileup due to the edge effect (1%).

The major backgrounds- synchrotron radiation from the spectrometer and beamline magnets, and transition radiation in the target- are expected to be small above 5 MeV. The synchrotron radiation and other beam related backgrounds have been measured in empty target holder runs, and found to be small, below 0.1% per electron. When significant (as for the 0.1% gold target), they have been subtracted from the corresponding target-in data.

The major difference between the Monte Carlo curves and the theoretical cross sections (Fig. 1) is that the Monte Carlo includes the effect of multi-photon emission, where a single electron passing through the target interacts twice, emitting two different photons. This depresses the low energy end of the spectrum, and increases the high energy end, tilting the otherwise fairly flat Bethe Heitler spectrum. The Monte Carlo curves also includes smearing to simulate calorimeter resolution and possible photon absorption in the target. At low energies, we include an additional suppression due to the longitudinal density effect

[11], following the prescription in Ref 2.

The Monte Carlo curves are normalized so that the LPM curves match the data in the region 450 < k < 500 MeV. Except for the $0.1\% X_0$ gold, the data are consistent with a common normalization factor of $0.94 \pm 0.01 \pm 0.032$. The 3.2% systematic error comes from uncertainties in the target thickness(2%), energy calibration (1%), electron flux (0.5%), Monte Carlo (excluding bremsstrahlung physics input) (1%), and the normalization technique (2%).

It is difficult to explain an overall 6% normalization error in terms of experimental biases alone. The error is likely due to the approximate nature of the Migdal formula (for example in neglecting the Z(Z+1)/3 term, which typically contributes 2% to the BH formula). The bremsstrahlung cross section may also be affected by the electron distribution in a solid, compared with the electron wave functions in a free atom.

In summary, we have found that the Migdal theory of the LPM effect accurately describes data taken on carbon and gold targets, once multi-photon pileup and edge effects are taken into account. The suppression varies with both target X_0 and electron energy as predicted by Migdal. For thin targets, edge effects reduce the suppression. For all of our data, the absolute bremsstrahlung rate is slightly below that predicted by current theory. More theoretical work is called for to accurately treat these edge effects, and the small but significant normalization difference between data and theory needs to be better understood.

We would like to thank the SLAC Experimental Facilities group for their assistance in setting up the experiment and the SLAC Accelerator Operators group for their efficient beam delivery. Walter Meyerhof provided the target holder. This work was supported by Department of Energy contracts DE-AC03-76SF00515 (SLAC), DE-AC03-76SF00098 (LBL), W-4705-ENG-48 (LLNL) and National Science Foundation grants NSF-PHY-9113428 (UCSC) and NSF-PHY-9114958 (American U.).

9

REFERENCES

- * Present address: Institut de Fisica d'Altes Energies, Universitat Autonima de Barcelona,
 08193 Bellaterra (Barcelona), Spain.
- L. D. Landau and I. J. Pomeranchuk, Dokl. Akad. Nauk. SSSR 92, 535 (1953); 92, 735 (1953). These two papers are available in English in L. Landau, *The Collected Papers of L. D. Landau*, Pergamon Press, 1965.
- [2] A. B. Migdal, Phys. Rev. **103**, 1811 (1956).
- [3] A. Varfolomeev et al., Sov. Phys. JETP 42, 218 (1976).
- [4] T. Stanev et al., Phys. Rev. D 25, 1291 (1982).
- [5] S. R. Klein et al., in Proc. XVI Int. Symp. Lepton and Photon Interactions at High Energies (Ithaca, 1993), Eds. P. Drell and D. Rubin, p. 172.
- [6] R. Becker-Szendy et al., in Proc. 21st SLAC Summer Institute on Particle Physics (Palo Alto, 1994), pg. 519.
- [7] M. Cavalli-Sforza et al., IEEE Trans. Nucl. Sci. 41, 1374 (1994).
 - [8] V. M. Galitsky and I. I. Gurevich, Il Nuovo Cimento 32, 396 (1964).
 - [9] M. L. Perl, in Proc. 1994 Les Rencontres de Physique de la Vallee D'Aoste, (Editions Frontieres, Gif-sur-Yvette, France, 1994). Ed. M. Grego, p. 567.
 - [10] Y.-S. Tsai, Rev. Mod. Phys. 46, 815 (1974).
 - M. L. Ter-Mikaelian, Dokl. Akad. Nauk. SSR 94, 1033 (1954). For a discussion in English, see M. L. Ter-Mikaelian, *High Energy Electromagnetic Processes in Condensed Media*, John Wiley & Sons, 1972.