SLAC–PUB–7573
June 1997

# A PRELIMINARY IMPROVED TEST OF THE FLAVOR INDEPENDENCE OF STRONG INTERACTIONS*

**The SLD Collaboration****

Stanford Linear Accelerator Center

Stanford University, Stanford, CA 94309

## ABSTRACT

We present an improved comparison of the strong couplings of gluons to light ($u$, $d$, and $s$), $c$, and $b$ quarks, determined from multijet rates in flavor-tagged samples of hadronic $Z^0$ decays recorded with the SLC Large Detector at the SLAC Linear Collider between 1993 and 1995. Flavor separation on the basis of lifetime and decay multiplicity differences among hadrons containing light, $c$, and $b$ quarks was made using the SLD precision tracking system, yielding tags with high purity and low bias against $\geq$ 3-jet final states. We find: $\alpha_s^{uds}/\alpha_s^{all} = 0.997 \pm 0.011(stat) \pm 0.011(syst) \pm 0.005(theory)$, $\alpha_s^c/\alpha_s^{all} = 0.984 \pm 0.042 \pm 0.053 \pm 0.022$, $\alpha_s^b/\alpha_s^{all} = 1.022 \pm 0.019 \pm 0.023 \pm 0.012$.

Contributed to the XVIII International Symposium on Lepton Photon Interactions, July 28 - August 1 1997, Hamburg, Germany, and to the International Europhysics Conference on High Energy Physics, 19-26 August 1997, Jerusalem, Israel; Ref: 289.

*Work supported by Department of Energy contract DE–AC03–76SF00515.

# 1. Introduction

A fundamental assumption of the theory of strong interactions, Quantum Chromodynamics (QCD), is that the strong coupling $\alpha_s$ is independent of quark flavor. This can be tested by measuring the strong coupling in events of the type $e^+e^- \to q\bar{q}(g)$ for specific quark flavors $q$. Although an absolute determination of $\alpha_s$ for each quark flavor would have large theoretical uncertainties [1], it is possible to test the flavor-independence of QCD precisely by measuring ratios of couplings in which most experimental errors and theoretical uncertainties are expected to cancel. Since it has recently been suggested [2] that a flavor-dependent anomalous quark chromomagnetic moment could modify the probability for the radiation of gluons, comparison of the strong coupling for different quark flavors may also provide information on physics beyond the Standard Model.

Comparisons of $\alpha_s$ for $b$ or $c$ quarks with $\alpha_s$ for all flavors made at PETRA [3] were limited in precision to $\pm 0.41$ ($c$) and $\pm 0.57$ ($b$) due to small data samples and limited heavy quark tagging capability. Measurements made at LEP of $\alpha_s^b/\alpha_s^{udsc}$ have reached precisions between $\pm 0.06$ and $\pm 0.013$ [4]. However, these tests make the simplifying assumption that $\alpha_s$ is independent of flavor for all the non-$b$ quarks, and are insensitive to differences between $\alpha_s$ for these flavors, especially a different $\alpha_s$ for $c$ quarks compared with either $b$ or light quarks. The OPAL Collaboration has measured $\alpha_s^f/\alpha_s^{all}$ for all five flavors $f$ with no assumption on the relative value of $\alpha_s$ for different flavors [5] to precisions of $\pm 0.026$ for $b$ and $\pm 0.09$ to $\pm 0.20$ for the other flavors. In that analysis the kinematic signatures used to tag $c$ and light quarks suffer from low efficiency and strong biases, due to preferential tagging of events without hard gluon radiation.

The SLC Large Detector (SLD) [6] at the SLAC Linear Collider (SLC) is an ideal environment in which to test the flavor independence of strong interactions. The tracking capability of the Central Drift Chamber (CDC) [7] and the precise CCD Vertex Detector (VXD) [8], combined with the stable, micron-sized beam interaction

point (IP), allows us to select $Z^0 \to b\bar{b}(g)$ and $Z^0 \to q_l\bar{q}_l(g)$ ($q_l = u, d, s$) events using their quark decay lifetime signatures with high efficiency and purity, and with low bias against 3-jet events, an important advantage of this analysis. Our previous measurement [9], based on the sample of roughly 50,000 $Z^0$ decay events collected in 1993, reached precisions of $\pm 0.04$ ($\alpha_s^{uds}/\alpha_s^{all}$), $\pm 0.17$ ($\alpha_s^c/\alpha_s^{all}$), and $\pm 0.06$ ($\alpha_s^b/\alpha_s^{all}$), and tested the flavor independence of the strong coupling making no assumptions about the relative values of $\alpha_s^b$, $\alpha_s^c$ and $\alpha_s^{uds}$. Here we present improved measurements of these quantities using the same technique, and based on the total sample of 150,000 $Z^0$ decay events collected by SLD between 1993 and 1995.

## 2. Event Selection and Flavor Tagging

This analysis is based on 5.4 $pb^{-1}$ of $e^+e^-$ annihilation data collected at a mean center-of-mass energy of $\sqrt{s} = 91.26$ GeV. The trigger and selection criteria for hadronic $Z^0$ decays are described in Ref. [1]. The efficiency for selecting a well-contained $Z^0 \to q\bar{q}(g)$ event was estimated to be above 96% independent of quark flavor. The selected sample comprised 78319 events, with an estimated $0.10 \pm 0.05\%$ background contribution dominated by $Z^0 \to \tau^+\tau^-$ events. This analysis used charged tracks measured in the CDC and in the VXD [1].

We used normalized impact parameters $d/\sigma_d$ as the basis for quark flavor tags, where $d$ is the signed distance of closest approach of a charged track to the IP in the ($x$–$y$) plane transverse to the beam axis, and $\sigma_d$ is the error on $d$. Tracks used for event flavor tagging were required to have: at least one VXD hit; at least 40 CDC hits, with the first hit at a radius less than 39 cm; a CDC fit quality ${\chi^2}_{CDC}/d.o.f < 5$; a combined CDC+VXD fit quality ${\chi^2}_{CDC+VXD}/d.o.f < 5$; momentum greater than 0.5 GeV/$c$; $\sigma_d < 250$ $\mu$m; and to miss the IP by less than 0.3 cm in the $x$–$y$ plane and by less than 1.5 cm in $z$. Tracks from candidate $K^0$ and $\Lambda$ decays and $\gamma$-conversions were removed [10]. For these selected tracks, $\sigma_d = 11 \oplus 70/(p_t(\mathrm{GeV}/c^2)\sqrt{\sin\theta})\mu$m has been measured using $Z^0 \to \mu^+\mu^-$ decays and hadronic events, where $p_t$ is the track momentum transverse to the beam axis, and $\theta$ is the polar angle with respect to

the beam axis. The spatial resolution on the average transverse IP position has been measured to be 7 $\mu$m [10]. The distributions of $d$ and $d/\sigma_d$ are modeled well by the SLD simulation [10].

Figure 1 shows the distribution of $N_{\rm sig}$, the number of tagging tracks per event with $d/\sigma_d \geq 3$. The data are well described by a Monte Carlo simulation of hadronic $Z^0$ decays [11] with parameter values tuned [12] to hadronic $e^+e^-$ annihilation data, combined with a simulation of the SLD. For the simulation, the contributions of events of different quark flavors are shown separately. The leftmost bin contains predominantly events containing primary $u$, $d$, or $s$ quarks, while the rightmost bins contain a pure sample of events containing primary $b$ quarks. The event sample was divided accordingly into five subsamples: (1) $N_{sig} = 0$, (2) $N_{sig} = 1$, (3) $N_{sig} = 2$, (4) $N_{sig} = 3$, and (5) $N_{sig} \geq 4$. We refer to subsample 1 as the uds-tagged sample, the union of subsamples 2, 3 and 4 as the c-tagged sample, and subsample 5 as the b-tagged sample. The hard $b$ tag yields a sample with very low contamination from charm events, maximizing the sensitivity of the three-flavor test. The light-quark tag does not change the relative flavor composition of the $uds$ sample. The efficiencies $\varepsilon$ for selecting events (after cuts) of type $i$ ($i = uds$, $c$, $b$) with tag $i$, and the fractions $\Pi$ of events of type $i$ in the $i$-tagged sample, were calculated from the Monte Carlo simulation to be: $(\varepsilon, \Pi)_{uds} = (84.2 \pm 0.1\%,\ 86.2 \pm 0.1\%)$; $(\varepsilon,\ \Pi)_c = (57.0 \pm 0.2\%,\ 32.3 \pm 0.1\%)$; $(\varepsilon,\ \Pi)_b = (43.4 \pm 0.2\%, 95.2 \pm 0.1\%)$; the errors are discussed below.

# 3. Jet Finding

Jets were then reconstructed using iterative clustering algorithms. We used the 'E', 'E0', 'P', and 'P0' variations of the JADE algorithm, as well as the 'Durham' ('D') and 'Geneva' ('G') algorithms [13]. We divided events into two categories: those containing: (1) two jets, and (2) three or more jets. The fraction of the event sample in category 2 was defined as the 3-jet rate $R_3$. This quantity is infrared- and collinear-safe and has been calculated to $\mathcal{O}(\alpha_s^2)$ in perturbative QCD [13, 14]. For each algorithm, the jet resolution parameter $y_c$ was chosen so as to minimize the combined statistical

and systematic errors on the $\alpha_s$ flavor ratios, as discussed below. This choice also avoids the 'Sudakov region' at low $y_c$ where multiple gluon emission requires that large logarithmic terms of $1/y_c$ be resummed in order to describe the data [1]. The chosen $y_c$ values are listed in Table 1.

The $R_3^j$ for each of the $j$ quark types ($j = uds$, $c$, $b$) was extracted from a simultaneous maximum likelihood fit to all $n_2^i$ and $n_3^i$, the number of 2-jet and 3-jet events, respectively, in subsample $i$ ($1 \leq i \leq 5$), using the relations:

$$\begin{aligned} n_2^i &= \sum_{j=1}^{3} \left( \varepsilon_{(2\to 2)}^{ij}(1 - R_3^j) + \varepsilon_{(3\to 2)}^{ij} R_3^j \right) f^j N \\ n_3^i &= \sum_{j=1}^{3} \left( \varepsilon_{(3\to 3)}^{ij} R_3^j + \varepsilon_{(2\to 3)}^{ij}(1 - R_3^j) \right) f^j N \ . \end{aligned} \tag{1}$$

Here $N$ is the total number of selected events corrected for the event selection efficiency, and $f^j$ is the Standard Model fractional hadronic width for $Z^0$ decays to quark type $j$. The $5 \times 3$ matrices $\varepsilon_{(2\to 2)}^{ij}$ and $\varepsilon_{(3\to 3)}^{ij}$ are the efficiencies for an event of type $j$, with 2- or 3-jets at the parton level, to pass all cuts and enter subsample $i$ as a 2- or 3-jet event, respectively. This formalism explicitly accounts for modifications of the parton-level 3-jet rate due to hadronization, detector effects, and tagging bias. These matrices were calculated from the Monte Carlo simulation. The efficiencies for correctly tagging a 2-jet event and a 3-jet event differ by an average over jet algorithms of 5.1%, 9.8%, and 19.6% for the $uds$, $c$, and $b$ tags, respectively.

Equations 1 were solved using 2- and 3-jet events defined by each of the six algorithms. The ratios $R_3^j/R_3^{all}$, where $R_3^{all}$ is the 3-jet rate in the total event sample, are shown in Table 1. Averaged over all six algorithms the correlation coefficients from the fit are: $uds$-$c$ : $-0.68$, $uds$-$b$ : $0.32$, $c$-$b$ : $-0.57$. The statistical errors were calculated using the full covariance matrix.

# 4. Comparison of $\alpha_s$ values

The 3-jet rate in heavy quark ($b$, $c$) events is expected to be modified relative to that in light quark events by the diminished phase-space for gluon emission due to the

quark masses. We evaluated the correction factors, $R_3^c/R_3^u$ and $R_3^b/R_3^d$, for each jet algorithm and $y_c$ value according to the JETSET7.4 parton shower simulation. These factors are listed in Table 1, and were used to correct the measured 3-jet rate ratios.

To $\mathcal{O}(\alpha_s^2)$ in perturbative QCD, $R_3(y_c) = A(y_c)\alpha_s + (B(y_c) + C(y_c))\alpha_s^2$, where the $\mathcal{O}(\alpha_s^2)$ coefficient includes a term $B(y_c)$ from 3-parton states calculated at next-to-leading order, and a term $C(y_c)$ from 4-parton states calculated at leading order. Hence, the ratio of the strong coupling of quark type $j$ to the mean coupling in the sample of all flavors, $\alpha_s^j/\alpha_s^{all}$, can be determined from:

$$\frac{R_3^j(y_c)}{R_3^{all}(y_c)} = \frac{A(y_c)\alpha_s^j + [B(y_c) + C(y_c)](\alpha_s^j)^2}{A(y_c)\alpha_s^{all} + [B(y_c) + C(y_c)](\alpha_s^{all})^2} \tag{2}$$

where $A(y_c)$, $B(y_c)$, and $C(y_c)$ for the different jet-finding algorithms were evaluated using Refs. [13, 14]. Equation 2 was solved to obtain $\alpha_s^j/\alpha_s^{all}$ for each jet algorithm. As an example, for the Geneva algorithm $R_3^i/R_3^{all}$ for each subsample $i$ ($1 \leq i \leq 5$) and the unfolded results $\alpha_s^j/\alpha_s^{all}$ ($j = uds$, $c$, $b$) are shown in Fig. 2 as a function of $y_c$. Fig. 3 summarises the results for all algorithms; the errors are statistical only.

We considered experimental systematic effects that could modify the tagging efficiencies. In each case the error was evaluated by varying the appropriate parameter in the Monte Carlo simulation, recalculating the matrices $\varepsilon$, performing a new fit of Eq. 1, and rederiving $\alpha_s^j/\alpha_s^{all}$. Suitable variation about the world average value of each parameter was considered [10]. The errors are summarized in Table 2, where averages over the six algorithms are shown. The dominant physics contributions in $\alpha_s^b/\alpha_s^{all}$ result from limited knowledge of the average $B$ decay multiplicity and the heavy quark fragmentation functions. The uncertainty in $BR(Z^0 \to c\bar{c})$ also produces variations in $\alpha_s^c/\alpha_s^{all}$ and $\alpha_s^{uds}/\alpha_s^{all}$. Contributions from $B$ hadron lifetimes, the fraction of $D^+$ in $B$ meson decays, $b$ baryon production rates, and the charm hadron decay multiplicity are small. The detector systematic error is dominated by the uncertainty in the charged track reconstruction efficiency. No systematic variation of the results was found when the event selection cuts, tag criteria, or $y_c$ values were changed.

We considered sources of uncertainty in the QCD predictions that affect the values

of $\alpha_s^j/\alpha_s^{all}$ derived from Eq. 2. For each jet algorithm these include variation of the QCD renormalization scale within the range allowed by our measurements of jet rates in the global sample [15] and variation of the heavy quark masses used in the phase-space correction factors by $\pm 0.25$ GeV/$c^2$. The rates of gluon splitting to heavy flavors were also considered. The variation of the results due to uncertainties in parton production and hadronization was investigated [1] by using JETSET [11] and was found to be small. These contributions are listed in Table 3.

There is some scatter among the $\alpha_s^j/\alpha_s^{all}$ values derived from the different jet algorithms. In order to quote a single $\alpha_s^j/\alpha_s^{all}$ value for each flavor $j$, we made the conservative assumption that the results are completely correlated, and we calculated the unweighted mean values and errors over all six algorithms. We obtained

$$\begin{aligned}
\frac{\alpha_s^{uds}}{\alpha_s^{all}} &= 0.997 \pm 0.011\,(\text{stat}) \pm 0.011\,(\text{syst}) \pm 0.005\,(\text{theory})\ , \\
\frac{\alpha_s^{c}}{\alpha_s^{all}} &= 0.984 \pm 0.042\,(\text{stat}) \pm 0.053\,(\text{syst}) \pm 0.022\,(\text{theory})\ , \\
\frac{\alpha_s^{b}}{\alpha_s^{all}} &= 1.022 \pm 0.019\,(\text{stat}) \pm 0.023\,(\text{syst}) \pm 0.012\,(\text{theory})\ , \qquad (3)
\end{aligned}$$

where the theoretical uncertainty is the sum in quadrature of the QCD uncertainty and the r.m.s. of the results over the six algorithms (Table 3). These average values are also shown in Fig. 3.

# 5. Conclusion

We have used hadron lifetime information to separate hadronic $Z^0$ decays into three flavor samples with high efficiency and purity, and small bias against events containing hard gluon radiation. From a comparison of the rates of multijet events in these samples, we find that the strong coupling is independent of quark flavor within our sensitivity. Our results are consistent with our previous measurements [9], but are substantially more precise, as well as with measurements performed at LEP using different flavor-tagging techniques [4, 5].

# Acknowledgements

We thank the personnel of the SLAC accelerator department and the technical staffs of our collaborating institutions for their outstanding efforts on our behalf.

*This work was supported by Department of Energy contracts: DE-FG02-91ER40676 (BU), DE-FG03-91ER40618 (UCSB), DE-FG03-92ER40689 (UCSC), DE-FG03-93ER40788 (CSU), DE-FG02-91ER40672 (Colorado), DE-FG02-91ER40677 (Illinois), DE-AC03-76SF00098 (LBL), DE-FG02-92ER40715 (Massachusetts), DE-FC02-94ER40818 (MIT), DE-FG03-96ER40969 (Oregon), DE-AC03-76SF00515 (SLAC), DE-FG05-91ER40627 (Tennessee), DE-FG02-95ER40896 (Wisconsin), DE-FG02-92ER40704 (Yale); National Science Foundation grants: PHY-91-13428 (UCSC), PHY-89-21320 (Columbia), PHY-92-04239 (Cincinnati), PHY-95-10439 (Rutgers), PHY-88-19316 (Vanderbilt), PHY-92-03212 (Washington); The UK Particle Physics and Astronomy Research Council (Brunel and RAL); The Istituto Nazionale di Fisica Nucleare of Italy (Bologna, Ferrara, Frascati, Pisa, Padova, Perugia); The Japan-US Cooperative Research Project on High Energy Physics (Nagoya, Tohoku); The Korea Science and Engineering Foundation (Soongsil).

# References

[1] SLD Collaboration: K. Abe et al., Phys. Rev. **D51** 962 (1995).

[2] T. G. Rizzo, Phys. Rev. **D50** 4478 (1994).

[3] TASSO Collaboration: W. Braunschweig et al., Z. Phys. **C42** 17 (1989); W. Braunschweig et al., Z. Phys. **C44** 365 (1989).

[4] DELPHI Collaboration: P. Abreu et al., Phys. Lett. **B307** 221 (1993); L3 Collaboration: B. Adeva et al., Phys. Lett. **B271** 461 (1991);

OPAL Collaboration: M.Z. Akrawy et al., Z. Phys. **C65** 31 (1995);
ALEPH Collaboration: D. Buskulic et al., Phys. Lett. **B355** 381 (1995).

[5] OPAL Collaboration: R. Akers et al., Z. Phys. **C60** 397 (1993).

[6] SLD Design Report, SLAC Report–273 (1984), unpublished.

[7] M. D. Hildreth et al., Nucl. Instr. Meth. **A367** 111 (1995).

[8] C.J.S. Damerell et. al., Nucl. Instr. Meth. **A288** 236 (1990).

[9] SLD Collaboration: K. Abe et al., Phys. Rev. **D53** 2271 (1996).

[10] SLD Collaboration: K. Abe et al., Phys. Rev. **D53** 1023 (1996)

[11] T. Sjöstrand, Comput. Phys. Commun. **43** 367 (1987).

[12] P. N. Burrows, Z. Phys. **C41** 375 (1988);
OPAL Collaboration: M. Z. Akrawy et al., Z. Phys. **C47** 505 (1990).

[13] S. Bethke et al., Nucl. Phys. **B370** 310 (1992).

[14] Z. Kunszt et al., CERN 89–08 Vol. I, 373 (1989).

[15] SLD Collaboration: K. Abe et al., Phys. Rev. Lett. **71** 2528 (1993).

## **List of Authors

K. Abe,[19] K. Abe,[30] T. Akagi,[28] N.J. Allen,[4] W.W. Ash,[28]† D. Aston,[28]
K.G. Baird,[24] C. Baltay,[34] H.R. Band,[33] M.B. Barakat,[34] G. Baranko,[9]
O. Bardon,[15] T. L. Barklow,[28] G.L. Bashindzhagyan,[18] A.O. Bazarko,[10]
R. Ben-David,[34] A.C. Benvenuti,[2] G.M. Bilei,[22] D. Bisello,[21] G. Blaylock,[16]
J.R. Bogart,[28] B. Bolen,[17] T. Bolton,[10] G.R. Bower,[28] J.E. Brau,[20]

M. Breidenbach,[28] W.M. Bugg,[29] D. Burke,[28] T.H. Burnett,[32] P.N. Burrows,[15] W. Busza,[15] A. Calcaterra,[12] D.O. Caldwell,[5] D. Calloway,[28] B. Camanzi,[11] M. Carpinelli,[23] R. Cassell,[28] R. Castaldi,[23][a] A. Castro,[21] M. Cavalli-Sforza,[6] A. Chou,[28] E. Church,[32] H.O. Cohn,[29] J.A. Coller,[3] V. Cook,[32] R. Cotton,[4] R.F. Cowan,[15] D.G. Coyne,[6] G. Crawford,[28] A. D'Oliveira,[7] C.J.S. Damerell,[25] M. Daoudi,[28] R. De Sangro,[12] R. Dell'Orso,[23] P.J. Dervan,[4] M. Dima,[8] D.N. Dong,[15] P.Y.C. Du,[29] R. Dubois,[28] B.I. Eisenstein,[13] R. Elia,[28] E. Etzion,[33] S. Fahey,[9] D. Falciai,[22] C. Fan,[9] J.P. Fernandez,[6] M.J. Fero,[15] R. Frey,[20] T. Gillman,[25] G. Gladding,[13] S. Gonzalez,[15] E.L. Hart,[29] J.L. Harton,[8] A. Hasan,[4] Y. Hasegawa,[30] K. Hasuko,[30] S. J. Hedges,[3] S.S. Hertzbach,[16] M.D. Hildreth,[28] J. Huber,[20] M.E. Huffer,[28] E.W. Hughes,[28] H. Hwang,[20] Y. Iwasaki,[30] D.J. Jackson,[25] P. Jacques,[24] J. A. Jaros,[28] Z. Y. Jiang,[28] A.S. Johnson,[3] J.R. Johnson,[33] R.A. Johnson,[7] T. Junk,[28] R. Kajikawa,[19] M. Kalelkar,[24] H. J. Kang,[26] I. Karliner,[13] H. Kawahara,[28] H.W. Kendall,[15] Y. D. Kim,[26] M.E. King,[28] R. King,[28] R.R. Kofler,[16] N.M. Krishna,[9] R.S. Kroeger,[17] J.F. Labs,[28] M. Langston,[20] A. Lath,[15] J.A. Lauber,[9] D.W.G.S. Leith,[28] V. Lia,[15] M.X. Liu,[34] X. Liu,[6] M. Loreti,[21] A. Lu,[5] H.L. Lynch,[28] J. Ma,[32] G. Mancinelli,[24] S. Manly,[34] G. Mantovani,[22] T.W. Markiewicz,[28] T. Maruyama,[28] H. Masuda,[28] E. Mazzucato,[11] A.K. McKemey,[4] B.T. Meadows,[7] R. Messner,[28] P.M. Mockett,[32] K.C. Moffeit,[28] T.B. Moore,[34] D. Muller,[28] T. Nagamine,[28] S. Narita,[30] U. Nauenberg,[9] H. Neal,[28] M. Nussbaum,[7]† Y. Ohnishi,[19] N. Oishi,[19] D. Onoprienko,[29] L.S. Osborne,[15] R.S. Panvini,[31] C.H. Park,[27] H. Park,[20] T.J. Pavel,[28] I. Peruzzi,[12][b] M. Piccolo,[12] L. Piemontese,[11] E. Pieroni,[23] K.T. Pitts,[20] R.J. Plano,[24] R. Prepost,[33] C.Y. Prescott,[28] G.D. Punkar,[28] J. Quigley,[15] B.N. Ratcliff,[28] T.W. Reeves,[31] J. Reidy,[17] P.L. Reinertsen,[6] P.E. Rensing,[28] L.S. Rochester,[28] P.C. Rowson,[10] J.J. Russell,[28] O.H. Saxton,[28] T. Schalk,[6] R.H. Schindler,[28] B.A. Schumm,[6] J. Schwiening,[28] S. Sen,[34]

V.V. Serbo,[33] M.H. Shaevitz,[10] J.T. Shank,[3] G. Shapiro,[14] D.J. Sherden,[28] K.D. Shmakov,[29] C. Simopoulos,[28] N.B. Sinev,[20] S.R. Smith,[28] M.B. Smy,[8] J.A. Snyder,[34] H. Staengle,[8] P. Stamer,[24] H. Steiner,[14] R. Steiner,[1] M.G. Strauss,[16] D. Su,[28] F. Suekane,[30] A. Sugiyama,[19] S. Suzuki,[19] M. Swartz,[28] A. Szumilo,[32] T. Takahashi,[28] F.E. Taylor,[15] E. Torrence,[15] A.I. Trandafir,[16] J.D. Turk,[34] T. Usher,[28] J. Va'vra,[28] C. Vannini,[23] E. Vella,[28] J.P. Venuti,[31] R. Verdier,[15] P.G. Verdini,[23] D.L. Wagner,[9] S.R. Wagner,[28] A.P. Waite,[28] S.J. Watts,[4] A.W. Weidemann,[29] E.R. Weiss,[32] J.S. Whitaker,[3] S.L. White,[29] F.J. Wickens,[25] D.C. Williams,[15] S.H. Williams,[28] S. Willocq,[28] R.J. Wilson,[8] W.J. Wisniewski,[28] M. Woods,[28] G.B. Word,[24] J. Wyss,[21] R.K. Yamamoto,[15] J.M. Yamartino,[15] X. Yang,[20] J. Yashima,[30] S.J. Yellin,[5] C.C. Young,[28] H. Yuta,[30] G. Zapalac,[33] R.W. Zdarko,[28] and J. Zhou,[20]

[1] *Adelphi University, Garden City, New York 11530*

[2] *INFN Sezione di Bologna, I-40126 Bologna, Italy*

[3] *Boston University, Boston, Massachusetts 02215*

[4] *Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom*

[5] *University of California at Santa Barbara, Santa Barbara, California 93106*

[6] *University of California at Santa Cruz, Santa Cruz, California 95064*

[7] *University of Cincinnati, Cincinnati, Ohio 45221*

[8] *Colorado State University, Fort Collins, Colorado 80523*

[9] *University of Colorado, Boulder, Colorado 80309*

[10] *Columbia University, New York, New York 10027*

[11] *INFN Sezione di Ferrara and Università di Ferrara, I-44100 Ferrara, Italy*

[12] *INFN Lab. Nazionali di Frascati, I-00044 Frascati, Italy*

[13] *University of Illinois, Urbana, Illinois 61801*

[14] *E.O. Lawrence Berkeley Laboratory, University of California, Berkeley, California*

*94720*

[15] *Massachusetts Institute of Technology, Cambridge, Massachusetts 02139*

[16] *University of Massachusetts, Amherst, Massachusetts 01003*

[17] *University of Mississippi, University, Mississippi 38677*

[18] *Moscow State University, Institute of Nuclear Physics 119899 Moscow, Russia*

[19] *Nagoya University, Chikusa-ku, Nagoya 464 Japan*

[20] *University of Oregon, Eugene, Oregon 97403*

[21] *INFN Sezione di Padova and Università di Padova, I-35100 Padova, Italy*

[22] *INFN Sezione di Perugia and Università di Perugia, I-06100 Perugia, Italy*

[23] *INFN Sezione di Pisa and Università di Pisa, I-56100 Pisa, Italy*

[24] *Rutgers University, Piscataway, New Jersey 08855*

[25] *Rutherford Appleton Laboratory, Chilton, Didcot, Oxon OX11 0QX United Kingdom*

[26] *Sogang University, Seoul, Korea*

[27] *Soongsil University, Seoul, Korea 156-743*

[28] *Stanford Linear Accelerator Center, Stanford University, Stanford, California 94309*

[29] *University of Tennessee, Knoxville, Tennessee 37996*

[30] *Tohoku University, Sendai 980 Japan*

[31] *Vanderbilt University, Nashville, Tennessee 37235*

[32] *University of Washington, Seattle, Washington 98195*

[33] *University of Wisconsin, Madison, Wisconsin 53706*

[34] *Yale University, New Haven, Connecticut 06511*

† *Deceased*

[a] *Also at the Università di Genova*

[b] *Also at the Università di Perugia*

**Table 1.** Results for $R_3^j/R_3^{\rm all}$, derived from Eq. 1; see text. The errors shown are statistical.

| Algorithm | $y_c$ | $R_3^{uds}/R_3^{all}$ | $R_3^c/R_3^{all}$ | $R_3^b/R_3^{all}$ | $R_3^c/R_3^u$ factor | $R_3^b/R_3^d$ factor |
|---|---|---|---|---|---|---|
| E | 0.030 | $0.996 \pm 0.014$ | $0.945 \pm 0.055$ | $1.056 \pm 0.024$ | 0.997 | 1.004 |
| E0 | 0.020 | $0.993 \pm 0.012$ | $0.975 \pm 0.046$ | $1.039 \pm 0.020$ | 0.993 | 1.026 |
| P | 0.020 | $1.001 \pm 0.012$ | $0.930 \pm 0.047$ | $1.054 \pm 0.021$ | 0.992 | 1.024 |
| P0 | 0.015 | $0.989 \pm 0.010$ | $0.999 \pm 0.040$ | $1.034 \pm 0.018$ | 0.992 | 1.031 |
| D | 0.010 | $1.006 \pm 0.016$ | $1.021 \pm 0.062$ | $0.966 \pm 0.026$ | 1.001 | 0.916 |
| G | 0.040 | $1.011 \pm 0.014$ | $0.999 \pm 0.065$ | $0.969 \pm 0.024$ | 0.998 | 0.944 |

**Table 2.** Systematic Erorrs

| source | Center Value | Variation | $\Delta\left(\frac{\alpha_s^{uds}}{\alpha_s^{all}}\right)$ | $\Delta\left(\frac{\alpha_s^{c}}{\alpha_s^{all}}\right)$ | $\Delta\left(\frac{\alpha_s^{b}}{\alpha_s^{all}}\right)$ |
|---|---|---|---|---|---|
| $B$ decay multiplicity $< n_{ch} >$ | 5.39 | $\pm$0.2trks | 0.001 | 0.022 | 0.014 |
| $B$ fragmentation $< x_b >$ | 0.700 | $\pm$0.008 | 0.002 | 0.008 | 0.012 |
| $B$ meson lifetime $\tau_B$ | 1.55$ps$ | $\pm$0.05$ps$ | 0.001 | 0.008 | 0.003 |
| $B$ baryon lifetime $\tau_B$ | 1.10$ps$ | $\pm$0.08$ps$ | 0.001 | 0.002 | 0.001 |
| $B$ baryon prod. rate $f_{\Lambda_b}$ | 7% | $\pm$4% | 0.001 | 0.004 | 0.006 |
| $B \to D^+ + X$ fraction | 0.15 | $\pm$0.05 | 0.001 | 0.003 | 0.002 |
| $R_b$ | 0.2216 | $\pm$0.0017 | 0.001 | 0.003 | 0.001 |
| $R_c$ | 0.17 | $\pm$0.010 | 0.0008 | 0.037 | 0.005 |
| $C$ fragmentation $< x_c >$ | 0.484 | $\pm$0.008 | 0.003 | 0.011 | 0.002 |
| $c\bar{c} \to D^+ + X$ fraction | 0.20 | $\pm$0.04 | 0.001 | 0.003 | 0.001 |
| $C$ decay multiplicity $< n_{ch} >$ | 2.34 | $\pm$0.2trks | 0.003 | 0.008 | 0.006 |
| tracking efficiency | correction | on/off | 0.002 | 0.013 | 0.005 |
| impact parameter resolution | smear | on/off | 0.001 | 0.005 | 0.001 |
| MC statistics | 0.8M events | | 0.005 | 0.020 | 0.008 |
| Total | | | 0.011 | 0.053 | 0.023 |

**Table 3.** Theoretical Uncertainties

| | | | | | |
|---|---|---|---|---|---|
| $\alpha_s^{all}$ | 0.120 | $\pm$0.01 | 0.001 | 0.001 | 0.002 |
| $m_b$ | 5.0GeV/$c^2$ | $\pm$0.25GeV/$c^2$ | 0.001 | 0.002 | 0.003 |
| $m_c$ | 1.35GeV/$c^2$ | $\pm$0.25GeV/$c^2$ | 0.001 | 0.002 | 0.003 |
| $g \to cc$ | $1.7 \times 10^{-2}$ | $\pm$50% | 0.001 | 0.002 | 0.002 |
| $g \to bb$ | $1.6 \times 10^{-3}$ | $\pm$50% | 0.001 | 0.001 | 0.003 |
| Jet Algorithm | average | r.m.s | 0.004 | 0.022 | 0.011 |
| Total | | | 0.005 | 0.022 | 0.012 |




Figure 1: The distribution of the the number of tracks per event that miss the interaction point by $\geq 3\sigma$.




Figure 2: a) Detector level $R_3^{subsample}/R_3^{all}$. b) Unfolded $\alpha_s^{flavor}/\alpha_s^{all}$ (Geneva algorithm). The error bars are statistical only.




Figure 3: Values of $\alpha_s^j/\alpha_s^{all}$ derived for each of the six jet algorithms for each of the quark flavors $j$ (see text). The error bars on the averages include only statistical errors.