

Probing the Geometry of the Universe at the NLC¹

Hooman Davoudiasl

Stanford Linear Accelerator Center, Stanford University, Stanford, California 94309, USA

Abstract.

The hierarchy problem in particle physics has recently been approached from a geometric point of view in different models. These approaches postulate the existence of extra dimensions with various geometric properties, to explain how the hierarchy between the apparent scale of gravity $\bar{M}_P \sim 10^{18}$ GeV and the weak scale $m_W \sim 100$ GeV can be generated. Generally, these models predict that the effects of gravity mediated interactions become strong at the weak scale. This fact makes the NLC a promising tool for testing such extra dimensional models.

I INTRODUCTION

The gravitational potential $V(r)$ of a test mass m_t at a distance r is observed to be

$$V(r) = -G_N \frac{m_t}{r}, \quad (1)$$

where G_N is the 4- d Newton's constant. Thus, gravitational interactions can be described by a non-renormalizable field theory, where the spin-2 graviton mediates the force, and couples to the energy momentum tensor with dimensionful coupling $1/\bar{M}_P$, where $\bar{M}_P \sim G_N^{-1/2} \sim 10^{18}$ GeV. However, the electroweak interactions have a typical scale of order the W mass $m_W \sim 100$ GeV. If the Higgs boson of the Standard Model (SM) is responsible for the electroweak symmetry breaking, then we expect that the mass of the Higgs $m_H \sim m_W$. Thus, m_H seems to be stable against $O(\bar{M}_P)$ quantum corrections. Explaining the origin of the large ratio $\bar{M}_P/m_H \sim 10^{16}$ is referred to as the hierarchy problem in particle physics.

¹⁾ Presented at the 5th International Linear Collider Workshop (LCWS 2000), Batavia, Illinois, October 24-28, 2000. Work supported by the Department of Energy, contract DE-AC03-76SF00515.

There have been a number of proposals for solving the hierarchy problem. An interesting and theoretically appealing proposal is low energy supersymmetry. In a supersymmetric solution, new fields are added to the Lagrangian, such that every known field will have a superpartner of weak scale mass. However, there is, as yet, no experimental evidence for this and other ideas requiring the discovery of new particles around the weak scale.

A new class of ideas approaches the hierarchy problem from a geometric point of view. Instead of postulating extra fields, such as in supersymmetry, one postulates the existence of extra dimensions in the universe. Here, we present two models that approach the question of hierarchy from an extra dimensional viewpoint. The first one, due to Arkani-Hamed, Dimopoulos, and Dvali (ADD) [1] uses $n > 1$ large extra dimensions; we only briefly discuss this model. The second model, due to Randall and Sundrum(RS) [2], assumes a warped 5- d universe, and is the main subject of what follows.

II LARGE EXTRA DIMENSIONS

In the ADD model, the assumption is that the fundamental scale of gravity in $(4+n)$ - d is M_F . The gravitational potential $V(r)$ at distances $r \gg R$, where R is the typical size of the extra dimensions, is given by Gauss' law

$$V(r) = -G_N \frac{m_t}{M_F^{(2+n)} R^n r}. \quad (2)$$

To recover the observed gravitational force, we must have

$$\bar{M}_P^2 \sim M_F^{(2+n)} R^n. \quad (3)$$

Now, if we require that the $M_F \sim m_W$, in order to eliminate the hierarchy between the two scales, we are forced to have large extra dimensions of size $1 \text{ fm} \lesssim R \lesssim 1 \text{ mm}$, for $2 \leq n \leq 6$. The case $n = 1$ is ruled out, since it requires $R \sim 1 \text{ AU}$, which would result in deviations in Newtonian gravity at the scale of the solar system.

In the ADD scenario with large extra dimensions, (i) there is a Kaluza-Klein tower of gravitons with mass $m_n \sim n/R$ with equal spacing; (ii) each KK mode couples with $1/\bar{M}_P$ in 4- d ; (iii) the KK tower at energies $\sqrt{s} \sim M_F \sim 1 \text{ TeV}$ interacts strongly, only suppressed by $1/M_F$, due to the KK multiplicity of $O(10^{16})$; (iv) the SM resides on a 4- d wall in a $(4+n)$ - d bulk; (v) the geometry is factorizable, and the n extra dimensions are flat, that is the metric is of the form

$$ds^2 = \eta_{\mu\nu} dx^\mu dx^\nu + \sum_{i=4}^{3+n} dx_i^2. \quad (4)$$

III THE RS MODEL

This model [2] is based on a 5- d spacetime of constant negative curvature, called AdS_5 , truncated by two 4- d Minkowski walls, separated by a fixed distance $L = \pi r_c$ with $r_c \sim \bar{M}_P^{-1}$ as the compactification scale; the 5th dimension y is parameterized by an angular variable $\phi \in [-\pi, \pi]$ and $y = \phi r_c$. The geometry is required to respect the Z_2 symmetry $\phi \rightarrow -\phi$. The ‘‘Planck wall’’ is at $\phi = 0$, whereas the ‘‘SM wall’’, corresponding to the visible 4- d universe, is at $\phi = \pi$. The energy density on the Planck wall V_P is equal and opposite to that on the SM wall and we have $V_P \sim M_5^3 k$, where $M_5 \sim \bar{M}_P$ is the fundamental 5- d scale, and $k \sim \bar{M}_P$ is the curvature scale. The 5- d cosmological constant is given by $\Lambda_5 = -kV_P$. Thus, we see that the parameters of the model do not establish new hierarchies.

The geometry of this model is warped and non-factorizable, with the metric

$$ds^2 = e^{-2\sigma(\phi)} \eta_{\mu\nu} dx^\mu dx^\nu + r_c^2 d\phi^2 ; \quad \sigma(\phi) = k r_c |\phi|, \quad (5)$$

where $e^{-2\sigma(\phi)}$ is the warp factor. This geometric warp factor offers a possible explanation of the hierarchy problem. Basically, if one writes down a 5- d action with Planckian mass parameters $m_5 \sim \bar{M}_P$, after a KK reduction to 4 dimensions, the 4- d fields with canonical 4- d kinetic terms will have mass parameters $m_4 = m_5 e^{-k r_c \pi}$. To have $m_4 \sim m_W$, we only need to require $kr_c \sim 10$, which has been shown to be easily realized in a mechanism that stabilizes the size of the 5th dimension [3]. In this way, numbers of $O(10)$ generate large hierarchies of $O(10^{16})$.

This model has features that are quite distinct from the ones of the ADD model. In the RS model [2,4] (i) the KK tower of gravitons starts at $m \sim 1$ TeV, the spacings between the tower masses $\Delta m \sim 1$ TeV are unequal and given by roots of Bessel functions; (ii) the zero mode (massless 4- d) graviton couples with $1/\bar{M}_P$ and the massive KK tower gravitons couple with $1/\Lambda_\pi \sim 1$ TeV⁻¹; this can be understood by noting that the wavefunction of the zero mode along the 5th dimension is localized near the Planck wall, characterized by \bar{M}_P , whereas the KK graviton wavefunctions are localized near the SM wall, characterized by Λ_π . The RS-type models are sometimes referred to as ‘‘Localized Gravity’’ models. (iii) In the original proposal by Randall and Sundrum [2], the SM fields are taken to reside only on the SM wall. With these features, the RS model predicts resonant production of KK gravitons at $\sqrt{s} \sim 1$ TeV at colliders such as the NLC.

There have been a number of generalizations and extensions of the original RS proposal. Some of these extensions study the possibility of having SM fields in the bulk and deriving the 4- d physics from the 5- d picture [5], since the SM scale of order 1 TeV can be generated on the SM wall through the warp factor. For various phenomenological reasons it is least problematic to keep the Higgs field on the SM wall [6–8]. However, as a first step, one can study the effect of placing the SM gauge fields in the bulk and keeping the fermions on the SM wall [9,10]. In this case, one finds that the fermions on the wall couple to the KK gauge fields $\sqrt{2kr_c\pi} \sim 10$ times more strongly than they couple to the zero mode gauge fields (γ, g, W^\pm, Z).

Here one expects to get strong constraints from data, and indeed agreement with precision electroweak data requires that the lightest KK gauge boson have a mass $m_1^{(A)} \gtrsim 23$ TeV. This value pushes the scale on the SM wall far above ~ 1 TeV, making this scenario disfavored in the context of the hierarchy problem.

The above bound can be somewhat relaxed, if the fermions also reside in the bulk [6]. In fact, by introducing bulk fermion 5- d masses m_Ψ , one can change the couplings of the fermion zero modes (observed SM fermions) to various KK fields, and place different bounds on the RS model, depending on the value of the bulk mass parameter $\nu \equiv m_\Psi/k$ [11,7,8]. The parameter ν controls the shape of the fermion zero mode wavefunction $f^{(0)} \sim e^{\nu\sigma(\phi)}$. Thus, negative values of ν localize $f^{(0)}$ near the Planck wall, whereas positive values of ν localize $f^{(0)}$ near the SM wall.

Avoiding large FCNC's may require keeping the value of ν nearly universal for all fermions. The 4- d Yukawa couplings of SM fermions depend on the value of ν . It can be shown that keeping the 5- d Yukawa couplings $\lambda_5 \sim 1$ requires that the $\nu \gtrsim -0.8$ (for the lightest fermion) [7,8]. Avoiding the generation of a new hierarchy, on the other hand, forces us to have $\nu \lesssim -0.3$. This can be understood by noting that as ν increases, the fermions get more and more localized toward the SM wall, and this takes us back to the case where leaving the fermions on the wall gave us a large lower bound on the mass of the first KK gauge field [8]. Although placing the SM gauge and fermion fields in the 5- d bulk results in a rich phenomenology [8], the range where the theory seems viable is rather narrow, and keeping the SM fields on the wall appears to be less problematic.

IV CONCLUSIONS

The hierarchy problem can be approached from a geometric $(4+n)$ - d point of view, in the ADD and the RS scenarios. Each scenario predicts a distinct set of signatures at $\sqrt{s} \sim 1$ TeV. Thus, an NLC with $\sqrt{s} \sim 1$ TeV can test these ideas and possibly yield information on the geometry of the extra dimensions of the universe by probing such features as their number, size, and curvature.

REFERENCES

1. N. Arkani-Hamed, S. Dimopoulos, and G. Dvali, Phys. Lett. **B429**, 263 (1998); Phys. Rev. **D59**, 086004 (1999).
2. L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999).
3. W. D. Goldberger and M. B. Wise, Phys. Rev. Lett. **83**, 4922 (1999).
4. H. Davoudiasl, J. L. Hewett and T. G. Rizzo, Phys. Rev. Lett. **84**, 2080 (2000).
5. W. D. Goldberger and M. B. Wise, Phys. Rev. D **60**, 107505 (1999).
6. S. Chang, J. Hisano, H. Nakano, N. Okada and M. Yamaguchi, Phys. Rev. D **62**, 084025 (2000).
7. T. Gherghetta and A. Pomarol, Nucl. Phys. **B586**, 141 (2000).

8. H. Davoudiasl, J. L. Hewett and T. G. Rizzo, hep-ph/0006041.
9. H. Davoudiasl, J. L. Hewett and T. G. Rizzo, Phys. Lett. **B473**, 43 (2000).
10. A. Pomarol, Phys. Lett. **B486**, 153 (2000).
11. Y. Grossman and M. Neubert, Phys. Lett. **B474**, 361 (2000).