

List of other measurements that are not included in the tables:

- $B^+ \rightarrow K^+\pi^-\pi^+\gamma$: LHCb has measured the up-down asymmetries in bins of the $K\pi\pi\gamma$ mass [1].
- In [2], LHCb has also measured the branching fraction of $B^+ \rightarrow K^+e^-e^+$ in the $m^2(\ell\ell)$ bin [1, 6] GeV^2/c^4 .
- In the $B^+ \rightarrow \pi^+\mu^+\mu^-$ paper [3], LHCb has also measured the differential branching fraction in bins of $m^2(\ell\ell)$.
- For $B \rightarrow K\ell^-\ell^+$, LHCb has measured F_H and A_{FB} in 17 (5) bins of $m^2(\ell\ell)$ for the K^+ (K_S^0) final state [4]. Belle has measured F_L and A_{FB} in 6 $m^2(\ell\ell)$ bins [64].
- For the $B \rightarrow K^*\ell^-\ell^+$ analyses, partial branching fractions and angular observables in bins of $m^2(\ell\ell)$ are also available:
 - $B^0 \rightarrow K^{*0}e^-e^+$: LHCb has measured F_L , $A_T^{(2)}$, A_T^{Im} , A_T^{Re} in the [0.002, 1.120] GeV^2/c^4 bin of $m^2(\ell\ell)$ [5], and has also determined the branching fraction in the dilepton mass region [10, 1000] MeV/c^2 [2].
 - $B \rightarrow K^*\ell^-\ell^+$: Belle has measured F_L , A_{FB} , isospin asymmetry in 6 $m^2(\ell\ell)$ bins [6] [41] and P'_4 , P'_5 , P'_6 , P'_8 in 4 $m^2(\ell\ell)$ bins [7]. In a more recent paper [8], they report measurements of P'_4 and P'_5 , separately for $\ell = \mu$ or e , in 4 $m^2(\ell\ell)$ bins and in the region [1, 6] GeV^2/c^4 bin of $m^2(\ell\ell)$. The measurements use both B^0 and B^+ decays. They also measure the LFV observables $Q_i = P_i^\mu - P_i^e$, for $i = 4, 5$. BABAR has measured F_L , A_{FB} , P_2 in 5 $m^2(\ell\ell)$ bins [9].
 - $B^0 \rightarrow K^{*0}\mu^-\mu^+$: LHCb has measured F_L , A_{FB} , $S_3 - S_9$, $A_3 - A_9$, $P_1 - P_3$, $P'_4 - P'_8$ in 8 $m^2(\ell\ell)$ bins [10]. CMS has measured F_L and A_{FB} in 7 $m^2(\ell\ell)$ bins [11], and P_1, P'_5 in [12]. ATLAS has measured F_L , $S_{3,4,5,7,8}$ and $P'_{1,4,5,6,8}$ in 6 $m^2(\ell\ell)$ bins [13].
- For $B \rightarrow X_s\ell^-\ell^+$ (X_s is a hadronic system with an s quark), Belle has measured A_{FB} in bins of $m^2(\ell\ell)$ with a sum of 10 exclusive final states [14].
- $B^0 \rightarrow K^+\pi^-\mu^+\mu^-$, with $1330 < m(K^+\pi^-) < 1530$ GeV/c^2 : LHCb has measured the partial branching fraction in bins of $m^2(\mu^-\mu^+)$ in the range [0.1, 8.0] GeV^2/c^4 , and has also determined angular moments [15].
- In [16], LHCb measures the phase difference between the short- and long-distance contributions to the $B^+ \rightarrow K^+\mu^+\mu^-$ decay. The measurement is based on the analysis of the dimuon mass distribution in the regions of the J/ψ and $\psi(2S)$ resonances and far from their poles, to probe long and short distance effects, respectively.
- In [17] LHCb performs a search for a hypothetical new scalar particle χ , assumed to have a narrow width, through the decay $B^+ \rightarrow \chi(\mu^+\mu^-)$ in the ranges of mass $250 < m(\chi) < 4700$ MeV/c^2 and lifetime $0.1 < \tau(\chi) < 1000$ ps. Upper limits are given as a function of $m(\chi)$ and $\tau(\chi)$.

References

- [1] R. Aaij *et al.*, (LHCb collaboration), Phys. Rev. Lett. **112**, 161801, (2014), [arXiv:1402.6852](https://arxiv.org/abs/1402.6852) [hep-ex].

- [2] R. Aaij *et al.*, (LHCb collaboration), JHEP **05**, 159, (2013), [arXiv:1304.3035 \[hep-ex\]](#).
- [3] R. Aaij *et al.*, (LHCb collaboration), JHEP **10**, 034, (2015), [arXiv:1509.00414 \[hep-ex\]](#).
- [4] R. Aaij *et al.*, (LHCb collaboration), JHEP **05**, 082, (2014), [arXiv:1403.8045 \[hep-ex\]](#).
- [5] R. Aaij *et al.*, (LHCb collaboration), JHEP **04**, 064, (2015), [arXiv:1501.03038 \[hep-ex\]](#).
- [6] J. T. Wei *et al.*, (Belle collaboration), Phys. Rev. Lett. **103**, 171801, (2009), [arXiv:0904.0770 \[hep-ex\]](#).
- [7] A. Abdesselam *et al.*, (Belle collaboration), [arXiv:1604.04042 \[hep-ex\]](#), (2016).
- [8] S. Wehle *et al.*, (Belle collaboration), Phys. Rev. Lett. **118**, no. 11, 111801, (2017), [arXiv:1612.05014 \[hep-ex\]](#).
- [9] J. P. Lees *et al.*, (BABAR collaboration), Phys. Rev. **D93**, 052015, (2016), [arXiv:1508.07960 \[hep-ex\]](#).
- [10] R. Aaij *et al.*, (LHCb collaboration), JHEP **02**, 104, (2016), [arXiv:1512.04442 \[hep-ex\]](#).
- [11] V. Khachatryan *et al.*, (CMS collaboration), Phys. Lett. **B753**, 424, (2016), [arXiv:1507.08126 \[hep-ex\]](#).
- [12] A. M. Sirunyan *et al.*, (CMS collaboration), (2017), [arXiv:1710.02846 \[hep-ex\]](#).
- [13] (ATLAS Collaboration collaboration), “Angular analysis of $B_d^0 \rightarrow K^* \mu^+ \mu^-$ decays in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Tech. Rep. ATLAS-CONF-2017-023, CERN, Geneva, Apr, 2017, <http://cds.cern.ch/record/2258146>.
- [14] Y. Sato *et al.*, (Belle collaboration), Phys. Rev. **D93**, 032008, (2016), [arXiv:1402.7134 \[hep-ex\]](#), Addendum *ibid.* **D93**, 059901, (2016).
- [15] R. Aaij *et al.*, (LHCb collaboration), JHEP **12**, 065, (2016), [arXiv:1609.04736 \[hep-ex\]](#).
- [16] R. Aaij *et al.*, (LHCb collaboration), Eur. Phys. J. **C77**, no. 3, 161, (2017), [arXiv:1612.06764 \[hep-ex\]](#).
- [17] R. Aaij *et al.*, (LHCb collaboration), Phys. Rev. **D95**, no. 7, 071101, (2017), [arXiv:1612.07818 \[hep-ex\]](#).