Measurement of the Time-Dependent CP Asymmetry in the $B^0 \rightarrow \phi K^0$ Decay

B. Aubert,¹ R. Barate,¹ D. Boutigny,¹ F. Couderc,¹ J.-M. Gaillard,¹ A. Hicheur,¹ Y. Karyotakis,¹ J. P. Lees,¹
 V. Tisserand,¹ A. Zghiche,¹ A. Palano,² A. Pompili,² J. C. Chen,³ N. D. Qi,³ G. Rong,³ P. Wang,³ Y. S. Zhu,³
 G. Eigen,⁴ I. Ofte,⁴ B. Stugu,⁴ G. S. Abrams,⁵ A. W. Borgland,⁵ A. B. Breon,⁵ D. N. Brown,⁵ J. Button-Shafer,⁵
 R. N. Cahn,⁵ E. Charles,⁵ C. T. Day,⁵ M. S. Gill,⁵ A. V. Gritsan,⁵ Y. Groysman,⁵ R. G. Jacobsen,⁵ R. W. Kadel,⁵
 J. Kadyk,⁵ L. T. Kerth,⁵ Yu. G. Kolomensky,⁵ G. Kukartsev,⁵ C. LeClerc,⁵ G. Lynch,⁵ A. M. Merchant,⁵
 L. M. Mir,⁵ P. J. Oddone,⁵ T. J. Orimoto,⁵ M. Pripstein,⁵ N. A. Roe,⁵ M. T. Ronan,⁵ V. G. Shelkov,⁵
 A. V. Telnov,⁵ W. A. Wenzel,⁵ K. Ford,⁶ T. J. Harrison,⁶ C. M. Hawkes,⁶ S. E. Morgan,⁶ A. T. Watson,⁶
 M. Fritsch,⁷ K. Goetzen,⁷ T. Held,⁷ H. Koch,⁷ B. Lewandowski,⁷ M. Pelizaeus,⁷ M. Steinke,⁷ J. T. Boyd,⁸
 N. Chevalier,⁸ W. N. Cottingham,⁸ M. P. Kelly,⁸ T. E. Latham,⁸ F. F. Wilson,⁸ T. Cuhadar-Donszelmann,⁹
 C. Hearty,⁹ T. S. Mattison,⁹ J. A. McKenna,⁹ D. Thiessen,⁹ P. Kyberd,¹⁰ L. Teodorescu,¹⁰ V. E. Blinov,¹¹
 A. D. Bukin,¹¹ V. P. Druzhinin,¹¹ V. B. Golubev,¹¹ V. N. Ivanchenko,¹¹ E. A. Kravchenko,¹¹ A. P. Onuchin,¹¹
 S. I. Serednyakov,¹¹ Yu. I. Skovpen,¹¹ E. P. Solodov,¹¹ A. N. Yushkov,¹¹ D. Best,¹² M. Bruinsma,¹² M. Chao,¹²
 I. Eschrich,¹² D. Kirkby,¹² A. J. Lankford,¹² M. Mandelkern,¹² R. K. Mommsen,¹² W. Roethel,¹² D. P. Stoker,¹²
 C. Buchanan,¹³ B. L. Hartfiel,¹³ J. W. Gary,¹⁴ B. C. Shen,¹⁴ K. Wang,¹⁴ D. del Re,¹⁵ H. K. Hadavand,¹⁵
 E. J. Hill,¹⁵ D. B. MacFarlane,¹⁵ H. P. Paar,¹⁵ Sh. Rahatlou,¹⁵ V. Sharma,¹⁵ J. W. Berryhill,¹⁶ C. Campagnari,¹⁶
 B. Dahmes,¹⁶ S. L. Levy,¹⁶ O. Long,¹⁶ A. Lu,¹⁶ M. A. Mazur,¹⁶ J. D. Richman,¹⁶ W. Verkerke,¹⁶ T. W. Beck,¹⁷
 A. M. Eisner,¹⁷ C. A. Heusch,¹⁷ W. S. Lockman,¹⁷ T. Schalk,¹⁷ R. E. Schmitz,¹⁷ B. A. Schumm,¹⁷ A. Seiden,¹⁷
 P. Spradlin,¹⁷ D. C. Williams,¹⁷ M. G. Wilson,¹⁷ J. Albert,¹⁸ E. Chen,¹⁸ G. P. Dubois-Felsmann,¹⁸ A. Dvoretzskii,¹⁸
 D. G. Hitlin,¹⁸ I. Narsky,¹⁸ T. Piatenko,¹⁸ F. C. Porter,¹⁸ A. Ryd,¹⁸ A. Samuel,¹⁸ S. Yang,¹⁸ S. Jayatilleke,¹⁹
 G. Mancinelli,¹⁹ B. T. Meadows,¹⁹ M. D. Sokoloff,¹⁹ T. Abe,²⁰ F. Blanc,²⁰ P. Bloom,²⁰ S. Chen,²⁰ P. J. Clark,²⁰
 W. T. Ford,²⁰ U. Nauenberg,²⁰ A. Olivas,²⁰ P. Rankin,²⁰ J. G. Smith,²⁰ L. Zhang,²⁰ A. Chen,²¹ J. L. Harton,²¹
 A. Soffer,²¹ W. H. Toki,²¹ R. J. Wilson,²¹ Q. L. Zeng,²¹ D. Altenburg,²² T. Brandt,²² J. Brose,²² T. Colberg,²²
 M. Dickopp,²² E. Feltresi,²² A. Hauke,²² H. M. Lacker,²² E. Maly,²² R. Müller-Pfefferkorn,²² R. Nogowski,²²
 S. Otto,²² A. Petzold,²² J. Schubert,²² K. R. Schubert,²² R. Schwierz,²² B. Spaan,²² J. E. Sundermann,²²
 D. Bernard,²³ G. R. Bonneaud,²³ F. Brochard,²³ P. Grenier,²³ S. Schrenk,²³ Ch. Thiebaux,²³ G. Vasileiadis,²³
 M. Verderi,²³ D. J. Bard,²⁴ A. Khan,²⁴ D. Lavin,²⁴ F. Muheim,²⁴ S. Playfer,²⁴ M. Andreotti,²⁵ V. Azzolini,²⁵
 D. Bettoni,²⁵ C. Bozzi,²⁵ R. Calabrese,²⁵ G. Cibinetto,²⁵ E. Luppi,²⁵ M. Negrini,²⁵ L. Piemontese,²⁵ A. Sarti,²⁵
 E. Treadwell,²⁶ R. Baldini-Ferroli,²⁷ A. Calcaterra,²⁷ R. de Sangro,²⁷ G. Finocchiaro,²⁷ P. Patteri,²⁷ M. Piccolo,²⁷
 A. Zallo,²⁷ A. Buzzo,²⁸ R. Capra,²⁸ R. Contri,²⁸ G. Crosetti,²⁸ M. Lo Vetere,²⁸ M. Macri,²⁸ M. R. Monge,²⁸
 S. Passaggio,²⁸ C. Patrignani,²⁸ E. Robutti,²⁸ A. Santroni,²⁸ S. Tosi,²⁸ S. Bailey,²⁹ G. Brandenburg,²⁹ M. Morii,²⁹
 E. Won,²⁹ R. S. Dubitzky,³⁰ U. Langenegger,³⁰ W. Bhimji,³¹ D. A. Bowerman,³¹ P. D. Dauncey,³¹ U. Egede,³¹
 J. R. Gaillard,³¹ G. W. Morton,³¹ J. A. Nash,³¹ G. P. Taylor,³¹ G. J. Grenier,³² U. Mallik,³² J. Cochran,³³
 H. B. Crawley,³³ J. Lamsa,³³ W. T. Meyer,³³ S. Prell,³³ E. I. Rosenberg,³³ J. Yi,³³ M. Davier,³⁴ G. Grosdidier,³⁴
 A. Höcker,³⁴ S. Laplace,³⁴ F. Le Diberder,³⁴ V. Lepeltier,³⁴ A. M. Lutz,³⁴ T. C. Petersen,³⁴ S. Plaszczynski,³⁴
 M. H. Schune,³⁴ L. Tantot,³⁴ G. Wormser,³⁴ C. H. Cheng,³⁵ D. J. Lange,³⁵ M. C. Simani,³⁵ D. M. Wright,³⁵
 A. J. Bevan,³⁶ J. P. Coleman,³⁶ J. R. Fry,³⁶ E. Gabathuler,³⁶ R. Gamet,³⁶ R. J. Parry,³⁶ D. J. Payne,³⁶
 R. J. Sloane,³⁶ C. Touramanis,³⁶ J. J. Back,³⁷ C. M. Cormack,³⁷ P. F. Harrison,³⁷ * G. B. Mohanty,³⁷
 C. L. Brown,³⁸ G. Cowan,³⁸ R. L. Flack,³⁸ H. U. Flaecher,³⁸ M. G. Green,³⁸ C. E. Marker,³⁸ T. R. McMahon,³⁸
 S. Ricciardi,³⁸ F. Salvatore,³⁸ G. Vaitsas,³⁸ M. A. Winter,³⁸ D. Brown,³⁹ C. L. Davis,³⁹ J. Allison,⁴⁰ N. R. Barlow,⁴⁰
 R. J. Barlow,⁴⁰ P. A. Hart,⁴⁰ M. C. Hodgkinson,⁴⁰ G. D. Lafferty,⁴⁰ A. J. Lyon,⁴⁰ J. C. Williams,⁴⁰ A. Farbin,⁴¹
 W. D. Hulsbergen,⁴¹ A. Jawahery,⁴¹ D. Kovalskyi,⁴¹ C. K. Lae,⁴¹ V. Lillard,⁴¹ D. A. Roberts,⁴¹ G. Blaylock,⁴²
 C. Dallapiccola,⁴² K. T. Flood,⁴² S. S. Hertzbach,⁴² R. Kofler,⁴² V. B. Koptchev,⁴² T. B. Moore,⁴² S. Saremi,⁴²
 H. Staengle,⁴² S. Willocq,⁴² R. Cowan,⁴³ G. Sciolla,⁴³ F. Taylor,⁴³ R. K. Yamamoto,⁴³ D. J. J. Mangeol,⁴⁴
 P. M. Patel,⁴⁴ S. H. Robertson,⁴⁴ A. Lazzaro,⁴⁵ F. Palombo,⁴⁵ J. M. Bauer,⁴⁶ L. Cremaldi,⁴⁶ V. Eschenburg,⁴⁶

R. Godang,⁴⁶ R. Kroeger,⁴⁶ J. Reidy,⁴⁶ D. A. Sanders,⁴⁶ D. J. Summers,⁴⁶ H. W. Zhao,⁴⁶ S. Brunet,⁴⁷ D. Côté,⁴⁷ P. Taras,⁴⁷ H. Nicholson,⁴⁸ N. Cavallo,⁴⁹ F. Fabozzi,^{49,†} C. Gatto,⁴⁹ L. Lista,⁴⁹ D. Monorchio,⁴⁹ P. Paolucci,⁴⁹ D. Piccolo,⁴⁹ C. Sciacca,⁴⁹ M. Baak,⁵⁰ H. Bulten,⁵⁰ G. Raven,⁵⁰ L. Wilden,⁵⁰ C. P. Jessop,⁵¹ J. M. LoSecco,⁵¹ T. A. Gabriel,⁵² T. Allmendinger,⁵³ B. Brau,⁵³ K. K. Gan,⁵³ K. Honscheid,⁵³ D. Hufnagel,⁵³ H. Kagan,⁵³ R. Kass,⁵³ T. Pulliam,⁵³ A. M. Rahimi,⁵³ R. Ter-Antonyan,⁵³ Q. K. Wong,⁵³ J. Brau,⁵⁴ R. Frey,⁵⁴ O. Igonkina,⁵⁴ C. T. Potter,⁵⁴ N. B. Sinev,⁵⁴ D. Strom,⁵⁴ E. Torrence,⁵⁴ F. Colecchia,⁵⁵ A. Dorigo,⁵⁵ F. Galeazzi,⁵⁵ M. Margoni,⁵⁵ M. Morandin,⁵⁵ M. Posocco,⁵⁵ M. Rotondo,⁵⁵ F. Simonetto,⁵⁵ R. Stroili,⁵⁵ G. Tiozzo,⁵⁵ C. Voci,⁵⁵ M. Benayoun,⁵⁶ H. Briand,⁵⁶ J. Chauveau,⁵⁶ P. David,⁵⁶ Ch. de la Vaissière,⁵⁶ L. Del Buono,⁵⁶ O. Hamon,⁵⁶ M. J. J. John,⁵⁶ Ph. Leruste,⁵⁶ J. Ocariz,⁵⁶ M. Pivk,⁵⁶ L. Roos,⁵⁶ S. T’Jampens,⁵⁶ G. Therin,⁵⁶ P. F. Manfredi,⁵⁷ V. Re,⁵⁷ P. K. Behera,⁵⁸ L. Gladney,⁵⁸ Q. H. Guo,⁵⁸ J. Panetta,⁵⁸ F. Anulli,^{27,59} M. Biasini,⁵⁹ I. M. Peruzzi,^{27,59} M. Pioppi,⁵⁹ C. Angelini,⁶⁰ G. Batignani,⁶⁰ S. Bettarini,⁶⁰ M. Bondioli,⁶⁰ F. Bucci,⁶⁰ G. Calderini,⁶⁰ M. Carpinelli,⁶⁰ V. Del Gamba,⁶⁰ F. Forti,⁶⁰ M. A. Giorgi,⁶⁰ A. Lusiani,⁶⁰ G. Marchiori,⁶⁰ F. Martinez-Vidal,^{60,‡} M. Morganti,⁶⁰ N. Neri,⁶⁰ E. Paoloni,⁶⁰ M. Rama,⁶⁰ G. Rizzo,⁶⁰ F. Sandrelli,⁶⁰ J. Walsh,⁶⁰ M. Haire,⁶¹ D. Judd,⁶¹ K. Paick,⁶¹ D. E. Wagoner,⁶¹ N. Danielson,⁶² P. Elmer,⁶² C. Lu,⁶² V. Miftakov,⁶² J. Olsen,⁶² A. J. S. Smith,⁶² F. Bellini,⁶³ G. Cavoto,^{62,63} R. Faccini,⁶³ F. Ferrarotto,⁶³ F. Ferroni,⁶³ M. Gaspero,⁶³ L. Li Gioi,⁶³ M. A. Mazzoni,⁶³ S. Morganti,⁶³ M. Pierini,⁶³ G. Piredda,⁶³ F. Safai Tehrani,⁶³ C. Voena,⁶³ S. Christ,⁶⁴ G. Wagner,⁶⁴ R. Waldi,⁶⁴ T. Adye,⁶⁵ N. De Groot,⁶⁵ B. Franek,⁶⁵ N. I. Geddes,⁶⁵ G. P. Gopal,⁶⁵ E. O. Olaiya,⁶⁵ R. Aleksan,⁶⁶ S. Emery,⁶⁶ A. Gaidot,⁶⁶ S. F. Ganzhur,⁶⁶ P.-F. Giraud,⁶⁶ G. Hamel de Monchenault,⁶⁶ W. Kozanecki,⁶⁶ M. Langer,⁶⁶ M. Legendre,⁶⁶ G. W. London,⁶⁶ B. Mayer,⁶⁶ G. Schott,⁶⁶ G. Vasseur,⁶⁶ Ch. Yèche,⁶⁶ M. Zito,⁶⁶ M. V. Purohit,⁶⁷ A. W. Weidemann,⁶⁷ F. X. Yumiceva,⁶⁷ D. Aston,⁶⁸ R. Bartoldus,⁶⁸ N. Berger,⁶⁸ A. M. Boyarski,⁶⁸ O. L. Buchmueller,⁶⁸ M. R. Convery,⁶⁸ M. Cristinziani,⁶⁸ G. De Nardo,⁶⁸ D. Dong,⁶⁸ J. Dorfan,⁶⁸ D. Dujmic,⁶⁸ W. Dunwoodie,⁶⁸ E. E. Elsen,⁶⁸ S. Fan,⁶⁸ R. C. Field,⁶⁸ T. Glanzman,⁶⁸ S. J. Gowdy,⁶⁸ T. Hadig,⁶⁸ V. Halyo,⁶⁸ C. Hast,⁶⁸ T. Hryn’ova,⁶⁸ W. R. Innes,⁶⁸ M. H. Kelsey,⁶⁸ P. Kim,⁶⁸ M. L. Kocian,⁶⁸ D. W. G. S. Leith,⁶⁸ J. Libby,⁶⁸ S. Luitz,⁶⁸ V. Luth,⁶⁸ H. L. Lynch,⁶⁸ H. Marsiske,⁶⁸ R. Messner,⁶⁸ D. R. Muller,⁶⁸ C. P. O’Grady,⁶⁸ V. E. Ozcan,⁶⁸ A. Perazzo,⁶⁸ M. Perl,⁶⁸ S. Petrak,⁶⁸ B. N. Ratcliff,⁶⁸ A. Roodman,⁶⁸ A. A. Salnikov,⁶⁸ R. H. Schindler,⁶⁸ J. Schwiening,⁶⁸ G. Simi,⁶⁸ A. Snyder,⁶⁸ A. Soha,⁶⁸ J. Stelzer,⁶⁸ D. Su,⁶⁸ M. K. Sullivan,⁶⁸ J. Va’vra,⁶⁸ S. R. Wagner,⁶⁸ M. Weaver,⁶⁸ A. J. R. Weinstein,⁶⁸ W. J. Wisniewski,⁶⁸ M. Wittgen,⁶⁸ D. H. Wright,⁶⁸ A. K. Yarritu,⁶⁸ C. C. Young,⁶⁸ P. R. Burchat,⁶⁹ A. J. Edwards,⁶⁹ T. I. Meyer,⁶⁹ B. A. Petersen,⁶⁹ C. Roat,⁶⁹ S. Ahmed,⁷⁰ M. S. Alam,⁷⁰ J. A. Ernst,⁷⁰ M. A. Saeed,⁷⁰ M. Saleem,⁷⁰ F. R. Wappler,⁷⁰ W. Bugg,⁷¹ M. Krishnamurthy,⁷¹ S. M. Spanier,⁷¹ R. Eckmann,⁷² H. Kim,⁷² J. L. Ritchie,⁷² A. Satpathy,⁷² R. F. Schwitters,⁷² J. M. Izen,⁷³ I. Kitayama,⁷³ X. C. Lou,⁷³ S. Ye,⁷³ F. Bianchi,⁷⁴ M. Bona,⁷⁴ F. Gallo,⁷⁴ D. Gamba,⁷⁴ C. Borean,⁷⁵ L. Bosio,⁷⁵ C. Cartaro,⁷⁵ F. Cossutti,⁷⁵ G. Della Ricca,⁷⁵ S. Dittongo,⁷⁵ S. Grancagnolo,⁷⁵ L. Lanceri,⁷⁵ P. Poropat,^{75,§} L. Vitale,⁷⁵ G. Vuagnin,⁷⁵ R. S. Panvini,⁷⁶ Sw. Banerjee,⁷⁷ C. M. Brown,⁷⁷ D. Fortin,⁷⁷ P. D. Jackson,⁷⁷ R. Kowalewski,⁷⁷ J. M. Roney,⁷⁷ H. R. Band,⁷⁸ S. Dasu,⁷⁸ M. Datta,⁷⁸ A. M. Eichenbaum,⁷⁸ J. J. Hollar,⁷⁸ J. R. Johnson,⁷⁸ P. E. Kutter,⁷⁸ H. Li,⁷⁸ R. Liu,⁷⁸ F. Di Lodovico,⁷⁸ A. Mihalysi,⁷⁸ A. K. Mohapatra,⁷⁸ Y. Pan,⁷⁸ R. Prepost,⁷⁸ S. J. Sekula,⁷⁸ P. Tan,⁷⁸ J. H. von Wimmersperg-Toeller,⁷⁸ J. Wu,⁷⁸ S. L. Wu,⁷⁸ Z. Yu,⁷⁸ and H. Neal⁷⁹

(The BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

²Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

³Institute of High Energy Physics, Beijing 100039, China

⁴University of Bergen, Inst. of Physics, N-5007 Bergen, Norway

⁵Lawrence Berkeley National Laboratory and University of California, Berkeley, CA 94720, USA

⁶University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁷Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁸University of Bristol, Bristol BS8 1TL, United Kingdom

⁹University of British Columbia, Vancouver, BC, Canada V6T 1Z1

¹⁰Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹¹Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹²University of California at Irvine, Irvine, CA 92697, USA

¹³University of California at Los Angeles, Los Angeles, CA 90024, USA

¹⁴University of California at Riverside, Riverside, CA 92521, USA

¹⁵University of California at San Diego, La Jolla, CA 92093, USA

¹⁶University of California at Santa Barbara, Santa Barbara, CA 93106, USA

- ¹⁷ *University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, CA 95064, USA*
- ¹⁸ *California Institute of Technology, Pasadena, CA 91125, USA*
- ¹⁹ *University of Cincinnati, Cincinnati, OH 45221, USA*
- ²⁰ *University of Colorado, Boulder, CO 80309, USA*
- ²¹ *Colorado State University, Fort Collins, CO 80523, USA*
- ²² *Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany*
- ²³ *Ecole Polytechnique, LLR, F-91128 Palaiseau, France*
- ²⁴ *University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom*
- ²⁵ *Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy*
- ²⁶ *Florida A&M University, Tallahassee, FL 32307, USA*
- ²⁷ *Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy*
- ²⁸ *Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy*
- ²⁹ *Harvard University, Cambridge, MA 02138, USA*
- ³⁰ *Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany*
- ³¹ *Imperial College London, London, SW7 2AZ, United Kingdom*
- ³² *University of Iowa, Iowa City, IA 52242, USA*
- ³³ *Iowa State University, Ames, IA 50011-3160, USA*
- ³⁴ *Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France*
- ³⁵ *Lawrence Livermore National Laboratory, Livermore, CA 94550, USA*
- ³⁶ *University of Liverpool, Liverpool L69 7ZE, United Kingdom*
- ³⁷ *Queen Mary, University of London, E1 4NS, United Kingdom*
- ³⁸ *University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom*
- ³⁹ *University of Louisville, Louisville, KY 40292, USA*
- ⁴⁰ *University of Manchester, Manchester M13 9PL, United Kingdom*
- ⁴¹ *University of Maryland, College Park, MD 20742, USA*
- ⁴² *University of Massachusetts, Amherst, MA 01003, USA*
- ⁴³ *Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, MA 02139, USA*
- ⁴⁴ *McGill University, Montréal, QC, Canada H3A 2T8*
- ⁴⁵ *Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy*
- ⁴⁶ *University of Mississippi, University, MS 38677, USA*
- ⁴⁷ *Université de Montréal, Laboratoire René J. A. Lévesque, Montréal, QC, Canada H3C 3J7*
- ⁴⁸ *Mount Holyoke College, South Hadley, MA 01075, USA*
- ⁴⁹ *Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy*
- ⁵⁰ *NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands*
- ⁵¹ *University of Notre Dame, Notre Dame, IN 46556, USA*
- ⁵² *Oak Ridge National Laboratory, Oak Ridge, TN 37831, USA*
- ⁵³ *Ohio State University, Columbus, OH 43210, USA*
- ⁵⁴ *University of Oregon, Eugene, OR 97403, USA*
- ⁵⁵ *Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy*
- ⁵⁶ *Universités Paris VI et VII, Lab de Physique Nucléaire H. E., F-75252 Paris, France*
- ⁵⁷ *Università di Pavia, Dipartimento di Elettronica and INFN, I-27100 Pavia, Italy*
- ⁵⁸ *University of Pennsylvania, Philadelphia, PA 19104, USA*
- ⁵⁹ *Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy*
- ⁶⁰ *Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy*
- ⁶¹ *Prairie View A&M University, Prairie View, TX 77446, USA*
- ⁶² *Princeton University, Princeton, NJ 08544, USA*
- ⁶³ *Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy*
- ⁶⁴ *Universität Rostock, D-18051 Rostock, Germany*
- ⁶⁵ *Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom*
- ⁶⁶ *DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France*
- ⁶⁷ *University of South Carolina, Columbia, SC 29208, USA*
- ⁶⁸ *Stanford Linear Accelerator Center, Stanford, CA 94309, USA*
- ⁶⁹ *Stanford University, Stanford, CA 94305-4060, USA*
- ⁷⁰ *State Univ. of New York, Albany, NY 12222, USA*
- ⁷¹ *University of Tennessee, Knoxville, TN 37996, USA*
- ⁷² *University of Texas at Austin, Austin, TX 78712, USA*
- ⁷³ *University of Texas at Dallas, Richardson, TX 75083, USA*
- ⁷⁴ *Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy*
- ⁷⁵ *Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy*
- ⁷⁶ *Vanderbilt University, Nashville, TN 37235, USA*
- ⁷⁷ *University of Victoria, Victoria, BC, Canada V8W 3P6*
- ⁷⁸ *University of Wisconsin, Madison, WI 53706, USA*
- ⁷⁹ *Yale University, New Haven, CT 06511, USA*
- (Dated: March 15, 2004)

We present a measurement of the time-dependent CP asymmetry for the neutral B -meson decay $B^0 \rightarrow \phi K^0$. We use a sample of approximately 114 million B -meson pairs taken at the $\Upsilon(4S)$ resonance with the *BABAR* detector at the PEP-II B -meson Factory at SLAC. We reconstruct the CP eigenstates ϕK_s^0 and ϕK_L^0 where $\phi \rightarrow K^+ K^-$, $K_s^0 \rightarrow \pi^+ \pi^-$, and K_L^0 is observed via its hadronic interactions. The other B meson in the event is tagged as either a B^0 or $\bar{B}^0$ from its decay products. The values of the CP -violation parameters are $S_{\phi K} = 0.47 \pm 0.34(\text{stat})_{-0.06}^{+0.08}(\text{syst})$ and $C_{\phi K} = 0.01 \pm 0.33(\text{stat}) \pm 0.10(\text{syst})$.

PACS numbers: 13.25.Hw, 11.30.Er, 12.15.Hh

Decays of B mesons into charmless hadronic final states with a ϕ meson are dominated by $b \rightarrow s\bar{s}s$ gluonic penguin amplitudes, possibly with smaller contributions from electroweak penguins, while other Standard Model (SM) amplitudes are strongly suppressed [1]. In the SM, CP violation arises from a single complex phase in the Cabibbo–Kobayashi–Maskawa (CKM) quark-mixing matrix [2]. Neglecting CKM-suppressed contributions, the time-dependent CP -violating asymmetries in the decays $B^0 \rightarrow \phi K^0$ and $B^0 \rightarrow J/\psi K^0$ are proportional to the same parameter $\sin 2\beta$ [3], where the latter decay is dominated by tree diagrams. Since many scenarios of physics beyond the SM introduce additional diagrams with heavy particles in the penguin loops and new CP -violating phases, comparison of CP -violating observables with SM expectations is a sensitive probe for new physics. Measurements of $\sin 2\beta$ in B decays to charmonium such as $B^0 \rightarrow J/\psi K_s^0$ have been reported by the *BABAR* [4] and Belle [5] collaborations, and the world average for $\sin 2\beta$ is 0.731 ± 0.056 [6]. The Belle collaboration measures $\sin 2\beta = -0.96 \pm 0.50_{-0.11}^{+0.09}$ in the decay $B^0 \rightarrow \phi K_s^0$ [7].

In this letter we report a measurement of the time-dependent CP asymmetry in the final state ϕK^0 based on an integrated luminosity of approximately 108 fb^{-1} collected at the $\Upsilon(4S)$ resonance with the *BABAR* detector [8] at the PEP-II asymmetric e^+e^- collider [9] located at the Stanford Linear Accelerator Center.

From a $B^0\bar{B}^0$ meson pair we fully reconstruct one meson, B_{CP} , in the final state ϕK^0 , and partially reconstruct the recoil B meson, B_{tag} . We examine B_{tag} for evidence that it decayed either as B^0 or $\bar{B}^0$ (flavor tag). The asymmetric beam configuration in the laboratory frame provides a boost of $\beta\gamma = 0.56$ to the $\Upsilon(4S)$, which allows the determination of the proper decay time difference $\Delta t = t_{CP} - t_{\text{tag}}$ from the vertex separation of the two neutral B mesons along the beam (z) axis. The decay rate f_+ (f_-) when the tagging meson is a B^0 ($\bar{B}^0$) is given by

$$f_{\pm}(\Delta t) = \frac{e^{-|\Delta t|/\tau_{B^0}}}{4\tau_{B^0}} [1 \pm S_{\phi K} \sin(\Delta m_d \Delta t) \mp C_{\phi K} \cos(\Delta m_d \Delta t)], \quad (1)$$

where τ_{B^0} is the neutral B meson mean lifetime, and Δm_d is the B^0 – $\bar{B}^0$ oscillation frequency. The time-dependent CP -violating asymmetry is defined as $A_{CP} \equiv$

$(f_+ - f_-)/(f_+ + f_-)$. In the SM, decays that proceed purely via the $b \rightarrow s\bar{s}s$ penguin transitions have CP parameters $S_{\phi K} = -\eta_f \sin 2\beta$ and $C_{\phi K} = 0$, where $\beta \equiv \arg[-V_{cd}V_{cb}^*/V_{td}V_{tb}^*]$. Here V_{ik} is the CKM matrix element for quarks i and k , and the CP eigenvalue is $\eta_f = -1$ ($+1$) for ϕK_s^0 (ϕK_L^0).

The B_{CP} candidate is reconstructed in the decay mode ϕK^0 with $\phi \rightarrow K^+ K^-$; the K^0 is either a K_L^0 or a $K_s^0 \rightarrow \pi^+ \pi^-$. We combine pairs of oppositely charged tracks extrapolated to a common vertex to form ϕ and K_s^0 candidates. For the charged tracks from the ϕ decay we require at least 12 measured drift-chamber (DCH) coordinates and a minimal transverse momentum p_T of $0.1 \text{ GeV}/c$. The tracks must also originate within 1.5 cm in xy and $\pm 10 \text{ cm}$ along the z -axis of the nominal beam spot. Tracks used to reconstruct the ϕ mesons are distinguished from pion and proton tracks via a requirement on a likelihood ratio that combines dE/dx information from the silicon vertex tracker (SVT) and the DCH for tracks with momentum $p < 0.7 \text{ GeV}/c$. For tracks with higher p , dE/dx in the DCH and the Cherenkov angle and the number of photons as measured by the internally reflecting ring-imaging Cherenkov detector are used in the likelihood. The two-kaon invariant mass must be within $16 \text{ MeV}/c^2$ of the nominal ϕ mass [6].

For tracks corresponding to K_s^0 and B_{tag} daughters our requirements are less restrictive. A $K_s^0 \rightarrow \pi^+ \pi^-$ candidate is accepted if its two-pion invariant mass is within $15 \text{ MeV}/c^2$ of the nominal K^0 mass [6], its reconstructed decay vertex is separated from the collision point by at least 3 standard deviations, and the angle between the line connecting the ϕ and K_s^0 decay vertices and the K_s^0 momentum direction is less than 45 mrad .

We identify a K_L^0 candidate as in our $B^0 \rightarrow J/\psi K_L^0$ analysis [10] either as a cluster of energy deposits in the electromagnetic calorimeter (EMC) or as a cluster of hits in two or more layers of the instrumented flux return (IFR) that cannot be associated with any charged track in the event. The K_L^0 energy is not well measured. Therefore, we determine the K_L^0 laboratory momentum from its flight direction as measured from the EMC or IFR cluster and the constraint that the invariant ϕK_L^0 mass agrees with the known B^0 mass. In those cases where the K_L^0 is detected in both the IFR and EMC we use the angular information from the EMC, as it has a higher precision. In order to reduce background from π^0 de-

cays, we reject an EMC K_L^0 candidate cluster if it forms an invariant mass between 100 and 150 MeV/ c^2 with any other cluster in the event under the $\gamma\gamma$ hypothesis, or if it has energy greater than 1 GeV and contains two shower maxima consistent with two photons from a π^0 decay. The remaining background of photons and overlapping showers is further reduced with the use of a neural network constructed from cluster shape variables, trained on Monte Carlo (MC) simulated $B^0 \rightarrow \phi K_L^0$ and measured radiative Bhabha events, and tested on measured $e^+e^- \rightarrow \phi(\rightarrow K_s^0 K_L^0)\gamma$ and $B^0 \rightarrow J/\psi K_L^0$ events. The final ϕK_L^0 sample consists of approximately equal numbers of IFR and EMC K_L^0 candidates.

The results are extracted from an extended unbinned maximum likelihood fit for which we parameterize the distributions of several kinematic and topological variables for signal and background events in terms of probability density functions (PDFs) [11]. The background arises primarily from random combinations of tracks produced in events of the type $e^+e^- \rightarrow q\bar{q}$, where $q = u, d, s, c$ (continuum). Background from other B decay final states with and without charm is estimated with MC simulations. Opposite CP contributions from the $K^+K^-K^0$ final state (e.g. K^+K^- S-wave) are estimated with an angular analysis on data to be less than 6.6% and treated as systematic error. The shapes of event variable distributions are obtained from signal and background MC samples and high statistics data control samples.

Each B_{CP} candidate is characterized by the energy difference $\Delta E = E_B^* - \frac{1}{2}\sqrt{s}$ and, except for $B^0 \rightarrow \phi K_L^0$, the beam-energy-substituted mass $m_{ES} = \sqrt{(\frac{1}{2}s + \vec{p}_0 \cdot \vec{p}_B)^2 / E_0^2 - p_B^2}$ [8]. The subscripts 0 and B refer to the initial $\Upsilon(4S)$ and the B_{CP} candidate, respectively, and the asterisk denotes the $\Upsilon(4S)$ rest frame. For signal events, ΔE is expected to peak at zero and m_{ES} at the nominal B mass. We require $\Delta E < 0.08$ GeV for $B^0 \rightarrow \phi K_L^0$ and $|\Delta E| < 0.2$ GeV and $m_{ES} > 5.2$ GeV/ c^2 for $B^0 \rightarrow \phi K_s^0$. In the fit we also use the helicity angle θ_H , which is defined as the angle between the directions of the K^+ and the parent B_{CP} in the K^+K^- rest frame. The $\cos\theta_H$ distribution for pseudoscalar-vector B decay modes is $\cos^2\theta_H$, and for the combinatorial background it is nearly uniform.

In continuum events, particles appear bundled into jets. This topology can be characterized with several variables computed in the CM frame. One such quantity is the angle θ_T between the thrust axis of the B_{CP} candidate and the thrust axis formed from the other charged and neutral particles in the event. We also use the angle θ_B between the B_{CP} momentum and the beam axis, and the sum of the momenta p_i of the other charged and neutral particles in the event weighted by the Legendre polynomials $L_n(\theta_i)$, $n = 0, 2$, where θ_i is the angle between the momentum of particle i and the thrust axis of the B_{CP} candidate. For $B^0 \rightarrow \phi K_s^0$ candidates, we

combine these variables into a Fisher discriminant $\mathcal{F}$ [12] after requiring $|\cos\theta_T| < 0.9$. In this mode background from other B decays is negligible, as demonstrated in MC simulation studies.

More stringent criteria must be applied to suppress backgrounds in the case of $B^0 \rightarrow \phi K_L^0$ candidates, and we require $|\cos\theta_T| < 0.8$ and $|\cos\theta_B| < 0.85$. We define the missing momentum $\vec{p}_{miss}$, calculated in the laboratory frame from the beam momentum and all tracks and EMC clusters, excluding the K_L^0 candidate. We require the polar angle θ_{miss} of the missing momentum with respect to the beam direction to be greater than 0.3 rad. The cosine of the angle between $\vec{p}_{miss}$ and the K_L^0 direction, θ_K , must satisfy $\cos\theta_K > 0.6$. In the plane transverse to the beam direction, the difference between the missing momentum projected along the K_L^0 direction and the calculated K_L^0 momentum must be greater than -0.75 GeV/ c . In the Fisher discriminant we replace $|\cos\theta_B|$ by the cosine of the angle between the missing momentum and the K^+ from the ϕ decay. In the ϕK_L^0 sample about 1.4% of the events originate from charm B decays, 0.7% originate from charmless B decays, and about 0.2% potentially have a CP asymmetry. The dominant contamination is the mode $B \rightarrow \phi K^*$, where the K^* decays to $K_L^0\pi$. In the likelihood fit we explicitly parameterize backgrounds from both charm and charmless B decays as derived from MC simulations.

All the other tracks and clusters in the event are used to form the B_{tag} , and its flavor is determined with a multivariate tagging algorithm [4]. The tagging efficiency ϵ and mistag probability w in four hierarchical and mutually exclusive categories is measured from fully reconstructed B^0 decays into the $D^{(*)}-X^+$ ($X^+ = \pi^+, \rho^+, a_1^+$) and $J/\psi K^{*0}$ ($K^{*0} \rightarrow K^+\pi^-$) flavor eigenstates (B_{flav} sample). The analyzing power $\epsilon(1-2w)^2$ is $(28.7 \pm 0.7)\%$.

A detailed description of the Δt reconstruction algorithm is given in Ref. [10]. The B_{CP} vertex resolution is dominated by the ϕ vertex. The average Δz resolution is 190 μm and is dominated by the tagging vertex in the event. Thus, we can characterize the resolution with the much larger B_{flav} sample, which we fit simultaneously with the CP samples. The amplitudes for the B_{CP} asymmetries and for the B_{flav} flavor oscillations are reduced by the same factor due to wrong tags. Both distributions are convoluted with a common Δt resolution function, and the backgrounds are accounted for by adding terms to the likelihood, incorporated with different assumptions about their Δt evolution and resolution function [10].

Since we measure the correlations among the observables to be small in the data samples entering the fit, we take the probability density function $\mathcal{P}_{i,c}^j$ for each event j to be a product of the PDFs for the separate observables. For each event hypothesis i (signal, background) and tagging category c , we define $\mathcal{P}_{i,c}^j = \mathcal{P}_i(m_{ES}) \cdot \mathcal{P}_i(\Delta E) \cdot \mathcal{P}_i(\mathcal{F}) \cdot \mathcal{P}_i(\cos\theta_H) \cdot \mathcal{P}_i(\Delta t; \sigma_{\Delta t}, c)$, where

for the ϕK_L^0 mode $\mathcal{P}_i(m_{ES}) = 1$ and for the flavor sample $\mathcal{P}_i(\mathcal{F}) \cdot \mathcal{P}_i(\cos \theta_H) = 1$. The $\sigma_{\Delta t}$ is the error on Δt for a given event. The likelihood function for each decay chain is then

$$\mathcal{L} = \prod_c \exp \left(- \sum_i N_{i,c} \right) \prod_j^{N_c} \left[\sum_i N_{i,c} \mathcal{P}_{i,c}^j \right], \quad (2)$$

where $N_{i,c}$ is the yield of events of hypothesis i found by the fitter in category c , and N_c is the number of category c events in the sample. The total sample consists of 86200 B_{flav} , 2138 ϕK_s^0 and 4730 ϕK_L^0 candidates. We find 70 ± 9 ϕK_s^0 and 52 ± 16 ϕK_L^0 signal events. The signal yields in both the ϕK^0 channels agree well with our determination of the branching fraction for $B^0 \rightarrow \phi K^0$ [13]. Fig. 1 shows the m_{ES} (ΔE) distribution of ϕK_s^0 (ϕK_L^0) events together with the result from the fit after a requirement on the likelihood (computed without the variable plotted) to enhance the sensitivity.

FIG. 1: Distribution of the event variable (a) m_{ES} for the ϕK_s^0 final state and (b) ΔE for the ϕK_L^0 final state after reconstruction and a requirement for the likelihood with total signal efficiency of 32% and 5%, respectively. The solid line represents the fit result for the total event yield and the dotted line for the background.

We determine the CP parameters $S_{\phi K}$ and $C_{\phi K}$ along with an additional 38 free parameters: the efficiency per tagging category (4 parameters), the average mistag fraction and the difference between B^0 and $\bar{B}^0$ mistags for each tagging category (8 parameters), the signal Δt resolution (9), and time dependence (6), Δt resolution (3) and mistag fractions (8) for the background. We fix τ_{B^0} and Δm_d to the world averages [6]. The determination of the mistag fractions and Δt -resolution parameters is dominated by the high-statistics B_{flav} sample. The fit was tested with a parameterized simulation of a large number of data-sized experiments and full detector simulated events for the different signal and background samples. The fit was also verified with our $J/\psi K_s^0$ data sample and a control sample of 232 ϕK^+ candidates where one expects $S_{\phi K^+} = C_{\phi K^+} = 0$. We measure $S_{\phi K^+} = 0.23 \pm 0.24$ and $C_{\phi K^+} = -0.14 \pm 0.18$ with statistical errors only. The simultaneous fit to the ϕK^0

FIG. 2: Plots a) and b) show the Δt distributions of B^0 - and $\bar{B}^0$ -tagged ϕK_s^0 events. The solid lines refer to the fit for all events; the dashed lines correspond to the background. Plot c) shows the asymmetry. A requirement for the event likelihood is applied. Plots d), e), and f) are the corresponding plots for ϕK_L^0 events.

and flavor decay modes yields:

$$\begin{aligned} S_{\phi K} &= 0.47 \pm 0.34(\text{stat})^{+0.08}_{-0.06}(\text{syst}), \\ C_{\phi K} &= 0.01 \pm 0.33(\text{stat}) \pm 0.10(\text{syst}). \end{aligned}$$

The result in the dominant channel $B^0 \rightarrow \phi K_s^0$ is $S_{\phi K} = 0.45 \pm 0.43$ and $C_{\phi K} = -0.38 \pm 0.37$ with statistical errors only. Fig. 2 shows the Δt distributions of the B^0 - and the $\bar{B}^0$ -tagged subsets together with the raw asymmetry for the ϕK_s^0 and ϕK_L^0 events with the result of the combined time-dependent CP -asymmetry fit superimposed.

We consider systematic uncertainties in the CP coefficients $S_{\phi K}$ and $C_{\phi K}$ due to the event-yield determination in the two channels (± 0.01 for $S_{\phi K}$, ± 0.05 for $C_{\phi K}$), contributions from B^0 final states with opposite CP ($+0.06$, ± 0.02), the parameterization of PDFs for the event yield in signal and background (± 0.02 , ± 0.05), composition and CP asymmetry of the background in the CP events (± 0.03 , ± 0.03), the assumed parameterization of the Δt resolution function (± 0.02 , ± 0.01), the m_{ES} background parameterization (± 0.02 , ± 0.05), a possible difference in the efficiency for B^0 and $\bar{B}^0$ (± 0.01 , ± 0.02), the fixed

values for Δm_d and τ_B (± 0.00 , ± 0.01), the beam-spot position (± 0.01 , ± 0.01), and uncertainties in the SVT alignment (± 0.01 , ± 0.01). The bias in the coefficients due to the fit procedure (± 0.03 , ± 0.01) is included in the uncertainty without making corrections to the final results. We estimate errors due to the effect of doubly CKM-suppressed decays [14] to be (± 0.01 , ± 0.03). We add these contributions in quadrature to obtain the total systematic uncertainty.

In summary, we have measured the time-dependent CP asymmetries in the combined B -meson final states ϕK_s^0 and ϕK_L^0 . We obtain values for the CP -violation parameters $S_{\phi K}$ and $C_{\phi K}$ that agree within one standard deviation with the ones measured in the charmonium channels [4, 5]; the central value of $S_{\phi K}$ is also consistent with no CP asymmetry at the 1.3σ level.

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from the A. P. Sloan Foundation, Research Corporation, and Alexander von Humboldt Foundation.

* Now at Department of Physics, University of Warwick, Coventry, United Kingdom

- [†] Also with Università della Basilicata, Potenza, Italy
- [‡] Also with IFIC, Instituto de Física Corpuscular, CSIC-Universidad de Valencia, Valencia, Spain
- [§] Deceased
- [1] N. G. Deshpande and J. Trampetic, Phys. Rev. D **41**, 895 (1990); N. G. Deshpande and G. He, Phys. Lett. B **336**, 471 (1994); R. Fleischer, Z. Phys. C **62**, 81 (1994); Y. Grossman *et al.*, Phys. Rev. D **68**, 015004 (2003).
- [2] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [3] A. B. Carter and A. I. Sanda, Phys. Rev. D **23**, 1567 (1981); I. I. Bigi and A. I. Sanda, Nucl. Phys. B **193**, 85 (1981); Y. Grossman and M. P. Worah, Phys. Lett. B **395**, 241 (1997); R. Fleischer, Int. J. Mod. Phys. A **12**, 2459 (1997); D. London and A. Soni, Phys. Lett. B **407**, 61 (1997).
- [4] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **89**, 201802 (2002).
- [5] Belle Collaboration, K. Abe *et al.*, hep-ex/0308036 (2003).
- [6] Particle Data Group, K. Hagiwara *et al.*, Phys. Rev. D **66**, 010001 (2002).
- [7] Belle Collaboration, K. Abe *et al.*, Phys. Rev. Lett. **91**, 261602 (2003).
- [8] *BABAR* Collaboration, B. Aubert *et al.*, Nucl. Instr. Meth. A **479**, 1 (2002).
- [9] PEP-II Conceptual Design Report, SLAC-R-418 (1993).
- [10] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. D **66**, 032003 (2002).
- [11] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **87**, 151801 (2001).
- [12] R. A. Fisher, Annals Eugen. **7**, 179 (1936).
- [13] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. D **69**, 011102 (2004).
- [14] O. Long, M. Baak, R. N. Cahn and D. Kirkby, Phys. Rev. D **68**, 034010 (2003).