Determination of the Branching Fraction for $B \rightarrow X_c \ell \nu$ Decays and of $|V_{cb}|$ from Hadronic-Mass and Lepton-Energy Moments

B. Aubert,¹ R. Barate,¹ D. Boutigny,¹ F. Couderc,¹ J.-M. Gaillard,¹ A. Hicheur,¹ Y. Karyotakis,¹ J. P. Lees,¹
 V. Tisserand,¹ A. Zghiche,¹ A. Palano,² A. Pompili,² J. C. Chen,³ N. D. Qi,³ G. Rong,³ P. Wang,³ Y. S. Zhu,³
 G. Eigen,⁴ I. Ofte,⁴ B. Stugu,⁴ G. S. Abrams,⁵ A. W. Borgland,⁵ A. B. Breon,⁵ D. N. Brown,⁵ J. Button-Shafer,⁵
 R. N. Cahn,⁵ E. Charles,⁵ C. T. Day,⁵ M. S. Gill,⁵ A. V. Gritsan,⁵ Y. Groysman,⁵ R. G. Jacobsen,⁵ R. W. Kadel,⁵
 J. Kadyk,⁵ L. T. Kerth,⁵ Yu. G. Kolomensky,⁵ G. Kukartsev,⁵ C. LeClerc,⁵ G. Lynch,⁵ A. M. Merchant,⁵
 L. M. Mir,⁵ P. J. Oddone,⁵ T. J. Orimoto,⁵ M. Pripstein,⁵ N. A. Roe,⁵ M. T. Ronan,⁵ V. G. Shelkov,⁵
 W. A. Wenzel,⁵ K. Ford,⁶ T. J. Harrison,⁶ C. M. Hawkes,⁶ S. E. Morgan,⁶ A. T. Watson,⁶ M. Fritsch,⁷ K. Goetzen,⁷
 T. Held,⁷ H. Koch,⁷ B. Lewandowski,⁷ M. Pelizaeus,⁷ M. Steinke,⁷ J. T. Boyd,⁸ N. Chevalier,⁸ W. N. Cottingham,⁸
 M. P. Kelly,⁸ T. E. Latham,⁸ F. F. Wilson,⁸ T. Cuhadar-Donszelmann,⁹ C. Hearty,⁹ N. S. Knecht,⁹
 T. S. Mattison,⁹ J. A. McKenna,⁹ D. Thiessen,⁹ A. Khan,¹⁰ P. Kyberd,¹⁰ L. Teodorescu,¹⁰ V. E. Blinov,¹¹
 A. D. Bukin,¹¹ V. P. Druzhinin,¹¹ V. B. Golubev,¹¹ V. N. Ivanchenko,¹¹ E. A. Kravchenko,¹¹ A. P. Onuchin,¹¹
 S. I. Serednyakov,¹¹ Yu. I. Skovpen,¹¹ E. P. Solodov,¹¹ A. N. Yushkov,¹¹ D. Best,¹² M. Bruinsma,¹² M. Chao,¹²
 I. Eschrich,¹² D. Kirkby,¹² A. J. Lankford,¹² M. Mandelkern,¹² R. K. Mommsen,¹² W. Roethel,¹² D. P. Stoker,¹²
 C. Buchanan,¹³ B. L. Hartfiel,¹³ J. W. Gary,¹⁴ B. C. Shen,¹⁴ K. Wang,¹⁴ D. del Re,¹⁵ H. K. Hadavand,¹⁵
 E. J. Hill,¹⁵ D. B. MacFarlane,¹⁵ H. P. Paar,¹⁵ Sh. Rahatlou,¹⁵ V. Sharma,¹⁵ J. W. Berryhill,¹⁶ C. Campagnari,¹⁶
 B. Dahmes,¹⁶ S. L. Levy,¹⁶ O. Long,¹⁶ A. Lu,¹⁶ M. A. Mazur,¹⁶ J. D. Richman,¹⁶ W. Verkerke,¹⁶ T. W. Beck,¹⁷
 A. M. Eisner,¹⁷ C. A. Heusch,¹⁷ W. S. Lockman,¹⁷ T. Schalk,¹⁷ R. E. Schmitz,¹⁷ B. A. Schumm,¹⁷ A. Seiden,¹⁷
 P. Spradlin,¹⁷ D. C. Williams,¹⁷ M. G. Wilson,¹⁷ J. Albert,¹⁸ E. Chen,¹⁸ G. P. Dubois-Felsmann,¹⁸ A. Dvoretzskii,¹⁹
 D. G. Hitlin,¹⁸ I. Narsky,¹⁸ T. Piatenko,¹⁸ F. C. Porter,¹⁸ A. Ryd,¹⁸ A. Samuel,¹⁸ S. Yang,¹⁸ S. Jayatilake,¹⁹
 G. Mancinelli,¹⁹ B. T. Meadows,¹⁹ M. D. Sokoloff,¹⁹ T. Abe,²⁰ F. Blanc,²⁰ P. Bloom,²⁰ S. Chen,²⁰ W. T. Ford,²⁰
 U. Nauenberg,²⁰ A. Olivas,²⁰ P. Rankin,²⁰ J. G. Smith,²⁰ J. Zhang,²⁰ L. Zhang,²⁰ A. Chen,²¹ J. L. Harton,²¹
 A. Soffer,²¹ W. H. Toki,²¹ R. J. Wilson,²¹ Q. L. Zeng,²¹ D. Altenburg,²² T. Brandt,²² J. Brose,²² T. Colberg,²²
 M. Dickopp,²² E. Feltresi,²² A. Hauke,²² H. M. Lacker,²² E. Maly,²² R. Müller-Pfefferkorn,²² R. Nogowski,²²
 S. Otto,²² A. Petzold,²² J. Schubert,²² K. R. Schubert,²² R. Schwierz,²² B. Spaan,²² J. E. Sundermann,²²
 D. Bernard,²³ G. R. Bonneaud,²³ F. Brochard,²³ P. Grenier,²³ S. Schrenk,²³ Ch. Thiebaux,²³ G. Vasileiadis,²³
 M. Verderi,²³ D. J. Bard,²⁴ P. J. Clark,²⁴ D. Lavin,²⁴ F. Muheim,²⁴ S. Playfer,²⁴ Y. Xie,²⁴ M. Andreotti,²⁵
 V. Azzolini,²⁵ D. Bettoni,²⁵ C. Bozzi,²⁵ R. Calabrese,²⁵ G. Cibinetto,²⁵ E. Luppi,²⁵ M. Negrini,²⁵ A. Sarti,²⁵
 E. Treadwell,²⁶ R. Baldini-Ferrolì,²⁷ A. Calcaterra,²⁷ R. de Sangro,²⁷ G. Finocchiaro,²⁷ P. Patteri,²⁷ M. Piccolo,²⁷
 A. Zallo,²⁷ A. Buzzo,²⁸ R. Capra,²⁸ R. Contri,²⁸ G. Crosetti,²⁸ M. Lo Vetere,²⁸ M. Macri,²⁸ M. R. Monge,²⁸
 S. Passaggio,²⁸ C. Patrignani,²⁸ E. Robutti,²⁸ A. Santroni,²⁸ S. Tosi,²⁸ S. Bailey,²⁹ G. Brandenburg,²⁹ M. Morii,²⁹
 E. Won,²⁹ R. S. Dubitzky,³⁰ U. Langenegger,³⁰ W. Bhimji,³¹ D. A. Bowerman,³¹ P. D. Dauncey,³¹ U. Egede,³¹
 J. R. Gaillard,³¹ G. W. Morton,³¹ J. A. Nash,³¹ G. P. Taylor,³¹ G. J. Grenier,³² U. Mallik,³² J. Cochran,³³
 H. B. Crawley,³³ J. Lamsa,³³ W. T. Meyer,³³ S. Prell,³³ E. I. Rosenberg,³³ J. Yi,³³ M. Davier,³⁴ G. Grosdidier,³⁴
 A. Höcker,³⁴ S. Laplace,³⁴ F. Le Diberder,³⁴ V. Lepeltier,³⁴ A. M. Lutz,³⁴ T. C. Petersen,³⁴ S. Plaszczynski,³⁴
 M. H. Schune,³⁴ L. Tantot,³⁴ G. Wormser,³⁴ C. H. Cheng,³⁵ D. J. Lange,³⁵ M. C. Simani,³⁵ D. M. Wright,³⁵
 A. J. Bevan,³⁶ J. P. Coleman,³⁶ J. R. Fry,³⁶ E. Gabathuler,³⁶ R. Gamet,³⁶ R. J. Parry,³⁶ D. J. Payne,³⁶
 R. J. Sloane,³⁶ C. Touramanis,³⁶ J. J. Back,³⁷ P. F. Harrison,³⁷ G. B. Mohanty,³⁷ C. L. Brown,³⁸ G. Cowan,³⁸
 R. L. Flack,³⁸ H. U. Flaecher,³⁸ M. G. Green,³⁸ C. E. Marker,³⁸ T. R. McMahon,³⁸ S. Ricciardi,³⁸ F. Salvatore,³⁸
 G. Vaitsas,³⁸ M. A. Winter,³⁸ D. Brown,³⁹ C. L. Davis,³⁹ J. Allison,⁴⁰ N. R. Barlow,⁴⁰ R. J. Barlow,⁴⁰ P. A. Hart,⁴⁰
 M. C. Hodgkinson,⁴⁰ G. D. Lafferty,⁴⁰ A. J. Lyon,⁴⁰ J. C. Williams,⁴⁰ A. Farbin,⁴¹ W. D. Hulsbergen,⁴¹
 A. Jawahery,⁴¹ D. Kovalskyi,⁴¹ C. K. Lae,⁴¹ V. Lillard,⁴¹ D. A. Roberts,⁴¹ G. Blaylock,⁴² C. Dallapiccola,⁴²
 K. T. Flood,⁴² S. S. Hertzbach,⁴² R. Kofler,⁴² V. B. Koptchev,⁴² T. B. Moore,⁴² S. Saremi,⁴² H. Staengle,⁴²
 S. Willocq,⁴² R. Cowan,⁴³ G. Sciolla,⁴³ F. Taylor,⁴³ R. K. Yamamoto,⁴³ D. J. J. Mangeol,⁴⁴ P. M. Patel,⁴⁴
 S. H. Robertson,⁴⁴ A. Lazzaro,⁴⁵ F. Palombo,⁴⁵ J. M. Bauer,⁴⁶ L. Cremaldi,⁴⁶ V. Eschenburg,⁴⁶ R. Godang,⁴⁶

R. Kroeger,⁴⁶ J. Reidy,⁴⁶ D. A. Sanders,⁴⁶ D. J. Summers,⁴⁶ H. W. Zhao,⁴⁶ S. Brunet,⁴⁷ D. Côté,⁴⁷ P. Taras,⁴⁷ H. Nicholson,⁴⁸ N. Cavallo,⁴⁹ F. Fabozzi,^{49,*} C. Gatto,⁴⁹ L. Lista,⁴⁹ D. Monorchio,⁴⁹ P. Paolucci,⁴⁹ D. Piccolo,⁴⁹ C. Sciacca,⁴⁹ M. Baak,⁵⁰ H. Bulten,⁵⁰ G. Raven,⁵⁰ L. Wilden,⁵⁰ C. P. Jessop,⁵¹ J. M. LoSecco,⁵¹ T. A. Gabriel,⁵² T. Allmendinger,⁵³ B. Brau,⁵³ K. K. Gan,⁵³ K. Honscheid,⁵³ D. Hufnagel,⁵³ H. Kagan,⁵³ R. Kass,⁵³ T. Pulliam,⁵³ A. M. Rahimi,⁵³ R. Ter-Antonyan,⁵³ Q. K. Wong,⁵³ J. Brau,⁵⁴ R. Frey,⁵⁴ O. Igonkina,⁵⁴ C. T. Potter,⁵⁴ N. B. Sinev,⁵⁴ D. Strom,⁵⁴ E. Torrence,⁵⁴ F. Colecchia,⁵⁵ A. Dorigo,⁵⁵ F. Galeazzi,⁵⁵ M. Margoni,⁵⁵ M. Morandin,⁵⁵ M. Posocco,⁵⁵ M. Rotondo,⁵⁵ F. Simonetto,⁵⁵ R. Stroili,⁵⁵ G. Tiozzo,⁵⁵ C. Voci,⁵⁵ M. Benayoun,⁵⁶ H. Briand,⁵⁶ J. Chauveau,⁵⁶ P. David,⁵⁶ Ch. de la Vaissière,⁵⁶ L. Del Buono,⁵⁶ O. Hamon,⁵⁶ M. J. J. John,⁵⁶ Ph. Leruste,⁵⁶ J. Ocariz,⁵⁶ M. Pivk,⁵⁶ L. Roos,⁵⁶ S. T'Jampens,⁵⁶ G. Therin,⁵⁶ P. F. Manfredi,⁵⁷ V. Re,⁵⁷ P. K. Behera,⁵⁸ L. Gladney,⁵⁸ Q. H. Guo,⁵⁸ J. Panetta,⁵⁸ F. Anulli,^{27,59} M. Biasini,⁵⁹ I. M. Peruzzi,^{27,59} M. Pioppi,⁵⁹ C. Angelini,⁶⁰ G. Batignani,⁶⁰ S. Bettarini,⁶⁰ M. Bondioli,⁶⁰ F. Bucci,⁶⁰ G. Calderini,⁶⁰ M. Carpinelli,⁶⁰ V. Del Gamba,⁶⁰ F. Forti,⁶⁰ M. A. Giorgi,⁶⁰ A. Lusiani,⁶⁰ G. Marchiori,⁶⁰ F. Martinez-Vidal,^{60,†} M. Morganti,⁶⁰ N. Neri,⁶⁰ E. Paoloni,⁶⁰ M. Rama,⁶⁰ G. Rizzo,⁶⁰ F. Sandrelli,⁶⁰ J. Walsh,⁶⁰ M. Haire,⁶¹ D. Judd,⁶¹ K. Paick,⁶¹ D. E. Wagoner,⁶¹ N. Danielson,⁶² P. Elmer,⁶² C. Lu,⁶² V. Miftakov,⁶² J. Olsen,⁶² A. J. S. Smith,⁶² A. V. Telnov,⁶² F. Bellini,⁶³ G. Cavoto,^{62,63} R. Faccini,⁶³ F. Ferrarotto,⁶³ F. Ferroni,⁶³ M. Gaspero,⁶³ L. Li Gioi,⁶³ M. A. Mazzoni,⁶³ S. Morganti,⁶³ M. Pierini,⁶³ G. Piredda,⁶³ F. Safai Tehrani,⁶³ C. Voena,⁶³ S. Christ,⁶⁴ G. Wagner,⁶⁴ R. Waldi,⁶⁴ T. Adye,⁶⁵ N. De Groot,⁶⁵ B. Franek,⁶⁵ N. I. Geddes,⁶⁵ G. P. Gopal,⁶⁵ E. O. Olaiya,⁶⁵ R. Aleksan,⁶⁶ S. Emery,⁶⁶ A. Gaidot,⁶⁶ S. F. Ganzhur,⁶⁶ P.-F. Giraud,⁶⁶ G. Hamel de Monchenault,⁶⁶ W. Kozanecki,⁶⁶ M. Langer,⁶⁶ M. Legendre,⁶⁶ G. W. London,⁶⁶ B. Mayer,⁶⁶ G. Schott,⁶⁶ G. Vasseur,⁶⁶ Ch. Yèche,⁶⁶ M. Zito,⁶⁶ M. V. Purohit,⁶⁷ A. W. Weidemann,⁶⁷ F. X. Yumiceva,⁶⁷ D. Aston,⁶⁸ R. Bartoldus,⁶⁸ N. Berger,⁶⁸ A. M. Boyarski,⁶⁸ O. L. Buchmueller,⁶⁸ M. R. Convery,⁶⁸ M. Cristinziani,⁶⁸ G. De Nardo,⁶⁸ D. Dong,⁶⁸ J. Dorfan,⁶⁸ D. Dujmic,⁶⁸ W. Dunwoodie,⁶⁸ E. E. Elsen,⁶⁸ S. Fan,⁶⁸ R. C. Field,⁶⁸ T. Glanzman,⁶⁸ S. J. Gowdy,⁶⁸ T. Hadig,⁶⁸ V. Halyo,⁶⁸ T. Hryn'ova,⁶⁸ W. R. Innes,⁶⁸ M. H. Kelsey,⁶⁸ P. Kim,⁶⁸ M. L. Kocian,⁶⁸ D. W. G. S. Leith,⁶⁸ J. Libby,⁶⁸ S. Luitz,⁶⁸ V. Luth,⁶⁸ H. L. Lynch,⁶⁸ H. Marsiske,⁶⁸ R. Messner,⁶⁸ D. R. Muller,⁶⁸ C. P. O'Grady,⁶⁸ V. E. Ozcan,⁶⁸ A. Perazzo,⁶⁸ M. Perl,⁶⁸ S. Petrak,⁶⁸ B. N. Ratcliff,⁶⁸ A. Roodman,⁶⁸ A. A. Salnikov,⁶⁸ R. H. Schindler,⁶⁸ J. Schwiening,⁶⁸ G. Simi,⁶⁸ A. Snyder,⁶⁸ A. Soha,⁶⁸ J. Stelzer,⁶⁸ D. Su,⁶⁸ M. K. Sullivan,⁶⁸ J. Va'vra,⁶⁸ S. R. Wagner,⁶⁸ M. Weaver,⁶⁸ A. J. R. Weinstein,⁶⁸ W. J. Wisniewski,⁶⁸ M. Wittgen,⁶⁸ D. H. Wright,⁶⁸ A. K. Yarritu,⁶⁸ C. C. Young,⁶⁸ P. R. Burchat,⁶⁹ A. J. Edwards,⁶⁹ T. I. Meyer,⁶⁹ B. A. Petersen,⁶⁹ C. Roat,⁶⁹ S. Ahmed,⁷⁰ M. S. Alam,⁷⁰ J. A. Ernst,⁷⁰ M. A. Saeed,⁷⁰ M. Saleem,⁷⁰ F. R. Wappler,⁷⁰ W. Bugg,⁷¹ M. Krishnamurthy,⁷¹ S. M. Spanier,⁷¹ R. Eckmann,⁷² H. Kim,⁷² J. L. Ritchie,⁷² A. Satpathy,⁷² R. F. Schwitters,⁷² J. M. Izen,⁷³ I. Kitayama,⁷³ X. C. Lou,⁷³ S. Ye,⁷³ F. Bianchi,⁷⁴ M. Bona,⁷⁴ F. Gallo,⁷⁴ D. Gamba,⁷⁴ C. Borean,⁷⁵ L. Bosisio,⁷⁵ C. Cartaro,⁷⁵ F. Cossutti,⁷⁵ G. Della Ricca,⁷⁵ S. Dittongo,⁷⁵ S. Grancagnolo,⁷⁵ L. Lanceri,⁷⁵ P. Poropat,^{75,‡} L. Vitale,⁷⁵ G. Vuagnin,⁷⁵ R. S. Panvini,⁷⁶ Sw. Banerjee,⁷⁷ C. M. Brown,⁷⁷ D. Fortin,⁷⁷ P. D. Jackson,⁷⁷ R. Kowalewski,⁷⁷ J. M. Roney,⁷⁷ H. R. Band,⁷⁸ S. Dasu,⁷⁸ M. Datta,⁷⁸ A. M. Eichenbaum,⁷⁸ M. Graham,⁷⁸ J. J. Hollar,⁷⁸ J. R. Johnson,⁷⁸ P. E. Kutter,⁷⁸ H. Li,⁷⁸ R. Liu,⁷⁸ F. Di Lodovico,⁷⁸ A. Mihalyi,⁷⁸ A. K. Mohapatra,⁷⁸ Y. Pan,⁷⁸ R. Prepost,⁷⁸ A. E. Rubin,⁷⁸ S. J. Sekula,⁷⁸ P. Tan,⁷⁸ J. H. von Wimmersperg-Toeller,⁷⁸ J. Wu,⁷⁸ S. L. Wu,⁷⁸ Z. Yu,⁷⁸ and H. Neal⁷⁹

(The BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

²Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

³Institute of High Energy Physics, Beijing 100039, China

⁴University of Bergen, Inst. of Physics, N-5007 Bergen, Norway

⁵Lawrence Berkeley National Laboratory and University of California, Berkeley, CA 94720, USA

⁶University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁷Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁸University of Bristol, Bristol BS8 1TL, United Kingdom

⁹University of British Columbia, Vancouver, BC, Canada V6T 1Z1

¹⁰Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹¹Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹²University of California at Irvine, Irvine, CA 92697, USA

¹³University of California at Los Angeles, Los Angeles, CA 90024, USA

¹⁴University of California at Riverside, Riverside, CA 92521, USA

¹⁵University of California at San Diego, La Jolla, CA 92093, USA

¹⁶University of California at Santa Barbara, Santa Barbara, CA 93106, USA

- ¹⁷University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, CA 95064, USA
¹⁸California Institute of Technology, Pasadena, CA 91125, USA
¹⁹University of Cincinnati, Cincinnati, OH 45221, USA
²⁰University of Colorado, Boulder, CO 80309, USA
²¹Colorado State University, Fort Collins, CO 80523, USA
²²Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany
²³Ecole Polytechnique, LLR, F-91128 Palaiseau, France
²⁴University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom
²⁵Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy
²⁶Florida A&M University, Tallahassee, FL 32307, USA
²⁷Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy
²⁸Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy
²⁹Harvard University, Cambridge, MA 02138, USA
³⁰Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany
³¹Imperial College London, London, SW7 2AZ, United Kingdom
³²University of Iowa, Iowa City, IA 52242, USA
³³Iowa State University, Ames, IA 50011-3160, USA
³⁴Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France
³⁵Lawrence Livermore National Laboratory, Livermore, CA 94550, USA
³⁶University of Liverpool, Liverpool L69 7ZE, United Kingdom
³⁷Queen Mary, University of London, E1 4NS, United Kingdom
³⁸University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom
³⁹University of Louisville, Louisville, KY 40292, USA
⁴⁰University of Manchester, Manchester M13 9PL, United Kingdom
⁴¹University of Maryland, College Park, MD 20742, USA
⁴²University of Massachusetts, Amherst, MA 01003, USA
⁴³Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, MA 02139, USA
⁴⁴McGill University, Montréal, QC, Canada H3A 2T8
⁴⁵Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy
⁴⁶University of Mississippi, University, MS 38677, USA
⁴⁷Université de Montréal, Laboratoire René J. A. Lévesque, Montréal, QC, Canada H3C 3J7
⁴⁸Mount Holyoke College, South Hadley, MA 01075, USA
⁴⁹Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy
⁵⁰NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands
⁵¹University of Notre Dame, Notre Dame, IN 46556, USA
⁵²Oak Ridge National Laboratory, Oak Ridge, TN 37831, USA
⁵³Ohio State University, Columbus, OH 43210, USA
⁵⁴University of Oregon, Eugene, OR 97403, USA
⁵⁵Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy
⁵⁶Universités Paris VI et VII, Lab de Physique Nucléaire H. E., F-75252 Paris, France
⁵⁷Università di Pavia, Dipartimento di Eletttronica and INFN, I-27100 Pavia, Italy
⁵⁸University of Pennsylvania, Philadelphia, PA 19104, USA
⁵⁹Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy
⁶⁰Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy
⁶¹Prairie View A&M University, Prairie View, TX 77446, USA
⁶²Princeton University, Princeton, NJ 08544, USA
⁶³Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy
⁶⁴Universität Rostock, D-18051 Rostock, Germany
⁶⁵Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom
⁶⁶DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France
⁶⁷University of South Carolina, Columbia, SC 29208, USA
⁶⁸Stanford Linear Accelerator Center, Stanford, CA 94309, USA
⁶⁹Stanford University, Stanford, CA 94305-4060, USA
⁷⁰State Univ. of New York, Albany, NY 12222, USA
⁷¹University of Tennessee, Knoxville, TN 37996, USA
⁷²University of Texas at Austin, Austin, TX 78712, USA
⁷³University of Texas at Dallas, Richardson, TX 75083, USA
⁷⁴Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy
⁷⁵Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy
⁷⁶Vanderbilt University, Nashville, TN 37235, USA
⁷⁷University of Victoria, Victoria, BC, Canada V8W 3P6
⁷⁸University of Wisconsin, Madison, WI 53706, USA
⁷⁹Yale University, New Haven, CT 06511, USA

(Dated: April 11, 2004)

We determine the inclusive $B \rightarrow X_c \ell \nu$ branching fraction, $\mathcal{B}_{c\ell\nu}$, the CKM matrix element $|V_{cb}|$, and other heavy-quark parameters from a simultaneous fit to moments of the hadronic-mass and lepton-energy distributions in semileptonic B -meson decays, measured as a function of the lower limit on the lepton energy, using data recorded with the BABAR detector. Using Heavy Quark Expansions (HQEs) to order $1/m_b^3$, we extract $\mathcal{B}_{c\ell\nu} = (10.61 \pm 0.16_{exp} \pm 0.06_{HQE})\%$ and $|V_{cb}| = (41.4 \pm 0.4_{exp} \pm 0.4_{HQE} \pm 0.6_{th}) \times 10^{-3}$. The stated errors refer to the experimental, HQE, and additional theoretical uncertainties.

PACS numbers: 12.15.Hh, 11.30.Er, 13.25.Hw

The CKM matrix element V_{cb} is one of the fundamental parameters of the Standard Model and thus its precise measurement with well understood uncertainties is important. At the parton level, the weak decay rate for $b \rightarrow c \ell \nu$ can be calculated accurately; it is proportional to $|V_{cb}|^2$ and depends on the quark masses, m_b and m_c . To relate measurements of the semileptonic B -meson decay rate to $|V_{cb}|$, the parton-level calculations have to be corrected for effects of strong interactions. Heavy-Quark Expansions (HQEs)[1] have become a useful tool for calculating perturbative and non-perturbative QCD corrections [2] and for estimating their uncertainties. In the kinetic-mass scheme [3] for example, these expansions in $1/m_b$ and $\alpha_s(m_b)$ (the strong coupling constant) to order $\mathcal{O}(1/m_b^3)$ contain six parameters: the running kinetic masses of the b - and c -quarks, $m_b(\mu)$ and $m_c(\mu)$, and four non-perturbative parameters. We determine these parameters from a fit to the moments of the hadronic-mass and electron-energy distributions in semileptonic B decays to charm particles, $B \rightarrow X_c \ell \nu$. This fit yields measurements of the inclusive branching fraction $\mathcal{B}_{c\ell\nu}$ and of $|V_{cb}|$, significantly improved compared to earlier BABAR measurements [4]. It also allows us to test the consistency of the data with the HQEs employed and to check for the possible impact of higher-order contributions. Moment measurements and the extraction of $|V_{cb}|$ based on HQEs [5] were first performed by the CLEO collaboration [6]. More recently, global fits to a variety of moments were presented [7–9], using HQEs in different mass schemes.

This analysis makes use of moments measured by the BABAR collaboration [10, 11]. The moments are derived from the inclusive hadronic-mass (m_X) and electron-energy (E_ℓ) distributions in $B \rightarrow X_c \ell \nu$ decays, averaged over charged and neutral B mesons produced at the $\Upsilon(4S)$ resonance. We have subtracted the charmless contributions based on the branching fraction $\mathcal{B}_{u\ell\nu} = (0.22 \pm 0.05)\%$ [12]. All moments are measured as functions of E_{cut} , a lower limit on the lepton energy (for energy moments we only use electrons, for mass moments we also use muons). The moments are corrected for detector effects and QED radiation [13]. The hadronic-mass distribution is measured in events tagged by the fully reconstructed hadronic decay of the second B meson. The hadronic-mass moments are defined as $M_n^X(E_{cut}) = \langle m_X^n \rangle_{E_\ell > E_{cut}}$ with $n = 1, 2, 3, 4$.

The electron-energy distribution is measured in events tagged by a high-momentum electron from the second B meson. We define the first energy moment as $M_1^\ell(E_{cut}) = \langle E_\ell \rangle_{E_\ell > E_{cut}}$ and the second and third moments as $M_n^\ell(E_{cut}) = \langle (E_\ell - M_1^\ell(E_{cut}))^n \rangle_{E_\ell > E_{cut}}$ with $n = 2, 3$. In addition, we use the partial branching fraction $M_0^\ell(E_{cut}) = \int_{E_{cut}}^{E_{max}} (d\mathcal{B}_{c\ell\nu}/dE_\ell) dE_\ell$. All measured moments are shown in Fig. 1.

In the kinetic-mass scheme the HQE to $\mathcal{O}(1/m_b^3)$ for the rate of $B \rightarrow X_c \ell \nu$ decays can be expressed as [14]

$$\Gamma_{c\ell\nu} = \frac{G_F^2 m_b^5}{192\pi^3} |V_{cb}|^2 (1 + A_{ew}) A_{pert}(r, \mu) \times \left[z_0(r) \left(1 - \frac{\mu_\pi^2 - \mu_G^2 + \frac{\rho_D^3 + \rho_{LS}^3}{m_b}}{2m_b^2} \right) - 2(1-r)^4 \frac{\mu_G^2 + \frac{\rho_D^3 + \rho_{LS}^3}{m_b}}{m_b^2} + d(r) \frac{\rho_D^3}{m_b^3} + \mathcal{O}(1/m_b^4) \right]. \quad (1)$$

The leading non-perturbative effects arise at $\mathcal{O}(1/m_b^2)$ and are parameterized by $\mu_\pi^2(\mu)$ and $\mu_G^2(\mu)$, the expectation values of the kinetic and chromomagnetic dimension-five operators. At $\mathcal{O}(1/m_b^3)$, two additional parameters enter, $\rho_D^3(\mu)$ and $\rho_{LS}^3(\mu)$, the expectation values of the Darwin (D) and spin-orbit (LS) dimension-six operators. These parameters depend on the scale μ that separates short-distance from long-distance QCD effects; the calculations are performed for $\mu = 1$ GeV [3]. Electroweak corrections are $1 + A_{ew} \cong (1 + \alpha/\pi \ln(M_Z/m_b))^2 = 1.014$ and perturbative QCD corrections are estimated to be $A_{pert}(r, \mu) \cong 0.91 \pm 0.01$ [14]. The ratio $r = m_c^2/m_b^2$ enters in the phase-space factor $z_0(r) = 1 - 8r + 8r^3 - r^4 - 12r^2 \ln r$ and the function $d(r) = 8 \ln r + 34/3 - 32r/3 - 8r^2 + 32r^3/3 - 10r^4/3$.

This analysis uses linearized expressions for the HQEs [15]. Specifically, the dependence of $|V_{cb}|$ on the true values of heavy-quark parameters, expanded around *a priori* estimates of these parameters, is

$$\begin{aligned} \frac{|V_{cb}|}{0.0417} &\cong \sqrt{\frac{\mathcal{B}_{c\ell\nu}}{(0.1032)} \frac{1.55 \text{ ps}}{\tau_B}} \\ &\times [1 + 0.30(\alpha_s(m_b) - 0.22)] \\ &\times [1 - 0.66(m_b - 4.60) + 0.39(m_c - 1.15) \\ &+ 0.013(\mu_\pi^2 - 0.40) + 0.09(\rho_D^3 - 0.20) \\ &+ 0.05(\mu_G^2 - 0.35) - 0.01(\rho_{LS}^3 + 0.15)]. \end{aligned} \quad (2)$$

FIG. 1: The measured hadronic-mass (a-d) and electron-energy (e-h) moments as a function of the cut-off energy, E_{cut} , compared with the result of the simultaneous fit (line), with the theoretical uncertainties [15] indicated as shaded bands. The solid data points mark the measurements included in the fit. The vertical bars indicate the experimental errors; in some cases they are comparable in size to the data points. Moment measurements for different E_{cut} are highly correlated.

Here m_b and all other parameters of the expansion are in $\text{GeV}^{(n)}$; τ_B refers to the average lifetime of B mesons produced at the $\Upsilon(4S)$. We use $\tau_B = f_0\tau_0 + (1-f_0)\tau_{\pm} = (1.608 \pm 0.012) \text{ ps}$, taking into account the lifetimes [16] of neutral and charged B mesons, τ_0 and $\tau_{\pm}$, and their relative production rates (defined in terms of $f_0 = 0.488 \pm 0.013$ [17], the fraction of $B^0\bar{B}^0$ pairs).

HQEs in terms of the same heavy-quark parameters are available for the hadronic-mass and electron-energy moments. The dependence on the heavy-quark parameters has again been linearized using the same *a priori* estimates of the parameters [14, 15]. We have verified that the differences between the linearized expressions and the full theoretical calculation are very small in all cases. We use these linear equations to determine the unknown heavy-quark parameters, the total branching fraction $\mathcal{B}_{cl\nu}$, and $|V_{cb}|$ from a simultaneous χ^2 fit to the measured moments and the partial branching fraction, all as a function of the cut-off lepton energy, E_{cut} .

In total, we have measured four hadronic-mass moments for each of seven different values of E_{cut} , ranging from 0.9 to 1.5 GeV, and three electron-energy moments plus the partial branching fraction at five values of E_{cut} , ranging from 0.6 to 1.5 GeV [10, 11]. Since many of these individual moments are highly correlated we select for the fitting procedure a set of moments for which the correlation coefficients do not exceed 95%. Thus we only use half of the 28 mass moments, and retain 13 of the 20 energy moments.

The global fit takes into account the statistical and

systematic errors and correlations of the individual measurements, as well as the uncertainties of the expressions for the individual moments. We assess the uncertainty of the calculated moments by varying, as suggested in [15], in the linearized expressions (given for $|V_{cb}|$ in Eq. 2) the *a priori* estimates for μ_{π}^2 and μ_G^2 by $\pm 20\%$ and for ρ_{LS}^3 and ρ_D^3 by $\pm 30\%$. We assume that for a given moment these changes are fully correlated for all values of E_{cut} , but uncorrelated for different moments. The resulting fit, shown in Fig. 1, describes the data well with $\chi^2 = 15.0$ for 20 degrees of freedom. Tables I and II list the fitted parameters, their errors and correlations. As expected, m_b and m_c are highly correlated. For the mass difference, we obtain $m_b - m_c = (3.436 \pm 0.025_{exp} \pm 0.018_{HQE} \pm 0.010_{\alpha_s}) \text{ GeV}$.

Beyond the uncertainties that are included in the fit, the limited knowledge of the expression for the decay rate, including various perturbative corrections and higher-order non-perturbative corrections, introduces an error in $|V_{cb}|$, assessed to be 1.5% [14]. By comparison, the impact of the uncertainty in α_s is estimated to be relatively small. For $M_n^l(E_{cut})$ moments, perturbative corrections of order α_s^2 are included with $\alpha_s(m_b) = 0.22 \pm 0.04$, whereas for $M_n^X(E_{cut})$ moments, they are calculated only to $\mathcal{O}(\alpha_s)$ with $\alpha_s(m_b) = 0.3 \pm 0.1$. We estimate the error on the perturbative corrections by varying α_s within the stated errors. The choice of the scale μ is estimated to have a very small impact on $|V_{cb}|$ and the branching fraction [14].

A series of tests has been performed to verify that the

TABLE I: Fit results and error contributions from the moment measurements, approximations to the HQEs, and additional theoretical uncertainties from α_s terms and other perturbative and non-perturbative terms contributing to $\Gamma_{cl\nu}$.

	$ V_{cb} (10^{-3})$	m_b (GeV)	m_c (GeV)	μ_π^2 (GeV ²)	ρ_D^3 (GeV ³)	μ_G^2 (GeV ²)	ρ_{LS}^3 (GeV ³)	$\mathcal{B}_{cl\nu}$ (%)
Results	41.390	4.611	1.175	0.447	0.195	0.267	-0.085	10.611
δ_{exp}	0.437	0.052	0.072	0.035	0.023	0.055	0.038	0.163
δ_{HQE}	0.398	0.041	0.056	0.038	0.018	0.033	0.072	0.063
δ_{α_s}	0.150	0.015	0.015	0.010	0.004	0.018	0.010	0.000
δ_Γ	0.620							
δ_{tot}	0.870	0.068	0.092	0.053	0.029	0.067	0.082	0.175

TABLE II: Correlation coefficients for the fit parameters.

	$ V_{cb} $	m_b	m_c	μ_π^2	ρ_D^3	μ_G^2	ρ_{LS}^3	$\mathcal{B}_{cl\nu}$
$ V_{cb} $	1.00	-0.49	-0.36	0.56	0.35	-0.37	0.64	0.61
m_b		1.00	0.97	-0.40	-0.13	0.16	-0.63	0.23
m_c			1.00	-0.38	-0.13	-0.04	-0.50	0.29
μ_π^2				1.00	0.82	0.08	0.46	0.16
ρ_D^3					1.00	0.08	0.23	0.12
μ_G^2						1.00	-0.43	-0.04
ρ_{LS}^3							1.00	0.09
$\mathcal{B}_{cl\nu}$								1.00

fit results are unbiased. Specifically, we enlarged and reduced the estimated theoretical uncertainties by a factor of two and verified that the changes in the fitted parameters were small compared to the errors of the standard fit. We have also checked that the choice of the set of moments that are used in the fit does not significantly affect the result. In particular, an energy cut-off above 1.2 GeV might have introduced larger theoretical uncertainties and a potential bias, and moments for lower values of E_{cut} might have been affected by higher backgrounds. We found no evidence for any such effects.

The fit results are fully compatible with independent estimates [15] of $\mu_G^2 = (0.35 \pm 0.07)$ GeV², based on the $B^* - B$ mass splitting, and of $\rho_{LS}^3 = (-0.15 \pm 0.10)$ GeV³, from heavy-quark sum rules [18].

Figure 2 shows the $\Delta\chi^2 = 1$ ellipses for $|V_{cb}|$ versus m_b and μ_π^2 , for a fit to all moments and separate fits to the electron-energy moments and the hadronic-mass moments, but including the partial branching fractions in both. The lepton-energy and hadronic-mass moments have slightly different sensitivity to the fit parameters, but the results for the separate fits, $|V_{cb}| = (41.4 \pm 0.7) \times 10^{-3}$ and $|V_{cb}| = (41.6 \pm 0.8) \times 10^{-3}$ are fully compatible with each other and with the global fit to all moments. Changes in the other fit parameters are also consistent within the stated errors. Since the expansions for the two sets of moments are sensitive to different theoretical uncertainties and assumptions, in particular the differences in the treatment of the perturbative corrections, the observed consistency of the separate fits indicates that such differences are small compared with the

FIG. 2: Fit results (crosses) with contours corresponding to $\Delta\chi^2 = 1$ for two pairs of the eight free parameters a) m_b and b) μ_π^2 versus $|V_{cb}|$, separately for fits using the hadronic-mass, the electron-energy, and all moments.

experimental and assumed theoretical uncertainties.

In conclusion, we have extracted $|V_{cb}|$, the semileptonic branching fraction, and the heavy-quark masses,

$$\begin{aligned}
 |V_{cb}| &= (41.4 \pm 0.4_{exp} \pm 0.4_{HQE} \pm 0.6_{th}) \times 10^{-3}, \\
 \mathcal{B}_{ce\nu} &= (10.61 \pm 0.16_{exp} \pm 0.06_{HQE})\%, \\
 m_b(1 \text{ GeV}) &= (4.61 \pm 0.05_{exp} \pm 0.04_{HQE} \pm 0.02_{th}) \text{ GeV}, \\
 m_c(1 \text{ GeV}) &= (1.18 \pm 0.07_{exp} \pm 0.06_{HQE} \pm 0.02_{th}) \text{ GeV},
 \end{aligned}$$

as well as the non-perturbative parameters in the kinetic-mass scheme up to order $(1/m_b^3)$ (see Table I). The total semileptonic branching fraction is $\mathcal{B}_{ce\nu} + \mathcal{B}_{ue\nu} = (10.83 \pm 0.16_{exp} \pm 0.06_{HQE})\%$. The errors refer to contributions from the experimental errors on the moment measurements, the HQE uncertainties included in the fit, and additional theoretical uncertainties, $\delta_{th} = \sqrt{\delta_{\alpha_s}^2 + \delta_\Gamma^2}$, derived from Refs. [14, 15].

Based on a large set of hadronic-mass and electron-energy moments and a consistent set of HQE calculations, we have also been able to assess the uncertainties in the $\mathcal{O}(1/m_b^3)$ terms from the data without constraints to any *a priori* values. The fitted values of the parameters are consistent with theoretical estimates [3, 14]. The uncertainties on the quark masses are much smaller than

those of previous measurements [16]. It would be interesting to compare the results of this analysis with fits based on recent calculations performed in the $1S$ mass scheme [19].

The result on $|V_{cb}|$ is in agreement with previous measurements using HQEs, either for a different mass scheme and with fixed terms of $\mathcal{O}(1/m_b^3)$ [8], or for the kinetic-mass scheme, but with external constraints on almost all HQE parameters [9], as well as with an analysis combining both of these measurements [7].

This work has greatly benefited from many interactions with N. Uraltsev and P. Gambino. We thank them as well as many other theorists for valuable discussions regarding this work. We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support BABAR. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from the A. P. Sloan Foundation, Research Corporation, and Alexander von Humboldt Foundation.

* Also with Università della Basilicata, Potenza, Italy

† Also with IFIC, Instituto de Física Corpuscular, CSIC-Universidad de Valencia, Valencia, Spain

‡ Deceased

- [1] M. Voloshin and M. Shifman, Sov. J. Nucl. Phys. **41**, 120 (1985); J. Chay, H. Georgi, and B. Grinstein, Phys. Lett. **B247**, 399 (1990); I. I. Bigi, and N. Uraltsev, Phys. Lett. **B280**, 271 (1992).
- [2] I. I. Bigi, N. Uraltsev, and A. I. Vainshtein, Phys. Lett. **B293**, 430 (1992); I. I. Bigi, M. Shifman, N. Uraltsev, and A. Vainshtein, Phys. Rev. Lett. **71**, 496 (1993); B. Blok, L. Koyrakh, M. Shifman, and A. Vainshtein, Phys. Rev. **D49**, 3356 (1994); A. V. Manohar and M. B. Wise, Phys. Rev. **D49**, 1310 (1994); M. Gremm and A. Kapustin, Phys. Rev. **D55**, 6924 (1997).
- [3] I. I. Bigi, M. Shifman, N. Uraltsev, and A. Vainshtein, Phys. Rev. **D56**, 4017 (1997).
- [4] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. **D67**, 031101 (2003).
- [5] M. Voloshin, Phys. Rev. **D51**, 4934 (1995); A. Falk, M. Luke, and M. Savage, Phys. Rev. **D53**, 2491 (2001); *ibid.*, **D53**, 6316 (1996).
- [6] D. Cronin-Hennessy *et al.* [CLEO Collaboration], Phys. Rev. Lett. **87**, 251808 (2001).
- [7] C. W. Bauer, Z. Ligeti, M. Luke and A. V. Manohar Phys. Rev. **D67**, 054012 (2003).
- [8] A. H. Mahmood *et al.* [CLEO Collaboration], Phys. Rev. **D67**, 072001 (2003).
- [9] M. Battaglia *et al.* [DELPHI Collaboration], Phys. Lett. **B556**, 41 (2003).
- [10] B. Aubert *et al.* [BABAR Collaboration], *Measurement of Moments of the Hadronic Mass Distribution in Semileptonic B Decays*, hep-ex/0403031, submitted to Phys. Rev. D (2004).
- [11] B. Aubert *et al.* [BABAR Collaboration], *Measurement of the Electron Energy Spectrum and its Moments in Inclusive $B \rightarrow X e \nu$ Decays*, hep-ex/0403030, submitted to Phys. Rev. D (2004).
- [12] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **92**, 071802 (2004).
- [13] E. Barbiero and Z. Was, Comput. Phys. Comm. **79**, 291 (1994).
- [14] D. Benson, I. I. Bigi, T. Mannel, and N. Uraltsev, Nucl. Phys. **B665**, 367 (2003).
- [15] P. Gambino and N. Uraltsev, *Moments of Semileptonic B Decay Distributions in the $1/m_b$ Expansion*, hep-ph/0401063, to be published in Eur. Phys. J. C (2004); the lepton-energy dependence of the perturbative corrections has been added for the hadronic-mass moments, based on N. Uraltsev, *Perturbative Corrections to Semileptonic B-Decay Moments*, hep-ph/0403166.
- [16] K. Hagiwara *et al.* Phys. Rev. **D66**, 010001 (2002).
- [17] J. P. Alexander *et al.* [CLEO Collaboration], Phys. Rev. Lett. **86**, 2737 (2001); B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. **D65**, 032001 (2002); S.B. Athar *et al.* [CLEO Collaboration], Phys. Rev. **D66**, 052003 (2002).
- [18] I. I. Bigi, M. A. Shifman, and N. G. Uraltsev, Ann. Rev. Nucl. Part. Sci., **47**, 591 (1997).
- [19] M. Trott, *Improving Extractions of $|V_{cb}|$ and m_b from the Hadronic Invariant Mass Moments of Semileptonic Inclusive B Decays*, hep-ph/0402120.