

# Soft QCD results from the ALICE experiment

*Luciano Ramello for the ALICE Collaboration*

*Di.S.T.A., Università del Piemonte Orientale, I-10121 Alessandria, ITALY*

## 1 Introduction

ALICE (A Large Ion Collider Experiment) is designed to study Heavy Ion collisions at the LHC (see [1]) and features in the barrel region a very robust tracking based on a large Time Projection Chamber (TPC) and an excellent vertexing provided by an Internal Tracking System (ITS) made of 6 layers of silicon detectors. The material budget up to the outer wall of the TPC (2.5 m) is very low ( $\approx 0.11 \cdot X_0$ ) implying a very low  $p_t$  cutoff of 100 MeV/c for pions. Particle identification is provided, for the results shown in the following, by the Time-of-Flight detector in addition to TPC and ITS detectors. More details are given in [1]. In the following, published and preliminary results obtained by ALICE for proton-proton collisions at LHC energies will be presented, covering diffraction and total cross-section, inclusive and identified particle spectra, correlations and fluctuations. Results from the first Pb-Pb run of 2010, for which pp data constitute the necessary baseline, are presented in [2].

## 2 Diffraction and total cross-section

Using the two-arm VZERO scintillator hodoscope for triggering and the van der Meer scan technique, cross-section have been measured for inelastic pp interaction at 2.76 and 7 TeV energies, see Table 1. The trigger acceptance for inelastic events was about 75% at both energies. In addition the cross section for a reference inelastic process, namely production of at least one charged particle with  $|\eta| < 0.8$  and  $p_t > 0.5$  GeV/c, has been measured and found in agreement with the results from ATLAS and CMS experiments. Single and double diffractive cross-sections have been measured as well<sup>1</sup>.

Energy (TeV)	Process	Cross-section (mb)
2.76	Inelastic	$62.1 \pm 1.6(\text{model}) \pm 4.3(\text{lumi})$
7.0	Inelastic	$72.7 \pm 1.1(\text{model}) \pm 5.1(\text{lumi})$
7.0	Inelastic, $N \geq 1$	$42.4 \pm 2.0$

Table 1: Preliminary cross-sections measured by ALICE for various processes.

---

<sup>1</sup>M. Poghosyan for the ALICE Coll., Proc. Quark Matter 2011, 23–28 May 2011, Annecy, France.

### 3 Inclusive production, identified particle spectra

Charged particle  $dN_{ch}/d\eta$  and multiplicity measured by ALICE in 0.9, 2.36 and 7 TeV pp collisions have been published [3], as well as charged particle  $p_t$  spectra at 0.9 TeV [4]. Preliminary  $p_t$  spectra at 2.76 and 7 TeV are shown in Fig. 1 (left), extending up to 100 GeV/c for the highest energy. Preliminary  $p_t$  spectra for identified  $\pi$ , K and p at midrapidity are shown in Fig. 1 (right), while spectra at 0.9 TeV are available in [5]. A fit with Lévy functions allowed us to estimate the yield and the average transverse momentum of each species with minimal extrapolation; the resulting mean  $p_t$  is shown in Fig. 2 (left), a modest increase with  $\sqrt{s}$  is seen. Fig. 2 (right) shows the  $K/\pi$  ratio evolution from SPS through RHIC to LHC energies; this ratio is approximately energy-independent between top RHIC energy and 7 TeV.




Figure 1:  $p_t$  spectra in pp collisions, for charged particles at three energies (left) and for pions, kaons and protons at 7 TeV (right).

ALICE has obtained preliminary spectra for multi-strange hadrons and resonances at 7 TeV, see Fig. 3. For  $\Omega$  and  $\Xi^2$ , the antiparticle/particle ratio measured over a wide  $p_t$  range is compatible with unity. The bottom panel of Fig. 3 (left) shows the ratio of ALICE data to the recent Z2 tune of PYTHIA 6, which is seen to underpredict multi-strange baryon production, with only a marginal improvement over the Perugia-0 tune (not shown).

The preliminary spectra of some resonances:  $\phi$ ,  $K^{*0}$  and  $\Xi^{*0}$  are shown in Fig. 3 (right). The  $\phi$  yield agrees with PYTHIA 6 (tune D6T, not shown) up to 2 GeV/c, while for higher  $p_t$  a better agreement with PHOJET is found.

<sup>2</sup>The mid-rapidity yield of  $\Xi$  measured by ALICE agrees with the one by CMS when one takes into account the different normalization (INEL vs. NSD).




Figure 2: (left) Average transverse momentum of pions, kaons and protons from RHIC to LHC energies; (right)  $K/\pi$  ratio in pp and  $\bar{p}p$  collisions vs. energy.




Figure 3: (left) Transverse momentum spectra of  $\Xi$ 's,  $\Omega$ 's and of the respective antibaryons in pp collisions at 7 TeV; (right) Transverse momentum spectra of some resonances in pp collisions at 7 TeV, with Lévy-Tsallis fits.

## 4 Correlations and fluctuations

The space-time characteristics of the particle production region have been studied in detail for the first time in pp collisions at 0.9, 2.76 and 7 TeV, as a function of pair transverse momentum  $k_t$  and  $dN_{ch}/d\eta$ , see Fig. 4 (left). The three HBT radii (out, side and long) at different  $\sqrt{s}$  are seen to scale vs.  $(dN_{ch}/d\eta)^{1/3}$ ; the side and long radii grow with  $dN_{ch}/d\eta$  at all  $k_t$ 's, while the out radius is flat and even decreasing at high  $k_t$ . At high  $dN_{ch}/d\eta$ , all three radii fall with  $k_t$ .

Event-by-event fluctuations of the mean  $p_t$  are measured via a two-particle correlator  $C_m$  which is zero for statistical fluctuations only; the relative fluctuations in pp collisions at the 3 LHC energies, shown in Fig. 4 (right), show a universal behaviour vs. uncorrected multiplicity, a good baseline for the same study in Pb-Pb collisions.




Figure 4: (left) Scaling of HBT radii in pp collisions at 3 different energies vs.  $\langle dN_{ch}/d\eta \rangle$ ; (right) Relative mean  $p_t$  fluctuations vs. uncorrected multiplicity.

## 5 Conclusions

A rich pp programme has been developed by ALICE at LHC energies. Inelastic, single and double diffraction cross-sections have been measured. The production of charged particles up to 100 GeV/c and of  $\pi$ ,  $K$ ,  $p$ ,  $\pi^0$ ,  $\eta$ , hyperons,  $\omega$ ,  $\phi$  and resonances up to 5-10 GeV/c has been measured as well. HBT radii vs.  $k_t$  and multiplicity, and mean  $p_t$  fluctuations vs. multiplicity have been measured at three LHC energies.  $J/\psi$  and  $D$  meson cross-sections and leptons from charm and beauty decays have been measured [6] both in the barrel and in the forward muon spectrometer. Extensive comparisons with pQCD and MonteCarlo models have been performed. Finally, a solid baseline for the study of Pb-Pb collisions at LHC has been established.

## References

- [1] J. Wessels, these Proceedings.
- [2] A. Morsch, these Proceedings.
- [3] K. Aamodt *et al.*, ALICE Coll., Eur. Phys. J. **C 68**, 89 (2010) and 345 (2010).
- [4] K. Aamodt *et al.*, ALICE Collaboration, Phys. Lett. **B 693**, 53 (2010).
- [5] K. Aamodt *et al.*, ALICE Collaboration, Eur. Phys. J. **C 71**, 1655 (2011).
- [6] C. Hadjidakis, these Proceedings; D. Caffarri, these Proceedings.