

D⁰ cross section in pp collisions at $\sqrt{s} = 7$ TeV, measured with the ALICE experiment

*Xianbao Yuan, for the ALICE Collaboration
 Institute of Particle Physics, Central China Normal University
 University and INFN, Padova, Italy
 Contact: yuanxb@iopp.cnu.edu.cn or xianbao.yuan@pd.infn.it*

The measurement of the cross-section for charm production in pp collisions at the LHC is not only a fundamental reference to investigate medium properties in heavy-ion collisions, but also a key test of pQCD predictions in a new energy domain.

The ALICE [1] experiment has measured the D meson production in pp collisions at $\sqrt{s} = 7$ TeV. We present the analysis procedure for $D^0 \rightarrow K^- \pi^+$ and for the calculation of efficiency and acceptance corrections. Finally, we show the preliminary results on D^0 cross section in pp collisions at $\sqrt{s} = 7$ TeV, measured in the region $2 < p_t < 12$ GeV/c at central rapidity $|\eta| < 0.5$. These results are compared to perturbative QCD predictions.

The analysis is based on an invariant mass analysis of opposite-charge pairs of reconstructed tracks that can represent a D^0 with a displaced vertex (the mean proper decay length of the D^0 is $c\tau \approx 123 \mu\text{m}$). The selection is based on topological cuts and particle identification via specific energy deposit and time-of-flight measurement. The cross section is calculated from the raw signal yield extracted with the invariant mass analysis, $N^{D^0 \text{ raw}}(p_t)$, using the following formula:

$$\left. \frac{d\sigma^{D^0}}{dp_t} \right|_{|\eta| < 0.5} = \frac{1}{2} \frac{1}{2 y_{\text{acc}} \Delta p_t} \frac{f_{\text{prompt}} \cdot N^{D^0 \text{ raw}}(p_t) \Big|_{|\eta| < y_{\text{acc}}}}{\epsilon_{\text{prompt}} \cdot \text{BR} \cdot L_{\text{int}}} \quad (1)$$

Here, ϵ_{prompt} means the efficiency of prompt mesons, which accounts for selection cuts, for track and primary vertex reconstruction efficiency, and for detector acceptance. The f_{prompt} is the prompt fraction of raw yield.

Figure 1 (Left) shows the invariant mass distribution for $p_t > 2$ GeV/c after applying the cuts, which corresponds to 1.1×10^8 minimum bias events collected by ALICE in 2010 at $\sqrt{s} = 7$ TeV. Figure 1 (Right) shows the efficiencies for $D^0 \rightarrow K^- \pi^+$ with all the decay particles in the acceptance $|\eta| < 0.9$. The efficiencies increase and flatten at about 0.1 at $p_t > 2$ GeV/c. The efficiency without particle identification selection, shown for comparison, is the same as with particle identification for $p_t > 2$ GeV/c, indicating that this selection is essentially fully efficient for the signal. The efficiencies for D^0 meson from B meson decay, also shown for comparison, are larger by a factor about 2, because this feed-down component is more displaced from the primary vertex, due to the longer B life time. The 10 – 15% feed-down from B decays is subtracted based on pQCD prediction [2].

Several sources of systematic uncertainties were considered, namely those affecting the signal extraction from the invariant mass spectra and all the correction factors applied to obtain the p_t -differential cross sections. A summary of the estimated relative systematic errors is shown in Fig 2 (Left).

The p_t -differential cross section for prompt D^0 , obtained from the yields extracted by fitting the invariant mass spectra and corrected for efficiency and B feed-down, is shown in Fig 2 (Right). The error bars represent the statistical errors, while the systematic errors are plotted as rectangle areas around the data points. The measured D^0 meson production cross section is compared to two theoretical predictions, namely FONLL [2] and GM-VFNS [3]. Our measurement of D^0 at $\sqrt{s} = 7$ TeV is reproduced by both models within their theoretical uncertainties.




Figure 1: Left: $p_T > 2$ GeV/c invariant mass distribution. Right: efficiencies for D^0 as a function of p_T .




Figure 2: Left: systematic errors summary plot. Right: p_T -differential cross section for prompt D^0 in pp collisions at $\sqrt{s} = 7$ TeV compared with FONLL [2] and GM-VFNS [3] theoretical predictions.

References

- [1] B. Abelev, A. Abrahantes Quintana, *et al.*, [ALICE Coll.], Arxiv: hep-ex/1111.1553v1.
- [2] M. Cacciari *et al.*, JHEP 0407 (2004) 033; Private communication.
- [3] B.A. Kniehl *et al.*, Phys. Rev. D77 (2008) 014011; Private communication.