

Particle Production and Diffraction in ATLAS

*James Robinson
The University of Manchester
Oxford Road, Manchester
M13 9PL, UK*

On behalf of the ATLAS Collaboration

1 Introduction

The study of soft QCD provides insight into the physics of hadronic cross-sections and hadron formation, while also enabling the validation and tuning of models of particle production and helping to describe and correct for pile-up or soft backgrounds to other processes. Soft QCD measurements in ATLAS [1] at the LHC mainly use the inner tracking detectors, sensitive to charged particles with $p_T > 100$ MeV and $|\eta| < 2.5$, and the electromagnetic and hadronic calorimeters, sensitive to electrons, photons and hadrons that have E_T greater than a few hundred MeV and $|\eta| < 4.9$. Most such measurements utilise Minimum Bias events: inclusive collisions which have been triggered by scintillators located in the region $2.1 < |\eta| < 3.8$.

2 Inclusive Two Particle Angular Correlations

Correlations in $\Delta\eta$ and $\Delta\phi$ were examined through a variable which takes into account both inter-event and intra-event correlations. Results, corrected using the HBOM unfolding technique [2] (repeatedly applying detector folding n times and extrapolating to the hypothetical $n = -1$ case), were compared to several PYTHIA 6 tunes [3], PYTHIA 8 and HERWIG++ [4], with the data not satisfactorily described by any of these. A large difference between PYTHIA and HERWIG++ is visible, reflecting the string/cluster hadronisation modelling difference [5].

3 Forward-backward Correlations

Correlations between the forward and backward regions were measured as a function of the multiplicity and $\sum p_T$ of charged particles in bins of $|\eta|$. The data were corrected for detector effects using multiple regression. This was the first measurement of $\sum p_T$ correlations, with the latest Monte Carlo tunes accurately reproducing the data [6].

4 Azimuthal ordering of charged hadrons

A spectral analysis of correlations between angular, energy and p_T orderings was performed. After correcting for detector effects, the comparison with Monte Carlo simulation showed too much correlation for high- p_T charged particles, but too little correlation at low- p_T (see Figure 1).

Increasing the level of underlying event in the Monte Carlo could improve the agreement at high p_T , while increasing the levels of initial state radiation could improve the agreement at low p_T . The net result is a problem for Monte Carlo tuning - simply changing the typical settings will not find a tune that can fit all of the data.




Figure 1: Angular/energy correlations at high p_T (left) and low p_T (right) [7].

The measured spectra show features consistent with helix-ordered gluon chain models [8], which impose correlations between angular and p_T orderings; including such effects could improve soft particle production and hadronisation models [7].

5 Inelastic cross-section as a function of forward rapidity gap size

Cross-sections, corrected to the hadron-level, were measured as a function of $\Delta\eta_F$, the largest η -region containing no particles with p_T above p_T^{cut} , counting inwards from the edge of the detector acceptance, at $|\eta| = 4.9$.

At low values of $\Delta\eta_F$, this measurement tests the reliability of hadronisation models in describing rapidity fluctuations in final-state particle production; at higher values it probes the diffractive cross-section. Figure 2 shows that the rise of the cross-section at the largest $\Delta\eta_F$ values is not reproduced in Monte Carlo (left) and also indicates that the data contain a higher proportion of low mass single-diffractive events than any of the theoretical models (right): this is shown by the steepness of the transition between ATLAS and TOTEM data [9].




Figure 2: Rapidity gap cross sections in PYTHIA 6, PYTHIA 8, PHOJET and data, with $p_T^{cut} = 200$ MeV (left). The total cross-section as a function of $\xi_X = M_X^2/s$, where ξ_{cut} is a minimum (right), comparing data against various theoretical predictions [9].

6 K_S^0 and Λ production

K_S^0 and Λ candidates were identified by fitting pairs of opposite-sign tracks to a common vertex and cutting on the transverse flight distance between primary and secondary vertices and on the angle between this vector and the particle momentum.

The observed distributions of p_T , y and multiplicity for K_S^0 and Λ hadrons were corrected to the hadron level and compared to Monte Carlo simulations. While the K_S^0 distributions agreed to within 15%, the Λ p_T distribution showed substantial discrepancies visible in both the HERWIG and PYTHIA predictions and across multiple generator tunes [10].

7 Underlying event in charged-particle jet events

Underlying event (UE) is used to describe any hadronic activity not associated with the hard scattering process. In typical measurements, the highest p_T jet in the event used to define a “transverse region”, the area satisfying $\pi/3 < |\phi^{particle} - \phi^{jet}| \leq 2\pi/3$, where ϕ is the azimuthal angle. The multiplicity and $\sum p_T$ of charged particles are used to characterise the UE.

Several Monte Carlo simulations are compared to the unfolded data, with PYTHIA 6 (AUET2B tune) showing the best agreement. The analysis is performed using a range of different anti- k_t R -parameters to identify the jets [11]. Figure 3 shows the ratio at different R -parameters of number of charged tracks identified in the transverse region as a function of the p_T of the jet. The number of tracks identified has a strong dependence on the R -parameter, both in data and Monte Carlo, indicating that the choice of hard scatter has a crucial effect on UE distributions [12].




Figure 3: Ratio of number of charged tracks for different anti- k_t R -parameters [12].

References

- [1] ATLAS Collaboration, J. Instrum. **3** (2008) S08003.
- [2] J. Monk and C. Oropeza-Barrera, arXiv:1111.4896 [hep-ex].
- [3] T. Sjostrand, S. Mrenna and P. Z. Skands, JHEP **0605** (2006) 026, arXiv:0603175 [hep-ph].
- [4] M. Bahr, S. Gieseke, M. A. Gigg, D. Grellscheid, K. Hamilton, O. Latund-Dada, S. Platzer and P. Richardson *et al.*, Eur. Phys. J. C **58** (2008) 639, arXiv:0803.0883 [hep-ph].
- [5] ATLAS Collaboration, JHEP **1205** (2012) 157, arXiv:1203.3549 [hep-ex].
- [6] ATLAS Collaboration, JHEP **1207** (2012) 019, arXiv:1203.3100 [hep-ex].
- [7] ATLAS Collaboration, arXiv:1203.0419 [hep-ex].
- [8] S. Todorova, arXiv:1204.2655v3 [hep-ph]
- [9] ATLAS Collaboration, Eur. Phys. J. **C72** (2012) 1926, arXiv:1201.2808 [hep-ex].
- [10] ATLAS Collaboration, Phys. Rev. **D85** (2012) 012001, arXiv:1111.1297 [hep-ex].
- [11] M. Cacciari, G. P. Salam and G. Soyez, JHEP **0804** (2008) 063, arXiv:0802.1189 [hep-ph].
- [12] ATLAS Collaboration, arXiv:1208.0563 [hep-ex].