arXiv:2203.00042v4 [physics.ins-det] 20 Apr 2022

A Call to Arms Control: Synergies between Nonproliferation Applications of Neutrino Detectors and Large-Scale Fundamental Neutrino Physics Experiments A Snowmass White Paper

Adam Bernstein,^{1, *} Tomi Akindele,^{1, †} Zelimir Djurcic,^{2, †} Logan Lebanowski,^{3, †} Jost Migenda,^{4, †} Anthony Papatyi,^{5, †} Mayly Sanchez,^{6, †} Michael Smy,^{7, †} and Mark Vagins^{7, †} (Editor and Contributors)

Z. Djurcic,² C. Mariani,⁸ A. Erlandson,⁹ R. Mahapatra,¹⁰ M. V. Garzelli,¹¹ E. Kemp,¹² R. Bruce Vogelaar,⁸
J. Pedro Ochoa-Ricoux,⁷ G. Pronost,¹³ S. Seo,¹⁴ F. Suekane,¹⁵ W. Wang,¹⁶ Y. Keung Hor,¹⁶ and M. Chen¹⁷ (Co-Signers from DUNE, SNO+, JUNO, Super-K, Hyper-K, and other collaborations)

T. Anderson,¹⁸ E. Anderssen,¹⁹ M. Askins,^{19,20} A. J. Bacon,³ Z. Bagdasarian,^{19,20} A. Baldoni,¹⁸ N. Barros,^{21,22} L. Bartoszek,²³ A. Bat,²⁴ M. Bergevin,¹ A. Bernstein,¹ E. Blucher,²⁵ J. Boissevain,²³ R. Bonventre,¹⁹ D. Brown,¹⁹ E. J. Callaghan,^{19,20} D. F. Cowen,¹⁸ S. Dazeley,¹ M. Diwan,²⁶ K. Frankiewicz,²⁷ C. Grant,²⁷ T. Kaptanoglu,^{19,20} J.R. Klein,³ C. Kraus,^{28,29} T. Kroupa,³ B. Land,^{19,20,3} L. Lebanowski,³ V. Lozza,^{21,22} A. Marino,³⁰ A. Mastbaum,³¹ C. Mauger,³ S. Naugle,³ M. Newcomer,³ A. Nikolica,³ G. D. Orebi Gann,^{19,20} L. Pickard,³² J. Saba,¹⁹ S. Schoppmann,^{19,20} J. Sensenig,³ M. Smiley,^{19,20} H. Steiger,^{33,34} R. Svoboda,³² E. Tiras,^{24,35} W. H. Trzaska,³⁶

R. van Berg,^{3,23} G. Wendel,¹⁸ M. Wetstein,⁶ M. Wurm,³³ G. Yang,^{19,20} M. Yeh,²⁶ and E.D. Zimmerman³⁰

(Co-Signers from Eos)

M. Askins,^{19,20} Z. Bagdasarian,^{19,20} N. Barros,^{3,21,22} E.W. Beier,³ A. Bernstein,¹ M. Böhles,³³ E. Blucher,²⁵ R. Bonventre,¹⁹ E. Bourret,¹⁹ E. J. Callaghan,^{20,19} J. Caravaca,^{20,19} M. Diwan,²⁶ S.T. Dye,³⁷ J. Eisch,³⁸ A. Elagin,²⁵ T. Enqvist,³⁶ U. Fahrendholz,³⁴ V. Fischer,³² K. Frankiewicz,²⁷ C. Grant,²⁷ D. Guffanti,³³ C. Hagner,³⁹ A. Hallin,⁴⁰ C. M. Jackson,⁵ R. Jiang,²⁵ T. Kaptanoglu,^{20,19} J.R. Klein,³ Yu. G. Kolomensky,^{20,19} C. Kraus,^{28,29} F. Krennrich,⁶ T. Kutter,⁴¹ T. Lachenmaier,⁴² B. Land,^{20,19,3} K. Lande,³ L. Lebanowski,³ J.G. Learned,³⁷ V.A. Li,¹ V. Lozza,^{21,22} L. Ludhova,^{43,44} M. Malek,⁴⁵ S. Manecki,^{29,46,28} J. Maneira,^{21,22} J. Maricic,³⁷ J. Martyn,³³ A. Mastbaum,³¹ C. Mauger,³ M. Mayer,³⁴ J. Migenda,⁴ F. Moretti,¹⁹ J. Napolitano,⁴⁷ B. Naranjo,⁴⁸ S. Naugle,³ M. Nieslony,³³ L. Oberauer,³⁴ G. D. Orebi Gann,^{20,19} J. Ouellet,⁴⁹ T. Pershing,³² S.T. Petcov,⁵⁰ L. Pickard,³² R. Rosero,²⁶ M. C. Sanchez,⁶ J. Sawatzki,³⁴ S. Schoppmann,^{20,19} S.H. Seo,⁵¹ M. Smiley,^{20,19} M. Smy,⁷ A. Stahl,⁴⁴ H. Steiger,^{33,34} M. R. Stock,³⁴ H. Sunej,²⁶ R. Svoboda,³² E. Tiras,^{24,35} W. H. Trzaska,³⁶ M. Tzanov,⁴¹ M. Vagins,⁷ C. Vilela,⁵² Z. Wang,⁵³ J. Wang,⁵⁴ M. Wetstein,⁶ M.J. Wilking,⁵² L. Winslow,⁴⁹ P. Wittich,⁵⁵ B. Wonsak,³⁹ E. Worcester,^{26,52} M. Wurm,³³ G. Yang,⁵² M. Yeh,²⁶ E.D. Zimmerman,³⁰ S. Zsoldos,^{20,19} and K. Zuber⁵⁶ (Co-Signers from Theia)

T. Akindele,¹ T. Anderson,¹⁸ M. Askins,^{19,20} A. Baldoni,¹⁸ A. Barna,³⁷ T. Benson,⁵⁷ M. Bergevin,¹ A. Bernstein,¹ B. Birrittella,⁵⁷ J. Boissevain,³ J. Borusinki,³⁷ D. Cowen,¹⁸ B. Crow,³⁷ F. Dalnoki-Veress,⁵⁸ D. Danielson,²⁵ S. Dazeley,¹ M. Diwan,²⁶ A. Druetzler,³⁷ S. Dye,³⁷ A. Fienberg,¹⁸ V. Fischer,³² K. (Kat) Frankiewicz,²⁷ D. Gooding,²⁷ C. Graham,⁵⁹ C. Grant,²⁷ J. Griskevich,⁷ J. He,³² J. Hecla,^{20,1} I. Jovanovic,⁵⁹ M. Keenan,¹⁸ P. Keener,³ P. Kunkle,²⁷ J. Learned,³⁷ V. Li,¹ J. Maricic,³⁷ P. Marr-Laundrie,⁵⁷ J. Moore,¹⁸ A. Mullen,^{20,1} E. Neights,¹⁸ K. Nishimura,³⁷ B. O'Meara,¹⁸ K. Ogren,⁵⁹ G. D. Orebi Gann,^{20,19} L. Oxborough,⁵⁷ A. Papatyi,⁵ B. Paulos,⁵⁷ T. Pershing,¹ L. Pickard,³² L. Sabarots,⁵⁷ V. Shebalin,³⁷ M. Smy,⁷ H. Song,²⁷ F. Sutanto,¹ R. Svoboda,³² M. Vagins,⁷ R. (Rick) Van Berg,³ G. Varner,³⁷ V. Veeraraghavan,⁶ S.

Ventura,³⁷ B. Walsh,²⁶ G. Wendel,¹⁸ D. Westphal,¹ M. Wetstein,⁶ A. Wilhelm,⁵⁹ S. Wolcott,⁵⁷ and M. Yeh²⁶

(Co-Signers from WATCHMAN)

¹Lawrence Livermore National Laboratory, Livermore, CA 94550, USA

²Argonne National Laboratory, Lemont, IL 60439, USA

³Department of Physics and Astronomy, University of Pennsylvania, Philadelphia, PA 19104-6396

⁴King's College London, Department of Physics, Strand Building, Strand, London WC2R 2LS, UK

⁵Pacific Northwest National Laboratory, Richland, WA 99352, USA

⁶Department of Physics and Astronomy, Iowa State University, Ames, IA 50011, USA

⁷University of California, Irvine, Department of Physics, CA 92697, Irvine, USA

⁸Center for Neutrino Physics, Virginia Tech, Blacksburg, Virginia 24061 USA

⁹Canadian Nuclear Laboratories Ltd., Chalk River, ON, Canada

¹⁰Texas A&M University, College Station, Texas

¹¹University of Hamburg, Hamburg, Germany

¹²Universidade de Campinas - UNICAMP, Cidade Universitaria "Zeferino Vaz" CEP 13083-970 - Campinas-SP

¹³Kamioka Observatory, ICRR, University of Tokyo

¹⁴IBS Center for Underground Physics, 34126, IBS, 55, Expo-ro, Yuseong-gu, Daejeon, Republic of Korea

¹⁵Research Center for Neutrino Science (RCNS), Tohoku University, Tohuku, Japan

¹⁶Sun Yat-sen University, Guangzhou, China

¹⁷Oueen's University, Kingston, ON K7L 3N6, Canada

¹⁸Pennsylvania State University, University Park, PA 16802, USA

¹⁹Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley, CA 94720-8153, USA

²⁰Physics Department, University of California at Berkeley, Berkeley, CA 94720-7300

²¹ Universidade de Lisboa, Faculdade de Ciências (FCUL), Departamento de Física, Campo Grande, Edifício C8, 1749-016 Lisboa, Portugal

²²Laboratório de Instrumentação e Física Experimental de Partículas (LIP), Av. Prof. Gama Pinto, 2, 1649-003, Lisboa, Portugal

²³Bartoszek Engineering, Aurora, IL 60506, USA

²⁴Department of Physics, Erciyes University, 38030, Kayseri, Turkey

²⁵The Enrico Fermi Institute and Department of Physics, The University of Chicago, Chicago, IL 60637, USA

²⁶Brookhaven National Laboratory, Upton, New York 11973, USA

²⁷Boston University, Department of Physics, Boston, MA 02215, USA

²⁸SNOLAB, Creighton Mine 9, 1039 Regional Road 24, Sudbury, ON P3Y 1N2, Canada

²⁹Laurentian University, Department of Physics, 935 Ramsey Lake Road, Sudbury, ON P3E 2C6, Canada

³⁰University of Colorado at Boulder, Department of Physics, Boulder, Colorado, USA

³¹Department of Physics and Astronomy, Rutgers, The State University

of New Jersey, 136 Frelinghuysen Road, Piscataway, NJ 08854-8019 USA ³²University of California, Davis, 1 Shields Avenue, Davis, CA 95616, USA

³³Institute of Physics and Excellence Cluster PRISMA, Johannes Gutenberg-Universität Mainz, 55099 Mainz, Germany

³⁴Physics Department, Technische Universität München, 85748 Garching, Germany

³⁵Department of Physics and Astronomy, The University of Iowa, Iowa City, Iowa, USA ³⁶Department of Physics, University of Jyväskylä, Finland

³⁷University of Hawai'i at Manoa, Honolulu, Hawai'i 96822, USA

³⁸Fermi National Accelerator Laboratory, Batavia, IL 60510, USA

³⁹Institut für Experimentalphysik, Universität Hamburg, 22761 Hamburg, Germany

⁴⁰University of Alberta, Department of Physics, 4-181 CCIS, Edmonton, AB T6G 2E1, Canada

⁴¹Department of Physics and Astronomy, Louisiana State University, Baton Rouge, LA 70803

⁴²Kepler Center for Astro and Particle Physics, Universität Tübingen, 72076 Tübingen, Germany

³Institut für Kernphysik, Forschungszentrum Jülich, 52425 Jülich, Germany

⁴⁴III. Physikalisches Institut B, RWTH Aachen University, 52062 Aachen, Germany

⁴⁵University of Sheffield, Physics & Astronomy, Western Bank, Sheffield S10 2TN, UK

⁴⁶Queen's University, Department of Physics, Engineering Physics & Astronomy, Kingston, ON K7L 3N6, Canada

⁴⁷Department of Physics, Temple University, Philadelphia, PA, USA

⁴⁸ University of California Los Angeles, Department of Physics & Astronomy, 475 Portola Plaza, Los Angeles, CA 90095-1547, USA

⁴⁹Massachusetts Institute of Technology, Department of Physics and Laboratory

for Nuclear Science, 77 Massachusetts Ave Cambridge, MA 02139, USA

⁵⁰SISSA/INFN, Via Bonomea 265, I-34136 Trieste, Italy

Kavli IPMU (WPI), University of Tokyo, 5-1-5 Kashiwanoha, 277-8583 Kashiwa, Japan

¹¹Center for Underground Physics, Institute for Basic Science, Daejeon 34126, Korea

⁵²State University of New York at Stony Brook, Department of Physics and Astronomy, Stony Brook, New York, USA

⁵³Department of Engineering Physics, Tsinghua University, Beijing 100084, China

⁵⁴Department of Physics, South Dakota School of Mines and Technology, Rapid City, SD 57701, USA ⁵⁵Cornell University, Ithaca, NY, USA

⁵⁶Institut für Kern und Teilchenphysik, TU Dresden, Zellescher Weg 19, 01069, Dresden, Germany

⁵⁷University of Wisconsin, Madison, Wisconsin 53706

⁵⁸Middlebury Institute of International Studies at Monterey, Monterey, California 93940, USA ⁵⁹University of Michigan, Ann Arbor, MI, USA

^{*} Editor.

[†] Primary Contributor.

EXECUTIVE SUMMARY

- The High Energy Physics community can benefit from a natural synergy in research activities into next-generation largescale water and scintillator neutrino detectors, now being studied for remote reactor monitoring, discovery and exclusion applications in cooperative nonproliferation contexts.
- Since approximately 2010, US nonproliferation researchers, supported by the National Nuclear Security Administration (NNSA), have been studying a range of possible applications of relatively large (100 ton) to very large (hundreds of kiloton) water and scintillator neutrino detectors.
- In parallel, the fundamental physics community has been developing detectors at similar scales and with similar design features for a range of high-priority physics topics, primarily in fundamental neutrino physics. These topics include neutrino oscillation studies at beams and reactors, solar, and geological neutrino measurements, supernova studies, and others.
- Examples of ongoing synergistic work at U.S. national laboratories and universities include prototype gadolinium-doped water and water-based and opaque scintillator test-beds and demonstrators, extensive testing and industry partnerships related to large area fast position-sensitive photomultiplier tubes, and the development of concepts for a possible underground kiloton-scale water-based detector for reactor monitoring and technology demonstrations.
- Some opportunities for engagement between the two communities include bi-annual Applied Antineutrino Physics conferences, collaboration with U.S. National Laboratories engaging in this research, and occasional NNSA funding opportunities supporting a blend of nonproliferation and basic science R&D, directed at the U.S. academic community.

I. INTRODUCTION

The High Energy Physics community can benefit from a natural synergy in research activities into next-generation large-scale water and scintillator neutrino detectors, now being studied for remote reactor monitoring, discovery and exclusion applications in cooperative nonproliferation contexts.

Since approximately 2010, US nonproliferation researchers, supported by the National Nuclear Security Administration (NNSA), have been studying a range of possible applications of relatively large (100 ton) to very large (hundreds of kiloton) water and scintillator neutrino detectors. Primarily due to investments by NNSA, but also with important HEP contributions, a significant R&D effort has arisen at US National Laboratories, working with academic collaborators to understand and advance the nonproliferation and fundamental science potential of such detectors. In this white paper, we offer a broad and non-exhaustive summary of areas of actual and potential overlap between HEP fundamental science priorities and NNSA nonproliferation priorities, and suggest ways to continue and formalize the exploitation of the synergies that are already in place.

In nonproliferation contexts, it is often difficult to reconcile the conflicting goals of non-intrusiveness and robust verification of the absence of illicit nuclear programs. One important area of interest is to confirm either the absence, or conversely the licit operation, of nuclear reactors - the sources of all the world's plutonium. Antineutrinos, whose penetrating signature of nuclear origin can provide insight into the operations and characteristics of nuclear reactors, offer a promising path towards remote and non-intrusive, but still robust and persistent verification of reactor operations. These applications have been explored in the last decade, and have resulted in ongoing US and international (primarily UK) collaborative efforts to advance relevant technologies, as well as a deeper understanding of the potential and limitations of those technologies for nonproliferation. In parallel, the high energy physics community is pursuing similar technical goals that advance fundamental physics, in the areas of MeV-scale solar, atmospheric and geological neutrinos, supernova detection, neutrinoless double beta decay, and the GeV-scale high energy neutrino oscillation and CP violation physics that are the main goals of the US's flagship Deep Underground Neutrino Experiment (DUNE) [1]. In this white paper, we provide an incomplete survey of ongoing work in this overlap area, and suggest possible approaches to facilitate cooperation and synergy among the two communities.

II. FUNDAMENTAL PHYSICS RELEVANCE

A range of fundamental physics applications make use of large unsegmented liquid detectors – from the 100 ton to the hundred kiloton scale. Water-based detectors such as Hyper-Kamiokande [2], scintillator detectors such as JUNO [3], and proposed hybrid or alternative media such as Theia [4] and Liquid-O [5] make use of broadly similar technologies. Science programs making use of these technologies are far-reaching and already popular in the HEP community. They include such goals as neutrinoless double beta decay, geoneutrino, solar neutrino and supernova physics at the MeV scale, and atmospheric neutrino and beam neutrino oscillation and CP-violation physics at the GeV scale. As presented in the 2019 DUNE Module of Opportunity workshop [6], such detectors are an attractive companion technology to the liquid argon detectors now envisioned for DUNE, with different systematic considerations for accelerator physics goals, and a different and lower energy regime made available by the hydrogenous target, relevant to a complementary set of physics goals.

III. NONPROLIFERATION APPLICATIONS

The monitoring of nuclear reactor operations, and the discovery or exclusion of unsanctioned plutonium production reactors are central concerns of the global nuclear nonproliferation regime. For convenience we define two standoff domains with their own technological approaches. In the mid-field, from 2 kilometers to 20 kilometers, 100 ton to tens of kiloton detectors can accumulate statistics sufficient to identify or exclude the existence of operating reactors, and to track basic features of their operation such as operational status, thermal power levels and fissile content. The interaction channel used is the inverse beta decay (IBD) process, in which a charged-current interaction of an antineutrino with a quasi-free proton generates a positron and a neutron, both of which can be detected in close time coincidence. While one must be mindful of potentially severe statistical and technology limitations, it may also be possible to extract an independent estimate of the range of the reactor based on a shape analysis of an acquired antineutrino spectrum, and of the direction of the reactor based on the collective displacements between reconstructed positions of the prompt positron emission and the delayed neutron capture in the IBD channel. This set of applications maps well into existing international protocols for monitoring reactors and other nuclear sites, in large part administered by the International Atomic Energy Agency.

In the far-field, from 20-200 kilometers, the small number of detectable IBD interactions in reasonable time periods restricts the applications to the exclusion or discovery of operating reactors, using detectors at the 10 kiloton scale, up to a limit of roughly megaton of target mass. Discovery and exclusion applications are relevant as confidence-building measures within the framework of existing treaties, and as more robust verification measures called for by potential future treaties such as the Fissile Material Cutoff Treaty [7] and the Treaty on the Prohibition of Nuclear Weapons [8].

IV. MID-FIELD AND FAR-FIELD TECHNOLOGY OPTIONS FOR NONPROLIFERATION OVERLAP WITH NEUTRINO PHYSICS EXPERIMENTAL NEEDS

Currently and for the foreseeable future, only liquid scintillator and water Cherenkov detectors (and possibly hybrids of these media) appear to have the potential to be practical at the hundreds of ton to hundreds of kiloton scale needed for mid-to-far-field monitoring. Kiloton-scale and larger detectors of both types have been built beginning in the 1970s, with increasing size and performance. The paradigm for reactor antineutrino detection is the kiloton-scale KamLAND liquid scintillator detector, which has operated for two decades, and has detected antineutrinos from reactors across Japan and South Korea, up to hundreds of kilometers away [9].

Attractions of scintillator detectors include high light output and consequent reduction in the number of sensors needed to capture the photons created by antineutrino interactions, and the ability to thoroughly purify the organic liquid of contaminating backgrounds. Disadvantages are the expense of the medium, in some cases the combustibility or toxicity of the organic materials comprising the scintillator, the need for transport or local manufacture of the scintillator at the deployment site, and the resultant additional complexity these imply for underground deployment and decommissioning of large-scale detectors. Moreover, since the scintillator absorbs light, it is difficult to build detectors with masses larger than a few tens of kilotons of the detector material before light absorption within the medium compromises detection. This constraining characteristic ultimately limits the effective standoff range for scintillator detectors, largely - though not entirely - relegating them to mid-field nonproliferation applications.

Detector masses ranging from tens of tons to tens of kilotons can address some mid-field and far-field applications. The largest currently planned liquid scintillator detector is JUNO [3], comprising 20 kilotons of scintillator immersed in a 35 kiloton water veto, and now being installed in an underground mine in China. JUNO, once operational, will therefore demonstrate a capability relevant for some mid-to-far-field applications.

For far-field deployments, with the detector located 20-200 kilometers away from the reactor, 10-1000 kiloton detectors are required to achieve reasonable dwell times to discover or exclude the presence of 50MWt reactors. At these scales, waterbased detectors would offer an attractive alternative to scintillator-based detectors. Water has the advantages of low cost per unit volume, ease of purification on large scales, and fewer adverse effects on health, safety, and the environment than liquid scintillator. The low cost of the raw material competes with the requirement for more high cost photosensors per unit of volume, needed to collect the smaller amount of light generated by particle interactions in water compared to scintillator.

Super-Kamiokande [10] is the largest water Cherenkov detector ever built, with a total mass of water of 50 kilotons. Hyper-Kamiokande[2] is approximately five times larger than Super-Kamiokande, and has been approved for deployment in Japan. Such a detector is on the scale required for remote exclusion and discovery of reactors at the 100 kilometer range. The SNO+ experiment[11] also anticipates collection of IBD events from distant reactors, providing further insight into sensitivity of large water-based detectors to the reactor IBD signal.

Pure water detectors provide sub-optimal spectral information about reactor antineutrinos. Two variants on water detectors show promise. First, one part per thousand quantities of the element gadolinium, a highly effective neutron capture agent, can be added to water, significantly enhancing sensitivity to the neutron generated in antineutrino interactions. This has been demonstrated in the EGADS 200-ton test-bed[12], and the SuperKamiokande experiment has recently added gadolinium in order to improve sensitivity to antineutrinos of astrophysical origin[13]. The WATCHMAN collaboration has proposed a design[14] for a Gd-H2O detector as a candidate for a demonstration of remote monitoring. Such a Water Cherenkov detector could provide the added potential benefit of direction reconstruction, via antineutrino-electron scattering. Solar Neutrino Electron Scattering has been demonstrated clearly in Super-Kamiokande, but this capability is difficult to achieve and as yet unproven for reactor antineutrinos. This is expected to be an extremely challenging measurement using a water-based target [15], due to as-yet undemonstrated radio-purity requirements.

Another promising approach is to try to combine the advantages of scintillator and water in a single medium. Water-based scintillators are water detectors in which part per hundred or greater quantities of scintillator are stably mixed, creating both scintillation and Cherenkov light and in principle combining the advantages of both materials and readout modes. Such detectors may hold promise as a mid-scale solution for reactor antineutrino detection. They have also been explored for close to a decade as one possible target material for the large-scale Theia detector for a range of fundamental physics experiments [4].

In addition to Cherenkov/scintillation light separation, it is also possible to exploit more traditional pulse-shape discrimination (PSD) methods in a suitable formulated WbLS cocktail. PSD has been used in pure liquid scintillators for decades to distinguish neutrons from gamma-rays with high reliability. Here, the fast and slow decay times of scintillation light arise from the excitation and decay of singlet and triplet states respectively. Neutrons and gamma-rays induce different triplet/singlet ratios in media due to their very different track ionization densities (DE/dx), with the result that pulse time profile differs substantially according to particle type. The use of scintillation PSD, possibly in combination with Cherenkov/scintillation separation, should allow for better particle identification for a larger range of particles. Another advantage of incorporating a 'standard' PSD capability into WbLS is that it may be possible to use slower-response light compared to methods relying on Cherenkov/scintillation discrimination. Recently, researchers at LLNL have developed and demonstrated a novel majority-water WbLS cocktail, which discriminates between gamma-rays and neutrons using scintillation-based PSD [16].

Another early-stage technology of interest is to use opaque scintillating oil (or possibly a water-scintillator combination) instrumented with solid-state photo-detectors coupled to wavelength shifting fibers immersed in the medium. This idea is being explored by the Liquid-O proto-collaboration [5]. The localized nature of the light collection may increase the fiducial-to-total mass ratio of the detector, and may offer significant reductions in the self-shielding and overburden requirements for reactor

antineutrino detection through efficient positron identification.

In sum, scintillator detectors have been demonstrated at the kiloton scale, scintillator and modified water detectors appear to hold promise up to the few tens of kiloton scale, suitable for the longer standoff distances in the mid-field, while large gadolinium-doped or scintillator-doped water designs could permit construction of the 100 kiloton and larger detectors that are required for the far-field.

V. OPPORTUNITIES FOR COLLABORATION

An excellent existing platform for exploration of the synergies described above is the Applied Antineutrino Physics workshop series[17]. Since 2004, these conferences have showcased developments in nonproliferation and in particle and nuclear physics that have the potential to influence the other domain. The conferences have been held in the U.S. and worldwide, and frequently include updates from representatives of particle physics experiments. The forum is an opportunity to learn the basic working concepts and priorities of the nuclear nonproliferation and disarmament communities, as well as providing nonproliferation experts with insight into cutting edge experimental physics projects worldwide.

Current examples of ongoing synergistic work at U.S. national laboratories and universities include prototype gadoliniumdoped water and water-based and opaque scintillator test-beds and demonstrators, extensive testing and industry partnerships related to large area fast position-sensitive photomultiplier tubes, and the development of concepts for a possible underground kiloton-scale water-based detector for reactor monitoring and technology demonstrations. The latter collaborative planning and research effort has served as a training ground for physics, nuclear engineering and policy students and postdoctoral fellows, who work side-by-side on technical problems and policy matters related to both nonproliferation and fundamental science.

NNSA also occasionally issues funding opportunities for academic collaborative efforts focused on nonproliferation and overlapping fundamental science efforts[18].

In the future, proposals that recognize and capitalize on the synergies between the nonproliferation and HEP communities could (and should) be considered for joint and combined funding —shared between the Office of High Energy Physics and NNSA. Pilot investments of this kind have been initiated in the past several years, and the horizon for large-scale neutrino and particle astrophysics experiments suggests that more collaborative opportunities may become available.

Specific areas of possible synergistic research include, but are not limited to:

- Joint technology demonstrations at the 0.1 to 10 kiloton scale;
- Further development of water-based scintillator, opaque scintillator, wavelength shifting chemicals added to water, and similar material advances;
- Development of high spatial and temporal resolution photosensors such as the Large Area Picosecond Photodetector (LAPPD)[19];
- Development of affordable instrumentation for large channel counts;
- Efficient and low-cost light collection across an increased wavelength regime extending into toward the IR, such as wavelength shifting plates;
- Multi-wavelength-discriminating sensors such as dichroicons[20], capable of manufacture and deployment on large scales.
- Development of new or improved scintillator materials, producing more fast light output as well as efficient particle identification (PSD).
- Development of new low energy detector types which permit some resolution of neutrino direction, including scaling up from ton scale to kiloton scale.
- Development of neutrino detectors capable of geoneutrino observation (<2.8 MeV), low-end reactor neutrino energies, and relic supernova neutrino studies.
- Detector development for neutrino spectral measurements extending down to the 1.8 MeV IBD threshold with few percent energy resolution enabling neutrino age measurement via spectrum modulation (neutrino ranging).
- Studies of cost effective very large IBD neutrino detectors, capable of long range reactor detection (>100 km), nucleon decay searches reaching to lifetimes in the 10³⁵ year range, and astrophysical neutrinos. Such detectors have important astrophysical observation capability (starburst galaxies and relic neutrinos), and supernova early warning capability via increased neutron appearance rate detection.
- Development of technology for large volume neutrino detectors employing replaceable optical modules without in-liquid wired connections.

- [1] R. Acciarri *et al.*, Long-Baseline Neutrino Facility (LBNF) and Deep Underground Neutrino Experiment (DUNE) Conceptual Design Report, volume 4 the DUNE detectors at LBNF, arXiv preprint arXiv:1601.02984 (2016), arXiv:1601.02984 [physics.ins-det].
- [2] K. Abe *et al.*, Letter of intent: The Hyper-Kamiokande Experiment Detector Design and Physics Potential (2011), arXiv:1109.3262 [hep-ex].
- [3] F. An et al. (JUNO), Neutrino Physics with JUNO, J. Phys. G 43, 030401 (2016).
- [4] G. D. O. G. (for the THEIA Interest Group), Physics Potential of an Advanced Scintillation Detector: Introducing THEIA (2015), arXiv:1504.08284 [physics.ins-det].
- [5] A. Cabrera et al., Neutrino physics with an opaque detector, Communications Physics 4, 10.1038/s42005-021-00763-5 (2021).
- [6] Dune Module of Opportunity Workshop, https://indico.fnal.gov/event/21535/timetable/?view=standard (2019).
- [7] Fissile Material Cut-off Treaty (FMCT) at a Glance, https://www.armscontrol.org/factsheets/fmct (2018).
- [8] The Treaty on the Prohibition of Nuclear Weapons, https://www.nti.org/education-center/treaties-and-regimes/treaty-on-the-prohibition-of-nuclear-we (2020).
- [9] A. Gando et al. (KamLAND), Reactor On-Off Antineutrino Measurement with KamLAND, Phys. Rev. D 88, 033001 (2013), arXiv:1303.4667 [hep-ex].
- [10] S. Fukuda et al., The Super-Kamiokande detector, Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Dete
- [11] L. Lebanowski and S. Andringa, Snowmass2021 Letter of Interest: Detecting Antineutrinos from Distant Reactors using Pure Water at SNO+, https://www.snowmass21.org/docs/files/summaries/NF/SNOWMASS21-NF7_NF0_SNOplus-186.pdf (2021).
- [12] L. Marti *et al.*, Evaluation of gadolinium's action on water Cherenkov detector systems with EGADS, Nucl. Instrum. Meth. A 959, 163549 (2020), arXiv:1908.11532 [physics.ins-det].
- [13] K. Abe et al., First gadolinium loading to Super-Kamiokande, Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectror
- [14] M. Askins *et al.*, The Physics and Nuclear Nonproliferation goals of WATCHMAN: a WATer CHerenkov Monitor for ANtineutrinos, arXiv preprint arXiv:1502.01132 (2015), arXiv:1502.01132 [physics.ins-det].
- [15] D. Hellfeld, A. Bernstein, S. Dazeley, and C. Marianno, Reconstructing the direction of reactor antineutrinos via electron scattering in Gd-doped water Cherenkov detectors, Nucl. Instrum. Meth. A 841, 130 (2017), arXiv:1512.00527 [physics.ins-det].
- [16] M. J. Ford, N. P. Zaitseva, M. L. Carman, S. A. Dazeley, A. Bernstein, A. Glenn, and O. A. Akindele, Pulse-shape discrimination in water-based scintillators, arXiv preprint arXiv:2202.07625 (2022), arXiv:2202.07625 [physics.ins-det].
- [17] Applied Antineutrino Physics Workshops, https://neutrinos.llnl.gov/workshops, https://indico.cern.ch/event/833568/ (2019).
- [18] https://www.energy.gov/nnsa/articles/nnsa-grants-50-million-university-consortia-supporting-nonproliferation-research-and (2019).
- [19] Adams, B. W. and Elagin, A. and Frisch, H. J. and Obaid, R. and Oberla, E. and Vostrikov, A. and Wagner, R. G. and Wang, J. and Wetstein, M., Timing characteristics of Large Area Picosecond Photodetectors, Nuclear Instruments and Methods in Physics Research. Section A, Accelerators, Spectrometers, Detectors and Associated Equipment **795**, 10.1016/j.nima.2015.05.027 (2015).
- [20] T. Kaptanoglu, M. Luo, and J. Klein, Cherenkov and Scintillation Light Separation Using Wavelength in LAB Based Liquid Scintillator, JINST 14 (05), T05001, arXiv:1811.11587 [physics.ins-det].