

---

---

Submitted to the Proceedings of the US Community Study  
on the Future of Particle Physics (Snowmass 2021)

---

---

IPPP/22/11, SLAC-PUB-17664, YITP-SB-2022-06

---

## Snowmass 2021 White Paper: Resummation for future colliders

---

**Melissa van Beekveld,<sup>a</sup> Sebastian Jaskiewicz,<sup>b</sup> Tao Liu,<sup>c,d</sup> Xiaohui Liu,<sup>e,f</sup> Duff Neill,<sup>g</sup>  
Alexander Penin,<sup>h</sup> Felix Ringer,<sup>i,j</sup> Robert Szafron,<sup>k</sup> Leonardo Vernazza,<sup>l,m</sup> Gherardo  
Vita,<sup>n</sup> Jian Wang<sup>o</sup>**

<sup>a</sup>*Rudolf Peierls Centre for Theoretical Physics, Clarendon Laboratory, Parks Road, University of Oxford, Oxford OX1 3PU, UK*

<sup>b</sup>*Institute for Particle Physics Phenomenology, Durham University, Durham DH1 3LE, United Kingdom*

<sup>c</sup>*Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049, China*

<sup>d</sup>*University of Chinese Academy of Sciences, Beijing 100049, China*

<sup>e</sup>*Center of Advanced Quantum Studies, Department of Physics, Beijing Normal University, Beijing 100875, China*

<sup>f</sup>*Center for High Energy Physics, Peking University, Beijing 100871, China*

<sup>g</sup>*Theoretical Division, MS B283, Los Alamos National Laboratory, Los Alamos, NM 87545, USA*

<sup>h</sup>*Department of Physics, University of Alberta, Edmonton AB T6G 2J1, Canada*

<sup>i</sup>*C.N. Yang Institute for Theoretical Physics, Stony Brook University, Stony Brook, NY 11794, USA*

<sup>j</sup>*Department of Physics and Astronomy, Stony Brook University, Stony Brook, NY 11794, USA*

<sup>k</sup>*Department of Physics, Brookhaven National Laboratory, Upton, N.Y., 11973, USA*

<sup>l</sup>*INFN, Sezione di Torino, Via P. Giuria 1, I-10125 Torino, Italy*

<sup>m</sup>*Nikhef, Science Park 105, NL-1098 XG Amsterdam, The Netherlands*

<sup>n</sup>*SLAC National Accelerator Laboratory, Stanford University, Stanford, CA 94039, USA*

<sup>o</sup>*School of Physics, Shandong University, Jinan, Shandong 250100, China*

*E-mail:* [melissa.vanbeekveld@physics.ox.ac.uk](mailto:melissa.vanbeekveld@physics.ox.ac.uk),  
[sebastian.jaskiewicz@durham.ac.uk](mailto:sebastian.jaskiewicz@durham.ac.uk), [liutao86@ihep.ac.cn](mailto:liutao86@ihep.ac.cn),  
[xiliu@bnu.edu.cn](mailto:xiliu@bnu.edu.cn), [duff.neill@gmail.com](mailto:duff.neill@gmail.com), [penin@ualberta.ca](mailto:penin@ualberta.ca),  
[felix.ringer@stonybrook.edu](mailto:felix.ringer@stonybrook.edu), [rszafron@bnl.gov](mailto:rszafron@bnl.gov),  
[leonardo.vernazza@to.infn.it](mailto:leonardo.vernazza@to.infn.it), [gvita@stanford.edu](mailto:gvita@stanford.edu), [j.wang@sdu.edu.cn](mailto:j.wang@sdu.edu.cn)

ABSTRACT: Resummation techniques are essential for high-precision phenomenology at current and future high-energy collider experiments. Perturbative computations of cross sections often suffer from large logarithmic corrections, which must be resummed to all orders to restore the reliability of predictions from first principles. The precise understanding of the all-order structure of field theories allows for fundamental tests of the Standard Model and new physics searches. In this white paper, we review recent progress in modern resummation techniques and outline future directions. In particular, we focus on the resummation beyond leading power, the joint resummation of different classes of logarithms relevant for jets and their substructure, small- $x$  resummation in the high-energy regime and the QCD fragmentation process in the small- $z_h$  limit.

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Resummation beyond leading power</b>	<b>3</b>
2.1	Diagrammatic methods for threshold resummation	3
2.2	Effective field theory approach for cross-sections	8
2.3	Diagrammatic methods for mass- and angle-suppressed amplitudes	11
2.4	Effective field theory approach for mass suppressed amplitudes	13
<b>3</b>	<b>Jets and their substructure</b>	<b>14</b>
<b>4</b>	<b>Small-<math>x</math> resummation</b>	<b>17</b>
<b>5</b>	<b>Fragmentation and small-<math>z_h</math> resummation</b>	<b>20</b>
<b>6</b>	<b>Conclusions</b>	<b>24</b>

---

# 1 Introduction

Tests of the Standard Model and New Physics searches at the Large Hadron Collider (LHC), Electron-Ion Collider (EIC) and other future colliders require extremely precise theoretical predictions. Correctly accounting for large quantum corrections will be indispensable for the High-Luminosity phase of the LHC, especially since the direct searches for new particles have not resulted in a discovery so far. Precise measurements of the Higgs boson and top quark properties, or tests of the Standard Model's flavor structure, rely on an excellent theoretical control over the QCD-induced corrections.

The accurate theoretical description of high-energy processes at colliders is among the most successful applications of perturbative QCD. The relevant QCD corrections are typically computed as an expansion in the strong coupling constant  $\alpha_s$ . Currently, state-of-the-art predictions include corrections up to the third order in  $\alpha_s$ , with most of the processes being known only up to the next-to-next-to-leading order (NNLO).

A typical process in QCD depends on various energy scales. When the scales are widely separated, the presence of the scale hierarchy manifests itself through logarithmically enhanced radiative corrections, which require an all-order resummation of the series in the strong coupling constant. One of the great simplifications of perturbative analysis at high energy is that the contributions suppressed by a power of a ratio of an infrared scale of the process to its center-of-mass energy are small and can be neglected in the first approximation, reducing the number of scales and hence the complexity of the calculations. Thus, the resummation techniques typically target the leading power (LP) terms in the power expansion. Before we review in more detail the recent progress in the most challenging fields, such as subleading power resummation, small- $x$  resummation, fragmentation and small- $z_h$  resummation, and the physics of jets and their substructure, let us briefly mention two of the well-established application of resummation techniques, which played a crucial role in the development of this research field.

Threshold resummation is one of the flagship applications of factorization and resummation for collider physics [1–5]. Typically, when the phase space available for radiation is limited, the corrections are dominated by soft gluons. Therefore, threshold resummation is relevant for the production of heavy states. In the context of LHC physics, the leading power threshold resummation is well understood. It has been applied to many processes, such as the production of the electroweak gauge and Higgs bosons and top quarks as well as production of new heavy particles, see e.g. Refs. [6–14].

A second important class of problems involves transverse momentum resummation. Substantial progress has been achieved in recent years, with the state-of-the-art results reaching next-to-next-to-next-to leading logarithmic ( $N^3LL$ ) accuracy combined with NNLO and even  $N^3LO$  fixed order computations, see e.g. [15–33]. These results are very important components for precision tests of the Standard Model at the LHC.

The need for a precise understanding of the all-order structure of QCD led to the development of new theoretical techniques and methods. Today the field of resummation is exploring new topics crucial for phenomenology and our understanding of gauge theories. This paper reviews the recent progress in the theoretical methods of resummation

and their relevance for collider phenomenology. We will discuss the most challenging problems currently under investigation, which are addressed with soft-collinear effective field theory (SCET) [34–37] and directly in QCD. In section 2, we discuss recent progress in understanding the all-order structure of subleading power terms using the traditional diagrammatic approach and SCET. Section 3 provides an overview of jet physics and jet substructure observables. Here the focus is in particular on the simultaneous resummation of several classes of large logarithmic corrections. This aspect is particularly relevant for jet substructure observables, where multiple energy scales appear. Next, in section 4, we discuss recent progress in understanding gluon saturation, which is relevant in the high-energy regime where the initial-state gluon momentum fraction is small. Section 5 focuses on the QCD fragmentation process in the limit where the energy fraction of the identified hadrons becomes small relative to the hard scale of the process. Finally, we conclude in section 6.

## 2 Resummation beyond leading power

In the leading-power approximation the structure of the logarithmic corrections is in general well understood and the relevant resummation techniques are well elaborated. However, in many crucial cases the continually improving accuracy of the experimental measurements requires the inclusion of power suppressed terms in the theoretical estimates. As a result, much effort is currently invested into the study of a diverse class of power corrections, see [38–58] and references therein. Incorporating logarithmically enhanced power-suppressed terms can significantly increase the accuracy and extend the region where the perturbative calculation is applicable. Besides the phenomenological importance, the power-suppressed contributions are very interesting from the general effective field theory point of view since the structure of the factorization and renormalization group evolution in this case becomes highly nontrivial already in the leading logarithmic (LL) approximation.

### 2.1 Diagrammatic methods for threshold resummation

**Large logarithms near threshold.** Power corrections play an important role for precise predictions of physical observables involving heavy final states. Because of a phenomenon of dynamical enhancement, these scattering processes receive a large contribution near their partonic threshold, where the dimensionless partonic cross takes the following schematic form

$$\Delta(\xi) = \sum_{n=0}^{\infty} \alpha_s^n \left\{ \sum_{m=0}^{2n-1} c_{nm}^{\text{LP}} \left( \frac{\ln^m \xi}{\xi} \right)_+ + d_n \delta(\xi) + \sum_{m=0}^{2n-1} c_{nm}^{\text{NLP}} \ln^m \xi + \dots \right\}. \quad (2.1)$$

In this equation  $\xi$  represents a variable, which approaches  $\xi \rightarrow 0$  near threshold; for instance,  $\xi = 1 - z \equiv 1 - Q^2/\hat{s}$  for the Drell-Yan process, where  $Q^2$  is the invariant mass of the final state, and  $\hat{s}$  the partonic center of mass energy squared. For  $\xi \rightarrow 0$  the partonic cross section in Eq. (2.1) develops large logarithms that need to be resummed to obtain precise predictions. The terms proportional to  $c_{nm}^{\text{LP}}$  contribute at leading power, and their resummation has been known for a long time to a high logarithmic accuracy,

since the seminal papers [1–5]. Later, LP threshold resummation has been reinterpreted and clarified using a wide variety of methods, including the use of Wilson lines [59, 60], the renormalization group [61], the connection to factorization theorems [62], and soft collinear effective theory [63–65]. The state-of-the-art for resummation at LP is next-to-next-to-next-to-leading logarithmic (N<sup>3</sup>LL) accuracy for color singlet final states, and next-to-next-to-leading logarithmic (NNLL) accuracy for processes involving colored particles in the final state.

The resummation of logarithms with  $c_{nm}^{\text{NLP}}$  coefficients, contributing at next-to-leading power (NLP), has been considered only more recently and it is still subject of intense study. Progress has been made by employing diagrammatic [51, 66–68], effective field theory [45, 48, 53, 69, 70] and renormalization group methods [71, 72]. In this section, we provide a brief overview of the diagrammatic approach, while the application of SCET to the resummation of large logarithms at NLP will be discussed in the following section.

**Resummation at NLP for diagonal partonic channels.** Near threshold, additional radiation is constrained to be soft and the partonic cross section develops a hierarchy of scales, with  $Q$  representing the energy involved in the hard interaction, and  $Q(1-z)$  with  $z \ll 1$  the energy of the soft radiation. It can be shown that soft radiation at LP is described in terms of uncorrelated eikonal emissions. As a consequence, the partonic cross section factorizes into a soft and a hard function, the latter representing the central hard scattering, while the former describes soft radiation in terms of a vacuum expectation value of Wilson lines along the directions of the hard scattering particles. The exponentiation of eikonal radiation together with the factorization of the associated phase space (in Mellin space) is the basis for resummation of the logarithms proportional to the  $c_{nm}^{\text{LP}}$  coefficients in Eq. (2.1).

Beyond LP, the factorization of the partonic cross section becomes more involved. In general, this happens because soft radiation becomes sensitive to the nature of the hard scattering particles and their hard interaction. For instance, soft gluons can be emitted through chromo-magnetic interactions, which are sensitive to the spin of the emitting particles; furthermore, at NLP one needs to take into account the emission of soft quarks, too. In this section we will focus on the emission of soft gluon radiation from collinear legs, which contributes at LP and NLP to the large logarithms in diagonal production channels, such as  $q\bar{q}$  in Drell-Yan, or  $gg$  in Higgs production. The emission of soft quarks from collinear legs contributes instead to the so-called off-diagonal partonic channels, such as the  $qg$  channel in Drell-Yan. We will briefly discuss these channels in the next subsection.

Besides being sensitive to the spin of the emitting particles, soft gluon radiation at NLP begins to reveal the structure of the hard scattering [73, 74], as well as the structure of virtual collinear radiation associated to the hard scattering particles of the process [75]. These effects can be described in terms of convolutions of soft and collinear matrix elements multiplying a hard function, which represents the central non-radiative scattering process. In a diagrammatic approach, the soft and collinear matrix elements are described in terms of QCD fields and Wilson lines. For instance, in the case of Drell-Yan, the  $q\bar{q}$  amplitude

with an emission of one additional soft gluon is written (up to NNLO) as [76, 77]

$$\mathcal{A}_{\mu,a}(p_j, k) = \sum_{i=1}^2 \left\{ \frac{1}{2} \tilde{\mathcal{S}}_{\mu,a}(p_j, k) + g \mathbf{T}_{i,a} G_{i,\mu}^\nu \frac{\partial}{\partial p_i^\nu} + [J_{\mu,a}(p_i, n_i, k) - \mathcal{J}_{\mu,a}(p_i, n_i, k)] \right\} \mathcal{A}(p_j), \quad (2.2)$$

where  $p_j$ ,  $j = 1, 2$  represent the momenta of the initial state partons, and

$$G_i^{\mu\nu} = g^{\mu\nu} - \frac{(2p_i - k)^\nu}{2p_i \cdot k - k^2} k^\mu. \quad (2.3)$$

In Eq. (2.2),  $\tilde{\mathcal{S}}_{\mu,a}(p_j, k)$  represents a generalized soft function, which accounts for soft emissions from the external particles beyond the eikonal approximation [78, 79]; the second term is given by the orbital angular momentum operator acting on the non-radiative amplitude  $\mathcal{A}(p_j)$ , and represents the so-called Low-Burnett-Kroll (LBK) theorem [73, 74]. In the last term,  $J_{\mu,a}(p_i, n_i, k)$  represents a radiative jet function, introduced first in [75], while  $\mathcal{J}_{\mu,a}(p_i, n_i, k)$  subtracts overlapping soft modes between the radiative and the soft function.

Starting from Eq. (2.2) it is possible to derive the corresponding factorized cross section, which has been explicitly verified up to NNLO in perturbation theory [76, 77]. The factorization theorem in Eq. (2.2) is valid up to this perturbative order, which means that it is not enough to develop a theory of resummation at NLP, at arbitrary logarithmic accuracy. While studies of the all-order factorization structure are in development (see e.g. [80–82] for recent progress), it has already been possible to exploit the result of Eq. (2.2) to obtain the resummation of NLP large logarithms at leading logarithmic accuracy. Indeed, it is possible to show that the second and third terms in Eq. (2.2) contribute only starting at next-to-leading logarithmic (NLL) accuracy, i.e., leading logarithms are generated only by momenta configurations which are maximally soft and collinear (see [51] for a detailed discussion), which are taken into account by the generalized soft function, i.e. the first term in Eq. (2.2). As a consequence, at LL accuracy the factorization theorem simplifies. In Mellin space, the bare partonic cross section takes the form


$$\widehat{\Delta}_{\text{NLP,LL}}^{(q\bar{q})}(N, Q^2, \epsilon) = |\mathcal{H}(Q^2)|^2 \mathcal{S}(N, Q^2, \epsilon). \quad (2.4)$$

By using tools such as the replica trick [83], the generalized soft function  $\mathcal{S}(N, Q^2, \epsilon)$  can be shown to exponentiate, [79], leading to [51]

$$\int_0^1 dz z^{N-1} \Delta_{\text{NLP,LL}}^{(q\bar{q})}(z) = \Delta_{\text{LO}}^{(q\bar{q})}(Q^2) \exp \left[ \frac{2\alpha_s C_F}{\pi} \left( \log^2 N + \frac{\log N}{N} \right) \right], \quad (2.5)$$

where the exponent of the one-loop soft function resums large leading logarithms, with the second term corresponding to the NLP correction.

It can be proven that the structure of the NLP correction in Eq. (2.5) is universal for all electroweak annihilation processes involving a colorless final state [51, 84]. The LL resummed cross section, up to NLP, takes the form of Eq. (2.5), with the LO partonic cross section multiplying a universal factor, whose color structure ( $C_F$  in Eq. (2.5)) depends on



**Figure 1.** Ratio plot of the total Higgs cross section (left) and the DY invariant mass distribution (right), normalized to the NNLL' + NNLO result. The NNLL' (+NLP LL) result is again shown by the orange solid (red dash-dotted) lines. The NNLL' NLP LL result is obtained by adding the N<sup>3</sup>LO NLP LL  $qg$  result, which is shown by the dotted light-blue line. Plots taken from [67].


the initial state particles. In general, in  $z$  space this leads to the resummation formula

$$\Delta_{aa}^{\text{dQCD, LP+NLP}}(N, Q^2/\mu^2) = g_0(\alpha_s) \exp \left\{ \int_0^1 dz z^{N-1} \left[ \frac{1}{1-z} D_{aa} \left( \alpha_s \left( \frac{(1-z)^2 Q^2}{z} \right) \right) + 2 \int_{\mu^2}^{(1-z)^2 Q^2/z} \frac{dk_T^2}{k_T^2} P_{aa}^{\text{LP+NLP}}(z, \alpha_s(k_T^2)) \right] \right\}, \quad (2.6)$$

where  $g_0(\alpha_s)$  collects the  $N$ -independent contributions. Large logarithms are resummed in the exponent, which contains the diagonal DGLAP splitting function  $P_{aa}(z, \alpha_s)$  and soft wide-angle contributions, collected in  $D_{aa}(\alpha_s)$ , where  $aa = q\bar{q}$  for quark-antiquark initiated processes, and  $aa = gg$  for gluon-gluon initiated processes. Evaluating these functions to a given order in  $\alpha_s$  resums the large logarithms to a corresponding logarithmic accuracy. To date,  $P_{aa}(z, \alpha_s)$  is known to fourth order in  $\alpha_s$  and  $D_{aa}(\alpha_s)$  to third order, which guarantees the resummation (at LP) up to N<sup>3</sup>LL accuracy. Both terms are process-independent, to the extent that they only depend on the color structure of the underlying hard-scattering process. The NLP correction in Eq. (2.5) is given by expanding  $P_{aa}(z, \alpha_s)$  up to NLP in the threshold variable. This ensures that all LLs are resummed at NLP. In principle, Eq. (2.6) resums also NLP logarithms beyond LL accuracy. These originates from keeping higher order contributions in  $P_{aa}(z, \alpha_s)$  at NLP, as well as from the argument of  $\alpha_s$  in the function  $D_{aa}$  and from the  $1/z$ -dependence of the upper limit of the  $k_T$  integral. However, these contributions give only part of the whole NLL contribution and beyond, and can be dropped for a consistent resummation at fixed logarithmic accuracy.

An extensive phenomenological analysis investigating the effects of resumming threshold LLs at NLP in physical observables of interest for the LHC has been presented in [67]. There, it has been shown that, in general, the NLP LL correction becomes competitive with





**Figure 2.** Left: ladder diagram contributing to the  $g\bar{q}$  channel in DY production; right: similar but for the  $qg$  channel in Higgs production.

the resummation of LP logarithm at NNLL accuracy, as can be seen for instance in Fig. 1. Therefore, any precise prediction in which LP resummation is performed at NNLL accuracy and beyond should include resummation of NLP logarithms. The analysis of [67] has also shown that the diagrammatic and the effective field theory (SCET) approach [48, 53], give almost identical results, when the same contributions at NLP are taken into account, thus removing earlier claims of numerical differences between the two approaches.

**Resummation at NLP for subleading partonic channels.** The resummation of large threshold logarithms in off-diagonal channels, such as the  $qg$  channel in Drell-Yan, are particularly interesting. These channels start contributing already at NLP, because near threshold they involve the emission of a soft quark, which is itself power-suppressed with respect to a soft gluon emission. Earlier conjectures for Deep Inelastic Scattering (DIS) and Drell-Yan [85, 86] showed that the resummation of large threshold logarithms in off-diagonal channels has a non-trivial exponentiation pattern, even at LL accuracy, which involves the color structure  $C_F - C_A$ . The reason for this non-trivial behavior can be understood, for instance, by investigating the factorization structure of the Drell-Yan  $qg$  channel by means of the method of regions, which shows that both virtual collinear and soft modes contribute at LL accuracy. Therefore, a simplification such as in Eq. (2.4) is not possible for off-diagonal channels. Despite this complication, it has been possible to derive an all-order resummation formula for NLP LLs in off diagonal channels of processes such as DIS, Drell-Yan and Higgs production [68, 70, 87]. To this end, a fundamental input is provided by deriving so-called consistency conditions, obtained by requiring the cancellation of leading poles in the total cross section. One parameterizes the leading poles in the total cross section in terms of unknown coefficients multiplying the relevant scales

of the process. Requiring pole cancellation in the bare total cross section provides a set of equations, whose solution determines the minimal number of parameters needed to fix all the poles – and thus the related large logarithms. The derivation is rather technical, and we refer to [68, 70] for further details. The relevant observation is that a minimal set of parameters is given entirely by those associated to a single scale in the problem – one parameter for each perturbative order. It is thus possible to obtain the entire cross section by determining the full tower of coefficients associated, for instance, to the hard scale, or to the soft scale in the problem. Determining the all-order coefficients associated to the hard scale can be done by means of effective field theory methods, and we refer to [70] for further details. The diagrammatic approach, instead, is particularly suitable to determine at all orders the coefficients associated to the leading poles in the soft region – in a suitable gauge, it can be shown that the soft region is determined by ladder diagrams such as those represented in Fig. 2 [68]. These can be calculated to all orders, and subsequently the consistency condition allow one to fix the entire cross section. For instance, in the case of the  $qg$  channel in Drell-Yan, the resummed NLP LL partonic cross section takes the form [68, 87]

$$\Delta_{qg}^{\text{dQCD, LP+NLP}} = \frac{T_R}{C_A - C_F} \frac{1}{2N \ln N} \left\{ e^{\frac{2C_F \alpha_s}{\pi} \ln^2 N} \mathcal{B}_0 \left[ \frac{\alpha_s}{\pi} (C_A - C_F) \ln^2 N \right] - e^{(2C_F + 6C_A) \frac{\alpha_s}{4\pi} \ln^2 N} \right\}, \quad (2.7)$$

where

$$\mathcal{B}_0(x) = \sum_{n=0}^{\infty} \frac{B_n}{(n!)^2} x^n, \quad (2.8)$$

and in turn  $B_n$  are Bernoulli numbers,  $B_0 = 1$ ,  $B_1 = -1/2$ ,  $\dots$ . Combining together Eq. (2.6) with Eq. (2.7) completes the resummation of LLs at NLP in all the production channels contributing to Drell-Yan. Analogous equations can be obtained for Higgs boson production in gluon fusion and DIS, and we refer to [68, 70] for a detailed derivation.

## 2.2 Effective field theory approach for cross-sections

The effective field theory approach can achieve a proper, systematic treatment of factorization and resummation beyond the leading power. SCET provides a framework to describe processes involving energetic and soft particles to all powers. It allows for an operator-based expansion, maintaining homogeneous power-counting and gauge invariance at every step of the computations. The EFT formulation provides further insights into the structure of well-established soft theorems [88–90], which are the basis for a systematic study of power suppressed effects. The EFT approach enables the study of subleading power terms in soft and collinear expansion not only in QCD but also in Gravity [91] and it would be interesting to explore further applications.

The SCET expansion parameter is typically denoted by  $\lambda$ . Its precise definition depends on the process under consideration, e.g.,  $\lambda \sim \sqrt{\xi}$  as in Eq. (2.1) above. The theory contains collinear and soft modes. Each collinear direction has its collinear mode, i.e., a generic  $N$ -jet process will be described by  $N$  collinear modes. Interactions within a given

collinear sector and interactions with the soft background are contained in the Lagrangian. The complete Lagrangian can then be written as a sum over collinear directions

$$\mathcal{L}_{\text{SCET}} = \sum_{i=1}^N \mathcal{L}_i + \mathcal{L}_s, \quad (2.9)$$

where  $\mathcal{L}_i$  is the  $i$ -collinear Lagrangian and  $\mathcal{L}_s$  is a pure soft Lagrangian. In principle, one also has to consider the Glauber Lagrangian [92]. However, for many observables, the Glauber contribution cancels and thus this term is often ignored. The  $i$ -collinear Lagrangian can then be written as a series in  $\lambda$

$$\mathcal{L}_i = \mathcal{L}_i^{(0)} + \mathcal{L}_i^{(1)} + \mathcal{L}_i^{(2)} + \mathcal{O}(\lambda^3). \quad (2.10)$$

Interactions between different collinear sectors are obtained after integrating out the hard modes. They are encoded in the currents, which are constructed from the collinear gauge-invariant building blocks [34, 93]. At leading power, we can only use a single building block


$$J_i^{A0}(t_i) \in \{\chi_i(t_i n_{i+}), \bar{\chi}_i(t_i n_{i+}), \mathcal{A}_{i\perp}(t_i n_{i+})\}, \quad (2.11)$$

for each collinear direction. Here  $\chi_i(x) = W_i^\dagger(x) \xi_i(x)$  is a fermionic building block, constructed out of the  $i$ -collinear field  $\xi_i(x)$  and the collinear Wilson line  $W_i$ . Similarly, the gluon building block is related to the collinear gluon field. Since the building blocks scale as  $\lambda$ , adding more  $i$ -collinear fields in one direction creates power-suppressed operators. Alternatively, power suppression can be introduced by the action of a derivative on the collinear fields. Details on the operatorial basis construction and renormalization of the subleading operators for various cases can be found in [89, 94–102].

While the study of power corrections in the context of flavor physics has a long history [103–110], the first resummation using SCET for event shape-type observables was achieved relatively recently for thrust distribution [45]. Subsequently, threshold resummation has been studied for Drell-Yan [48] and Higgs production [53]. Each of these studies considers only diagonal channels, i.e., not involving soft quarks and achieved leading logarithmic accuracy. The LL results have a relatively simple structure for all these processes. For example, the Drell-Yan threshold resummation leads to the following expression

$$\begin{aligned} \Delta_{\text{NLP}}^{\text{LL}}(z, \mu) &= \frac{\hat{\sigma}_{\text{NLP}}^{\text{LL}}(z, \mu)}{z} = \exp \left[ -2 \frac{\alpha_s C_F}{\pi} \ln^2 \frac{\mu}{\mu_h} \right] \exp \left[ +2 \frac{\alpha_s C_F}{\pi} \ln^2 \frac{\mu}{\mu_s} \right] \\ &\quad \times (-4) \frac{\alpha_s C_F}{\pi} \ln \frac{\mu_s}{\mu} \theta(1-z). \end{aligned} \quad (2.12)$$

One can recognize the subleading power logarithm appearing at NLO in the second line, dressed by Sudakov exponents originating from the hard and soft functions. Interestingly, even these simple results required the development of new techniques allowing to perform the resummation. These are related to the observation that the leading logarithms appear due to operator mixing under renormalization [45, 48].



**Figure 3.** Soft scale dependence of the NNLL LP and LL NLP resummed Higgs production cross section. Left panel:  $\mu_h^2 = m_H^2$ , right panel:  $\mu_h^2 = -m_H^2$ . Letters *A* and *B* refer to different possible choice of initial conditions, see [53].

The result for the Higgs threshold production [53] can be obtained from the Drell-Yan result (2.12) after substituting color factors  $C_F \leftrightarrow C_A$ . Phenomenological study for 13TeV proton-proton collisions shows that the LL NLP corrections can exceed 30% of the NNLL LP resummed cross section. However, as shown in Fig. 3, the dependence on the soft scale is substantial and indicates a need for extension for the NLP resummation beyond LL accuracy, and possibly even further terms in the power expansion [111].

While the general procedure of deriving subleading power factorization theorems is well understood [54, 102], the resummation beyond leading logarithmic accuracy for diagonal channels is still an open problem. The so-called endpoint divergent problem hinders the extension of the above results beyond LL accuracy or to the off-diagonal contributions involving soft quarks. The factorization theorems at NLP generically are a sum over several terms, where each term is a convolution between the hard, collinear, and soft functions. While these convolution integrals are well defined for the bare factorization theorem in  $d$ -dimensions, the limit  $d \rightarrow 4$  does not exist as was explicitly demonstrated in [54, 112]. This problem originates, for example, due to the overlap between soft and collinear modes. In [101], it has been observed that for certain cases protected by reparameterization invariance, one can redefine the operator basis to remove the endpoint divergences. In [69] the result for the Sudakov factor involving soft quarks has been conjectured and it has been shown that endpoint divergences are absent for  $\mathcal{N} = 1$  QCD, i.e. when the color charge of quark and gluon are equal. The conjectured soft quark Sudakov factor has been subsequently proven and generalized in [70]. This has been possible thanks to the development of the additional endpoint factorization of the divergent contributions and the introduction of new auxiliary modes, which parameterize the overlap between soft and collinear contributions. Using this result, the short-distance coefficient in the  $\overline{\text{MS}}$  scheme in off-diagonal Higgs induced DIS can be obtained in closed form, yielding the following result

$$\begin{aligned} \tilde{C}_{\phi,q}^{NLP,LL} \Big|_{\epsilon \rightarrow 0} &= \frac{1}{2N \ln N} \frac{C_F}{C_F - C_A} \left( \mathcal{B}_0(a) \exp \left[ C_A \frac{\alpha_s}{\pi} \left( \frac{1}{2} \ln^2 N + \ln N \ln \frac{\mu^2}{Q^2} \right) \right] \right. \\ &\quad \left. - \exp \left[ \frac{\alpha_s C_F}{\pi} \left( \frac{1}{2} \ln^2 N + \ln N \ln \frac{\mu^2}{Q^2} \right) \right] \right), \end{aligned} \quad (2.13)$$

where  $a = \frac{\alpha_s}{\pi} (C_F - C_A) \ln^2 N$  and  $\mathcal{B}_0(x)$  is defined in (2.8). The above result agrees with

Eq. (29) of [85] for  $\mu = Q$ , and generalizes it to  $\mu \neq Q$ .

Dealing with the endpoint divergences is currently at the forefront of EFT studies. Once a systematic treatment is established, we can expect the accuracy of resummation to be improved. Not much is known about the properties of the subleading soft and jet functions. Further progress will require a better understanding of the renormalization properties of these complicated non-local objects. Beyond the applications of SCET<sub>I</sub> that are discussed here, there are new interesting areas of study in SCET<sub>II</sub> involving subleading power rapidity divergences for transverse momentum distributions [50], Energy-Energy correlators [113], Regge kinematics and forward scattering [114, 115], and azimuthal asymmetries in semi-inclusive deep inelastic scattering [116].

Progress on the application of the EFT methods to power expansion for collider observables will be crucial for achieving the precision required by the High-Luminosity LHC, EIC and future colliders. Applications of a systematic power expansion and resummation with the help of the renormalization group equations will enable us to achieve better precision and has a potential to impact the development of subtraction methods for fixed order computations [43, 46, 50, 117–121].

### 2.3 Diagrammatic methods for mass- and angle-suppressed amplitudes

Two of the most phenomenologically and conceptually interesting examples of power corrections are given by the mass-suppressed amplitudes in processes with massive fermions in the fixed-angle *Sudakov* limit and the amplitudes suppressed by the ratio of a characteristic momentum transfer to the total energy in the small-angle or *Regge* limit.

In processes with massive fermions, the origin of the logarithmic corrections and the asymptotic behavior of the amplitudes at next-to-leading power (NLP) drastically differ from the classical leading-power Sudakov result [122]. The double-logarithmic terms in this case are related to the effect of the eikonal color charge nonconservation in the process with soft fermion exchange and result in asymptotic exponential enhancement for a wide class of amplitudes and a breakdown of a formal power counting [38, 42].

To illustrate the general properties of such processes, we consider the amplitude of light quark mediated Higgs boson production in gluon fusion which is suppressed by the quark to Higgs boson mass ratio  $m_q^2/m_H^2$  relative to the leading contribution of the top quark loop. With the targeted precision of the QCD prediction for the cross section in a percent range [123] the contributions of bottom quark loop cannot be ignored and the problem of its theoretical description ultimately goes beyond the finite order perturbative calculation [124, 125]. Indeed, the radiative corrections in this case are enhanced by the second power of the large logarithm  $\ln(m_H/m_q)$  with the effective expansion parameter being  $\alpha_s \ln^2(m_H/m_b) \approx 40\alpha_s$ . In the leading (double) logarithmic approximation, the light quark mediated  $gg \rightarrow H$  amplitude reads [42, 47]

$$\mathcal{M}_{gg \rightarrow H}^{qLL} = -\frac{3}{2} Z_g^2 g(z) \frac{m_q^2}{m_H^2} \ln^2 \left( \frac{m_H^2}{m_q^2} \right) \mathcal{M}_{gg \rightarrow H}^{t(0)}, \quad (2.14)$$

where  $\mathcal{M}_{gg \rightarrow H}^{t(0)}$  is the leading order heavy top quark mediated amplitude. Eq. (2.14) reveals a number of characteristic features of mass-suppressed processes. It includes the standard

infrared divergent Sudakov factor  $Z_g^2$  for the external on-shell gluon states and the function  $g(z)$  of the variable  $z = (C_A - C_F) \frac{\alpha_s}{4\pi} \ln^2(m_H^2/m_q^2)$  which accounts for the non-Sudakov mass logarithms to all orders in  $\alpha_s$ . It can be solved in terms of the generalized hypergeometric function  $g(z) = {}_2F_2(1, 1; 3/2, 2; z/2)$  and has the following asymptotic behavior at  $z \rightarrow \infty$

$$g(z) \sim \left( \frac{2\pi e^z}{z^3} \right)^{1/2}, \quad (2.15)$$

i.e. in contrast to the Sudakov factor it results in the exponential enhancement of the amplitude in the high-energy (small quark mass) limit. The difference  $C_A - C_F$  of the quadratic Casimir operators of the gluon and quark color group representations in the definition of the variable  $z$  accounts for the color charge variation of the highly energetic parton after emission of a soft quark, which constitutes the physical origin of the non-Sudakov double logarithms.

Though the leading logarithmic result determines the qualitative behavior of the amplitude, it cannot be used for a reliable quantitative estimates since it does not account for the numerically important effects such as the renormalization scale dependence of the quark mass and coupling constants, which requires the inclusion of subleading logarithms. In Ref. [55], the next-to-leading logarithmic corrections to Eq. (2.14) of the form  $\alpha_s^n \ln^{2n-1}(m_H/m_q)$  for all  $n$  have been computed and applied to the analysis of the Higgs boson production in threshold approximation. For the yet unknown next-to-next-to-leading and next-to-next-to-next-to-leading bottom quark contribution to the total cross section this analysis gives  $-0.12 \text{ pb}$  and  $-0.02 \text{ pb}$ , respectively. With a rather conservative assessment of the accuracy of the next-to-leading logarithmic and the threshold approximations it provides a rough estimate of the bottom quark mediated contribution to the total cross section of Higgs boson production in gluon fusion beyond NLO to be in the range from  $-0.32$  to  $0.08 \text{ pb}$ , thereby reducing the previous uncertainty estimate [123] by a factor of two.

The actual accuracy of the logarithmic and threshold approximations, however, is difficult to estimate, and the above interval has to be further reduced by evaluating the next-to-next-to-leading logarithmic contribution and getting an approximation valid beyond the threshold region. The latter requires the analysis of the logarithmically enhanced corrections to the hard real emission which is currently not available even in the leading logarithmic approximation (only the abelian part of the double-logarithmic corrections for the  $gg \rightarrow Hg$  amplitude of Higgs plus jet production has been obtained in Ref. [39]). The extension of the method [42, 47, 55] beyond the next-to-leading logarithms and to processes with hard real radiation is one of big challenges for the modern effective field theory and is of primary phenomenological importance for the high-precision Higgs physics program at the LHC. At the same time the high-order next-to-next-to-leading power terms are well under control [58].

The Regge limit of high-energy scattering describes a kinematical configuration with vanishing ratio of the characteristic momentum transfer to the total energy of the process. The relevant expansion parameter is given by the ratio of the Mandelstam variables  $t/s$ . Despite a crucial simplification due to decoupling of the light-cone and transversal degrees

of freedom, the gauge interactions in this limit possess highly nontrivial dynamics giving a rigorous quantum field theory realization of the Regge concept for high-energy scattering. Major progress has been achieved in the analysis of the leading-power amplitudes which culminated in the evaluation of the next-to-leading QCD corrections to the theory of BFKL pomeron [126].

The expansion in  $t/s$  is also mandatory for the evaluation of the nonfactorizable QCD corrections to the Higgs boson production via vector boson fusion. The leading order of this expansion is equivalent to the eikonal approximation, which reduces the problem to the effective theory analysis in the two-dimensional transversal space. This analysis has been performed at next-to-next-to-leading order in QCD for the cross sections of a single Higgs boson [127] and Higgs boson pair production [128].

At the same time, very little is known about the structure of the logarithmic corrections and the asymptotic behavior of the power suppressed amplitudes which are crucial to control the accuracy and validity of the analysis based on the eikonal approximation. Only recently the leading logarithmic result for the power-suppressed part of the fermion scattering amplitude has been obtained in an abelian gauge theory within an effective field theory framework [52]. In contrast to the single-logarithmic Regge behavior of the leading-power amplitude, the radiative corrections to the power-suppressed term exhibit a double-logarithmic enhancement and the result for the amplitude has the following form

$$\mathcal{M}^{LL} = |t/s| z g(2z) \mathcal{M}^{\text{Born}}, \quad (2.16)$$

where  $z = \frac{\alpha}{2\pi} \ln(|t/s|)$  and  $g(z)$  is, amazingly, the same function as in Eq. (2.14) describing the non-Sudakov corrections to the mass suppressed amplitude. Due to the exponential growth of this function at large values of its argument, at  $|s/t| \approx e^{2\pi/\alpha}$ , the formally power-suppressed contribution to the scattering cross section becomes comparable to the leading-power result and the small-angle expansion breaks down. This, in particular, may be considered as a possible solution of the long standing unitarity problem of the Regge analysis of the leading-power scattering amplitudes in QED [129]. The generalization of the above result to the nonabelian gauge groups and more complex amplitudes is therefore of primary theoretical interest.

## 2.4 Effective field theory approach for mass suppressed amplitudes

In parallel to the direct QCD approach, the mass suppressed effects on the amplitude level have been studied in the SCET<sub>II</sub> framework. It has been shown that these effects give rise to unexpectedly large QED corrections for the leptonic decays of  $B_s$  mesons [130]. They have been resummed, in part, in [131]. The resummed corrections were due to a contribution which is not endpoint divergent thanks to the stronger suppression of the endpoint region at the hard scale.

The breakthrough has been achieved in the study of the  $h \rightarrow \gamma\gamma$  amplitude mediated by the light quark. The condition on cancellation of the rapidity divergences has been used to derive refactorization conditions [132] and perform the LL resummation. The renormalization properties of the NLP jet and soft functions have been subsequently studied

in [133–136]. These objects have a close connection to functions appearing earlier in the flavor applications of SCET. Generally, they are simpler than the corresponding objects defined at the level of the cross section in SCET<sub>1</sub>, as their dependence on the convolution variables can be traced back to a particular external state which defines the amplitude.

The refactorization conditions were then rigorously derived in [56] as operatorial identities. This allowed proving that the reshuffling of the factorization theorem removes end-point divergences. Using the previously computed anomalous dimensions of soft and jet functions led to the first complete NLL resummation [137] for mass suppressed effects.

Resummation of power suppressed amplitudes will have a substantial impact on the precision in studies of the top and Higgs sectors at the LHC. Applications of the EFT methods will impact also flavor studies and in particular the evaluation of the QED corrections, which are a necessary ingredient for the resolution of flavor anomalies.

### 3 Jets and their substructure

Jets are collimated sprays of particles which are observed in the detectors of high-energy scattering experiments such as the LHC, RHIC, LEP, HERA and the future EIC. They reflect the underlying quark and gluon degrees of freedom which emit radiation approximately collinear to their initial direction. In recent years, significant progress has been made in improving the perturbative precision of perturbative QCD calculations. For example, fixed order calculations for inclusive jet production are now available at NNLO [138, 139]. In addition, logarithmic corrections at threshold and in the jet radius have been resummed at NLL' accuracy [140–142]. In the future, it will be critical to extend these calculations to N<sup>3</sup>LO and next-to-next-to leading logarithmic (NNLL) accuracy in order to match the experimental precision. In particular, the field of jet substructure has received a growing attention over the last decade [143–145]. Here the goal is to characterize and utilize the radiation pattern inside jets. Jet substructure observables are relevant for searches of BSM physics and various precision measurements of Standard Model processes such as Transverse Momentum Dependent (TMD) PDFs and fragmentation functions [146–151], the strong coupling constant  $\alpha_s$  [152, 153], quarkonia [154, 155], and the top quark mass [156] and to extract medium properties in heavy-ion collisions. The need for precision calculations will increase during the high-luminosity era of the LHC, where the detectors will be optimized for the first time to perform precision jet substructure measurements. In addition, the EIC is expected to perform precise measurements of low-energy jets. The clean environment of the EIC will allow for novel measurements complimentary to the LHC. In addition, RHIC and LEP data are analyzed to measure jet energy spectra and jet substructure observables. Jet grooming techniques [157–161] have been developed to systematically remove soft radiation from identified jets which can be difficult to account for from first principles in QCD. These techniques can reduce nonperturbative effects and allow for direct and precise comparisons between theoretical calculations and data. Jet substructure observables at collider experiments typically involve the measurement of multi-differential cross sections. This necessitates the joint resummation of several large logarithmic corrections to all orders in the QCD strong coupling constant, which is the main theoretical challenge. Examples




include logarithms in the jet mass, jet radius and grooming parameters. These different large logarithmic corrections are generally not independent and require a careful identification of relevant energy scales and corresponding factorization theorems. Depending on the relative scaling of the involved variables, different factorization theorems need to be derived. Eventually, the different results need to be merged to allow for a meaningful comparison to experimental data. Moreover, observables like the groomed momentum sharing fraction  $z_g$  are Sudakov safe [162] and require, in addition, the joint resummation of an auxiliary variable like the groomed jet radius  $\theta_g$ . To illustrate the complexity of factorization theorems for jet substructure observables, we consider the momentum sharing fraction  $z_g$  [162]. It is of considerable phenomenological interest as it allows for the most direct measurement of the QCD splitting function. For inclusive jet production, jet functions  $\mathcal{G}_{i=q,g}$  that depend on the jet substructure measurement can be factored out, which are then multiplied by appropriate quark/gluon fractions. In order to resum all relevant large logarithmic corrections, the jet functions need to be refactorized as [163]

$$\begin{aligned} \mathcal{G}_i &= \Theta(1/2 > z_g > z_{\text{cut}}\theta_g^\beta) \tilde{H}_i(p_T R, \mu) C_i^{\text{Egr}}(\theta_g p_T R, \mu) S_i^{\text{Egr}}(z_{\text{cut}} p_T R, \beta, \mu) \\ &\tilde{\mathcal{S}}_G(z_{\text{cut}}\theta_g^{1+\beta} p_T R, \beta, \mu) S_i^{\text{NG}}(z_{\text{cut}}) \left[ \frac{d}{dz_g} \frac{d}{d\theta_g} \tilde{\mathcal{S}}_{z_g}(z_g \theta_g p_T R, \mu) \right. \\ &\left. + \tilde{\mathcal{S}}_{i,1}^{\text{NG}}(z_g \theta_g, z_g) + \tilde{\mathcal{S}}_{i,2}^{\text{NG}}\left(z_g \theta_g, \frac{z_g}{z_{\text{cut}}\theta_g^\beta}\right) \right]. \end{aligned} \quad (3.1)$$

Here,  $z_{\text{cut}}, \beta$  denote parameters of the soft drop grooming procedure and the functions  $\tilde{\mathcal{S}}_{i,j}^{\text{NG}}$  account for non-global contributions. The factorization in Eq. (3.1) enables the joint resummation of four classes of large logarithmic corrections in the jet radius  $R$ , the grooming parameter  $z_{\text{cut}}$  and the variables  $z_g, \theta_g$ . Currently, the achieved precision using the factorization in Eq. (3.1) is NLL' accuracy, which can be extended to yet higher accuracy in the future. These results allow percent level comparisons to the available data without the need to model nonperturbative effects. As an example, we show the comparison to ATLAS data [164] in the left panel of Fig. 4.

In recent years, significant progress has been made in developing factorization formulas and improving the theoretical precision of theoretical calculations using both direct perturbative QCD methods and SCET. Several hallmark observables in the field of jet substructure at the LHC have now been extended systematically to NLL' and partially NNLL accuracy. Examples include the jet mass [165–169], jet angularities [167, 170–174], the groomed jet radius [175], the momentum sharing fraction  $z_g$  [162, 163], the jet shape [176], the angle between different jet axes [177], the jet pull [178, 178, 179], and the primary Lund plane [180]. A new jet grooming technique called dynamical grooming has been developed in Ref. [181, 182]. Dedicated observables to distinguish boosted Higgs,  $Z$ ,  $W$  or BSM jets have been studied analytically in Refs. [183–185]. These observables make use of the multi-prong structure of jets originating from boosted resonances compared to single-prong QCD jets. In Refs. [186–188], energy-energy correlators in the collinear limit were discussed with are particularly suitable for extensions to higher accuracy. Recently, jet tagging has also been proposed as a useful tool to constrain collinear PDFs at the LHC [189]. While jet grooming has been introduced initially to remove soft wide-angle radiation from the



**Figure 4.** Left: Comparison of numerical results for the momentum sharing fraction  $z_g$  at NLL' accuracy [163] to ATLAS data [164]. Figure taken from Ref. [163]. Right: Prediction of the Siverts asymmetry at the EIC in back-to-back electron-jet production for different bins of the jet charge  $Q_k$ . Figure taken from Ref. [196]

jet, it has recently been utilized to design Infrared-Collinear Safe observables which are particularly sensitive to soft radiation [177, 190]. These observables may lead to a better understanding of the underlying event contribution and hadronization effects. In the future it will be essential to extend these calculations to NNLL' or even N<sup>3</sup>LL accuracy. This requires progress in fixed order calculations of relevant collinear and soft functions. For example, in Ref. [191], the exclusive collinear jet function for anti- $k_T$  jets [192] was computed at NNLO which is an essential ingredient to extend the precision of existing jet substructure calculations. See also Refs. [193, 194]. Moreover, future developments are necessary to resum non-global logarithms. In Ref. [195], the NLL resummation of NGLs was achieved for the  $e^+e^-$  hemisphere case which sets the starting point for extending the precision of relevant jet substructure observables.

At the EIC, precision calculations of jets and their substructure will be critical to explore the 3D structure of nucleons and nuclei and study aspect of the QCD hadronization process. Inclusive jet production has been calculated up to NNLO in unpolarized and polarized  $ep$  scattering which can provide important constraints for collinear PDFs. See Refs. [197–201] for recent developments. Jets are expected to play an important role to constrain TMD PDFs and fragmentation functions. Different than Semi-Inclusive Deep-Inelastic Scattering (SIDIS), jet observables allow for a decorrelation of initial and final-state TMD dynamics. Back-to-back electron-jet correlations measured in the laboratory frame [202] can be used to study TMD PDFs which is independent of TMD fragmentation. In Ref. [196], it was proposed to use the jet charge [203, 204] to separate up and down quark contributions to the Siverts asymmetry in transversely polarized  $ep$  scattering. The numerical predictions for the EIC are shown in the right panel of Fig. 4. In the Breit frame, jets can be clustered in terms of energy instead of transverse energy or momentum as in the laboratory frame [205]. Since the photon virtuality  $Q$  provides an additional reference scale, the jet energy spectrum can be accessed. This provides a unique opportunity to not only measure inclusive jets, but also the energy loss of leading jets in vacuum and

the nuclear environment [142]. Breit frame jets also allow for measurements which are sensitive to TMD PDFs [206]. Chiral-odd TMDs such as the transversity can be probed by time-reversal odd jets [207]. The observable and factorization structure is similar to SIDIS but the final-state TMD fragmentation function is replaced with a jet function that can be calculated perturbatively. Moreover, gluon TMD PDFs can be accessed using high transverse momentum di-jets in the Breit frame [208, 209]. TMD fragmentation can be studied independent of TMD PDFs using jet substructure observables. This includes both the unpolarized case and for example the Collins effect which can be accessed when the initial proton is transversely polarized [209–211]. Energy-energy correlators in the back-to-back region can also provide important new insights into TMD dynamics [212]. Moreover, EIC jets provide a powerful tool to probe cold nuclear matter effects in electron-nucleus collisions [202, 213–216]. In order to match the precision of traditional processes such as SIDIS, higher perturbative accuracy is needed at leading and subleading power.

#### 4 Small- $x$ resummation

Gluon saturation has attracted a lot of attention in recent years and is one of the major scientific goals of the future EIC [217]. Gluon saturation plays the key role in understanding proton and heavy nuclei collisions in the high-energy limit, where the gluon momentum fraction  $x$  is very small. In this small- $x$  region, the gluon density grows dramatically and enters the nonlinear regime where gluon recombination becomes equally important to the splitting process, and the color glass condensate (CGC) effective theory [218–221] is the proper framework to describe this regime. The nonlinear B-JIMWLK (or its infinite color approximation, the BK equation) equation [222–229] replaces the linear parton evolution equations [230], which inevitably leads to gluon saturation [231, 232] with a characteristic scale  $Q_s$ . The saturation scale  $Q_s$  features the typical transverse momentum of the gluons inside the proton or the nucleus and grows as  $x$  decreases.

There are experimental results [233–240] that are compatible with saturation-model predictions, but there still exists no definitive evidence to claim a discovery. In the future, dedicated measurements at the EIC will provide further information on the regime of gluon saturation.

In order to faithfully and unambiguously establish gluon saturation and its onset, reliable theoretical predictions for small- $x$  collider phenomena are crucial. In the small- $x$  regime, the typical semi-hard saturation scale is of the order of a few GeV and, therefore,  $\alpha_s(Q_s)$  is typically not small enough. As a consequence, calculations beyond the leading order (LO) are generally required to ensure the convergence of the perturbative results. Recently, some attempts have been made in realizing the next-to-leading order (NLO) calculations for small- $x$  physics [241–254]. However, systematic calculations remain challenging and often cut-offs need to be introduced that violate the small- $x$  factorization theorem and mix non-perturbative and perturbative quantities that can invalidate the predictions. Meanwhile, there also lacks a systematic way to resum the large logarithms in the semi-hard NLO coefficient of the small- $x$  factorization theorem, which leads to the infamous

negative cross section problem [255] and hinders practical phenomenological applications of the NLO calculations.

One recent proposal suggests to use power counting arguments and to couple SCET to the CGC formalism as a possible approach to combine systematic fixed order results and resummation [256, 257]. The idea is applied to single-inclusive hadron production in proton-nucleus collisions,  $pA \rightarrow hX$ , at forward rapidity. The CGC factorization for this observable is confirmed at NLO [243]. However, the observed negative cross section when the hadron transverse momentum  $p_{h,\perp}$  becomes a bit larger has been quite puzzling to the community [255]. The NLO partonic cross section contains contributions of the form


$$\begin{aligned} \frac{d^2\hat{\sigma}^{(1)}}{dzd^2p'_\perp} &\propto -\frac{\alpha_s}{2\pi}\mathbf{T}_i^2P_{i\rightarrow i}(z)\ln\frac{r_\perp^2\mu^2}{c_0^2}\left(1+\frac{1}{z^2}e^{i\frac{1-z}{z}p'_\perp\cdot r_\perp}\right) \\ &- \frac{\alpha_s}{\pi}\mathbf{T}_i^a\mathbf{T}_j^{a'}\int\frac{dx_\perp}{\pi}\left\{\frac{1}{z}\frac{2}{(1-z)_+}e^{i\frac{1-z}{z}p'_\perp\cdot r'_\perp}\frac{r'_\perp\cdot r''_\perp}{r'_\perp{}^2r''_\perp{}^2}\right. \\ &\left.+\delta(1-z)\ln\frac{X_f}{X_A}\left[\frac{r_\perp^2}{r'_\perp{}^2r''_\perp{}^2}\right]_+\right\}W_{aa'}(x_\perp)+\dots, \end{aligned} \quad (4.1)$$

where  $p'_\perp$  is the transverse momentum of the fragmenting parton. Here,  $W_{aa'}$  is the CGC Wilson line in the adjoint representation,  $X_A$  is the momentum fraction carried by the gluon from the nucleus and  $X_f$  is the scale due to the rapidity divergence [92, 251, 256, 258]. Moreover,  $P_{i\rightarrow i}(z)$  is the QCD splitting function. Other conventions of the notation can be found in [257]. The terms in the second line are due to the interference between initial and final state radiation. At forward rapidity, the interference is not power suppressed. It was explicitly demonstrated in [257] that as the hadron transverse momentum  $p_{h,\perp}$  increases,  $z$  quickly approaches 1, and the threshold logarithms of the form  $\frac{1}{(1-z)_+}$  overwhelmingly dominate over the other contributions and drive the cross section to negative values. Therefore, an appropriate threshold resummation is required to ensure a reliable prediction.

It is worthwhile to point out that the threshold resummation in the small- $x$  regime is different from the threshold resummation within the collinear factorization, as is obvious from Eq. (4.1) where the color structure  $\mathbf{T}_i^a\mathbf{T}_j^{a'}W_{aa'}$  of the single logarithmic interference term proposes new challenges and can hardly be produced by the known threshold resummation techniques. Numerically, it is found that it is this interference term that makes up the dominant bulk of the threshold contribution [257]. Therefore, a new resummation formalism is required.

In [257], the couplings between the CGC Wilson line  $W_{aa'}(x)$  and the SCET soft and collinear fields are put in by hand. Invoking SCET power counting and factorization, it is shown that the cross section for this process can be factorized as

$$\begin{aligned} \frac{d^2\sigma}{dy_h d^2p_{h\perp}} &= \sum_{i,j=g,q} \frac{1}{4\pi^2} \int \frac{d\xi}{\xi^2} \frac{dx}{x} z x f_{i/P}(x, \mu) D_{h/j}(\xi, \mu) \\ &\times \int d^2b_\perp d^2b'_\perp e^{ip'_\perp\cdot r_\perp} \left\langle \langle \mathcal{M}_0(b'_\perp) | \mathcal{J}(z, \mu, \nu, b_\perp, b'_\perp) \mathcal{S}(\mu, \nu, b_\perp, b'_\perp) | \mathcal{M}_0(b_\perp) \rangle \right\rangle_\nu. \end{aligned} \quad (4.2)$$



**Figure 5.** Comparison of the CGC threshold resummation prediction with the LHC [260] and RHIC [234] data. Figure taken from Ref. [257]

where  $\mathcal{J}$  and  $\mathcal{S}$  represent the collinear and soft contributions, respectively. The LO color space notation  $|\mathcal{M}_0(b_\perp)\rangle$  [259] is used and encodes the CGC (Glauber) Wilson line  $W_{i_c j_c}(b_\perp)$  where  $i_c$  and  $j_c$  are the color indices for the incoming and the outgoing partons using the fundamental for quarks and the adjoint representation for gluons.

It is shown in [257] that the formalism

- reproduces the NLO results in covariant gauge. The soft function reproduces the so-called kinematic constraint, which was put in by hand in the previous calculations, but shows up naturally and automatically within the SCET/CGC hybrid formalism.
- systematically regulates all the divergences by dimensional regularization or by the rapidity regulator. And the final results are consistent with the  $\overline{\text{MS}}$  scheme; No cutoffs that violate the SCET power counting are introduced;
- naturally assigns the IR/rapidity poles to either the PDF/FFs or the nucleus CGC dipole distribution. The scale dependence correctly reproduces the DGLAP evolution for the proton and the final state hadron, as well as the small- $x$  BK equation for the colliding nucleus.
- allows for the resummation of the threshold logarithms, which formally can be achieved through the evolution factor of the jet and the soft function

$$U_J U_S = \exp \left[ -\frac{\alpha_s}{\pi} \int \frac{dx_\perp}{\pi} \left( \ln \frac{\nu_S}{\nu_J} I_{BK,r} + \ln \frac{X_f}{X_A} I_{BK} \right) \mathbf{T}_i^a \mathbf{T}_j^{a'} W_{aa'}(x_\perp) \right]. \quad (4.3)$$

Here we see clearly that the resummation is not a conventional Sudakov evolution but it contains a complicated color structures and is highly non-linear in color. The exponentiation of the CGC Wilson line  $W_{aa'}$  suggests that multiple structures will arise at higher orders which, again, cannot be reproduced by conventional threshold resummation. After the threshold resummation, the problem of the negative cross section is resolved and the theoretical predictions are found to agree with the known

data [257]. A different approach to the negative cross section problem can be found in [261].

The success of handling the threshold logarithms in forward  $pA$  collisions suggests plausible future applications to other small- $x$  phenomena by combining the SCET and CGC formalism. For instance to apply the formalism to the forward di-jet/di-hadron production at the EIC. At the moment, the hybrid formalism is performed in a brute force way by coupling the CGC shock wave (Glauber Wilson line) to the SCET soft and collinear fields. In the future, a first principle and more systematic treatment is required along this line.

## 5 Fragmentation and small- $z_h$ resummation

Another critical observable requiring resummation is fragmentation. Fragmentation is simply the inclusive cross-section for a particle to carry a certain amount of momentum in the event, while ignoring the rest of the process. These cross-sections have long been foundational to our understanding of both the perturbative and non-perturbative structure of high energy scattering. Perturbative physics since, at asymptotically high energies, the scaling properties with the hard momentum scale of the collision can be calculated, but non-perturbative physics plays a critical role since the observable is defined directly on the hadrons exiting the hard interaction.

The canonical example of fragmentation is  $e^+e^- \rightarrow h + X$ , where one asks how many hadrons in the event carry a fraction of the total energy of the event. This example also enjoys one of the oldest factorization theorems of QCD, as well as the corresponding processes  $ep \rightarrow h + X$  and  $pp \rightarrow h + X$ , see Refs. [262–265]. If  $Q$  is the center of mass energy of the electron-positron pair, and  $p_h$  is the momentum of the hadron, the cross-section is:

$$z_h = \frac{2Q \cdot p_h}{Q^2}, \text{ with } 0 \leq z_h \leq 1, \quad (5.1)$$

$$\frac{1}{\sigma_0} \frac{d\sigma}{dz_h} = \sum_{i=q,\bar{q},g} \int_{z_h}^1 \frac{dz}{z} C_i(z, \mu^2, Q^2) d_{h/i}\left(\frac{z_h}{z}, \mu^2, \Lambda_{QCD}^2\right) + \mathcal{O}\left(\frac{\Lambda_{QCD}^2}{Q^2}\right). \quad (5.2)$$

Here  $C_i$  is the coefficient function that describes how the  $e^+e^-$  pair excite the QCD vacuum at high energies, and from this exits a parton of flavor  $i$  carrying energy fraction  $z$ . From this parton, a hadron is formed carrying energy fraction  $z_h$ , and this process is captured by the fragmentation function  $d_{h/i}$ . The fragmentation function depends upon the scale of confinement  $\Lambda_{QCD}$ , and both functions depend on the factorization scale  $\mu$ . The cross-section or number density  $d\sigma/\sigma_0$ , (normalized by the total inclusive cross-section) does not depend on the factorization scale  $\mu$ , leading to the renormalization group equations:

$$\bar{C}_i(n, \mu^2, Q^2) = \int_0^1 \frac{dz}{z} z^{n+1} C_i(z, \mu^2, Q^2), \quad (5.3)$$

$$\bar{d}_{h/i}(n, \mu^2, Q^2) = \int_0^1 \frac{dz}{z} z^{n+1} d_{h/i}(z, \mu^2, \Lambda_{QCD}^2), \quad (5.4)$$

$$\mu^2 \frac{d}{d\mu^2} \bar{C}_i(n, \mu^2, Q^2) = - \sum_j \gamma_{ji}^T(n) \bar{C}_j(n, \mu^2, Q^2), \quad (5.5)$$

$$\mu^2 \frac{d}{d\mu^2} \bar{d}_{h/i}(n, \mu^2, Q^2) = \sum_j \gamma_{ij}^T(n) \bar{d}_{h/j}(n, \mu^2, Q^2). \quad (5.6)$$

All processes have the same fragmentation function  $d_{h/i}$ , an early and testable prediction of factorization: the universality of the hadronization process when the cross-section is sufficiently inclusive. Moreover, the fragmentation to hadrons can be generalized to fragmentation to jets, where one replaces the intrinsic cut-off scale of confinement with the jet radius  $R$ , as was done in Refs. [266–268]. At fixed order,  $C_i$  has been calculated to order  $\alpha_s^2$  in Ref. [269–271], with partial results at the same order for the DIS fragmentation process Refs. [272, 273], and  $\gamma_{ij}^T$  has been calculated to order  $\alpha_s^3$  Refs. [274–277].

The cross-section is largest in the  $z_h \rightarrow 0$  limit. That is, the bulk of hadrons produced have low energy. This has its origins in the fact that QCD enjoys an enhanced infra-red region of phase-space, and the  $z_h \rightarrow 0$  limit probes the soft singularities of QCD. The importance of this region was recognized early on, Refs. [278–280], and a resummation of the logarithms of  $z_h$  in the anomalous dimension was accomplished at leading and next-to leading order, however, with a scheme dependence which did not interface well with the minimal subtraction schemes of dimensional regularization, making it difficult to compare to fixed order calculations. At leading log order in the  $z_h \rightarrow 0$  limit, which in mellin space corresponds to  $n \rightarrow 0$ , we have:

$$\gamma_{gg}^T(n) = \frac{n}{4} \left( -1 + \sqrt{1 + 8 \frac{\alpha_s C_A}{\pi n^2}} \right) + \dots \quad (5.7)$$

After performing the inverse mellin transform, one can see then that this is double logarithmic series, *in the anomalous dimension itself*.

The resummation was further developed to embody the concepts of angular-ordering and hard cut-offs in transverse momentum, culminating in the so-called mixed leading log (MLL) approximation, see Refs. [281–283]. The MLL has seen surprising phenomenological success (Ref. [284]) when coupled to the hypothesis of parton-hadron duality (*very crudely*, each parton becomes a hadron), see Ref. [285], but with no clear connection to traditional fixed order calculations of the anomalous dimension or the coefficient function.

More recently, two other approaches to small- $z_h$  resummation have been developed, which have been pushed beyond leading logarithmic order, and include a clear relationship to fixed order calculations.<sup>1</sup> The first approach of Refs. [287, 288], which has also been used to resum transverse-momentum dependent fragmentation functions Ref. [26], makes use of a set of recursion relations based on an ansatz that the energy fraction  $z_h$  must ultimately cut-off soft divergences in the calculation. This leads to an over-determined set of linear relations, whose unique solution enables N<sup>2</sup>LL resummation of the coefficient function, and

---

<sup>1</sup>Though see also Ref. [286], which clarified the role of scheme-dependence in small- $z_h$  resummations, thus enabling the first successful matching of fixed-order to resummed perturbation theory in the context of fragmentation.

$N^3$ LL resummation of the anomalous dimension with arbitrary flavor dependence. Parton-to-pion fragmentation functions have been fit using these resummed coefficient functions and anomalous dimensions Ref. [289].

The second approach of Ref. [290] seeks to put the notion of an angular-ordered cascade on firmer theoretical grounds, by explicitly writing an evolution equation for the partonic coefficient function. The evolution equation seeks to make use of a universal kernel that in principle should be calculated from the BFKL equation that governs forward scattering and the resummation of the DIS cross-section as Bjorken  $x$  goes to zero. Consistency of the BFKL equation [230, 291–294] with the collinear factorization of DIS cross-section has been known to give the resummation of space-like anomalous dimensions of twist-two operators, where the consistency in dimensional regularization was first worked out in Ref. [295, 296]. However, work on the resummation of the soft physics of NGLs in Refs. [297–301] established an essential connection between forward scattering physics and the resummation of non-global-logs (NGLs), based on the correspondence between the BK and BMS equations [224, 225, 302]. While the NGL appear in semi-exclusion processes where radiation is vetoed, in certain angular regions Ref. [290] makes the connection between forward scattering and soft parton cascades explicit for the case of inclusive fragmentation, writing a deformation of the BFKL equation which resums the *time-like* anomalous dimensions in complete analogy to the space-like case. It is worth noting that Refs. [303, 304] also examined a form of the BFKL equation that could be used for fragmentation, but never explicitly connected the results with the traditional collinear factorization approach.

In addition to developing a BFKL approach to soft fragmentation, the authors also showed how one may write DGLAP-style evolution equations that resum the small- $z_h$  logs. Focusing on the case of pure Yang-Mills, one postulates that in  $4 - 2\epsilon$  dimensions, the cross-section  $d(z_h, \mu^2, Q^2, R^2)$  for fragmenting a gluon of energy fraction  $z_h$  can be defined with an angular cut-off: the descendant parton cannot approach within angle  $R$  of their parent or siblings. This regulates the collinear divergence. Then one takes the limit  $R \rightarrow 0$  using the evolution equation:

$$R^2 \frac{\partial}{\partial R^2} z_h^{1+2\epsilon} d(z_h, \mu^2, Q^2, R^2) = \rho\left(\frac{\mu^2}{R^2 Q^2}\right) z_h^{1+2\epsilon} d(z_h, \mu^2, Q^2, R^2) + \int_{z_h}^1 \frac{dz}{z} P\left(\frac{z_h}{z}; \frac{\mu^2}{z^2 R^2 Q^2}\right) z^{1+2\epsilon} d(z, \mu^2, Q^2, R^2), \quad (5.8)$$


$$P\left(\frac{z_h}{z}; \frac{\mu^2}{z^2 R^2 Q^2}\right) = \sum_{\ell=1}^{\infty} P^{(\ell-1)}\left(\frac{z_h}{z}; \alpha_s; \epsilon\right) \left(\frac{\mu^2}{z^2 R^2 Q^2}\right)^{\ell\epsilon}, \quad (5.9)$$

$$\rho\left(\frac{\mu^2}{R^2 Q^2}\right) = \sum_{\ell=1}^{\infty} \rho^{(\ell-1)}(\alpha_s; \epsilon) \left(\frac{\mu^2}{R^2 Q^2}\right)^{\ell\epsilon}. \quad (5.10)$$

$P$  is derived from the space-like anomalous dimension in  $4 - 2\epsilon$  dimensions of Refs. [305–307], and  $\rho$  is connected to the beta-function. After taking  $R \rightarrow 0$ , we recover the bare perturbative cross-section in  $4 - 2\epsilon$  dimensions, so:

$$\bar{d}(n, \mu^2, \mu^2, 0) = \exp\left(\int_0^{\alpha_s} \frac{d\alpha}{\beta(\alpha, \epsilon)} \gamma^T(\alpha, n)\right) \mathcal{R}^T(\alpha_s, n). \quad (5.11)$$






**Figure 6.** The fragmentation spectrum for  $e^+e^- \rightarrow h^\pm + X$  at center of mass energies  $Q = 91$  and  $206$  GeV. The gray region is excluded from the  $\chi^2/\text{d.o.f}$  determination based on hadron mass corrections or  $z_h \sim O(1)$ , and the quoted  $\chi^2/\text{d.o.f}$  is the goodness-of-fit for that data set. All curves are fitted at  $91$  GeV and evolved to  $206$  GeV. Compared to the NLL curve (red) is the evolved 3 flavor scheme MLLA limiting spectrum (purple LS) from Ref. [284], and a 5 flavor scheme evolving the moments of a Gaussian according to the MLLA anomalous dimension, as in Ref. [281] (orange G).

Both  $\gamma^T$  and  $\mathcal{R}^T$  are resummed automatically in the soft limit, and indeed,  $\mathcal{R}^T$  is the soft resummed coefficient function of Eq. (5.2). The evolution equation (5.8) can be viewed as an appropriate generalization of the reciprocity relations of Refs. [308, 309]

Going into the future, there are multiple directions to study. Firstly, it is necessary to extend the resummation of soft fragmentation beyond the well-studied case of  $e^+e^- \rightarrow h + X$  to processes that include initial state hadrons like  $ep \rightarrow h + X$  and  $pp \rightarrow h + X$ . Secondly, there was a history of so-called fractal observables that were resummed in the unsystematic MLL approximation, many of which are outlined in Ref. [283]. It would be important to revisit these observables with an improved understanding of the resummation of the soft fragmentation region. Additionally, fragmentation in a constrained jet (Ref. [146, 310]) is critical for many experimental analyses, as is following the flow of charge within the jet Ref. [188, 311, 312]. The extension of the soft resummation to these cases would bring insights into the flow of charge and momentum in QCD cascades. Lastly, the approach of Ref. [290] should be extended to include the full flavor structure of QCD, and the exact connection to forward scattering physics should move beyond conjectures and verification order-by-order, but involve full-scale proofs and examples showing the limits of the correspondence between time-like and space-like branching processes.

On a more phenomenological note, Ref. [315] initiated a study of parton-hadron duality based off of the improved understanding of the resummation of the soft physics from Ref. [290], resumming all logarithms of  $z_h$  to NLL order. The results showed a marked improvement on the older MLL approximation, see Fig. 6, where fragmentation to charged hadrons is compared to theory predictions making use of parton-hadron-duality hypothesis. Perhaps even more surprising was the ability to describe the data in regions outside the



**Figure 7.** The fragmentation spectrum extrapolated into the large  $z_h$  region for  $e^+e^- \rightarrow h^\pm + X$  at center of mass energies  $Q = 91$  and  $189$  GeV. The region below  $\ln \frac{1}{z_h} < 0.5$  was excluded from the fit. Data taken from the OPAL, SLD, and ALEPH collaborations Refs. [284, 313, 314]. The NLL result was a global fit to data from  $Q = 35$  to  $206$  GeV.

soft limit, see Fig. 7. With improved analytical control of the perturbative structure of the theory, the time seems ripe to revisit parton-hadron duality, and the ultimate limits of perturbation theory.

## 6 Conclusions

The perturbative approach to calculate QCD scattering processes from first principles can be spoiled by the presence of large logarithmic corrections that appear in the power expansion of cross sections in the QCD strong coupling constant  $\alpha_s$ . These large logarithmic corrections need to be resummed to all orders in  $\alpha_s$ , which restores the predictive power of perturbative QCD. In this work, we reviewed recent progress in QCD resummation techniques. An improved understanding of the all-order structure of QCD is relevant for the high-luminosity era at the LHC as well as future experiments such as the Electron-Ion Collider. First, we considered the resummation at subleading power, which has become possible for several processes. For example, the threshold resummation for the Drell-Yan process and Higgs production via gluon fusion has been now been performed at subleading power and leading logarithmic accuracy. It was found that the numerical impact at the LHC can be comparable to high-precision resummation at leading power. We discussed techniques both in soft collinear effective theory and direct QCD. Both methods still face conceptual challenges which hinder systematic evaluation of the higher logarithmic orders. Second, we reviewed progress in the calculation of jet and jet substructure observables. Here the challenging aspect is the simultaneous resummation of multiple classes of large logarithmic corrections. Especially, jet substructure observables can be sensitive to several disparate energy scales that require careful treatment to identify the relevant energy scales and establish the corresponding factorization theorems. We discussed jet measurements at the LHC and outlined how jet physics can provide new insights into the structure of hadrons at the future EIC. Third, we focused on small- $x$  resummation, which is relevant

in the high-energy limit where the momentum fraction carried by the initial-state gluon becomes small. We discussed recent work that combines resummation techniques in SCET and the Color Glass Condensate formalism. The additional resummation of threshold logarithms allows for the description of forward  $pA$  scattering data. Lastly, we focused on the QCD fragmentation process. The inclusive hadron cross section peaks in the limit where the momentum fraction of the identified hadrons becomes small  $z_h \rightarrow 0$ . Large logarithmic corrections in the coefficient function and the time-like anomalous dimension need to be taken into account to achieve a reliable prediction within perturbative QCD. We discussed recent progress that employs space-time reciprocity relations, which connect the description of final-state hadrons to BFKL dynamics. The resummation of small- $z_h$  logarithms allows for a precise comparison to the available data from LEP and SLD colliders. The topics discussed here provide a starting point for future studies at high perturbative accuracy both at leading and subleading power. QCD resummation is a critical ingredient to stress test the Standard Model of particle physics, better understanding QCD itself, and to improve searches for physics beyond the Standard Model.

## Acknowledgments

MvB acknowledges the UK Science and Technology Facilities Council (STFC) grant 452 number ST/T000864/1 and the Royal Society Research Professorship (RP\R1\180112). SJ is supported by the UK Science and Technology Facilities Council (STFC) under grant ST/T001011/1. XL is supported by the National Natural Science Foundation of China under Grant No. 12175016. DN is supported by the U.S. Department of Energy under Contract No. 89233218CNA000001 and by the LDRD program at LANL. FR is supported by the Simons Foundation under the Simons Bridge program for Postdoctoral Fellowships at SCGP and YITP, award number 815892 and the NSF, award number 1915093. RS is supported by the United States Department of Energy under Grant Contract DE-SC0012704. LV is supported by Fellini, Fellowship for Innovation at INFN, funded by the European Union's Horizon 2020 research programme under the Marie Skłodowska-Curie Cofund Action, grant agreement no. 754496. GV is supported by the United States Department of Energy, Contract DE-AC02-76SF00515. JW is supported by the National Natural Science Foundation of China under Grant No. 12005117.

## References

- [1] G. Parisi, *Summing Large Perturbative Corrections in QCD*, *Phys. Lett. B* **90** (1980) 295.
- [2] G. Curci and M. Greco, *Large Infra-red Corrections in QCD Processes*, *Phys. Lett. B* **92** (1980) 175.
- [3] G.F. Sterman, *Summation of Large Corrections to Short Distance Hadronic Cross-Sections*, *Nucl. Phys. B* **281** (1987) 310.
- [4] S. Catani and L. Trentadue, *Resummation of the QCD Perturbative Series for Hard Processes*, *Nucl. Phys. B* **327** (1989) 323.

- [5] S. Catani and L. Trentadue, *Comment on QCD exponentiation at large  $x$* , *Nucl. Phys. B* **353** (1991) 183.
- [6] S. Moch and A. Vogt, *Higher-order soft corrections to lepton pair and Higgs boson production*, *Phys. Lett. B* **631** (2005) 48 [[hep-ph/0508265](#)].
- [7] E. Laenen and L. Magnea, *Threshold resummation for electroweak annihilation from DIS data*, *Phys. Lett. B* **632** (2006) 270 [[hep-ph/0508284](#)].
- [8] A. Idilbi, X.-d. Ji, J.-P. Ma and F. Yuan, *Threshold resummation for Higgs production in effective field theory*, *Phys. Rev. D* **73** (2006) 077501 [[hep-ph/0509294](#)].
- [9] T. Becher, M. Neubert and G. Xu, *Dynamical Threshold Enhancement and Resummation in Drell-Yan Production*, *JHEP* **07** (2008) 030 [[0710.0680](#)].
- [10] A. Kulesza and L. Motyka, *Threshold resummation for squark-antisquark and gluino-pair production at the LHC*, *Phys. Rev. Lett.* **102** (2009) 111802 [[0807.2405](#)].
- [11] M. Beneke, P. Falgari, S. Klein and C. Schwinn, *Hadronic top-quark pair production with NNLL threshold resummation*, *Nucl. Phys. B* **855** (2012) 695 [[1109.1536](#)].
- [12] M. Bonvini, S. Forte, M. Ghezzi and G. Ridolfi, *Threshold Resummation in SCET vs. Perturbative QCD: An Analytic Comparison*, *Nucl. Phys. B* **861** (2012) 337 [[1201.6364](#)].
- [13] M. Bonvini and S. Marzani, *Resummed Higgs cross section at  $N^3LL$* , *JHEP* **09** (2014) 007 [[1405.3654](#)].
- [14] P. Hinderer, F. Ringer, G. Sterman and W. Vogelsang, *Threshold Resummation at NNLL for Single-particle Production in Hadronic Collisions*, *Phys. Rev. D* **99** (2019) 054019 [[1812.00915](#)].
- [15] Y. Li and H.X. Zhu, *Bootstrapping Rapidity Anomalous Dimensions for Transverse-Momentum Resummation*, *Phys. Rev. Lett.* **118** (2017) 022004 [[1604.01404](#)].
- [16] W. Bizon, P.F. Monni, E. Re, L. Rottoli and P. Torrielli, *Momentum-space resummation for transverse observables and the Higgs  $p_{\perp}$  at  $N^3LL+NNLO$* , *JHEP* **02** (2018) 108 [[1705.09127](#)].
- [17] F. Coradeschi and T. Cridge, *reSolve — A transverse momentum resummation tool*, *Comput. Phys. Commun.* **238** (2019) 262 [[1711.02083](#)].
- [18] X. Chen, T. Gehrmann, E.W.N. Glover, A. Huss, Y. Li, D. Neill et al., *Precise QCD Description of the Higgs Boson Transverse Momentum Spectrum*, *Phys. Lett. B* **788** (2019) 425 [[1805.00736](#)].
- [19] S. Camarda et al., *DYTurbo: Fast predictions for Drell-Yan processes*, *Eur. Phys. J. C* **80** (2020) 251 [[1910.07049](#)].
- [20] A. Bacchetta, V. Bertone, C. Bissolotti, G. Bozzi, F. Delcarro, F. Piacenza et al., *Transverse-momentum-dependent parton distributions up to  $N^3LL$  from Drell-Yan data*, *JHEP* **07** (2020) 117 [[1912.07550](#)].
- [21] M.-x. Luo, T.-Z. Yang, H.X. Zhu and Y.J. Zhu, *Quark Transverse Parton Distribution at the Next-to-Next-to-Next-to-Leading Order*, *Phys. Rev. Lett.* **124** (2020) 092001 [[1912.05778](#)].
- [22] M.A. Ebert, B. Mistlberger and G. Vita, *Transverse momentum dependent PDFs at  $N^3LO$* , *JHEP* **09** (2020) 146 [[2006.05329](#)].

- [23] S. Kallweit, E. Re, L. Rottoli and M. Wiesemann, *Accurate single- and double-differential resummation of colour-singlet processes with MATRIX+RADISH: WW production at the LHC*, *JHEP* **12** (2020) 147 [[2004.07720](#)].
- [24] M. Wiesemann, L. Rottoli and P. Torrielli, *The  $Z\gamma$  transverse-momentum spectrum at NNLO+N<sup>3</sup>LL*, *Phys. Lett. B* **809** (2020) 135718 [[2006.09338](#)].
- [25] M.A. Ebert, J.K.L. Michel, I.W. Stewart and F.J. Tackmann, *Drell-Yan  $q_T$  resummation of fiducial power corrections at N<sup>3</sup>LL*, *JHEP* **04** (2021) 102 [[2006.11382](#)].
- [26] M.-x. Luo, T.-Z. Yang, H.X. Zhu and Y.J. Zhu, *Unpolarized quark and gluon TMD PDFs and FFs at N<sup>3</sup>LO*, *JHEP* **06** (2021) 115 [[2012.03256](#)].
- [27] T. Becher and T. Neumann, *Fiducial  $q_T$  resummation of color-singlet processes at N<sup>3</sup>LL+NNLO*, *JHEP* **03** (2021) 199 [[2009.11437](#)].
- [28] M.A. Ebert, B. Mistlberger and G. Vita, *TMD Fragmentation Functions at N<sup>3</sup>LO*, *JHEP* **07** (2021) 121 [[2012.07853](#)].
- [29] M.A. Ebert, B. Mistlberger and G. Vita, *The Energy-Energy Correlation in the back-to-back limit at N<sup>3</sup>LO and N<sup>3</sup>LL'*, *JHEP* **08** (2021) 022 [[2012.07859](#)].
- [30] W.-L. Ju and M. Schönherr, *The  $q_T$  and  $\Delta\phi$  spectra in W and Z production at the LHC at N<sup>3</sup>LL'+N<sup>2</sup>LO*, *JHEP* **10** (2021) 088 [[2106.11260](#)].
- [31] T. Neumann, *The diphoton  $q_T$  spectrum at N<sup>3</sup>LL' + NNLO*, *Eur. Phys. J. C* **81** (2021) 905 [[2107.12478](#)].
- [32] G. Billis, B. Dehnadi, M.A. Ebert, J.K.L. Michel and F.J. Tackmann, *Higgs  $p_T$  Spectrum and Total Cross Section with Fiducial Cuts at Third Resummed and Fixed Order in QCD*, *Phys. Rev. Lett.* **127** (2021) 072001 [[2102.08039](#)].
- [33] X. Chen, T. Gehrmann, E.W.N. Glover, A. Huss, P. Monni, E. Re et al., *Third order fiducial predictions for Drell-Yan at the LHC*, [2203.01565](#).
- [34] C.W. Bauer, S. Fleming, D. Pirjol and I.W. Stewart, *An Effective field theory for collinear and soft gluons: Heavy to light decays*, *Phys. Rev. D* **63** (2001) 114020 [[hep-ph/0011336](#)].
- [35] C.W. Bauer, D. Pirjol and I.W. Stewart, *Soft collinear factorization in effective field theory*, *Phys. Rev. D* **65** (2002) 054022 [[hep-ph/0109045](#)].
- [36] M. Beneke, A.P. Chapovsky, M. Diehl and T. Feldmann, *Soft collinear effective theory and heavy to light currents beyond leading power*, *Nucl. Phys. B* **643** (2002) 431 [[hep-ph/0206152](#)].
- [37] M. Beneke and T. Feldmann, *Multipole expanded soft collinear effective theory with nonAbelian gauge symmetry*, *Phys. Lett. B* **553** (2003) 267 [[hep-ph/0211358](#)].
- [38] A.A. Penin, *High-Energy Limit of Quantum Electrodynamics beyond Sudakov Approximation*, *Phys. Lett. B* **745** (2015) 69 [[1412.0671](#)].
- [39] K. Melnikov and A. Penin, *On the light quark mass effects in Higgs boson production in gluon fusion*, *JHEP* **05** (2016) 172 [[1602.09020](#)].
- [40] A.A. Penin and N. Zerf, *Two-loop Bhabha Scattering at High Energy beyond Leading Power Approximation*, *Phys. Lett. B* **760** (2016) 816 [[1606.06344](#)].
- [41] T. Liu, A.A. Penin and N. Zerf, *Three-loop quark form factor at high energy: the leading mass corrections*, *Phys. Lett. B* **771** (2017) 492 [[1705.07910](#)].

- [42] T. Liu and A.A. Penin, *High-Energy Limit of QCD beyond the Sudakov Approximation*, *Phys. Rev. Lett.* **119** (2017) 262001 [[1709.01092](#)].
- [43] R. Boughezal, A. Isgro and F. Petriello, *Next-to-leading-logarithmic power corrections for  $N$ -jettiness subtraction in color-singlet production*, *Phys. Rev. D* **97** (2018) 076006 [[1802.00456](#)].
- [44] R. Brüser, S. Caron-Huot and J.M. Henn, *Subleading Regge limit from a soft anomalous dimension*, *JHEP* **04** (2018) 047 [[1802.02524](#)].
- [45] I. Moulton, I.W. Stewart, G. Vita and H.X. Zhu, *First Subleading Power Resummation for Event Shapes*, *JHEP* **08** (2018) 013 [[1804.04665](#)].
- [46] M.A. Ebert, I. Moulton, I.W. Stewart, F.J. Tackmann, G. Vita and H.X. Zhu, *Power Corrections for  $N$ -Jettiness Subtractions at  $\mathcal{O}(\alpha_s)$* , *JHEP* **12** (2018) 084 [[1807.10764](#)].
- [47] T. Liu and A. Penin, *High-Energy Limit of Mass-Suppressed Amplitudes in Gauge Theories*, *JHEP* **11** (2018) 158 [[1809.04950](#)].
- [48] M. Beneke, A. Broggio, M. Garny, S. Jaskiewicz, R. Szafron, L. Vernazza et al., *Leading-logarithmic threshold resummation of the Drell-Yan process at next-to-leading power*, *JHEP* **03** (2019) 043 [[1809.10631](#)].
- [49] T. Engel, C. Gnendiger, A. Signer and Y. Ulrich, *Small-mass effects in heavy-to-light form factors*, *JHEP* **02** (2019) 118 [[1811.06461](#)].
- [50] M.A. Ebert, I. Moulton, I.W. Stewart, F.J. Tackmann, G. Vita and H.X. Zhu, *Subleading power rapidity divergences and power corrections for  $q_T$* , *JHEP* **04** (2019) 123 [[1812.08189](#)].
- [51] N. Bahjat-Abbas, D. Bonocore, J. Sinninghe Damsté, E. Laenen, L. Magnea, L. Vernazza et al., *Diagrammatic resummation of leading-logarithmic threshold effects at next-to-leading power*, *JHEP* **11** (2019) 002 [[1905.13710](#)].
- [52] A.A. Penin, *Regge Limit of Gauge Theory Amplitudes beyond Leading Power Approximation*, *JHEP* **04** (2020) 156 [[1909.02013](#)].
- [53] M. Beneke, M. Garny, S. Jaskiewicz, R. Szafron, L. Vernazza and J. Wang, *Leading-logarithmic threshold resummation of Higgs production in gluon fusion at next-to-leading power*, *JHEP* **01** (2020) 094 [[1910.12685](#)].
- [54] M. Beneke, A. Broggio, S. Jaskiewicz and L. Vernazza, *Threshold factorization of the Drell-Yan process at next-to-leading power*, *JHEP* **07** (2020) 078 [[1912.01585](#)].
- [55] C. Anastasiou and A. Penin, *Light Quark Mediated Higgs Boson Threshold Production in the Next-to-Leading Logarithmic Approximation*, *JHEP* **07** (2020) 195 [[2004.03602](#)].
- [56] Z.L. Liu, B. Mécaj, M. Neubert and X. Wang, *Factorization at subleading power and endpoint divergences in  $h \rightarrow \gamma\gamma$  decay. Part II. Renormalization and scale evolution*, *JHEP* **01** (2021) 077 [[2009.06779](#)].
- [57] A. Broggio, S. Jaskiewicz and L. Vernazza, *Next-to-leading power two-loop soft functions for the Drell-Yan process at threshold*, *JHEP* **10** (2021) 061 [[2107.07353](#)].
- [58] T. Liu, S. Modi and A.A. Penin, *Higgs boson production and quark scattering amplitudes at high energy through the next-to-next-to-leading power in quark mass*, *JHEP* **02** (2022) 170 [[2111.01820](#)].
- [59] G.P. Korchemsky and G. Marchesini, *Structure function for large  $x$  and renormalization of Wilson loop*, *Nucl. Phys. B* **406** (1993) 225 [[hep-ph/9210281](#)].

- [60] G.P. Korchemsky and G. Marchesini, *Resummation of large infrared corrections using Wilson loops*, *Phys. Lett. B* **313** (1993) 433.
- [61] S. Forte and G. Ridolfi, *Renormalization group approach to soft gluon resummation*, *Nucl. Phys. B* **650** (2003) 229 [[hep-ph/0209154](#)].
- [62] H. Contopanagos, E. Laenen and G.F. Sterman, *Sudakov factorization and resummation*, *Nucl. Phys. B* **484** (1997) 303 [[hep-ph/9604313](#)].
- [63] T. Becher and M. Neubert, *Threshold resummation in momentum space from effective field theory*, *Phys. Rev. Lett.* **97** (2006) 082001 [[hep-ph/0605050](#)].
- [64] M.D. Schwartz, *Resummation and NLO matching of event shapes with effective field theory*, *Phys. Rev. D* **77** (2008) 014026 [[0709.2709](#)].
- [65] C.W. Bauer, S.P. Fleming, C. Lee and G.F. Sterman, *Factorization of  $e+e-$  Event Shape Distributions with Hadronic Final States in Soft Collinear Effective Theory*, *Phys. Rev. D* **78** (2008) 034027 [[0801.4569](#)].
- [66] M. van Beekveld, W. Beenakker, E. Laenen and C.D. White, *Next-to-leading power threshold effects for inclusive and exclusive processes with final state jets*, *JHEP* **03** (2020) 106 [[1905.08741](#)].
- [67] M. van Beekveld, E. Laenen, J. Sinninghe Damsté and L. Vernazza, *Next-to-leading power threshold corrections for finite order and resummed colour-singlet cross sections*, *JHEP* **05** (2021) 114 [[2101.07270](#)].
- [68] M. van Beekveld, L. Vernazza and C.D. White, *Threshold resummation of new partonic channels at next-to-leading power*, *JHEP* **12** (2021) 087 [[2109.09752](#)].
- [69] I. Moutl, I.W. Stewart, G. Vita and H.X. Zhu, *The Soft Quark Sudakov*, *JHEP* **05** (2020) 089 [[1910.14038](#)].
- [70] M. Beneke, M. Garry, S. Jaskiewicz, R. Szafron, L. Vernazza and J. Wang, *Large- $x$  resummation of off-diagonal deep-inelastic parton scattering from  $d$ -dimensional refactorization*, *JHEP* **10** (2020) 196 [[2008.04943](#)].
- [71] A.H. Ajjath, P. Mukherjee and V. Ravindran, *On next to soft corrections to Drell-Yan and Higgs Boson productions*, [2006.06726](#).
- [72] A.H. Ajjath, P. Mukherjee, V. Ravindran, A. Sankar and S. Tiwari, *On next to soft threshold corrections to DIS and SIA processes*, *JHEP* **04** (2021) 131 [[2007.12214](#)].
- [73] F.E. Low, *Bremsstrahlung of very low-energy quanta in elementary particle collisions*, *Phys. Rev.* **110** (1958) 974.
- [74] T.H. Burnett and N.M. Kroll, *Extension of the low soft photon theorem*, *Phys. Rev. Lett.* **20** (1968) 86.
- [75] V. Del Duca, *High-energy Bremsstrahlung Theorems for Soft Photons*, *Nucl. Phys. B* **345** (1990) 369.
- [76] D. Bonocore, E. Laenen, L. Magnea, S. Melville, L. Vernazza and C.D. White, *A factorization approach to next-to-leading-power threshold logarithms*, *JHEP* **06** (2015) 008 [[1503.05156](#)].
- [77] D. Bonocore, E. Laenen, L. Magnea, L. Vernazza and C.D. White, *Non-abelian factorisation for next-to-leading-power threshold logarithms*, *JHEP* **12** (2016) 121 [[1610.06842](#)].

- [78] E. Laenen, G. Stavenga and C.D. White, *Path integral approach to eikonal and next-to-eikonal exponentiation*, *JHEP* **03** (2009) 054 [[0811.2067](#)].
- [79] E. Laenen, L. Magnea, G. Stavenga and C.D. White, *Next-to-Eikonal Corrections to Soft Gluon Radiation: A Diagrammatic Approach*, *JHEP* **01** (2011) 141 [[1010.1860](#)].
- [80] H. Gervais, *Soft Photon Theorem for High Energy Amplitudes in Yukawa and Scalar Theories*, *Phys. Rev. D* **95** (2017) 125009 [[1704.00806](#)].
- [81] E. Laenen, J. Sinninghe Damsté, L. Vernazza, W. Waalewijn and L. Zoppi, *Towards all-order factorization of QED amplitudes at next-to-leading power*, *Phys. Rev. D* **103** (2021) 034022 [[2008.01736](#)].
- [82] D. Bonocore and A. Kulesza, *Soft photon bremsstrahlung at next-to-leading power*, [2112.08329](#).
- [83] E. Gardi, E. Laenen, G. Stavenga and C.D. White, *Webs in multiparton scattering using the replica trick*, *JHEP* **11** (2010) 155 [[1008.0098](#)].
- [84] V. Del Duca, E. Laenen, L. Magnea, L. Vernazza and C.D. White, *Universality of next-to-leading power threshold effects for colourless final states in hadronic collisions*, *JHEP* **11** (2017) 057 [[1706.04018](#)].
- [85] A. Vogt, *Leading logarithmic large- $x$  resummation of off-diagonal splitting functions and coefficient functions*, *Phys. Lett. B* **691** (2010) 77 [[1005.1606](#)].
- [86] N.A. Lo Presti, A.A. Almasy and A. Vogt, *Leading large- $x$  logarithms of the quark-gluon contributions to inclusive Higgs-boson and lepton-pair production*, *Phys. Lett. B* **737** (2014) 120 [[1407.1553](#)].
- [87] M. Beneke, M. Garry, S. Jaskiewicz, J. Strohm, R. Szafron, L. Vernazza et al., unpublished.
- [88] A.J. Larkoski, D. Neill and I.W. Stewart, *Soft Theorems from Effective Field Theory*, *JHEP* **06** (2015) 077 [[1412.3108](#)].
- [89] M. Beneke, M. Garry, R. Szafron and J. Wang, *Subleading-power  $N$ -jet operators and the LBK amplitude in SCET*, *PoS RADCOR2017* (2017) 048 [[1712.07462](#)].
- [90] M. Beneke, P. Hager and R. Szafron, *Gravitational soft theorem from emergent soft gauge symmetries*, [2110.02969](#).
- [91] M. Beneke, P. Hager and R. Szafron, *Soft-collinear gravity beyond the leading power*, [2112.04983](#).
- [92] I.Z. Rothstein and I.W. Stewart, *An Effective Field Theory for Forward Scattering and Factorization Violation*, *JHEP* **08** (2016) 025 [[1601.04695](#)].
- [93] C.W. Bauer and I.W. Stewart, *Invariant operators in collinear effective theory*, *Phys. Lett. B* **516** (2001) 134 [[hep-ph/0107001](#)].
- [94] I. Moulton, I.W. Stewart, F.J. Tackmann and W.J. Waalewijn, *Employing Helicity Amplitudes for Resummation*, *Phys. Rev. D* **93** (2016) 094003 [[1508.02397](#)].
- [95] D.W. Kolodrubetz, I. Moulton and I.W. Stewart, *Building Blocks for Subleading Helicity Operators*, *JHEP* **05** (2016) 139 [[1601.02607](#)].
- [96] M. Beneke, M. Garry, R. Szafron and J. Wang, *Anomalous dimension of subleading-power  $N$ -jet operators*, *JHEP* **03** (2018) 001 [[1712.04416](#)].



- [97] I. Moulton, I.W. Stewart and G. Vita, *A subleading operator basis and matching for  $gg \rightarrow H$* , *JHEP* **07** (2017) 067 [[1703.03408](#)].
- [98] C.-H. Chang, I.W. Stewart and G. Vita, *A Subleading Power Operator Basis for the Scalar Quark Current*, *JHEP* **04** (2018) 041 [[1712.04343](#)].
- [99] I. Feige, D.W. Kolodrubetz, I. Moulton and I.W. Stewart, *A Complete Basis of Helicity Operators for Subleading Factorization*, *JHEP* **11** (2017) 142 [[1703.03411](#)].
- [100] M. Beneke, M. Garny, R. Szafron and J. Wang, *Anomalous dimension of subleading-power  $N$ -jet operators. Part II*, *JHEP* **11** (2018) 112 [[1808.04742](#)].
- [101] M. Beneke, M. Garny, R. Szafron and J. Wang, *Violation of the Kluberg-Stern-Zuber theorem in SCET*, *JHEP* **09** (2019) 101 [[1907.05463](#)].
- [102] I. Moulton, I.W. Stewart and G. Vita, *Subleading Power Factorization with Radiative Functions*, *JHEP* **11** (2019) 153 [[1905.07411](#)].
- [103] M. Beneke and T. Feldmann, *Factorization of heavy to light form-factors in soft collinear effective theory*, *Nucl. Phys. B* **685** (2004) 249 [[hep-ph/0311335](#)].
- [104] R.J. Hill, T. Becher, S.J. Lee and M. Neubert, *Sudakov resummation for subleading SCET currents and heavy-to-light form-factors*, *JHEP* **07** (2004) 081 [[hep-ph/0404217](#)].
- [105] K.S.M. Lee and I.W. Stewart, *Factorization for power corrections to  $B \rightarrow X(s)$  gamma and  $B \rightarrow X(u) l$  anti-nu*, *Nucl. Phys. B* **721** (2005) 325 [[hep-ph/0409045](#)].
- [106] M. Beneke, Y. Kiyo and D.s. Yang, *Loop corrections to subleading heavy quark currents in SCET*, *Nucl. Phys. B* **692** (2004) 232 [[hep-ph/0402241](#)].
- [107] M. Beneke, F. Campanario, T. Mannel and B.D. Pecjak, *Power corrections to anti- $B \rightarrow X(u) l$  anti-nu ( $X(s)$  gamma) decay spectra in the 'shape-function' region*, *JHEP* **06** (2005) 071 [[hep-ph/0411395](#)].
- [108] G. Paz, *Subleading Jet Functions in Inclusive  $B$  Decays*, *JHEP* **06** (2009) 083 [[0903.3377](#)].
- [109] M. Benzke, S.J. Lee, M. Neubert and G. Paz, *Factorization at Subleading Power and Irreducible Uncertainties in  $\bar{B} \rightarrow X_s \gamma$  Decay*, *JHEP* **08** (2010) 099 [[1003.5012](#)].
- [110] M. Beneke, *Soft-collinear factorization in  $B$  decays*, *Nucl. Part. Phys. Proc.* **261-262** (2015) 311 [[1501.07374](#)].
- [111] C. Anastasiou, C. Duhr, F. Dulat, F. Herzog and B. Mistlberger, *Higgs Boson Gluon-Fusion Production in QCD at Three Loops*, *Phys. Rev. Lett.* **114** (2015) 212001 [[1503.06056](#)].
- [112] S. Jaskiewicz, *Next-to-leading power threshold factorization for Drell-Yan production*, in *14th International Symposium on Radiative Corrections: Application of Quantum Field Theory to Phenomenology*, 12, 2019, DOI [[1912.08882](#)].
- [113] I. Moulton, G. Vita and K. Yan, *Subleading power resummation of rapidity logarithms: the energy-energy correlator in  $\mathcal{N} = 4$  SYM*, *JHEP* **07** (2020) 005 [[1912.02188](#)].
- [114] I. Moulton, M.P. Solon, I.W. Stewart and G. Vita, *Fermionic Glauber Operators and Quark Reggeization*, *JHEP* **02** (2018) 134 [[1709.09174](#)].
- [115] A. Bhattacharya, A.V. Manohar and M.D. Schwartz, *Quark-gluon backscattering in the Regge limit at one-loop*, *JHEP* **02** (2022) 091 [[2110.02978](#)].
- [116] M.A. Ebert, A. Gao and I.W. Stewart, *Factorization for Azimuthal Asymmetries in SIDIS at Next-to-Leading Power*, [2112.07680](#).

- [117] I. Moulton, L. Rothen, I.W. Stewart, F.J. Tackmann and H.X. Zhu, *Subleading Power Corrections for  $N$ -Jettiness Subtractions*, *Phys. Rev. D* **95** (2017) 074023 [[1612.00450](#)].
- [118] R. Boughezal, X. Liu and F. Petriello, *Power Corrections in the  $N$ -jettiness Subtraction Scheme*, *JHEP* **03** (2017) 160 [[1612.02911](#)].
- [119] I. Moulton, L. Rothen, I.W. Stewart, F.J. Tackmann and H.X. Zhu,  *$N$ -jettiness subtractions for  $gg \rightarrow H$  at subleading power*, *Phys. Rev. D* **97** (2018) 014013 [[1710.03227](#)].
- [120] A. Bhattacharya, I. Moulton, I.W. Stewart and G. Vita, *Helicity Methods for High Multiplicity Subleading Soft and Collinear Limits*, *JHEP* **05** (2019) 192 [[1812.06950](#)].
- [121] R. Boughezal, A. Isgro and F. Petriello, *Next-to-leading power corrections to  $V + 1$  jet production in  $N$ -jettiness subtraction*, *Phys. Rev. D* **101** (2020) 016005 [[1907.12213](#)].
- [122] V.V. Sudakov, *Vertex parts at very high-energies in quantum electrodynamics*, *Sov. Phys. JETP* **3** (1956) 65.
- [123] C. Anastasiou, C. Duhr, F. Dulat, E. Furlan, T. Gehrmann, F. Herzog et al., *High precision determination of the gluon fusion Higgs boson cross-section at the LHC*, *JHEP* **05** (2016) 058 [[1602.00695](#)].
- [124] J.M. Lindert, K. Melnikov, L. Tancredi and C. Wever, *Top-bottom interference effects in Higgs plus jet production at the LHC*, *Phys. Rev. Lett.* **118** (2017) 252002 [[1703.03886](#)].
- [125] M.L. Czakon and M. Niggetiedt, *Exact quark-mass dependence of the Higgs-gluon form factor at three loops in QCD*, *JHEP* **05** (2020) 149 [[2001.03008](#)].
- [126] V.S. Fadin and L.N. Lipatov, *BFKL pomeron in the next-to-leading approximation*, *Phys. Lett. B* **429** (1998) 127 [[hep-ph/9802290](#)].
- [127] T. Liu, K. Melnikov and A.A. Penin, *Nonfactorizable QCD Effects in Higgs Boson Production via Vector Boson Fusion*, *Phys. Rev. Lett.* **123** (2019) 122002 [[1906.10899](#)].
- [128] F.A. Dreyer, A. Karlberg and L. Tancredi, *On the impact of non-factorisable corrections in VBF single and double Higgs production*, *JHEP* **10** (2020) 131 [[2005.11334](#)].
- [129] G.V. Frolov, V.N. Gribov and L.N. Lipatov, *On Regge poles in quantum electrodynamics*, *Phys. Lett. B* **31** (1970) 34.
- [130] M. Beneke, C. Bobeth and R. Szafron, *Enhanced electromagnetic correction to the rare  $B$ -meson decay  $B_{s,d} \rightarrow \mu^+ \mu^-$* , *Phys. Rev. Lett.* **120** (2018) 011801 [[1708.09152](#)].
- [131] M. Beneke, C. Bobeth and R. Szafron, *Power-enhanced leading-logarithmic QED corrections to  $B_q \rightarrow \mu^+ \mu^-$* , *JHEP* **10** (2019) 232 [[1908.07011](#)].
- [132] Z.L. Liu and M. Neubert, *Factorization at subleading power and endpoint-divergent convolutions in  $h \rightarrow \gamma\gamma$  decay*, *JHEP* **04** (2020) 033 [[1912.08818](#)].
- [133] Z.L. Liu, B. Mecaj, M. Neubert, X. Wang and S. Fleming, *Renormalization and Scale Evolution of the Soft-Quark Soft Function*, *JHEP* **07** (2020) 104 [[2005.03013](#)].
- [134] G.T. Bodwin, J.-H. Ee, J. Lee and X.-P. Wang, *Analyticity, renormalization, and evolution of the soft-quark function*, *Phys. Rev. D* **104** (2021) 016010 [[2101.04872](#)].
- [135] Z.L. Liu and M. Neubert, *Two-Loop Radiative Jet Function for Exclusive  $B$ -Meson and Higgs Decays*, *JHEP* **06** (2020) 060 [[2003.03393](#)].
- [136] Z.L. Liu, M. Neubert, M. Schnubel and X. Wang, *Radiative quark jet function with an external gluon*, *JHEP* **02** (2022) 075 [[2112.00018](#)].

- [137] Z.L. Liu, B. Mecaj, M. Neubert and X. Wang, *Factorization at subleading power, Sudakov resummation, and endpoint divergences in soft-collinear effective theory*, *Phys. Rev. D* **104** (2021) 014004 [[2009.04456](#)].
- [138] J. Currie, E.W.N. Glover and J. Pires, *Next-to-Next-to Leading Order QCD Predictions for Single Jet Inclusive Production at the LHC*, *Phys. Rev. Lett.* **118** (2017) 072002 [[1611.01460](#)].
- [139] M. Czakon, A. van Hameren, A. Mitov and R. Poncelet, *Single-jet inclusive rates with exact color at  $\mathcal{O}(\alpha_s^4)$* , *JHEP* **10** (2019) 262 [[1907.12911](#)].
- [140] M. Dasgupta, F. Dreyer, G.P. Salam and G. Soyez, *Small-radius jets to all orders in QCD*, *JHEP* **04** (2015) 039 [[1411.5182](#)].
- [141] X. Liu, S.-O. Moch and F. Ringer, *Threshold and jet radius joint resummation for single-inclusive jet production*, *Phys. Rev. Lett.* **119** (2017) 212001 [[1708.04641](#)].
- [142] D. Neill, F. Ringer and N. Sato, *Leading jets and energy loss*, *JHEP* **07** (2021) 041 [[2103.16573](#)].
- [143] A.J. Larkoski, I. Moult and B. Nachman, *Jet Substructure at the Large Hadron Collider: A Review of Recent Advances in Theory and Machine Learning*, *Phys. Rept.* **841** (2020) 1 [[1709.04464](#)].
- [144] R. Kogler et al., *Jet Substructure at the Large Hadron Collider: Experimental Review*, *Rev. Mod. Phys.* **91** (2019) 045003 [[1803.06991](#)].
- [145] S. Marzani, G. Soyez and M. Spannowsky, *Looking inside jets: an introduction to jet substructure and boosted-object phenomenology*, vol. 958, Springer (2019), [10.1007/978-3-030-15709-8](#), [[1901.10342](#)].
- [146] M. Procura and I.W. Stewart, *Quark Fragmentation within an Identified Jet*, *Phys. Rev. D* **81** (2010) 074009 [[0911.4980](#)].
- [147] R. Bain, Y. Makris and T. Mehen, *Transverse Momentum Dependent Fragmenting Jet Functions with Applications to Quarkonium Production*, *JHEP* **11** (2016) 144 [[1610.06508](#)].
- [148] Z.-B. Kang, X. Liu, F. Ringer and H. Xing, *The transverse momentum distribution of hadrons within jets*, *JHEP* **11** (2017) 068 [[1705.08443](#)].
- [149] D. Neill, I. Scimemi and W.J. Waalewijn, *Jet axes and universal transverse-momentum-dependent fragmentation*, *JHEP* **04** (2017) 020 [[1612.04817](#)].
- [150] D. Neill, A. Papaefstathiou, W.J. Waalewijn and L. Zoppi, *Phenomenology with a recoil-free jet axis: TMD fragmentation and the jet shape*, *JHEP* **01** (2019) 067 [[1810.12915](#)].
- [151] D. Gutierrez-Reyes, Y. Makris, V. Vaidya, I. Scimemi and L. Zoppi, *Probing Transverse-Momentum Distributions With Groomed Jets*, *JHEP* **08** (2019) 161 [[1907.05896](#)].
- [152] *Les Houches 2017: Physics at TeV Colliders Standard Model Working Group Report*, 3, 2018.
- [153] S. Marzani, D. Reichelt, S. Schumann, G. Soyez and V. Theeuwes, *Fitting the Strong Coupling Constant with Soft-Drop Thrust*, *JHEP* **11** (2019) 179 [[1906.10504](#)].
- [154] R. Bain, L. Dai, A. Leibovich, Y. Makris and T. Mehen, *NRQCD Confronts LHCb Data on Quarkonium Production within Jets*, *Phys. Rev. Lett.* **119** (2017) 032002 [[1702.05525](#)].

- [155] Z.-B. Kang, J.-W. Qiu, F. Ringer, H. Xing and H. Zhang, *J/ψ production and polarization within a jet*, *Phys. Rev. Lett.* **119** (2017) 032001 [[1702.03287](#)].
- [156] A.H. Hoang, S. Mantry, A. Pathak and I.W. Stewart, *Extracting a Short Distance Top Mass with Light Grooming*, *Phys. Rev. D* **100** (2019) 074021 [[1708.02586](#)].
- [157] D. Krohn, J. Thaler and L.-T. Wang, *Jet Trimming*, *JHEP* **02** (2010) 084 [[0912.1342](#)].
- [158] S.D. Ellis, C.K. Vermilion and J.R. Walsh, *Recombination Algorithms and Jet Substructure: Pruning as a Tool for Heavy Particle Searches*, *Phys. Rev. D* **81** (2010) 094023 [[0912.0033](#)].
- [159] M. Dasgupta, A. Fregoso, S. Marzani and G.P. Salam, *Towards an understanding of jet substructure*, *JHEP* **09** (2013) 029 [[1307.0007](#)].
- [160] M. Cacciari, G.P. Salam and G. Soyez, *SoftKiller, a particle-level pileup removal method*, *Eur. Phys. J. C* **75** (2015) 59 [[1407.0408](#)].
- [161] A.J. Larkoski, S. Marzani, G. Soyez and J. Thaler, *Soft Drop*, *JHEP* **05** (2014) 146 [[1402.2657](#)].
- [162] A.J. Larkoski, S. Marzani and J. Thaler, *Sudakov Safety in Perturbative QCD*, *Phys. Rev. D* **91** (2015) 111501 [[1502.01719](#)].
- [163] P. Cal, K. Lee, F. Ringer and W.J. Waalewijn, *The soft drop momentum sharing fraction  $z_g$  beyond leading-logarithmic accuracy*, [2106.04589](#).
- [164] ATLAS collaboration, *Measurement of soft-drop jet observables in pp collisions with the ATLAS detector at  $\sqrt{s} = 13$  TeV*, *Phys. Rev. D* **101** (2020) 052007 [[1912.09837](#)].
- [165] C. Frye, A.J. Larkoski, M.D. Schwartz and K. Yan, *Factorization for groomed jet substructure beyond the next-to-leading logarithm*, *JHEP* **07** (2016) 064 [[1603.09338](#)].
- [166] S. Marzani, L. Schunk and G. Soyez, *A study of jet mass distributions with grooming*, *JHEP* **07** (2017) 132 [[1704.02210](#)].
- [167] Z.-B. Kang, K. Lee, X. Liu and F. Ringer, *Soft drop groomed jet angularities at the LHC*, *Phys. Lett. B* **793** (2019) 41 [[1811.06983](#)].
- [168] A.J. Larkoski, *Improving the understanding of jet grooming in perturbation theory*, *JHEP* **09** (2020) 072 [[2006.14680](#)].
- [169] K. Benkendorfer and A.J. Larkoski, *Grooming at the cusp: all-orders predictions for the transition region of jet groomers*, *JHEP* **11** (2021) 188 [[2108.02779](#)].
- [170] S.D. Ellis, C.K. Vermilion, J.R. Walsh, A. Hornig and C. Lee, *Jet Shapes and Jet Algorithms in SCET*, *JHEP* **11** (2010) 101 [[1001.0014](#)].
- [171] A.J. Larkoski, D. Neill and J. Thaler, *Jet Shapes with the Broadening Axis*, *JHEP* **04** (2014) 017 [[1401.2158](#)].
- [172] Z.-B. Kang, K. Lee and F. Ringer, *Jet angularity measurements for single inclusive jet production*, *JHEP* **04** (2018) 110 [[1801.00790](#)].
- [173] S. Caletti, O. Fedkevych, S. Marzani, D. Reichelt, S. Schumann, G. Soyez et al., *Jet angularities in Z+jet production at the LHC*, *JHEP* **07** (2021) 076 [[2104.06920](#)].
- [174] D. Reichelt, S. Caletti, O. Fedkevych, S. Marzani, S. Schumann and G. Soyez, *Phenomenology of jet angularities at the LHC*, [2112.09545](#).
- [175] Z.-B. Kang, K. Lee, X. Liu, D. Neill and F. Ringer, *The soft drop groomed jet radius at NLL*, *JHEP* **02** (2020) 054 [[1908.01783](#)].

- [176] P. Cal, F. Ringer and W.J. Waalewijn, *The jet shape at NLL'*, *JHEP* **05** (2019) 143 [[1901.06389](#)].
- [177] P. Cal, D. Neill, F. Ringer and W.J. Waalewijn, *Calculating the angle between jet axes*, *JHEP* **04** (2020) 211 [[1911.06840](#)].
- [178] A.J. Larkoski, S. Marzani and C. Wu, *Theory Predictions for the Pull Angle*, *Phys. Rev. D* **99** (2019) 091502 [[1903.02275](#)].
- [179] Y. Bao and A.J. Larkoski, *Calculating Pull for Non-Singlet Jets*, *JHEP* **12** (2019) 035 [[1910.02085](#)].
- [180] A. Lifson, G.P. Salam and G. Soyez, *Calculating the primary Lund Jet Plane density*, *JHEP* **10** (2020) 170 [[2007.06578](#)].
- [181] Y. Mehtar-Tani, A. Soto-Ontoso and K. Tywoniuk, *Dynamical grooming of QCD jets*, *Phys. Rev. D* **101** (2020) 034004 [[1911.00375](#)].
- [182] P. Caucal, A. Soto-Ontoso and A. Takacs, *Dynamical Grooming meets LHC data*, *JHEP* **07** (2021) 020 [[2103.06566](#)].
- [183] A.J. Larkoski, I. Moult and D. Neill, *Factorization and Resummation for Groomed Multi-Prong Jet Shapes*, *JHEP* **02** (2018) 144 [[1710.00014](#)].
- [184] D. Napoletano and G. Soyez, *Computing  $N$ -subjettiness for boosted jets*, *JHEP* **12** (2018) 031 [[1809.04602](#)].
- [185] L. Cavallini, A. Coccaro, C.K. Khosa, G. Manco, S. Marzani, F. Parodi et al., *Tagging the Higgs boson decay to bottom quarks with colour-sensitive observables and the Lund jet plane*, [2112.09650](#).
- [186] H. Chen, I. Moult, X. Zhang and H.X. Zhu, *Rethinking jets with energy correlators: Tracks, resummation, and analytic continuation*, *Phys. Rev. D* **102** (2020) 054012 [[2004.11381](#)].
- [187] H. Chen, I. Moult and H.X. Zhu, *Quantum Interference in Jet Substructure from Spinning Gluons*, *Phys. Rev. Lett.* **126** (2021) 112003 [[2011.02492](#)].
- [188] Y. Li, I. Moult, S.S. van Velzen, W.J. Waalewijn and H.X. Zhu, *Extending Precision Perturbative QCD with Track Functions*, [2108.01674](#).
- [189] S. Caletti, O. Fedkevych, S. Marzani and D. Reichelt, *Tagging the initial-state gluon*, *Eur. Phys. J. C* **81** (2021) 844 [[2108.10024](#)].
- [190] Y.-T. Chien and I.W. Stewart, *Collinear Drop*, *JHEP* **06** (2020) 064 [[1907.11107](#)].
- [191] H.-y. Liu, X. Liu and S.-O. Moch, *Anti- $k_T$  jet function at next-to-next-to-leading order*, *Phys. Rev. D* **104** (2021) 014016 [[2103.08680](#)].
- [192] M. Cacciari, G.P. Salam and G. Soyez, *The anti- $k_t$  jet clustering algorithm*, *JHEP* **04** (2008) 063 [[0802.1189](#)].
- [193] A. Kardos, A.J. Larkoski and Z. Trócsányi, *Two- and three-loop data for the groomed jet mass*, *Phys. Rev. D* **101** (2020) 114034 [[2002.05730](#)].
- [194] A. Kardos, A.J. Larkoski and Z. Trócsányi, *Groomed jet mass at high precision*, *Phys. Lett. B* **809** (2020) 135704 [[2002.00942](#)].
- [195] A. Banfi, F.A. Dreyer and P.F. Monni, *Next-to-leading non-global logarithms in QCD*, *JHEP* **10** (2021) 006 [[2104.06416](#)].

- [196] Z.-B. Kang, X. Liu, S. Mantry and D.Y. Shao, *Jet Charge: A Flavor Prism for Spin Asymmetries at the EIC*, *Phys. Rev. Lett.* **125** (2020) 242003 [2008.00655].
- [197] P. Hinderer, M. Schlegel and W. Vogelsang, *Single-Inclusive Production of Hadrons and Jets in Lepton-Nucleon Scattering at NLO*, *Phys. Rev. D* **92** (2015) 014001 [1505.06415].
- [198] G. Abelo, R. Boughezal, X. Liu and F. Petriello, *Single-inclusive jet production in electron-nucleon collisions through next-to-next-to-leading order in perturbative QCD*, *Phys. Lett. B* **763** (2016) 52 [1607.04921].
- [199] R. Boughezal, F. Petriello and H. Xing, *Inclusive jet production as a probe of polarized parton distribution functions at a future EIC*, *Phys. Rev. D* **98** (2018) 054031 [1806.07311].
- [200] I. Borsa, D. de Florian and I. Pedron, *Inclusive-jet and dijet production in polarized deep inelastic scattering*, *Phys. Rev. D* **103** (2021) 014008 [2010.07354].
- [201] I. Borsa, D. de Florian and I. Pedron, *Dijet Production in Neutral Current and Charged Current Polarized Deep Inelastic Scattering*, [2112.08223](#).
- [202] X. Liu, F. Ringer, W. Vogelsang and F. Yuan, *Lepton-jet Correlations in Deep Inelastic Scattering at the Electron-Ion Collider*, *Phys. Rev. Lett.* **122** (2019) 192003 [1812.08077].
- [203] R.D. Field and R.P. Feynman, *A Parametrization of the Properties of Quark Jets*, *Nucl. Phys. B* **136** (1978) 1.
- [204] D. Krohn, M.D. Schwartz, T. Lin and W.J. Waalewijn, *Jet Charge at the LHC*, *Phys. Rev. Lett.* **110** (2013) 212001 [1209.2421].
- [205] M. Arratia, Y. Song, F. Ringer and B.V. Jacak, *Jets as precision probes in electron-nucleus collisions at the future Electron-Ion Collider*, *Phys. Rev. C* **101** (2020) 065204 [1912.05931].
- [206] D. Gutierrez-Reyes, I. Scimemi, W.J. Waalewijn and L. Zoppi, *Transverse momentum dependent distributions with jets*, *Phys. Rev. Lett.* **121** (2018) 162001 [1807.07573].
- [207] X. Liu and H. Xing, *The Time-reversal Odd Side of a Jet*, [2104.03328](#).
- [208] R.F. del Castillo, M.G. Echevarria, Y. Makris and I. Scimemi, *Transverse momentum dependent distributions in dijet and heavy hadron pair production at EIC*, [2111.03703](#).
- [209] Z.-B. Kang, K. Lee, D.Y. Shao and F. Zhao, *Spin asymmetries in electron-jet production at the future electron ion collider*, *JHEP* **11** (2021) 005 [2106.15624].
- [210] F. Yuan, *Azimuthal asymmetric distribution of hadrons inside a jet at hadron collider*, *Phys. Rev. Lett.* **100** (2008) 032003 [0709.3272].
- [211] M. Arratia, Z.-B. Kang, A. Prokudin and F. Ringer, *Jet-based measurements of Sivers and Collins asymmetries at the future electron-ion collider*, *Phys. Rev. D* **102** (2020) 074015 [2007.07281].
- [212] A. Gao, H.T. Li, I. Moutl and H.X. Zhu, *Precision QCD Event Shapes at Hadron Colliders: The Transverse Energy-Energy Correlator in the Back-to-Back Limit*, *Phys. Rev. Lett.* **123** (2019) 062001 [1901.04497].
- [213] M.D. Sievert and I. Vitev, *Quark branching in QCD matter to any order in opacity beyond the soft gluon emission limit*, *Phys. Rev. D* **98** (2018) 094010 [1807.03799].
- [214] J.a. Barata, Y. Mehtar-Tani, A. Soto-Ontoso and K. Tywoniuk, *Revisiting transverse momentum broadening in dense QCD media*, *Phys. Rev. D* **104** (2021) 054047 [2009.13667].

- [215] H.T. Li and I. Vitev, *Nuclear Matter Effects on Jet Production at Electron-Ion Colliders*, *Phys. Rev. Lett.* **126** (2021) 252001 [[2010.05912](#)].
- [216] Y.-Y. Zhang and X.-N. Wang, *Multiple parton scattering and gluon saturation in dijet production at EIC*, [2104.04520](#).
- [217] A. Accardi et al., *Electron Ion Collider: The Next QCD Frontier: Understanding the glue that binds us all*, *Eur. Phys. J. A* **52** (2016) 268 [[1212.1701](#)].
- [218] L.D. McLerran and R. Venugopalan, *Computing quark and gluon distribution functions for very large nuclei*, *Phys. Rev. D* **49** (1994) 2233 [[hep-ph/9309289](#)].
- [219] L.D. McLerran and R. Venugopalan, *Gluon distribution functions for very large nuclei at small transverse momentum*, *Phys. Rev. D* **49** (1994) 3352 [[hep-ph/9311205](#)].
- [220] L.D. McLerran and R. Venugopalan, *Green's functions in the color field of a large nucleus*, *Phys. Rev. D* **50** (1994) 2225 [[hep-ph/9402335](#)].
- [221] F. Gelis, E. Iancu, J. Jalilian-Marian and R. Venugopalan, *The Color Glass Condensate*, *Ann. Rev. Nucl. Part. Sci.* **60** (2010) 463 [[1002.0333](#)].
- [222] A.H. Mueller, *Small  $x$  Behavior and Parton Saturation: A QCD Model*, *Nucl. Phys. B* **335** (1990) 115.
- [223] A.H. Mueller, *Soft gluons in the infinite momentum wave function and the BFKL pomeron*, *Nucl. Phys. B* **415** (1994) 373.
- [224] I. Balitsky, *Operator expansion for high-energy scattering*, *Nucl. Phys. B* **463** (1996) 99 [[hep-ph/9509348](#)].
- [225] Y.V. Kovchegov, *Small  $x$   $F(2)$  structure function of a nucleus including multiple pomeron exchanges*, *Phys. Rev. D* **60** (1999) 034008 [[hep-ph/9901281](#)].
- [226] J. Jalilian-Marian, A. Kovner, A. Leonidov and H. Weigert, *The BFKL equation from the Wilson renormalization group*, *Nucl. Phys. B* **504** (1997) 415 [[hep-ph/9701284](#)].
- [227] J. Jalilian-Marian, A. Kovner, A. Leonidov and H. Weigert, *The Wilson renormalization group for low  $x$  physics: Towards the high density regime*, *Phys. Rev. D* **59** (1998) 014014 [[hep-ph/9706377](#)].
- [228] E. Iancu, A. Leonidov and L.D. McLerran, *Nonlinear gluon evolution in the color glass condensate. 1.*, *Nucl. Phys. A* **692** (2001) 583 [[hep-ph/0011241](#)].
- [229] E. Ferreiro, E. Iancu, A. Leonidov and L. McLerran, *Nonlinear gluon evolution in the color glass condensate. 2.*, *Nucl. Phys. A* **703** (2002) 489 [[hep-ph/0109115](#)].
- [230] I. Balitsky and L. Lipatov, *The Pomeron Singularity in Quantum Chromodynamics*, *Sov.J.Nucl.Phys.* **28** (1978) 822.
- [231] L.V. Gribov, E.M. Levin and M.G. Ryskin, *Semihard Processes in QCD*, *Phys. Rept.* **100** (1983) 1.
- [232] A.H. Mueller and J.-w. Qiu, *Gluon Recombination and Shadowing at Small Values of  $x$* , *Nucl. Phys. B* **268** (1986) 427.
- [233] K.J. Golec-Biernat and M. Wusthoff, *Saturation effects in deep inelastic scattering at low  $Q^2$  and its implications on diffraction*, *Phys. Rev. D* **59** (1998) 014017 [[hep-ph/9807513](#)].
- [234] BRAHMS collaboration, *On the evolution of the nuclear modification factors with rapidity*

- and centrality in  $d + Au$  collisions at  $s(NN)^{1/2} = 200\text{-GeV}$ , *Phys. Rev. Lett.* **93** (2004) 242303 [[nucl-ex/0403005](#)].
- [235] PHENIX collaboration, *Nuclear modification factors for hadrons at forward and backward rapidities in deuteron-gold collisions at  $s(NN)^{1/2} = 200\text{-GeV}$* , *Phys. Rev. Lett.* **94** (2005) 082302 [[nucl-ex/0411054](#)].
- [236] STAR collaboration, *Forward neutral pion production in  $p+p$  and  $d+Au$  collisions at  $s(NN)^{1/2} = 200\text{-GeV}$* , *Phys. Rev. Lett.* **97** (2006) 152302 [[nucl-ex/0602011](#)].
- [237] PHENIX collaboration, *Jet structure from dihadron correlations in  $d+Au$  collisions at  $s(NN)^{1/2} = 200\text{-GeV}$* , *Phys. Rev. C* **73** (2006) 054903 [[nucl-ex/0510021](#)].
- [238] STAR collaboration, *Two Particle Correlations at Forward Rapidity in STAR*, *Nucl. Phys. A* **854** (2011) 168 [[1008.3989](#)].
- [239] ALICE collaboration, *Neutral pion and  $\eta$  meson production in  $p\text{-Pb}$  collisions at  $\sqrt{s_{NN}} = 5.02\text{ TeV}$* , *Eur. Phys. J. C* **78** (2018) 624 [[1801.07051](#)].
- [240] A. Morreale and F. Salazar, *Mining for Gluon Saturation at Colliders*, *Universe* **7** (2021) 312 [[2108.08254](#)].
- [241] A. Dumitru, A. Hayashigaki and J. Jalilian-Marian, *The Color glass condensate and hadron production in the forward region*, *Nucl. Phys. A* **765** (2006) 464 [[hep-ph/0506308](#)].
- [242] T. Altinoluk and A. Kovner, *Particle Production at High Energy and Large Transverse Momentum - 'The Hybrid Formalism' Revisited*, *Phys. Rev. D* **83** (2011) 105004 [[1102.5327](#)].
- [243] G.A. Chirilli, B.-W. Xiao and F. Yuan, *One-loop Factorization for Inclusive Hadron Production in  $pA$  Collisions in the Saturation Formalism*, *Phys. Rev. Lett.* **108** (2012) 122301 [[1112.1061](#)].
- [244] G. Beuf, *NLO corrections for the dipole factorization of DIS structure functions at low  $x$* , *Phys. Rev. D* **85** (2012) 034039 [[1112.4501](#)].
- [245] G. Beuf, *Dipole factorization for DIS at NLO: Loop correction to the  $\gamma_{T,L}^* \rightarrow q\bar{q}$  light-front wave functions*, *Phys. Rev. D* **94** (2016) 054016 [[1606.00777](#)].
- [246] G. Beuf, *Dipole factorization for DIS at NLO: Combining the  $q\bar{q}$  and  $q\bar{q}g$  contributions*, *Phys. Rev. D* **96** (2017) 074033 [[1708.06557](#)].
- [247] R. Boussarie, A.V. Grabovsky, L. Szymanowski and S. Wallon, *On the one loop  $\gamma^{(*)} \rightarrow q\bar{q}$  impact factor and the exclusive diffractive cross sections for the production of two or three jets*, *JHEP* **11** (2016) 149 [[1606.00419](#)].
- [248] R. Boussarie, A.V. Grabovsky, D.Y. Ivanov, L. Szymanowski and S. Wallon, *Next-to-Leading Order Computation of Exclusive Diffractive Light Vector Meson Production in a Saturation Framework*, *Phys. Rev. Lett.* **119** (2017) 072002 [[1612.08026](#)].
- [249] I. Balitsky and G.A. Chirilli, *Photon impact factor and  $k_T$ -factorization for DIS in the next-to-leading order*, *Phys. Rev. D* **87** (2013) 014013 [[1207.3844](#)].
- [250] K. Roy and R. Venugopalan, *Inclusive prompt photon production in electron-nucleus scattering at small  $x$* , *JHEP* **05** (2018) 013 [[1802.09550](#)].
- [251] H.-Y. Liu, Y.-Q. Ma and K.-T. Chao, *Improvement for Color Glass Condensate factorization: single hadron production in  $pA$  collisions at next-to-leading order*, *Phys. Rev. D* **100** (2019) 071503 [[1909.02370](#)].



- [252] K. Roy and R. Venugopalan, *Extracting many-body correlators of saturated gluons with precision from inclusive photon+dijet final states in deeply inelastic scattering*, *Phys. Rev. D* **101** (2020) 071505 [[1911.04519](#)].
- [253] K. Roy and R. Venugopalan, *NLO impact factor for inclusive photon+dijet production in  $e + A$  DIS at small  $x$* , *Phys. Rev. D* **101** (2020) 034028 [[1911.04530](#)].
- [254] P. Caucal, F. Salazar and R. Venugopalan, *Dijet impact factor in DIS at next-to-leading order in the Color Glass Condensate*, *JHEP* **11** (2021) 222 [[2108.06347](#)].
- [255] A.M. Stasto, B.-W. Xiao and D. Zaslavsky, *Towards the Test of Saturation Physics Beyond Leading Logarithm*, *Phys. Rev. Lett.* **112** (2014) 012302 [[1307.4057](#)].
- [256] Z.-B. Kang and X. Liu, *Power Counting the Small- $x$  Observables*, [1910.10166](#).
- [257] H.-Y. Liu, Z.-B. Kang and X. Liu, *Threshold resummation for hadron production in the small- $x$  region*, *Phys. Rev. D* **102** (2020) 051502 [[2004.11990](#)].
- [258] S. Fleming, *The role of Glauber exchange in soft collinear effective theory and the Balitsky–Fadin–Kuraev–Lipatov Equation*, *Phys. Lett. B* **735** (2014) 266 [[1404.5672](#)].
- [259] S. Catani and M. Grazzini, *Infrared factorization of tree level QCD amplitudes at the next-to-next-to-leading order and beyond*, *Nucl. Phys. B* **570** (2000) 287 [[hep-ph/9908523](#)].
- [260] ATLAS collaboration, *Transverse momentum, rapidity, and centrality dependence of inclusive charged-particle production in  $\sqrt{s_{NN}} = 5.02$  TeV  $p + Pb$  collisions measured by the ATLAS experiment*, *Phys. Lett. B* **763** (2016) 313 [[1605.06436](#)].
- [261] Y. Shi, L. Wang, S.-Y. Wei and B.-W. Xiao, *Pursuing the Precision Study for Color Glass Condensate in Forward Hadron Productions*, [2112.06975](#).
- [262] G. Altarelli, R.K. Ellis, G. Martinelli and S.-Y. Pi, *Processes Involving Fragmentation Functions Beyond the Leading Order in QCD*, *Nucl. Phys. B* **160** (1979) 301.
- [263] A.H. Mueller, *Cut vertices and their renormalization: A generalization of the wilson expansion*, *Phys. Rev. D* **18** (1978) 3705.
- [264] J.C. Collins and D.E. Soper, *Parton Distribution and Decay Functions*, *Nucl. Phys. B* **194** (1982) 445.
- [265] W. Furmanski and R. Petronzio, *Lepton - Hadron Processes Beyond Leading Order in Quantum Chromodynamics*, *Z. Phys. C* **11** (1982) 293.
- [266] M. Dasgupta, F.A. Dreyer, G.P. Salam and G. Soyez, *Inclusive jet spectrum for small-radius jets*, *JHEP* **06** (2016) 057 [[1602.01110](#)].
- [267] Z.-B. Kang, F. Ringer and I. Vitev, *The semi-inclusive jet function in SCET and small radius resummation for inclusive jet production*, *JHEP* **10** (2016) 125 [[1606.06732](#)].
- [268] L. Dai, C. Kim and A.K. Leibovich, *Fragmentation of a Jet with Small Radius*, *Phys. Rev. D* **94** (2016) 114023 [[1606.07411](#)].
- [269] P.J. Rijken and W.L. van Neerven,  *$O(\alpha_s^{**2})$  contributions to the longitudinal fragmentation function in  $e^+ e^-$  annihilation*, *Phys. Lett. B* **386** (1996) 422 [[hep-ph/9604436](#)].
- [270] P.J. Rijken and W.L. van Neerven, *Higher order QCD corrections to the transverse and longitudinal fragmentation functions in electron - positron annihilation*, *Nucl. Phys. B* **487** (1997) 233 [[hep-ph/9609377](#)].

- [271] A. Mitov and S.-O. Moch, *QCD Corrections to Semi-Inclusive Hadron Production in Electron-Positron Annihilation at Two Loops*, *Nucl. Phys. B* **751** (2006) 18 [[hep-ph/0604160](#)].
- [272] M. Abele, D. de Florian and W. Vogelsang, *Approximate NNLO QCD corrections to semi-inclusive DIS*, *Phys. Rev. D* **104** (2021) 094046 [[2109.00847](#)].
- [273] D. Anderle, D. de Florian and Y. Rotstein Habarnau, *Towards semi-inclusive deep inelastic scattering at next-to-next-to-leading order*, *Phys. Rev. D* **95** (2017) 034027 [[1612.01293](#)].
- [274] A. Mitov, S. Moch and A. Vogt, *Next-to-Next-to-Leading Order Evolution of Non-Singlet Fragmentation Functions*, *Phys. Lett.* **B638** (2006) 61 [[hep-ph/0604053](#)].
- [275] S. Moch and A. Vogt, *On third-order timelike splitting functions and top-mediated Higgs decay into hadrons*, *Phys. Lett.* **B659** (2008) 290 [[0709.3899](#)].
- [276] A.A. Almasy, S. Moch and A. Vogt, *On the Next-to-Next-to-Leading Order Evolution of Flavour-Singlet Fragmentation Functions*, *Nucl. Phys.* **B854** (2012) 133 [[1107.2263](#)].
- [277] H. Chen, T.-Z. Yang, H.X. Zhu and Y.J. Zhu, *Analytic Continuation and Reciprocity Relation for Collinear Splitting in QCD*, *Chin. Phys. C* **45** (2021) 043101 [[2006.10534](#)].
- [278] A.H. Mueller, *On the Multiplicity of Hadrons in QCD Jets*, *Phys. Lett.* **B104** (1981) 161.
- [279] A.H. Mueller, *Multiplicity and Hadron Distributions in QCD Jets: Nonleading Terms*, *Nucl. Phys.* **B213** (1983) 85.
- [280] A. Bassetto, M. Ciafaloni, G. Marchesini and A.H. Mueller, *Jet Multiplicity and Soft Gluon Factorization*, *Nucl. Phys.* **B207** (1982) 189.
- [281] C.P. Fong and B.R. Webber, *One and two particle distributions at small  $x$  in QCD jets*, *Nucl. Phys. B* **355** (1991) 54.
- [282] Y.L. Dokshitzer, V.A. Khoze and S.I. Troian, *Inclusive particle spectra from QCD cascades*, *Int. J. Mod. Phys. A* **7** (1992) 1875.
- [283] V.A. Khoze and W. Ochs, *Perturbative QCD approach to multiparticle production*, *Int. J. Mod. Phys. A* **12** (1997) 2949 [[hep-ph/9701421](#)].
- [284] OPAL collaboration, *A Study of coherence of soft gluons in hadron jets*, *Phys. Lett. B* **247** (1990) 617.
- [285] Y.I. Azimov, Y.L. Dokshitzer, V.A. Khoze and S.I. Troyan, *Similarity of Parton and Hadron Spectra in QCD Jets*, *Z. Phys. C* **27** (1985) 65.
- [286] S. Albino, P. Bolzoni, B.A. Kniehl and A. Kotikov, *Fully double-logarithm-resummed cross sections*, *Nucl. Phys.* **B851** (2011) 86 [[1104.3018](#)].
- [287] C.H. Kom, A. Vogt and K. Yeats, *Resummed small- $x$  and first-moment evolution of fragmentation functions in perturbative QCD*, *JHEP* **10** (2012) 033 [[1207.5631](#)].
- [288] A. Vogt, *Resummation of small- $x$  double logarithms in QCD: semi-inclusive electron-positron annihilation*, *JHEP* **10** (2011) 025 [[1108.2993](#)].
- [289] D.P. Anderle, T. Kaufmann, M. Stratmann and F. Ringer, *Fragmentation Functions Beyond Fixed Order Accuracy*, *Phys. Rev.* **D95** (2017) 054003 [[1611.03371](#)].
- [290] D. Neill and F. Ringer, *Soft Fragmentation on the Celestial Sphere*, *JHEP* **06** (2020) 086 [[2003.02275](#)].

- [291] E.A. Kuraev, L.N. Lipatov and V.S. Fadin, *Multi - Reggeon Processes in the Yang-Mills Theory*, *Sov.Phys.JETP* **44** (1976) 443.
- [292] L. Lipatov, *Reggeization of the Vector Meson and the Vacuum Singularity in Nonabelian Gauge Theories*, *Sov.J.Nucl.Phys.* **23** (1976) 338.
- [293] E. Kuraev, L. Lipatov and V.S. Fadin, *The Pomeranchuk Singularity in Nonabelian Gauge Theories*, *Sov.Phys.JETP* **45** (1977) 199.
- [294] L. Lipatov, *The Bare Pomeron in Quantum Chromodynamics*, *Sov.Phys.JETP* **63** (1986) 904.
- [295] S. Catani, M. Ciafaloni and F. Hautmann, *High-energy factorization in QCD and minimal subtraction scheme*, *Phys. Lett.* **B307** (1993) 147.
- [296] S. Catani and F. Hautmann, *High-energy factorization and small  $x$  deep inelastic scattering beyond leading order*, *Nucl. Phys.* **B427** (1994) 475 [[hep-ph/9405388](#)].
- [297] H. Weigert, *Nonglobal jet evolution at finite  $N(c)$* , *Nucl.Phys.* **B685** (2004) 321 [[hep-ph/0312050](#)].
- [298] Y. Hatta, *Relating  $e+ e-$  annihilation to high energy scattering at weak and strong coupling*, *JHEP* **0811** (2008) 057 [[0810.0889](#)].
- [299] E. Avsar, Y. Hatta and T. Matsuo, *Soft gluons away from jets: Distribution and correlation*, *JHEP* **0906** (2009) 011 [[0903.4285](#)].
- [300] S. Caron-Huot, *Resummation of non-global logarithms and the BFKL equation*, *JHEP* **03** (2018) 036 [[1501.03754](#)].
- [301] S. Caron-Huot and M. Herranen, *High-energy evolution to three loops*, [1604.07417](#).
- [302] A. Banfi, G. Marchesini and G. Smye, *Away from jet energy flow*, *JHEP* **0208** (2002) 006 [[hep-ph/0206076](#)].
- [303] G. Marchesini and A. Mueller, *BFKL dynamics in jet evolution*, *Phys.Lett.* **B575** (2003) 37 [[hep-ph/0308284](#)].
- [304] G. Marchesini and E. Onofri, *Exact solution of BFKL equation in jet-physics*, *JHEP* **0407** (2004) 031 [[hep-ph/0404242](#)].
- [305] R.D. Ball and S. Forte, *All order running coupling BFKL evolution from GLAP (and vice-versa)*, *Nucl. Phys.* **B742** (2006) 158 [[hep-ph/0601049](#)].
- [306] M. Ciafaloni and D. Colferai, *Dimensional regularisation and factorisation schemes in the BFKL equation at subleading level*, *JHEP* **09** (2005) 069 [[hep-ph/0507106](#)].
- [307] M. Ciafaloni, D. Colferai, G.P. Salam and A.M. Stasto, *Minimal subtraction vs. physical factorisation schemes in small- $x$  QCD*, *Phys. Lett.* **B635** (2006) 320 [[hep-ph/0601200](#)].
- [308] Y.L. Dokshitzer, G. Marchesini and G.P. Salam, *Revisiting parton evolution and the large- $x$  limit*, *Phys. Lett.* **B634** (2006) 504 [[hep-ph/0511302](#)].
- [309] B. Basso and G.P. Korchemsky, *Anomalous dimensions of high-spin operators beyond the leading order*, *Nucl. Phys.* **B775** (2007) 1 [[hep-th/0612247](#)].
- [310] A. Jain, M. Procura and W.J. Waalewijn, *Fully-Unintegrated Parton Distribution and Fragmentation Functions at Perturbative  $k_T$* , *JHEP* **04** (2012) 132 [[1110.0839](#)].
- [311] W.J. Waalewijn, *Calculating the Charge of a Jet*, *Phys. Rev. D* **86** (2012) 094030 [[1209.3019](#)].

- [312] H.-M. Chang, M. Procura, J. Thaler and W.J. Waalewijn, *Calculating Track-Based Observables for the LHC*, *Phys. Rev. Lett.* **111** (2013) 102002 [[1303.6637](#)].
- [313] SLD collaboration, *Production of  $\pi^+$ ,  $\pi^-$ ,  $K^+$ ,  $K^-$ ,  $p$  and anti- $p$  in light ( $uds$ ),  $c$  and  $b$  jets from  $Z^0$  decays*, *Phys. Rev. D* **69** (2004) 072003 [[hep-ex/0310017](#)].
- [314] ALEPH collaboration, *Studies of QCD at  $e^+e^-$  centre-of-mass energies between 91-GeV and 209-GeV*, *Eur. Phys. J. C* **35** (2004) 457.
- [315] D. Neill, *The Fragmentation Spectrum from Space-Time Reciprocity*, *JHEP* **03** (2021) 081 [[2010.02934](#)].