Submitted to the Proceedings of the US Community Study on the Future of Particle Physics (Snowmass 2021)

Electroweak Precision Physics and Constraining New Physics

Conveners: Alberto Belloni^{*1}, Ayres Freitas^{†2}, Junping Tian^{‡3},

Juan Alcaraz Maestre⁴, Aram Apyan⁵, Bianca Azartash-Namin⁶, Paolo Azzurri⁷, Swagato Banerjee⁸, Jakob Beyer⁹, Saptaparna Bhattacharya¹⁰, Jorge de Blas¹¹, Alain Blondel¹², Daniel Britzger¹³, Mogens Dam¹⁴, Yong Du¹⁵, David d'Enterria¹⁶, Keisuke Fujii¹⁷, Christophe Grojean^{9,18}, Jiayin Gu¹⁹, Tao Han², Michael Hildreth²⁰, Adrián Irles²¹, Patrick Janot¹⁶, Daniel Jeans¹⁷, Mayuri Kawale⁶, Elham E Khoda²², Samuel Lane²³, Ian Lewis²³, Zhijun Liang²⁴, Jenny List⁹, Zhen Liu²⁵, Yang Ma², Victor Miralles²⁶, Takahiro Mizuno^{17,27}, Chilufya Mwewa²⁸, Meenakshi Narain²⁹, Luka Nedic³⁰, Mark S. Neubauer³¹, Chaitanya Paranjape^{32,33}, Michael Peskin³⁴, Roberto Petti³⁵, Marc-André Pleier²⁸, Roman Poeschl³⁶, Karolos Potamianos³⁰, Laura Reina³⁷, Jürgen Reuter⁹, Tania Robens³⁸, Philipp Roloff¹⁶, Michael Roney³⁹, Manqi Ruan²⁴, Richard Ruiz⁴⁰, Alex Schuy²², William Shepherd⁴¹, Daniel Stolarski³³, John Stupak⁶, Taikan Suehara⁴², Roberto Tenchini⁷, Alessandro Tricoli²⁸, Eleni Vryonidou⁴³, Marcel Vos²¹, Connor Waits⁶, Graham Wilson²³, Yongcheng Wu⁴⁴, Keping Xie², Siqi Yang⁴⁵, Tianyi Yang⁴⁶, Keita Yumino²⁷, Shuang-Yong Zhou^{47,48}, and Junjie Zhu⁴⁹

¹University of Maryland, College Park, MD, USA

²University of Pittsburgh, Pittsburgh, PA, USA

³ICEPP, University of Tokyo, Japan

⁴CIEMAT, Madrid, Spain

⁵Brandeis University, Waltham, MA, USA

⁶The University of Oklahoma, Norman, OK, USA

⁷INFN Pisa, Italy

⁸University of Louisville, Louisville, KY, USA

⁹Deutsches Elektronen-Synchrotron DESY, 22607 Hamburg, Germany

¹⁰Northwestern University, Evanston, IL, USA

¹¹CAFPE and Departamento de Física Teórica y del Cosmos, Universidad de Granada, Spain

¹²LPNHE, Paris, France

¹³Max-Planck-Institut für Physik, Munich, Germany

¹⁴Niels Bohr Institute, Copenhagen University, Denmark

¹⁵Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing, China

¹⁶CERN, Geneva, Switzerland

¹⁷IPNS, KEK, Tsukuba, Japan ¹⁸Humboldt-Universität zu Berlin, 12489 Berlin, Germany ¹⁹Fudan University, Shanghai, China ²⁰University of Notre Dame, Notre Dame, IN, USA ²¹IFIC (UV/CSIC) Valencia, 46980 Paterna, Spain ²²University of Washington, Seattle, WA, USA ²³University of Kansas, Lawrence, KS, USA ²⁴Institute of High Energy Physics, Chinese Academy of Sciences, Beijing, China ²⁵University of Minnesota, Minneapolis, MN, USA ²⁶INFN Rome. Italy ²⁷The Graduate University for Advanced Studies, SOKENDAI, Japan ²⁸Brookhaven National Laboratory, Upton, NY, USA ²⁹Brown University, Providence, RI, USA ³⁰Oxford University, Oxford OX1 3PJ, United Kingdom ³¹University of Illinois at Urbana-Champaign, IL, USA ³²Indian Institute of Technology (Indian School of Mines) Dhanbad, Jharkhand-826004, India ³³Ottawa-Carleton Institute for Physics, Carleton University, Ottawa, ON, Canada ³⁴SLAC National Accelerator Lab., Menlo Park, CA, USA ³⁵University of South Carolina, Columbia, SC, USA ³⁶IJCLab, CNRS / Université Paris-Saclay / Université Paris Cité, 91405 Orsay cedex, France ³⁷Florida State University, Tallahassee, FL, USA ³⁸Institut Ruder Boskovic, 10000 Zagreb, Croatia ³⁹University of Victoria, Victoria, BC, Canada ⁴⁰Institute of Nuclear Physics, Polish Academy of Sciences (IFJ PAN), Kraków, Poland ⁴¹Sam Houston State University, Huntsville TX, USA ⁴²Kvushu University, Fukuoka 819Ž0130395, Japan ⁴³University of Manchester, Manchester M13 9PL, United Kingdom ⁴⁴Oklahoma State University, Stillwater, OK, USA ⁴⁵University of Science & Technology, Hefei, China ⁴⁶Peking University, Beijing, China ⁴⁷ICTS, University of Science & Technology, Hefei, China ⁴⁸Peng Huanwu Center for Fundamental Theory, Hefei, China ⁴⁹University of Michigan, Ann Arbor, MI, USA

Version 2.4; October 24, 2022

 $^{{\}rm *Alberto.Belloni@cern.ch}$

 $^{^{\}dagger}$ afreitas@pitt.edu

[‡]tian@icepp.s.u-tokyo.ac.jp

Contents

| 1 | Inti | roduction | 3 | | |
|----------|---|---|----|--|--|
| 2 | Electroweak precision tests at future colliders | | | | |
| | 2.1 | Current status of electroweak precision tests | 5 | | |
| | 2.2 | Electroweak precision measurements at future e^+e^- colliders | 8 | | |
| | 2.3 | Electroweak precision measurements at other facilities | 15 | | |
| | 2.4 | Theory needs for the interpretation of Electroweak precision data | 16 | | |
| 3 | Mu | lti-boson processes at high-energy colliders | 18 | | |
| | 3.1 | Theory studies on anomalous couplings | 19 | | |
| | 3.2 | Multi-boson processes at future lepton colliders | 20 | | |
| | 3.3 | Multi-boson processes at future hadron colliders | 22 | | |
| 4 | Glo | bal fits of new physics | 24 | | |
| | 4.1 | Framework and scope | 26 | | |
| | 4.2 | Collider scenarios and observables | 28 | | |
| | 4.3 | Results | 29 | | |
| 5 | Cor | nclusions | 42 | | |

1 Introduction

The precise measurement of physics observables and the test of their consistency within the standard model (SM) are an invaluable approach, complemented by direct searches for new physics, to determine the existence of physics beyond the standard model (BSM). Historically, the discovery of new particles (e.g., the W and Z bosons by the UA1 and UA2 collaborations [1–4]) has been followed by the construction of accelerator machines dedicated to the in-depth study of the new particles' features. After the discovery of a Higgs boson in 2012, there is no compelling theoretical argument or measurement result that predicts the mass scale of any BSM physics. The indirect search for new physics, which exploits off-shell and loop contributions of new particles, allows one to explore a much wider range of energy scales than those probed by direct searches in specific BSM scenarios. Such indirect BSM effects are typically inversely proportional to some power of the mass scale of the new degrees of freedom [5], so that high precision is crucial for probing large energy scales. The

achievable precision of an experiment is determined by the statistics of the collected data sample, the experimental and theoretical systematic uncertainties, and their correlations.

Studies of massive electroweak gauge bosons (W and Z bosons) are a promising target for indirect BSM searches, since the interactions of photons and gluons are strongly constrained by the unbroken gauge symmetries. They can be divided into two categories:

• Fermion scattering processes mediated by s- or t-channel W/Z bosons. These are known as *electroweak precision measurements*, since large-statistics samples can be produced at e^+e^- and $pp/p\bar{p}$ colliders. These measurements are sensitive to modifications of the gauge-boson–fermion couplings and the gauge-boson masses.

Electroweak precision tests at e^+e^- colliders benefit from the clean and controlled initial state, whereas hadron colliders are affected by large systematic uncertainties due to parton distributions functions and other QCD effects. Thus e^+e^- colliders have the potential to have a better sensitivity for electroweak precision measurements than hadron colliders, but a large integrated luminosity is crucial for that purpose.

Electroweak precision measurements will be covered in more detail in section 2, in particular the interplay of statistical, experimental systematic and theory uncertainties. It should be noted that it is very difficult to realistically predict the systematic uncertainties (both experimental and theory) of a future facility, since any uncertainty estimate is based on assumptions that can only be tested with data or by carrying out a certain theoretical calculation. Nevertheless, to fairly compare the potential of different proposed e^+e^- colliders, the systematic uncertainties should be based on the same assumptions for all these machines. Such a consistent treatment of systematic error estimates has been attempted in this document.

 Multi-boson processes, which include production of two or more vector bosons in fermion-antifermion annihilation, as well as vector boson scattering (VBS) processes. These processes can test modifications of gauge-boson self-interactions, and the sensitivity is typically improved with increased collision energy, so that hadron colliders tend to provide the strongest limits, although a future multi-TeV electron-positron or muon collider would also be very competitive.

A more extensive discussion of multi-boson physics at the high-luminosity run of the LHC (HL-LHC), at future higher-energy pp colliders, and at high-energy e^+e^- and $\mu^+\mu^-$ colliders is the topic of section 3.

A model-independent description of indirect BSM effects is given by an extension of the SM with higher-dimensional operators. The most common effective theory framework for this purpose is the Standard Model Effective Field Theory (SMEFT), which has the same field content and symmetries as the SM. The leading contributions to electroweak observables stem from operators of dimension 6, which are suppressed by Λ^{-2} , where Λ indicates an effective new physics scale.

Generally, even at the dimension-6 level, there are more operators than independent observables, so that additional assumptions (e.g., about flavor symmetries) are needed to

constrain the operator coefficient from the data. On the other hand, some of these operators also contribute to other phenomenological sectors of the SM, i.e., to Higgs physics or top physics, and measurements in these different sectors can help to break some parameter degeneracies. Thus it is advantageous to perform a *qlobal fit* of a large number of operators to a large number of observables from different sectors. In particular, such a global fit can be used to evaluate and compare the new physics reach of future experimental facilities.

Various global SMEFT fits of different scope are presented in section 4. Compared to previous such studies in the literature, the analysis in section 4 uses updated inputs for the expected statistical and systematic uncertainties of key measurements at future colliders. Furthermore, it also extends previous studies by including 4-fermion operators, which generate contact interactions contributing to processes like $e^+e^- \to ff$, and which can also modify the non-resonant background in Z-pole precision studies.

$\mathbf{2}$ Electroweak precision tests at future colliders

Precision measurements of the properties of W and Z bosons can be used to test the SM at the quantum level and to indirectly constrain potential BSM physics. The masses, widths and effective couplings of these gauge bosons can be modified through many different extensions of the SM, including new gauge interactions, extended Higgs sectors, composite Higgs scenarios, vector-like fermion fields, etc. (e.g., see section 10 of Ref. [6] for an overview).

2.1Current status of electroweak precision tests

An important class of electroweak precision measurements focuses on fermion-pair production processes, $e^+e^- \to f\bar{f}$ and $pp \to \ell^+\ell^-$. For electron-positron colliders with center-ofmass energies near the Z-boson mass, the dominant contribution to the cross section follows from the Z resonance, which can be approximately written as

$$\frac{d\sigma}{d\Omega}[e^{+}e^{-} \to f\overline{f}] \approx \frac{N_{c}^{f}s}{64\pi^{2}} \times \frac{(1 - P_{+}P_{-})[G_{1}(1 + c_{\theta}^{2}) + 2G_{3}c_{\theta}] + (P_{+} - P_{-})[H_{1}(1 + c_{\theta}^{2}) + 2H_{3}c_{\theta}]}{(s - m_{Z}^{2})^{2} + s^{2}\Gamma_{Z}^{2}/m_{Z}^{2}}, \quad (1)$$

where

$$G_1 = (v_e^2 + a_e^2)(v_f^2 + a_f^2),$$
 $G_3 = 4v_e a_e v_f a_f,$ (2)

$$G_1 = (v_e^2 + a_e^2)(v_f^2 + a_f^2), G_3 = 4v_e a_e v_f a_f, (2)$$

$$H_1 = 2v_e a_e (v_f^2 + a_f^2), H_3 = 2(v_e^2 + a_e^2) v_f a_f, (3)$$

where v_f and a_f are the effective vector and axial-vector couplings of the Z-boson to the fermion type f, respectively, $N_c^f = 1 \, (3)$ for leptons (quarks), and $P_{+/-}$ is the degree of longitudinal polarization of the positron/electron beam. Furthermore, $c_{\theta} \equiv \cos \theta$, where θ is the scattering angle.

The partial and total Z-boson widths can also be expressed in terms of these effective couplings,

$$\Gamma_Z = \sum_f \Gamma_Z^f, \qquad \Gamma_Z^f \approx \frac{N_c^f m_Z}{12\pi} (v_f^2 + a_f^2), \qquad (4)$$

which in turn lead to the following expression for the total peak cross-section:

$$\sigma[e^+e^- \to f\overline{f}]_{s=m_Z^2} \approx \frac{12\pi}{m_Z^2} \frac{\Gamma_Z^e \Gamma_Z^f}{\Gamma_Z^2}.$$
 (5)

The Z mass and total width can be determined from measurements of the cross-section lineshape at a few center-of-mass energies near the resonance peak. From measurements of cross sections with different final states one can determine ratios of the Z-boson partial widths. It is customary to express them in terms of

$$\sigma_{\rm had}^0 \equiv \sigma[e^+e^- \to {\rm had.}]_{s=m_Z^2}, \quad R_q \equiv \frac{\Gamma_q}{\Gamma_{\rm had}} \ (q=b,c), \quad R_\ell \equiv \frac{\Gamma_{\rm had}}{\Gamma_\ell} \ (\ell=e,\mu,\tau), \quad (6)$$

where "had" refers to all hadronic final states (i.e., the sum over u, d, c, s, b final states at the partonic level).

Ratios of the vector and axial-vector couplings can be extracted from the forward-backward asymmetry, the average polarization degree of produced τ leptons (for $f = \tau$), and the left-right asymmetry (for a polarized electron beam):

$$A_{\rm FB} \equiv \frac{\sigma_F - \sigma_B}{\sigma_F + \sigma_B} \approx \frac{3}{4} A_e A_f,$$

$$\langle P_\tau \rangle = A_\tau, \qquad A_f \equiv \frac{2v_f a_f}{v_f^2 + a_f^2}.$$

$$A_{\rm LR} \equiv \frac{\sigma_L - \sigma_R}{\sigma_L + \sigma_R} \approx A_e,$$
(7)

Here σ_F and σ_B refer to the cross section for only positive and negative values of $\cos \theta$, respectively; whereas σ_L and σ_R denote the cross section for left-handed $(P_- < 0)$ and right-handed $(P_- > 0)$ electrons (assuming $P_+ = 0$). The ratio v_f/a_f is also related to the effective weak mixing angle

$$\sin^2 \theta_{\text{eff}}^f \equiv \frac{1}{4|Q_f|} \left(1 - \frac{v_f}{a_f} \right). \tag{8}$$

The expressions above do not include the contributions stemming from photon-exchange and box diagrams and from radiative corrections that cannot be absorbed into the effective couplings, in particular initial-state and final-state radiation. These effects need to be predicted from theory and subtracted from the data in order to extract "measured" values of the quantities in eqs. (6). The latter are therefore known as electroweak *pseudo-observables* (EWPOs).

| EWPO Uncertainties | Current | HL-LHC |
|--|----------------------|--------|
| $\Delta m_W \; ({ m MeV})$ | $12 / 9.4^{\dagger}$ | 5 |
| $\Delta m_Z \; ({ m MeV})$ | 2.1 | |
| $\Delta\Gamma_Z \; ({ m MeV})$ | 2.3 | |
| $\Delta m_t \; ({\rm GeV})$ | 0.6* | 0.2 |
| $\Delta \sin^2 \theta_{\text{eff}}^{\ell} \times 10^{5}$ | 13 | < 10 |
| $\delta R_{\mu} \ (\times 10^3)$ | 1.6 | |
| $\delta R_b \ (\times 10^3)$ | 3.1 | |

[†] The recent W mass measurement from CDF with 9.4 MeV precision [13] has not yet been included in the global average [6].

Table 1: The current precision of a few selected EWPOs, based on data from LEP, SLC, TeVatron and LHC [6], and expected improvements from the HL-LHC [15]. Δ (δ) stands for absolute (relative) uncertainty.

Other EWPOs include the W-boson mass (m_W) and branching ratios, as well as the Fermi constant of muon decay, G_F . The latter is a key ingredient for predicting m_W in the SM, based on the relation

$$\frac{G_F}{\sqrt{2}} = \frac{\pi \alpha}{2m_W^2 (1 - m_W^2 / m_Z^2)} (1 + \Delta r),\tag{9}$$

where Δr describes higher-order corrections. G_F is currently known with a precision of 0.5 ppm [6], which may be further improved in the future, and thus it is a negligible source of uncertainty.

Moreover, when comparing experimental values for the EWPOs to theory predictions in the SM, other SM parameters are needed as inputs for the latter. Specifically, the mass of the top quark and the Higgs boson play an important role, as well as the strong coupling constant α_s and the shift due to the running of the fine structure constant from the Thomson limit to the Z scale, $\Delta\alpha \equiv 1 - \frac{\alpha(0)}{\alpha(m_Z)}$. $\Delta\alpha$ receives contributions from leptons, which can be computed perturbatively [7], and from hadronic states. The hadronic part can be split into non-perturbative and perturbative contributions. The non-perturbative contributions can be extracted from data for $e^+e^- \to \text{had}$. [8–10] or from lattice QCD simulations [11,12] using a dispersive approach. The data-driven methods are currently more precise, with an uncertainty for $\Delta\alpha_{\text{had}}$ of about 10^{-4} [8–10].

Reducing the uncertainty of $\Delta\alpha_{\rm had}$ requires improved measurements of $e^+e^-\to {\rm had.}$ for energies below 2 GeV (e.g., with ongoing measurements at VEPP-2000 and BEPC-II), 4-loop perturbative QCD corrections, and more precise determinations of the charm and bottom quark masses. With these improvements, an uncertainty of $< 0.5 \times 10^{-4}$ appears within reach [8]. Similarly, the lattice QCD evaluation of $\Delta\alpha_{\rm had}$ is expected to continue to improve, but quantitative estimates are currently not available.

^{*} This value includes an additional uncertainty due to ambiguities in the top mass definition (see EF TOPHF report [14] for more details).

The current precision for a few selected EWPOs is listed in Tab. 1. Most of these results stem from measurements at the e^+e^- colliders LEP and SLC, but the hadron colliders TeVatron and LHC contribute important results for $\sin^2 \theta_{\text{eff}}^{\ell}$, m_W and m_t .

The HL-LHC with integrated luminosity of 3000 fb⁻¹ can make improved measurements of certain EWPOs. The effective weak mixing angle can be extracted from measurements of the forward-backward asymmetry in Drell-Yan production, $pp \to \ell^+\ell^-$ ($\ell = e, \mu$). The measurement precision is mostly limited by uncertainties of the parton distribution functions (PDFs), but the PDFs can be constrained simultaneously with the weak mixing angle determination through Drell-Yan data. In particular, the $m_{\ell\ell}$ distribution can be useful in disentangling the effect of PDFs from the weak mixing angle determination [16]. It is estimated that the total uncertainty of $\sin^2 \theta_{\text{eff}}^{\ell}$ can be reduced below 10^{-4} at HL-LHC [15].

Similarly, the W-boson mass can be extracted from measurements of $pp \to \ell\nu$, by performing fits to the lepton p_T and transverse mass distributions. This measurement benefits from a dedicated run with low instantaneous luminosity to improve the accurate reconstruction of the missing transverse momentum. Again PDF uncertainties are expected to dominate, and an ultimate precision of about 5 MeV appears achievable [15].

It should be noted that these precision measurements at the HL-LHC will rely on detailed theory input, including higher-order EW and mixed QCD×EW corrections [17–19], as well as resummation for low p_T (e.g., see Ref. [20] and references therein). Moreover, the extraction of $\sin^2\theta_{\rm eff}^{\ell}$ assumes that the dependence of $\sin^2\theta_{\rm eff}^f$ on different fermion flavor f is small and as predicted in the SM*.

2.2 Electroweak precision measurements at future e^+e^- colliders

Future high-luminosity e^+e^- colliders proposed as Higgs factories can also be used to study the masses and interactions of electroweak bosons to much higher precision than before. We here focus on four collider proposals: ILC [21–23], CLIC [24, 25], FCC-ee [26, 27], and CEPC [28–30]. Table 2 summarizes the run scenarios considered for these colliders within the Snowmass 2021 study. The 50 MW upgrade of CEPC [30] is assumed for all quantitative analyses throughout this document. The recent Cool Copper Collider (C³) [31,32] proposal has parameters very similar to ILC and will not be discussed separately in what follows.

The linear collider projects ILC and CLIC feature polarized electron beams (and also a polarized positron beam in the case of ILC). Two options are considered for ILC, the default option with center-of-mass energies of 250 GeV and above, and the "GigaZ" option that includes a run at the Z pole. [Note that a Z-pole run is also considered as a possible option for CLIC [34].] The ILC and CLIC runs with 500 GeV and above are irrelevant for "canonical" electroweak precision studies (*i.e.*, not considering multi-gauge-boson processes).

The circular colliders (FCC-ee and CEPC) can deliver very large integrated luminosities on the Z pole, yielding samples of $\mathcal{O}(10^{12})$ events. On the other hand, the standard run

^{*}In other words, this measurement can serve as a high-precision consistency check of the SM, but it will be difficult to interpret an observed deviation from the SM without model assumptions.

| Collider | \sqrt{s} | P [%] | $L_{ m int}$ |
|-----------|-----------------|-------------------|--------------|
| | | e^-/e^+ | ab^{-1} |
| ILC | $250~{\rm GeV}$ | $\pm 80 / \pm 30$ | 2 |
| | 350 GeV | $\pm 80/ \pm 30$ | 0.2 |
| | 500 GeV | $\pm 80/ \pm 30$ | 4 |
| | 1 TeV | $\pm 80/ \pm 20$ | 8 |
| ILC-GigaZ | m_Z | $\pm 80 / \pm 30$ | 0.1 |
| CLIC | $380~{\rm GeV}$ | $\pm 80/0$ | 1 |
| | 500 GeV | $\pm 80/0$ | 2.5 |
| | 1 TeV | $\pm 80/0$ | 5 |
| CEPC | m_Z | | 60 / 100 |
| | $2m_W$ | | 3.6 / 6 |
| | $240~{\rm GeV}$ | | 12 / 20 |
| | $2m_t$ | | - / 1 |
| FCC-ee | m_Z | | 150 |
| | $2m_W$ | | 10 |
| | $240~{\rm GeV}$ | | 5 |
| | $2m_t$ | | 1.5 |

Table 2: Electron-positron collider run scenarios used for the Snowmass 2021 study. The two sets of numbers for CEPC refer to the 30 MW baseline and 50 MW upgrade for the beam power. Also see section 1.3 and table 1-1 in the main EF report [33].

scenarios for ILC (without the GigaZ option) and CLIC do not include any run at the Z pole. Instead, precision studies of the Z boson are possible through the radiative return method, i.e., by producing Z bosons together with one or more initial-state photons, $e^+e^- \to Z + n\gamma$. The photons are emitted predominantly at low angles and lost in the beam pipe. However, the requirement of hard photon emission reduces the event yield and thus the achievable precision.

Given the large statistics of these future e^+e^- colliders, systematic uncertainties may have a significant impact on the achievable precision. In the following we discuss the dominant machine-specific systematic error sources, as well as uncertainties that are common to all machines. It should be emphasized that these systematic error evaluations are just order-of-magnitude estimates, while a more precise assessment would require instrumentation detail and tools that are not available at this time.

Common systematics: Uncertainties due to the physics modeling affect all collider proposals equally. Previous publications by the collider collaborations [28, 35–37] have made differing assumptions for the size and relevance of these common uncertainties. This situation creates problems for the comparison of the new-physics reach between different machines. Therefore, as part of the Snowmass 2021 process, a consistent set of assumptions

is being used and applied uniformly to all e^+e^- collider proposals.

For branching ratios of heavy-quark (b and c) final states, the tagging efficiency can be controlled in situ by comparing single and double tag rates. However, the simple scaling $\epsilon_{2\text{tag}} = (\epsilon_{1\text{tag}})^2$ gets modified by so-called hemisphere correlations. These correlations can be produced by detector effects, vertex fitting, and QCD effects. The first two sources can be reduced to a sub-dominant level through the availability of large-statistics calibration samples and the increased vertex precision of modern vertex detectors. The most important QCD effect is gluon splitting into a heavy-flavor $q\bar{q}$ pair. The contamination from gluon splitting can be reduced with acolinearity cuts between the two tagged jets. Moreover, the large available data sets can be used to dramatically improve the modeling of gluon splitting, but this will require parallel improvements in Monte-Carlo (MC) simulation tools. Here it is assumed that the QCD uncertainty can be improved by about one order magnitude compared to LEP [38], leading to relative uncertainties of 0.2×10^{-3} for R_b and 1×10^{-3} for R_c , respectively.

Similarly, QCD effects are a dominant source of uncertainty for determinations of A_b (A_c) from the forward-backward asymmetry of $e^+e^- \to b\bar{b}$ ($c\bar{c}$). QCD radiation can change the angular distributions and correlations of the heavy-quark jets, which in turn modified the observable asymmetry. This has been studied in detail in Ref. [39], where it was found that the impact of QCD effects can be substantially reduced with an acolinearity cut. With a moderate acolinearity cut and assuming NNLO QCD corrections, the relative error on $A_{b,c}$ due to missing higher-order perturbative QCD contributions is estimated to be about 3×10^{-4} (see Tab. 9 in Ref. [39]). With future work on QCD calculations this can likely be reduced to the level of 1×10^{-4} . In addition, one needs to consider non-perturbative hadronization and showering uncertainties (see also Ref. [40]). Due to wealth of available data at any of the proposed colliders, a significant improvement of the hadronization and showering models should be possible. Assuming an improvement of a factor 5 compared to currently available MC tunes (see Tab. 9 in Ref. [39]), this leads to an estimated relative error of 2×10^{-4} . Combining perturbative and non-perturbative uncertainties, the total absolute error due to QCD effects amounts to 2.1×10^{-4} for A_b and 1.5×10^{-4} for A_c .

Experimental systematic for linear colliders: For electroweak precision measurements at ILC250 or CLIC380 using the radiative return method, signal events need to be selected based on the invariant mass m_{ff} of two fermions from $Z \to f\bar{f}$. m_{ff} can be reconstructed using the polar angles of the fermions [41], which can be measured very precisely, so that this becomes a negligible source of systematic uncertainty. Note that multi-photon emission produces a tail in the reconstructed m_{ff} distribution, but this dilution does not diminish the precision of the overall energy scale calibration.

Combining this method with a precise calibration of the tracker momentum scale using large samples of kaon and Λ baryon decays, it may be possible to determine the Z mass and width at ILC250 with a systematic uncertainty of 2 ppm [42].

Many measurements at linear e^+e^- colliders profit from polarized beams, which in turn makes the polarization calibration a leading source of systematic uncertainties. Both

the ILC and CLIC designs expect that the luminosity-weighted long-term average of the polarization can be controlled to 0.1% [23,24]. However, as demonstrated in Ref. [43], the impact of the polarization uncertainty can be further reduced by treating the polarization values as nuisance parameters in the actual extraction of physics parameters from a set of observables.

The asymmetry parameter A_e can be determined from the left-right asymmetry A_{LR} (see Ref. [44] for a full simulation study), while A_f for other fermion types $(f = \mu, \tau, b, c)$ can be obtained from the left-right-forward-backward asymmetry for the process $e^+e^- \to f\overline{f}$,

$$A_{\rm LR,FB} = \frac{\sigma_{LF} - \sigma_{LB} - \sigma_{RF} + \sigma_{RB}}{\sigma_{LF} + \sigma_{LB} + \sigma_{RF} + \sigma_{RB}} \approx \frac{3}{4} A_f \tag{10}$$

The polarization calibration leads to a relative systematic uncertainty of 3×10^{-4} . Other important systematic uncertainties include the control of the luminosity and detector acceptance between runs with different polarization, which are estimated to be subdominant.

For the branching ratios R_i , the dominant source of uncertainty stems from the flavor identification, which is estimated at the level of 0.1% [23].

Measurements of $e^+e^- \to W^+W^-$ at ILC250 or CLIC380 can yield information about a variety of properties of the W bosons, including anomalous gauge-boson couplings. The W mass can be determined from a variety of kinematic final-state observables [23]: (1) constrained reconstruction of $qq\ell\nu$ events; (2) di-jet invariant mass for semi-leptonic and all-hadronic final states; (3) endpoints of the lepton energy spectrum for di-lepton ($\ell\nu\ell\nu$) and semi-leptonic ($qq\ell\nu$) final states; and (4) approximate kinematic reconstruction of di-lepton events by assuming that the event has a planar topology ("pseudo-mass" method). With an integrated luminosity of a few ab⁻¹ at ILC250, a statistical uncertainty of 0.5 MeV on m_W can be achieved [23]. The systematic uncertainty was estimated in the Snowmass 2013 study with 2.4 MeV [45], which receives comparable contributions from the beamenergy calibration, luminosity spectrum, modeling of hadronization, modeling of radiative corrections, and detector energy calibration. With improved detectors and methods for addressing the other systematic issues, a total error of 1 MeV at ILC250 may well be feasible.

An ILC run on the Z pole (ILC-GigaZ) would yield a higher-statistics sample of clean Z events, thus leading to improved overall precision for EWPOs. For the asymmetry parameters A_f , the systematic errors are again dominated by the polarization uncertainty, whereas the acceptance is the leading source of systematics for the branching ratios R_f [35].

Experimental systematic for circular colliders: The beam energy at circular colliders can be controlled with high accuracy using resonant depolarization, leading to an absolute precision of 100 keV (for $\sqrt{s} \sim 100$ GeV) and a point-to-point precision of 25 keV (i.e. the accuracy with which the energy difference between two nearby center-of-mass energies can be determined). These two numbers are the leading systematic uncertainties for the determination of the Z mass and width, respectively.

For the determination of A_e , it is advantageous to consider the forward-backward tau

polarization in $e^+e^- \to \tau^+\tau^-$,

$$\langle P_{\tau, \text{FB}} \rangle \equiv \frac{\langle P_{\tau} \rangle_F - \langle P_{\tau} \rangle_B}{\langle P_{\tau} \rangle_F + \langle P_{\tau} \rangle_B} \approx \frac{3}{4} A_e \,.$$
 (11)

This quantity is independent of the tau polarization distributions and of hemisphere migration effects. It would only be affected by correlations between these two effects, which are expected to be very small. As a result, the dominant systematic uncertainty would instead stem from non-tau backgrounds. These are estimated from the statistics of control samples used for calibrating the background, leading to an error estimate of 2×10^{-5} .

Other A_f parameters can be determined from $A_{\rm FB}$ for $e^+e^- \to f\overline{f}$. The main systematic uncertainty for A_μ is point-to-point control of the luminosity and detector acceptance. For A_c and A_b the dominant systematic error stems from QCD effects (see "Common systematics" above). While A_τ could also be obtained from the forward-backward asymmetry, a more precise determination is possible from the tau polarization. The main systematic uncertainty for the polarization measurement is due to the modeling of the hadronic tau decay modes. Since these are expected to be substantially improved by using the large available calibration samples at FCC-ee/CEPC, it is expected that this error is reduced by a factor 10 compared to LEP [38], leading to uncertainty of 2×10^{-4} .

The measurement of the total peak cross-section, $\sigma_{\rm had}^0$, is limited by the luminosity calibration. Using low-angle Bhabha events, a relatively precision of 10^{-4} or better should be achievable. For the hadronic branching fractions $R_{b,c}$, QCD uncertainties from gluon splitting are the dominant error source (see "Common systematics" above). For the leptonic branching fractios $R_{e,\mu,\tau}$ the lepton acceptance and beam energy control will be important factors. R_e is additionally affected by the subtraction of Bhabha backgrounds.

The W mass and width can be extracted with high precision from measurements at a few energy points near the WW threshold. As mentioned above, the dominant systematic uncertainty due to the beam energy calibration can be controlled using resonant depolarization, but with a slightly lower precision (0.0002%) at this center-of-mass energy compared to the Z pole.

Summary: A summary of projected statistical and systematic uncertainties for the different proposed e^+e^- colliders is given in Tab. 3. This table also serves as an input for the global fits presented in section 4.

Measurements of leptonic branching ratios R_{ℓ} ($\ell = e, \mu, \tau$) can be used to extract a precise value for the strong coupling constant α_s , which enters through final-state radiative corrections in Γ_{had} . Given that R_{ℓ} is a highly inclusive quantity, this determination of α_s is essentially free of non-perturbative QCD effects, so that a robust $\mathcal{O}(10^{-4})$ precision is achievable. However, it should be noted that this method assumes the validity of the SM, but the Z decay ratios may in general be modified by BSM physics. Similar considerations apply to the determination of α_s from the leptonic branching fraction of W bosons. For more information about future determinations of α_s , see Ref. [48].

As mentioned in the previous subsection, the shift $\Delta \alpha$ between the $\alpha(m_Z)$ and $\alpha(0)$

| Quantity | current | ILC250 | ILC-GigaZ | FCC-ee | CEPC | CLIC380 |
|---|---------|-----------|-----------|---|-------------------|---|
| $\Delta \alpha(m_Z)^{-1} \ (\times 10^3)$ | 17.8* | 17.8* | | 3.8 (1.2) | 17.8* | |
| $\Delta m_W \; ({ m MeV})$ | 12* | 0.5(2.4) | | 0.25(0.3) | 0.35(0.3) | |
| $\Delta m_Z \; ({ m MeV})$ | 2.1* | 0.7(0.2) | 0.2 | 0.004(0.1) | 0.005(0.1) | 2.1* |
| $\Delta m_H \; ({ m MeV})$ | 170* | 14 | | 2.5(2) | 5.9 | 78 |
| $\Delta\Gamma_W \; ({ m MeV})$ | 42* | 2 | | 1.2(0.3) | 1.8 (0.9) | |
| $\Delta\Gamma_Z \; ({ m MeV})$ | 2.3* | 1.5(0.2) | 0.12 | $0.004 \ (0.025)$ | $0.005 \ (0.025)$ | 2.3* |
| $\Delta A_e \times 10^5$ | 190* | 14(4.5) | 1.5(8) | 0.7(2) | 1.5(2) | $\begin{bmatrix} -60 \\ (15) \end{bmatrix}$ |
| $\Delta A_{\mu} \ (\times 10^5)$ | 1500* | 82 (4.5) | 3 (8) | 2.3(2.2) | 3.0 (1.8) | 390 (14) |
| $\Delta A_{\tau} \ (\times 10^5)$ | 400* | 86 (4.5) | 3 (8) | 0.5(20) | 1.2 (20) | 550 (14) |
| $\Delta A_b \ (\times 10^5)$ | 2000* | 53 (35) | 9 (50) | 2.4(21) | 3 (21) | 360 (92) |
| $\Delta A_c \ (\times 10^5)$ | 2700* | 140 (25) | 20(37) | 20 (15) | 6 (30) | 190 (67) |
| $\Delta \sigma_{\rm had}^{\bar{0}}$ (pb) | 37* | | | $\begin{bmatrix} -0.035(4) \end{bmatrix}$ | 0.05(2) | $[-\bar{3}\bar{7^*}^-]$ |
| $\delta R_e \ (\times 10^3)$ | 2.4* | 0.5(1.0) | 0.2(0.5) | 0.004(0.3) | 0.003(0.2) | 2.5 (1.0) |
| $\delta R_{\mu} \ (\times 10^3)$ | 1.6* | 0.5(1.0) | 0.2(0.2) | $0.003 \ (0.05)$ | 0.003(0.1) | 2.5 (1.0) |
| $\delta R_{\tau} \ (\times 10^3)$ | 2.2* | 0.6 (1.0) | 0.2(0.4) | 0.003(0.1) | 0.003(0.1) | 3.3 (5.0) |
| $\delta R_b \ (\times 10^3)$ | 3.1* | 0.4 (1.0) | 0.04(0.7) | 0.0014 (< 0.3) | 0.005 (0.2) | 1.5 (1.0) |
| $\delta R_c(\times 10^3)$ | 17* | 0.6(5.0) | 0.2 (3.0) | 0.015 (1.5) | 0.02 (1) | 2.4(5.0) |

Table 3: EWPOs at future e^+e^- colliders: statistical error (estimated experimental systematic error). Δ (δ) stands for absolute (relative) uncertainty, while * indicates inputs taken from current data [6]. See Refs. [23, 30, 35, 36, 46, 47].

is also an important ingredient for precision electroweak studies. Future e^+e^- Higgs factories could in principle provide data for the dispersive approach using the radiative return method, $e^+e^- \to \text{had.} + n\gamma$. While no detailed studies have been performed, it is not expected that this will lead to an improvement compared to data from lower-energy e^+e^- colliders. On the other hand, with sufficient amounts of luminosity spent at two center-of-mass energy a few GeV below and above the Z peak, it is possible to determine $\alpha(m_Z)$ directly, since the γ -Z interference contribution is sensitive to this quantity [49]. However, this method crucially depends on multi-loop theory calculations for the process $e^+e^- \to \mu^+\mu^-$.

Since the list of EWPOs in Tab. 3 is an over-constrained set of inputs for a SM fit, it can be used to indirectly determine the Higgs-boson and top-quark masses, which only appear within loop corrections. This is illustrated in Fig. 1, which demonstrates that all future e^+e^- colliders will tremendously improve the precision of this indirect test compared to the currently available data. The increased precision for the indirect determination of m_H and m_t at CEPC/FCC-ee compared to ILC is driven by the higher expected precision for the EWPOs themselves and for the strong coupling constant α_s . For ILC we assume $\Delta \alpha_s = 0.0005$, while for CEPC/FCC-ee we use $\Delta \alpha_s = 0.0002$ [48]. The difference in contours between CEPC and FCC-ee is mostly due to different assumptions about the precision of $\alpha(m_Z)$, where for FCC-ee we consider the direct determination according to Ref. [49] with $\Delta \alpha(m_Z) \sim 3 \times 10^{-5}$. On the other hand, we take the present-day uncertainty $\Delta \alpha(m_Z) \sim 1 \times 10^{-4}$ for CEPC, which is excessively consersative but serves to illustrate the impact of $\Delta \alpha(m_Z)$ in the electroweak precision fit.




Figure 1: Indirect sensitivity to m_H and m_t for a fit of SM theory predictions to current and projected future data for electroweak precision tests (W mass and Z-pole quantities). For comparison, the direct measurement precision is also shown (on the scale of the plot the width of the m_H band is not visible). The light (dark) shaded areas depict 95% (68%) confidence level regions. For the future collider scenarios it is assumed that the central values coincide with the SM expectations. The plot was made using SM theory predictions from Refs. [50, 51].

Further improvements: Some of the statistical and systematic uncertainties discussed above may be further improved with new advances of detector design and reconstruction techniques. Ref. [52] studied the prospects for precision measurements of the tau polarization, which can be used for measuring Z-boson coupling ratios, as mentioned in section 2.1. Due to the unobserved neutrino in the tau decays, a direct measurement of the tau polarization is not possible. Approximate polarization observables can be defined using only the visible decay products of individual tau decays. A better approximation of the true polarization may be obtained by using the full visible event information in di-tau events, $e^+e^- \rightarrow \tau^+\tau^-$. In this case it is possible to kinematically reconstruct the invisible neutrino momenta up to a two-fold ambiguity. However, all of the aforementioned polarization measurement methods require knowledge of the collision energy, and thus they suffer from ISR and beamstrahlung.

A new reconstruction method, which is much less sensitive to ISR, also makes use of the impact parameter of the visible tau decay products with the beam axis [52]. This information allows one to reconstruct the tau momenta exactly in the presence of a single ISR photon. For tau decays, the observable impact parameters are typically below 1 mm and thus they require precise vertex detectors. The impact parameter method appears promising to achieve 70–80% efficiency for the tau momentum and polarization reconstruction [52].

2.3 Electroweak precision measurements at other facilities

Besides high-energy pp and e^+e^- colliders, other experiments can also perform interesting precision measurements of the electroweak sector.

At electron-positron colliders with $\sqrt{s} \ll m_Z$, the process $e^+e^- \to f\bar{f}$ is dominated by photon exchange, and thus it is less sensitive to electroweak physics. However, the Belle II experiment at the SuperKEKB collider with $\sqrt{s} = 10.58 \,\text{GeV}$ will benefit from the very large integrated luminosity to obtain some competitive constraints. In particular, an upgrade SuperKEKB with polarized electron beams would open up the possibility of measuring the left-right asymmetry of the process $e^+e^- \to f\bar{f}$ [53]. For $\sqrt{s} \ll m_Z$, this process is mainly sensitive to v_f , the vector coupling of the Z-boson to $f\bar{f}$. With 40 ab⁻¹ integrated luminosity, the precision of v_f for $f = \mu, b, c$ could be improved by a factor of 4–7 compared to the current world average. For most final states, the precision is statistics limited, except for the $b\bar{b}$ final state. The dominant systematic error sources are the polarization measurement (0.3%) and subtraction of backgrounds (which include the Υ resonances) [53].

One also can interpret the results for v_f as a determination of the running weak mixing angle in the \overline{MS} scheme, $\sin^2 \overline{\theta}(\mu)$. The achievable precision is comparable to the combined LEP+SLC precision, but at a lower scale $\mu \approx 10 \,\text{GeV}$, thus providing a non-trivial test of the running of $\sin^2 \overline{\theta}(\mu)$.

Similarly, various low-energy precision experiments can determine the running weak mixing angle at very small scales, $\mu \lesssim 1\,\mathrm{GeV}$, through measurements of parity violation in fixed-target electron scattering and in atomic physics, see Ref. [54] for a brief overview.

On the other hand, new information on the running of $\sin^2 \overline{\theta}(\mu)$ at larger scales, $10 \,\text{GeV} < \mu < 60 \,\text{GeV}$ will be accessible at the Electron-Ion Collider (EIC), using scattering of electron and positron beams on proton and deuteron beams [55,56].

While e^+e^- colliders can deliver the best precision for many EWPOs, it is difficult to disentangle individual couplings of gauge bosons to light quarks, due to the low sensitivity of tagging light-quark flavor and charge. Lepton-proton colliders are ideally suited to overcome this difficulty. With the possibility of switching between e^-p and e^+p runs and with polarized e^+ beams, it is possible to individually determine the vector and axial-vector couplings of the Z-boson to light quarks (v_d, a_d, v_u, a_u) and simulaneously constrain the relevant PDFs. This is achieved by measuring neutral-current deep-inelastic scattering (DIS), $e^{\pm}p \rightarrow e^{\pm} + X$, where photons and Z-bosons appear in the t-channel. The relatively contribution of Z-boson increases with higher energies, so that a future high-energy ep collider will have substantially higher sensitivity to these couplings than previous experiments.

Two proposals for such a collider utilize the proton beam from the LHC (called LHeC) [57] or from FCC (called FCC-eh) [58], with center-of-mass energies of 1.3 TeV and 3.5 TeV, respectively. By performing a simultaneous coupling and PDF fit, it was found the LHeC can determine all for couplings (v_d, a_d, v_u, a_u) with $\mathcal{O}(\%)$ precision [59], while the precision can be improved by another factor 2–3 at FCC-eh [60], see Fig. 2.

A muon collider with center-of-mass energy $\sqrt{s} \approx 91 \text{ GeV}$ [61] could perform electroweak




Figure 2: Projected precision for the Z-boson vector and axial-vector couplings to light quarks at LHeC and FCC-eh, compared to the current precision from LEP, SLC, TeVatron and HERA (figure taken from Ref. [60]). Here g_V^f and g_A^f are rescaled versions of the vector and axial-vector couplings introduced in eqs. (2),(3): $v_f = g/(2\cos\theta_w)g_V^f$, $a_f = g/(2\cos\theta_w)g_A^f$.

measurements with a precision that greatly exceeds that currently available data from LEP/SLC. More studies for electroweak physics at muon colliders would be important to more thoroughly assess its potential.

2.4 Theory needs for the interpretation of Electroweak precision data

To fully exploit the potential of electroweak precision measurements to test the SM and possible new physics effects, theory inputs are needed in multiple places:

• Most of the quantities in Tab. 3 are not real observables, but pseudo-observables. The pseudo-observables are defined without backgrounds, initial-state radiation (ISR), the impact of final-state QED/QCD radiation on distributions, and detector smearing and acceptance effects. Various corrections factors and subtraction terms are needed to translate real observables to pseudo-observables. While it is possible to extract some of these terms with data-driven methods, theory input is needed in many instances, either for calibration or because the data-driven methods do not capture all relevant effects. The current state of the art are NLO results for the irreducible background contributions, MC tools with full NLO and partial higher-order QED radiation, and higher-order initial-state photon radiation in a leading-log approximation (see e.g. Ref. [62] for a review). For the expected precision of future e^+e^- Higgs factories, one more order of perturbation theory (NNLO) will likely be needed for the background contributions, and one or more orders of improvement are required for the simulation of QED effects in MC tools, which may require novel frameworks for the architecture of MC programs [63–65].

| EWPO | Current | Projected | Current | Projected p | error error |
|--|--------------|--------------|--------------|-------------|-------------|
| uncertainties | theory error | theory error | param. error | Scenario 1 | Scenario 2 |
| $\Delta m_W \; ({ m MeV})$ | 4 | 1 | 5 | 2.8 | 0.6 |
| $\Delta\Gamma_Z~({ m MeV})$ | 0.4 | 0.1 | 0.5 | 0.3 | 0.1 |
| $\Delta \sin^2 \theta_{\mathrm{eff}}^{\ell} (\times 10^5)$ | 4.5 | 1.5 | 4.2 | 3.7 | 1.1 |
| $\Delta A_{\ell} \ (\times 10^5)$ | 32 | 11 | 30 | 25 | 7.5 |
| $\delta R_{\ell} \ (\times 10^3)$ | 6 | 1.5 | 6 | 3.2 | 1.3 |

Table 4: Impact of theory and parametric uncertainties on the prediction of a few selected EWPOs (see Ref. [66]). For the theory errors, the uncertainty estimates from currently available calculations are compared to the projected improvement when assuming the availability of N³LO corrections and leading N⁴LO corrections. For the parametric errors, current uncertainties are compared to two future scenarios, see eq. (12).

- For the interpretation of measured values of the pseudo-observables, they need to be compared to precise predictions within the SM. For Z-pole EWPOs, full NNLO and partial higher-order corrections are currently known, while NLO plus partial higher orders are available for most other processes (such as $e^+e^- \to WW$). The estimated theory uncertainties are subdominant compared to current experimental accuracies, but are significantly larger than the anticipated precision of future e^+e^- collider, cf. Tabs. 1, 3, 4. The dominant missing contributions are 3-loop corrections with at least one closed fermion loop[†] and leading 4-loop corrections enhanced by powers of the top Yukawa coupling [66]. It is not possible to provide a reliable projection for how much the availability of these corrections would reduce the overall theory uncertainty, but a very rough estimate has been attempted in Ref. [66], using a combination of methods (extrapolation of the perturbation series, counting of known prefactors, scheme comparisons). As shown in Tab. 4, these corrections will likely be needed to match the precision of future e^+e^- colliders (Tab. 3), and in some cases even higher orders may be necessary. Fortunately, there is continuous progress in the development of new calculational techniques for loop diagrams [67–70].
- Furthermore, as already mentioned above, SM theory predictions of EWPOs require various SM parameters as inputs, most notably the top mass m_t and Higgs mass m_H , the strong coupling constant α_s , and the shift of the fine structure constant, $\Delta \alpha$. While the latter has been discussed above on pages 7 and 12, information about the other parameters can be found in Ref. [48] and the EF Higgs and TOPHF reports [14,71]. The impact of SM parameter uncertainties are illustrated in Tab. 4 for current results for these parameters and two future scenarios:

| | $\Delta m_t \; [{\rm GeV}]$ | $\Delta m_H \; [{ m GeV}]$ | $\Delta m_Z \; [{ m MeV}]$ | $\Delta(\Delta \alpha)$ | $\Delta \alpha_s$ | |
|------------|-----------------------------|----------------------------|----------------------------|-------------------------|--------------------|------|
| Current | 0.6 | 0.17 | 2.1 | 10^{-4} | 9×10^{-4} | (19) |
| Scenario 1 | 0.3 | 0.02 | 0.8 | 10^{-4} | 5×10^{-4} | (12) |
| Scenario 2 | 0.05 | 0.01 | 0.1 | 3×10^{-5} | 2×10^{-4} | |

[†]Corrections with fermion loops are enhanced due to the large top Yukawa coupling and the large fermion multiplicity in the SM.

Scenario 1 approximately corresponds to a Higgs factory with a Giga-Z Z-pole run and no data taking at the $t\bar{t}$ threshold. Scenario is more similar to a Higgs factory with a Tera-Z Z-pole run (FCC-ee, CEPC) and including $t\bar{t}$ threshold run. In particular, the improvement in the m_t precision is crucial for reducing the parametric uncertainties in Scenario 2, to a level that is roughly comparable to the target precision for these EWPOs shown in Tab. 3.

Note that the dependence of the predictions for Γ_Z and R_ℓ on α_s are to a certain extent circular, since these quantities would be used for the extraction of the strong coupling constant at future e^+e^- colliders [48].

3 Multi-boson processes at high-energy colliders

The SM predicts the existence of multi-boson interactions, which give rise to final states with two or three bosons. Anomalies in the rate and kinematic of these final states can be indicative of new physics not currently described in the SM. Such anomalies can be parametrized through modifications of the strength or form of the SM multi-boson vertices. A newer approach consists in using EFT operators of dimension six or above, where measurements of multi-boson processes can be recast as direct determinations of the Wilson coefficients of these operators.

It shall be noted that the sensitivity to BSM effects, or, in other terms, the upper limits to the Wilson coefficients of new operators, scale with a power of the c.o.m. energy, thus making multi-TeV colliders the ideal tools for studying these final states. At this time, the most promising avenues for reaching multi-TeV energies are proton-proton colliders or $\mu^+\mu^-$ colliders.

Di-boson final states can be produced directly through annihilation of the colliding particles or partons, or indirectly through vector-boson fusion (VBF) / vector-boson scattering (VBS) processes. These processes can give important clues about the origin of the electroweak symmetry breaking, and whether the Higgs mechanism is the only source of it. Hadron colliders also offer the possibility to study same-sign WW production through VBS. Other interesting final states contain three bosons, such as WWW, or the as-of-yet unobserved ZZZ. These final states can be produced via quartic-gauge couplings, and allow one to unveil one of the ostensibly least known sector of the SM.

As noted above, discrepancies between the expected total and differential cross sections for each of the multi-boson final states and their SM predictions can be studied with two different approaches. A modification of the strength of the SM couplings constitute the premise of the searches for triple- and quartic-gauge-coupling anomalies (TGC and QGC, respectively). The fundamental assumption is that there are no additional types of interaction among SM particles than the ones already included in the SM Lagrangian. The adoption of EFT operators allows one to eliminate this constraint, and bestows the freedom to obtain a model-independent extension of the SM Lagrangian, under the assumption that there are no additional fields. The SMEFT approach is described in more detail in Section 4.1.

Plentiful experimental results with multi-boson final states are available. Both the ATLAS and CMS collaborations have measured di-boson [72–80], tri-boson processes [81–84], as well as VBF/VBS processes [85–95], which are characterized by a VVjj final state. Di-boson final states include W^+W^- , same-sign $W^\pm W^\pm$, WZ, ZZ, $Z\gamma$. Tri-boson final states include $W\gamma\gamma$, $Z\gamma\gamma$, $WV\gamma$ (where V=W,Z), and WVV' (where V,V'=W,Z). A summary of the expected sensitivities of multi-boson cross-section measurements for HL-LHC is reported in Ref. [15].

Bounds on new physics have been determined in the language of anomalous gauge-boson couplings (aGCs) [72,75–77,80] and effective operators [73,77–79,84,91–94]. The latter is theoretically preferred since it provides a consistent power counting and allows one to implement theoretical consistency constraints. In these studies, only one or two aGCs/operators are allowed to be non-zero at the same time, i.e., no full aGC/SMEFT analysis has been performed.

The most up-to-date limits on gauge-coupling anomalies are available at Refs. [96, 97]. Expected limits at the end of the HL-LHC and HE-LHC runs are reported in Refs. [98, 99].

3.1 Theory studies on anomalous couplings

It is well know that diverging from the SM predictions of the TGC and QGC causes the growth of scattering amplitudes, up to the point at which unitarity is lost. Various methods have been implemented in TGC and QGC searches to enforce the conservation of unitarity, and the consistency of the analyses. A study of the different methods employed by experimental collaborations is presented, e.g., in Ref. [100]. A new direction of research toward the imposition of constraints dictated by the necessity that SMEFT admits a UV completion is explored in Ref. [101]. The conventional approach to the derivation of positivity bounds uses 2-to-2 scattering amplitudes, showing that one can obtain a set of homogeneous inequalities for the dim-8 Wilson coefficients. The limit of this technique is that it requires one to consider scattering amplitudes between arbitrarily superimposed particle states, which has not been done systematically. The new approach draws a connection between the positivity bounds of EFT Wilson coefficients and the solution of a geometric problem, i.e., finding the extremal rays of a spectrahedron, built from the crossing symmetries and the SM symmetries of an interaction amplitude. The bounds obtained with new method are compared to the ones from the elastic positivity bounds and shown to be more stringent. A concise survey of the recent advances in constraining the SMEFT parameter space from the UV considerations can be found in Ref. [102], section 2.5.

A review of the usage and potential pitfalls of SMEFT is presented in [103]. As indicated before in this report, EFT is the leading tool employed to determine, in a model-independent way (with certain symmetry assumptions), the impact of SM measurements on new physics. Two energy regimes, in which SMEFT studies are currently performed, are identified and separately discussed: resonant and near-threshold processes at low energy; distribution tails at high energy. In the former case, it is possible to show that the number of ways in which SMEFT can contribute is finite; this allows to identify combinations of Wilson coefficients as contributors to the process in question, and effectively obtain a resummation

of all orders in the SMEFT expansion that affect the process of interest. In the latter case, it is not possible to consider the full EFT effect on a process, and one needs to cut the expansion at dimension 6. As hinted above, one then must consider the effect of truncating the expansion by neglecting higher-order operators, such as dim-8 ones, as well as the limits of validity of the EFT approximation. The choice of a method to fit any model data is also discussed. A global fit, where all measurements are considered on equal footing, is ideal, but requires significant work to properly combine and compare the fit inputs. A sequential fit is presented as a quicker alternative, in which intermediate fits are performed by adding measurements divided in subsets, in order of decreasing precision. Directions for future progress are summarized at the end of the paper, and involve studies of SMEFT at higher order for on-shell and near-threshold observables, the adoption by experiments of an error estimation scheme for high-energy observables, and the development of sequential fits toward the definition of a fully-global fitting framework.

3.2 Multi-boson processes at future lepton colliders

While limited in energy reach compared to hadron colliders, lepton colliders with $\sqrt{s} \gtrsim 1$ TeV have advantages for measurement of vector-boson scattering (VBS), due to the well-defined initial state, complete coverage of final states, and the possibility to separate spin, isospin and CP quantum numbers. Particle flow algorithms enable very good particle ID (to reduce photon-induced background) and W/Z discrimination from hadronic decays [104].

An e^+e^- collider like ILC or CLIC can cover energies of a few TeV, while a muon collider or more speculative proposals such as plasma wakefield accelerators (e.g., Ref [105]) can reach tens of TeV [106]. In the latter case, VBS can be described with good accuracy by factorizing the full process $\ell\ell \to VV\ell'\ell'$ into a $V'V' \to VV$ hard process and V' radiation in the initial state described by electroweak PDFs [107, 108].

Significant backgrounds arise from a number of processes (including $\ell^+\ell^- \to VV$ without VBS), but they can be reduced with suitable cuts or machine learning techniques, and they also become less important relative to the signal process when going to higher values of \sqrt{s} [109–111]. However, the achievable constraints on SMEFT coefficients do not always improve when increased center-of-mass energy [107].

Reference [107] presents a review of how electroweak vector boson fusion/scattering processes become the dominant production modes of vector bosons as the center-of-mass energy of a lepton collider enters the few-TeV range. The size and growth of VBF cross sections for numerous SM and new physics processes are investigated. The key observation is that s-channel production rates decrease, with collider energy, as 1/s, while VBF rates grow as $\log s$, eventually becoming the most dominant process.

A comprehensive review of VBS processes at current and future colliders is presented in Ref. [112]. This paper also discusses the importance of adopt the proper formalism to describe initial-and final-state radiation (electroweak parton distribution functions, and resummation of fragmentation functions, respectively). This latter topic is presented also in Ref. [113] for lepton colliders (and similarly, in the scenario of a high-energy hadron

collider, in Ref. [114], which is discussed later in section 3.3). Multi-TeV lepton colliders are effectively weak-boson colliders, which suggests that EW bosons should be treated as constituents of high-energy leptons. The paper reviews the validity of W and Z parton distribution functions, investigates power-law and logarithmic corrections that arise in the derivation of weak boson PDFs in the collinear limit, and reports an implementation of the Effective W/Z and Weizsäker-Williams approximations into the Monte Carlo generator MadGraph_aMC@LNO. The key question is how factorization and resummation work in the weak sector, and how it differs from QED and perturbative QCD. This question is critically important, as in multi-TeV muon colliders, and at 100 TeV hadron colliders, typical parton collisions satisfy the criteria for collinear factorization of weak bosons. It will furthermore be important to extend the factorization framework to higher orders (see e.g. Ref. [115]).

The future lepton colliders obviously offer the opportunity to investigate, in detail, interesting experimental signatures. Three such studies are presented below, and touch two specific aspects of lepton colliders: the precision study of Higgs physics, and the unique advantage (high-energy, high-statistics, clean environment) offered by muon colliders as weak-boson colliders.

Reference [116] reports a study of the Vh process that is relevant for both the HL-LHC and future lepton colliders, in which the signal to background ratio is significantly higher than at the LHC. A particularly interesting aspect of the analysis is the ability to check whether the Higgs couplings to W and Z, κ_W and κ_Z have the same sign; models in which they do not include scalars which have higher isospin representations. The idea is to exploit the tree-level destructive interference between the W and Z mediated processes that contribute to the production of a Higgs boson in association with a vector boson via vector-boson fusion. The Vh matrix element contains in fact a term that grows with energy and is proportional to $\lambda_{WZ} - 1$, where $\lambda_{WZ} = \kappa_W/\kappa_Z$ (i.e., $\lambda_{WZ} = 1$ in the SM). The future lepton collider being considered is CLIC, at 1.5 TeV and 3 TeV center-of-mass energy. The achievable sensitivity at a lepton collider is shown in Fig. 3. It is reported that the point $(\kappa_W, \kappa_Z) = (1, -1)$ is excluded at more than two standard deviation at the end of the HL-LHC run[‡], while 3.4 fb⁻¹ (14.1 fb⁻¹) are enough at CLIC 3 TeV (1.5 TeV) to exclude that point at 95% CL against the SM case.

Prospects for searches for anomalous quartic gauge couplings at a high-energy muon collider are presented in Ref. [117]. A multi-TeV muon collider is effectively a high-luminosity weak boson collider, and allows for the measurement, in a relatively clean environment, of vector boson scattering processes. The study of W pair production, in association with two muons or two neutrinos, is presented in the reference. Deviations of the proposed measurements with respect to the SM predictions could indicate the presence of an anomalous quartic gauge coupling. Figure 4 shows the distribution of the mass of the W pair in the WW $\nu\nu$ and WW $\mu\mu$ channels, using a simulation with full matrix elements (rather than the effective W-boson approximation or EW PDFs). It also includes an example of a signal prediction for one illustrative aQCG parameter. The limits on anomalous quartic gauge

[‡]Note that a more precise determination of the magnitude of $\kappa_{W,Z}$ can be achieved with a global fit of HL-LHC measurements, but the discussion here focuses only on the direct determination of the sign of these couplings.




Figure 3: Left: 1- and 2- σ sensitivity of the measurement to κ_W and κ_Z at the HL-LHC. Right: constraints in the $\kappa_W - \kappa_Z$ plane for the total rate measurement at CLIC. (figures taken from Ref. [116]).

couplings that can be set with a luminosity of 4/ab of muon collisions at a center-of-mass energy of 6 TeV are about two orders of magnitude tighter than the current limits. It shall be noted that the effects of beam-induced background have not been included in the analysis, and that the limits on aQGCs are set under the assumption that triple gauge couplings are not modified.

Muon colliders also offers the opportunity to study in detail the topic of unitarity restoration, for example, by measuring longitudinally polarized vector boson scattering. It is shown in Ref. [118] that such a study could surpass the end-of-life HL-LHC results in the ZZ channel. The study utilizes a Boosted Decision Tree and shows that, even with a conservative estimation, a 5 standard deviation discovery of longitudinally polarized ZZ scattering can be achieved with 3/ab of data collected at a 14 TeV muon collider. This results outperforms the expected results of the end-of-life HL-LHC, which expects to have a sensitivity of about 2 standard deviations. The paper also reports the study of a 6 TeV muon-collider case, and shows that its sensitivity is comparable to the HL-LHC one.

3.3 Multi-boson processes at future hadron colliders

Multi-TeV future hadron colliders provide a unique laboratory to explore the nature of EW symmetry breaking, and its restoration as energies increase above the EW scale. The most compelling direction of investigation entails the study of vector bosons produced in association with a Higgs boson, or the longitudinal polarization of pair-produced vector bosons. Both these lines of investigations are discussed below.

A discussion of the prospects for physics measurements at future hadron colliders requires a suitable description of the physics in the high-energy regime. Reference [114] discusses the definition of a consistent theoretical treatment to describe the physics processes that take place in particle collisions at multi-TeV energies. In that regime, beyond the weak mass scale, all SM particles, including the gauge bosons, can be considered to be




Figure 4: Distribution of m(WW) in the WW $\nu\nu$ and WW $\mu\mu$ channels, after event selection. The dashed lines show the signal prediction for one illustrative aQGC parameter. (figure taken from Ref. [117]).

massless. Collinear splitting becomes the dominant phenomenon. The proper description of parton distribution functions, initial state radiations, final state radiations and fragmentation functions is needed. The focus of the paper is the resummation of final-state radiation up to leading-log accuracy, and show the effect of high-energy splitting at a 100 TeV collider. The formalism developed in the paper is applicable also to the case of multi-TeV muon colliders.

As indicated earlier, the study of Higgs production in association with a vector boson at a high-energy hadron collider offers a test stand to probe the restoration of EW symmetry. Reference [119] presents a new test to study the restoration of EW symmetry at high energy. The two colliders under consideration are the 14 TeV HL-LHC and the 27 TeV HE-LHC. The main assumption of the analysis is that at those energies the EW vector bosons become massless, and one can replace their longitudinal modes with the associated Goldstone bosons. The conclusion is that while the VV' production is dominantly transversely polarized, up to very high energies, the Vh channel is longitudinally dominated starting at relatively low energies (e.g., at a c.o.m. energy of 14 TeV, the W boson is longitudinally polarized in $\approx 90\%$ of the Wh events with a Higgs transverse momentum above 200 GeV). Figure 5 demonstrates this effect, by showing the fraction of polarized gauge-boson production as a function of the boson transverse momentum. It is ultimately demonstrated that the EW restoration can be confirmed with a precision of 40% at the HL-LHC, and 6% at the HE-LHC.

The second compelling direction of investigation previously identified is the study of longitudinal polarization of pair-produced vector bosons. The sensitivity to longitudinal vector boson scattering at a multi-TeV proton-proton collider, using same-sign WW pairs produced in association with two jets, is presented in Ref. [120]. Vector boson scattering processes are important probes of the non-Abelian structure of electroweak interactions, as the unitarity of the tree-level amplitude of longitudinally polarized gauge boson scattering




Figure 5: Ratio of transverse momentum distributions of polarized gauge boson production to the total distribution summed over polarizations. (figures taken from Ref. [119]).

could be restored at high energies by the Higgs boson. Extensions of the SM introduce new resonances or modifications of the Higgs boson couplings that modify the cross sections of processes involving the scattering of longitudinally polarized gauge bosons. The same-sign WW state is of particular interest because the requirement of two same-sign leptons in the final states greatly reduces the backgrounds from other SM processes. The study reported in the paper assumes 30/ab collected at a center-of-mass energy of 27, 50, and 100 TeV. The result of the analysis is that a cut-and-count method is sufficient to measure with a relative precision of 39%, 22%, and 17% the fraction of the purely longitudinal contribution to same-sign WWjj production, using the fully leptonic decay mode (at a center-of-mass energy of 27, 50, and 100 TeV, respectively). The purely transverse and mixed longitudinal-transverse contributions are measured with a relative precision of 2% and 4%, respectively, at 100 TeV. Figure 6 shows the distribution of the signal (W_LW_L) and background components, as a function of the pseudorapidity difference between the two jets. As expected, the signal-to-background ratio is higher at a large pseudorapidity difference.

4 Global fits of new physics

Assuming new physics scales are significantly higher than the EW scale, Effective Field Theories (EFT) provide a model-independent prescription that allows us to put generic constraints on new physics and to study and combine large sets of experimental data in a systematically improvable quantum field theory approach. All new physics effects are represented by a set of higher-dimensional operators which consist of only the SM fields and respect the SM gauge symmetries. Depending on whether the SU(2)×U(1) gauge symmetries are realized linearly or nonlinearly, there are two classes of formalism popular for studying EW physics at colliders, the Standard Model Effective Field Theory (SMEFT) [121, 122] or the Higgs Effective Field Theory (HEFT) [123,124]; see Ref. [125] for a pedagogical review. The EFT approach has some features that are of particular interest for studying precision EW physics, for instance: it provides a well-defined theoretical framework that enables the inclusion of radiative corrections for both the SM and BSM parts; and the synergies between different precision EW measurements can be explored globally so that a




Figure 6: The distribution of events as a function of $\Delta \eta(jj)$, the pseudorapidity difference between the two leading jets, after full event selection, for $\sqrt{s} = 100$ TeV (figure taken from Ref. [120]).

comprehensive picture of the constraints on new physics can be drawn. However, the EFT approach also has some practical limitations since it has in principle an infinite number of degrees of freedom, and it is only an adequate description if the new physics scales are larger than the experimentally reachable energies. In a realistic global EFT fit, various flavor assumptions and truncations to the lowest order of relevant operators often have to be applied, to limit the number of parameters to a manageable level. The HEFT allows considerably more parameter freedom than the SMEFT. A subset of BSM physics that couples only to SM gauge bosons can also be represented by the so-called oblique parameters [126, 127]. Two of the oblique parameters (S and T) are directly related to dimension-6 SMEFT operators, while the U parameter corresponds to a dimension-8 operator.

Most of the global EFT fits are currently performed based on the SMEFT, which will also be the focus here. An up-to-date global SMEFT fit at future colliders has been performed for the European Study Group (ESG), which combines measurements of EWPOs, Higgs production and decay rates at LHC and future colliders, and $e^+e^- \to WW$ [46]. The results of global EFT fit may give important implications for proposals of future colliders which otherwise would not get recognized. Just to name two examples from ESG: Z-pole and WW runs at circular e^+e^- colliders can help improve significantly the Higgs coupling precisions with respect to what can be obtained using only ZH runs; and beam polarization at linear e^+e^- colliders can help lift degeneracies of different new physics effects, as a result of which similar Higgs coupling precision can be achieved at both linear and circular e^+e^- colliders, in spite of the difference in integrated luminosity.

For Snowmass 2021, the global EFT fit for ESG has been extended in a few directions [128]: consistent implementation of full EFT treatment in $e^+e^- \to WW$ using optimal observables; new inclusion of a large set of 4-fermion operators; more complete set of operators that are related to top-quark. The projections of the uncertainties of required input observables are provided by the Topical Group EF01 for Higgs related observables [71], EF03 for top-quark related observables [14], EF04 for W/Z related observables (see section 2), and the Rare Process and Precision Frontier (RF) for a set of low-energy measurements. The projections for various future e^+e^- colliders are made to be as consistent as possible, for instance: by applying common systematic errors as explained in Section 2.2; by extrapolating from one collider to another whenever there is any important missing input. More details about the considerations of inputs can be found in Ref. [128]. The global fits are performed with respect to various run scenarios for each collider. The results of global fits are provided as bounds on the Wilson coefficients as well as uncertainties of effective H/Z/Wcouplings. The intrinsic and parametric theory errors of the SM predictions of observables are not included in the global fits by default, instead their impact is quantified separately in terms of how much change the theory error would bring on the fitted uncertainties on various effective couplings. The computation of EFT contributions to various observables is done at tree level only except for the loop contribution from the triple Higgs coupling in the single Higgs processes.

4.1 Framework and scope

The SMEFT takes a form of Effective Lagrangian from the SM part up to dimension-4 operators plus an infinite tower of higher-dimensional (d > 4) operators (O_i^d) which respect Lorentz and the SM gauge symmetries and are suppressed by the corresponding inverse powers of the cut-off scale Λ ,

$$\mathcal{L}_{\text{SMEFT}} = \mathcal{L}_{\text{SM}} + \sum_{d=5}^{\infty} \sum_{i} \frac{C_i^{(d)}}{\Lambda^{d-4}} \mathcal{O}_i^{(d)}.$$
 (13)

The information of new physics is encoded in the series of Wilson Coefficients $C_i^{(d)}$. The number of non-redundant operators at d=5,6,7,8 is known [121,122,129–134]. For the global fits presented here, we restrict ourselves to operators of dimension 6 (d=6) that preserve baryon and lepton numbers. A complete basis of such operators contains 2499 operators without flavor assumptions. It reduces to 84 if only one generation of fermions are considered, and further to 59 if only for CP-even operators. Three main global fits are performed for Snowmass 2021, called Fit-1/2/3, which each consider a different subset of operators to parametrize EW physics at future colliders. Independent operators for more than one generation of fermions are considered, with a general assumption that the flavor structure is diagonal for simplicity, but without assuming lepton-flavor universality. The operators for the 3rd-generation of quarks are always treated separately, while universality for the 1st- and 2nd-generation of quarks is assumed in Fit-1. Bounds on Wilson Coefficients are given in terms of the original Warsaw basis [122]. More details can be found in Ref. [128].

Fit-1 is mainly focused on the Higgs and EW sectos. It explores the interplay among




Figure 7: The contribution from the same operator $\mathcal{O}_{\phi e}$ (defined in the text) in three different processes: double Higgs production (left), single Higgs production (middle) and Z-pole production (right).

measurements for Higgs production and decay rate, EWPOs and di-boson processes; the roles played by energy, luminosity and beam polarizations; and the synergy between LHC and future lepton colliders. There are around 20 operators that contribute to those measurements, which is the complete set given the assumptions as mentioned above. Let us consider one of the operators as an example to illustrate why Higgs and EW measurements are inherently related: $\mathcal{O}_{\phi e} = (\phi^{\dagger}iD_{\mu}\phi)(\overline{e}_{R}\gamma^{\mu}e_{R})$. As sketched in Fig. 7, this operator will directly generate a five-point interaction that contributes to $e^{+}e^{-} \rightarrow ZHH$. By replacing the Higgs field by v, it will also generate a four-point interaction that contributes to $e^{+}e^{-} \rightarrow ZH$ which is one of the leading Higgs production channels. Furthermore, replacing the other Higgs field by v, it will result in a vertex correction to Z pole observables. Therefore, the interplay of Higgs measurements and EWPOs at the Z-pole together will be advantageous for probing the effects from new physics.

Fit-1 results are given mainly in terms of effective couplings [46,135] which are defined by pseudo-observables and thus are independent of the operator basis one would have chosen for the fit. The Higgs effective couplings (g_{HX}^{eff}) are defined as

$$g_{HX}^{\text{eff}} = \frac{\Gamma_{H \to X}}{\Gamma_{H \to X}^{\text{SM}}},$$
 (14)

where $\Gamma_{H\to X}$ is the decay partial width of $H\to X^\S$ The electroweak effective couplings, g_L^f and g_R^f for each fermion f, are defined similarly through the partial decay widths of $Z\to f_L\overline{f}_L$ and $Z\to f_R\overline{f}_R$ where f_L and f_R are the left- and right-handed fermion, respectively.

Fit-2 is focused on probing 4-fermion interactions which are present in many BSM models with new gauge bosons that couple to the SM fermions. The framework is based on the study in Ref. [136] and extends it by including measurements at future colliders. It involves both 4-fermion operators and 2-fermion operators that modify W/Z vertices, around 60 parameters in total. While the vertex part has significant overlap with Fit-1, Fit-2 extends the scope by relaxing the universality assumption that was imposed in Fit-1 for the first and second generation quarks. Essentially Fit-2 does not apply any flavor assumption besides focusing only on the diagonal part (i.e. ignoring flavor violating effects). The effect of 4-

[§]Each Higgs effective coupling is related to one κ parameter in the κ -formalism [46].

[¶]See Ref. [137] for an earlier study of a global fit to data with generic flavor structure.

fermion operators grows quadratically as energy increases, so that this global fit can provide a very good measure of the merit of running at high energies at future colliders. There are degeneracies which can only be lifted by low-energy measurements; thus this fit is also a place to study the synergy between future colliders and low-energy experiments. The renormalization group evolution of the relevant operators at different scales are properly taken into account. The results from Fit-2 are presented in terms of $1-\sigma$ bounds directly on the Wilson Coefficients of 4-fermion operators as well as the precision on electroweak effective couplings. Fit-2 results can also be interpreted in terms of the bounds on O_{2W} and O_{2B} operators (defined in [138]) which correspond to the oblique parameters W and Y [139].

The top-quark sector has essentially been excluded in the scope of Fit-1 and Fit-2. Fit-3 is thus focused on top-quark electroweak couplings and eett 4-fermion operators, by considering around 20 operators that are directly related to top-quark or the third generation quarks. The top quark has currently not yet been directly produced at any lepton colliders, so that top-quark measurements at hadron colliders play an essential role. Fit-3 also allows one to study the synergy between HL-LHC and future lepton colliders where top-quark can be directly produced. There have already been global SMEFT fits to current data including the interplay between the top-quark sector and Higgs/EW sectors [140,141]. The interplay may become more subtle when loop effects from top-quark operators in the Higgs/EW observables are included [142–144]. The Fit-3 for Snowmass 2021 extends these studies by focusing on future colliders, but it is restricted to a more limited set of observables and operators since not all ingredients that are needed for such a combined top-quark/Higgs/EW fit are technically ready. Nevertheless, it is useful for studying the interplay of the top-Yukawa coupling with the Higgs/EW sector.

A few independent sets of codes have been developed to do the global fits, using HEP-fit [145], Mathematica and C++, each using different statistical models**. Cross checks has been performed extensively and excellent consistency has been achieved for the fit results. Since the focus here is on the projected precision and new-physics reach, the central values of all input observables are by default set to the SM expectations. Nevertheless it has been confirmed in Fit-2 that identical uncertainties are obtained if the central values of input observables take their current measurement values from the PDG.

4.2 Collider scenarios and observables

The colliders scenarios that are considered in the global fits include the HL-LHC and future e^+e^- colliders as shown in Tab. 2. In addition, future muon colliders are also included in Fit-1 with three scenarios: 1 ab⁻¹ at 3 TeV; 10 ab⁻¹ at 10 TeV; 10 ab⁻¹ at 10 TeV

Using the equations of motion, these operators lead to charged and neutral current four-fermion contact interactions with flavor universality.

^{**}The global fit code by HEPfit performs a Bayesian analysis following Markov Chain Monte Carlo procedures with the logarithm of the likelihood function built from measurement projections; the Mathematica code relies on a χ^2 constructed from all observables; and the C++ code obtains the fitting parameter uncertainties directly from their covariance matrix.

plus 20 fb⁻¹ at 125 GeV^{††}. No muon collider scenarios have been considered for Fit-2 and Fit-3, partly due to a lack of suitable studies of input observables^{‡‡}. The projected uncertainties of various input observables are mainly supplied by the corresponding collider collaborations [21–28,30,147–149]. The list of input observables in each fit is too lengthy to be included in this report, and we will show here only a few typical examples. The complete list and details can be found in [128].

The input observables for Fit-1 include: EWPOs as shown in Tab. 3; Higgs production and decay rates, as shown partially in Tab. 5 for HL-LHC and Tab. 6 for FCC-ee and CEPC*; optimal observables for $e^+e^- \to W^+W^-$ [150]. The input observables for Fit-2 include: EWPOs as above; cross section and forward-backward asymmetry in $e^+e^- \to f\bar{f}$ off the Z-pole[†]; low-energy observables as shown in Tab. 7. Fit-3 includes the following observables: cross section or differential cross section for $t\bar{t}$, single-top, $t\bar{t}Z$ and $t\bar{t}\gamma$ production at (HL-) LHC; cross section and forward-backward asymmetry in $e^+e^- \to b\bar{b}$; optimal observables in $e^+e^- \to t\bar{t} \to bW^+\bar{b}W^-$ [151]; cross section for $e^+e^- \to t\bar{t}H$.

It is important to ensure the consistency among inputs provided by different collider collaborations. One example about common systematic errors in A_b measurements has been elaborated in Section 2.2. Another example is shown in Tab. 6, where we can compare the direct inputs by one collaboration with extrapolated inputs from another collaboration, as illustrated for FCC-ee240 direct inputs and ILC extrapolations (numbers in brackets) in the second column. Even though the $BR_{\gamma Z}$ input was missing in the FCC-ee documents, the extrapolated projection is included in the global fit since this observable turns out to play a sensitive role. More examples about the procedures that were taken to ensure consistency on inputs can be found in Ref. [128].

4.3 Results

The results of Fit-1 are shown in Fig. 8 for electroweak and Higgs effective couplings as defined in Sec. 4.1. They are plotted as 1σ relative uncertainties for two cases of global fits: the wider (narrower) bars assume that the Higgs total width is constrained (free)[‡]. The grey bars represent the expectation from HL-LHC measurements while colored bars are for various future lepton colliders as indicated in the legend. The input measurements from HL-LHC are always included in the fits for future colliders. For each future collider, results are shown for various running scenarios with measurements in earlier stages always combined with the later ones (denoted by the "+" symbol in the legend), except for the muon colliders. The electroweak effective couplings include: Z couplings to left- and right-handed

^{††}Another scenario which combines a 10 TeV muon collider with a future e^+e^- machine is in preparation. ^{‡‡}However, a study of the sensitivity of muon colliders to new 4-fermion interactions was performed in Ref. [146] in the framework of the W/Y parameters, which is more constrained than the SMEFT framework.

^{*}There are many more tables for Higgs inputs at other energies and other colliders [128] which are not explicitly listed here for conciseness.

[†]Due to insufficient inputs from community, a common analysis was performed to obtain those uncertainties for all future e^+e^- which however included only statistical errors.

[‡]Allowing the Higgs total width to be a free parameter accounts for the possibility of non-standard Higgs decays into BSM particles.

| HL-LHC | $3 \text{ ab}^{-1} \text{ ATLAS+CMS}$ | | | | |
|-----------------------------------|---------------------------------------|------|------|------|------|
| Prod. | ggH | VBF | WH | ZH | ttH |
| σ | - | - | _ | - | _ |
| $\sigma \times BR_{bb}$ | 19.1 | - | 8.3 | 4.6 | 10.2 |
| $\sigma \times BR_{cc}$ | - | - | _ | - | _ |
| $\sigma \times BR_{gg}$ | - | - | _ | - | _ |
| $\sigma \times BR_{ZZ}$ | 2.5 | 9.5 | 32.1 | 58.3 | 15.2 |
| $\sigma \times BR_{WW}$ | 2.5 | 5.5 | 9.9 | 12.8 | 6.6 |
| $\sigma \times BR_{\tau\tau}$ | 4.5 | 3.9 | _ | _ | 10.2 |
| $\sigma \times BR_{\gamma\gamma}$ | 2.5 | 7.9 | 9.9 | 13.2 | 5.9 |
| $\sigma \times BR_{\gamma Z}$ | 24.4 | 51.2 | _ | - | _ |
| $\sigma \times BR_{\mu\mu}$ | 11.1 | 30.7 | _ | - | _ |
| $\sigma \times BR_{inv.}$ | - | 2.5 | _ | - | _ |
| Δm_H | 30 MeV | - | - | - | - |

Table 5: Projected uncertainties of Higgs observables at HL-LHC for the leading five production channels and various decay modes. Numbers are in %, except for m_H .

| | $FCCee240 5ab^{-1}$ | | CEP | $C240 \ 20ab^{-1}$ |
|-----------------------------------|---------------------|-------------|------|--------------------|
| Prod. | ZH | $\nu \nu H$ | ZH | $\nu \nu H$ |
| σ | 0.5(0.537) | - | 0.26 | - |
| $\sigma \times BR_{bb}$ | 0.3(0.380) | 3.1(2.78) | 0.14 | 1.59 |
| $\sigma \times BR_{cc}$ | 2.2(2.08) | - | 2.02 | - |
| $\sigma \times BR_{gg}$ | 1.9(1.75) | - | 0.81 | - |
| $\sigma \times BR_{ZZ}$ | 4.4(4.49) | - | 4.17 | - |
| $\sigma \times BR_{WW}$ | 1.2(1.16) | - | 0.53 | - |
| $\sigma \times BR_{\tau\tau}$ | 0.9(0.822) | - | 0.42 | - |
| $\sigma \times BR_{\gamma\gamma}$ | 9(8.47) | - | 3.02 | - |
| $\sigma \times BR_{\gamma Z}$ | (17^*) | - | 8.5 | - |
| $\sigma \times BR_{\mu\mu}$ | 19(17.9) | - | 6.36 | - |
| $\sigma \times BR_{inv}$. | 0.3(0.226) | - | 0.07 | - |

Table 6: Projected uncertainties of Higgs observables at FCCee240 and CEPC240 in the two leading production channels and various decay modes. Numbers are in %. The numbers in brackets are extrapolated from the projections at ILC250.




Figure 8: Precision reach on Higgs and electroweak effective couplings from a SMEFT global analysis of the Higgs and EW measurements at various future colliders. The wide (narrow) bars correspond to the results from the constrained- Γ_H (free- Γ_H) fit. The HL-LHC and LEP/SLD measurements are combined with all future lepton collider scenarios. For e^+e^- colliders, the high-energy runs are always combined with the low energy ones. For the ILC, the (upper edge of the) triangle mark shows the results for which a Giga-Z run is also included. For the muon collider, three separate scenarios are considered. The subscripts in the collider scenarios denote the corresponding integrated luminosity of the run in ab⁻¹. Note the Higgs total width measurement from the off-shell Higgs processes at the HL-LHC is not included in the global fit.

| Process | Observable | Experimental value | Ref. | SM prediction |
|---|--|--|--|--|
| (-) | $g_{LV}^{ u_{\mu}e}$ | -0.035 ± 0.017 | CHARM-II [152] | -0.0396 [153] |
| $\begin{array}{ccc} \begin{pmatrix} - \\ \nu \\ \mu \end{pmatrix} - e^{-} \text{ scattering} \end{array}$ | $g_{LA}^{ u_{\mu}e}$ | -0.503 ± 0.017 | OHARM-H [102] | -0.5064 [153] |
| τ decay | $\frac{G_{\tau e}^2}{G_F^2}$ | 1.0029 ± 0.0046 | PDG2014 [154] | 1 |
| / decay | $\frac{G_{Te}^2}{G_F^2}$ $\frac{G_{T\mu}^2}{G_F^2}$ $\frac{G_{\tau\mu}^2}{G_F^2}$ | 0.981 ± 0.018 | 1 DG2014 [104] | 1 |
| | $R_{\nu_{\mu}}$ | 0.3093 ± 0.0031 | CHARM $(r = 0.456)$ [155] | 0.3156 [155] |
| | $R_{\overline{\nu}_{\mu}}$ | 0.390 ± 0.014 | CHARM (7 = 0.400) [100] | 0.370 [155] |
| Neutrino scattering | $R_{ u_{\mu}}$ | 0.3072 ± 0.0033 | CDHS $(r = 0.393)$ [156] | 0.3091 [156] |
| reduino scattering | $R_{\overline{\nu}_{\mu}}$ | 0.382 ± 0.016 | CDIIO (7 = 0.555) [100] | 0.380 [156] |
| | κ | 0.5820 ± 0.0041 | CCFR [157] | 0.5830 [157] |
| | $R_{ u_e\overline{ u}_e}$ | $0.406^{+0.145}_{-0.135}$ | CHARM [158] | 0.33 [159] |
| | $(s_w^2)^{\text{Møller}}$ | 0.2397 ± 0.0013 | SLAC-E158 [160] | 0.2381 ± 0.0006 [161] |
| | $Q_W^{\text{Cs}}(55, 78)$ | -72.62 ± 0.43 | PDG2016 [159] | $-73.25 \pm 0.02 $ [159] |
| | $Q_W^{\mathrm{p}}(1,0)$ | 0.064 ± 0.012 | QWEAK [162] | 0.0708 ± 0.0003 [159] |
| D : | A_1 | $(-91.1 \pm 4.3) \times 10^{-6}$ | PVDIS [163] | $(-87.7 \pm 0.7) \times 10^{-6} $ [163] |
| Parity-violating scattering | A_2 | $(-160.8 \pm 7.1) \times 10^{-6}$ | 1 (100) | $(-158.9 \pm 1.0) \times 10^{-6}$ [163] |
| | $g_{VA}^{eu} - g_{VA}^{ed}$ | -0.042 ± 0.057 | SAMPLE ($\sqrt{Q^2} = 200 \text{MeV}$) [164] | -0.0360 [159] |
| | g_{VA} g_{VA} | -0.12 ± 0.074 | SAMPLE ($\sqrt{Q^2} = 125 \text{MeV}$) [164] | 0.0265 [159] |
| | $b_{ m SPS}$ | $-(1.47\pm0.42)\times10^{-4}\mathrm{GeV^{-2}}$ | SPS ($\lambda = 0.81$) [165] | $-1.56 \times 10^{-4} \mathrm{GeV^{-2}}$ [165] |
| | 9515 | $-(1.74\pm0.81)\times10^{-4}\mathrm{GeV^{-2}}$ | SPS ($\lambda = 0.66$) [165] | $-1.57 \times 10^{-4} \mathrm{GeV^{-2}}$ [165] |
| τ polarization | $\mathcal{P}_{	au}$ | 0.012 ± 0.058 | VENUS [166] | 0.028 [166] |
| , poterización | $\mathcal{A}_{\mathcal{P}}$ | 0.029 ± 0.057 | , 111.00 [100] | 0.021 [166] |
| Neutrino trident production | $\frac{\frac{\sigma}{\sigma^{\rm SM}}(\nu_{\mu}\gamma^* \to \nu_{\mu}\mu^+\mu^-)}$ | 0.82 ± 0.28 | CCFR [167–169] | 1 |
| $d_I \to u_J \ell \overline{\nu}_\ell(\gamma)$ | $\epsilon_{L,R,S,P,T}^{de_J}$ | See Ref. [128] | [170] | 0 |

Table 7: Low-energy observables included in Fit-2.

leptons (for e, μ, τ) and quarks (for u, d, b^{\S}); W couplings to leptons. The Higgs effective couplings include Higgs couplings to $ZZ, WW, \gamma\gamma, Z\gamma, gg, cc, bb, \tau\tau, \mu\mu$ as well as the Higgs total width. Z couplings to top quarks and the top-Yukawa coupling will be discussed in Fit-3 results. In addition to the above effective couplings, which are independent of the operator basis, the figure also shows three anomalous triple gauge couplings (aTGCs): $g_{1,Z}, \kappa_{\gamma}, \lambda_{Z}$. The exact definition of each parameter or Wilson coefficient shown in this and following plots can be found in Ref. [128].

From the Fit-1 results one can deduce that future e^+e^- colliders can improve our knowledge of electroweak effective couplings by a few orders of magnitude. The improvement will mainly come from dedicated runs at the Z-pole and WW-threshold for circular e^+e^- , and lower-energy stages at linear e^+e^- via the radiative return process. The higher energy stages of any e^+e^- have little impact on most of the effective couplings except for: Z cou-

Note that the universality assumption for 1st and 2nd generation quarks implies that couplings for c(s) quarks are as same as for u(d) quarks

 $[\]P Z$ couplings to neutrinos and W couplings to quarks are not listed separately since they are related to the other couplings for operators up to dimension 6 in SMEFT.




Figure 9: Ratios of the measurement precision (shown in Figure 8) to the one assuming perfect EW measurements (Z pole + W mass/width) in the constrained- Γ_H fit. Results are only shown for Higgs couplings and aTGCs with ratios significantly larger than one. For CEPC/FCC-ee, we also show (with the thin bars) the results without their Z-pole measurements.

plings to electrons, which as illustrated in Fig. 7 are related to eeZH contact interactions which increase quadratically with energy; W couplings to leptons, simply due to increased statistics from WW production. At muon colliders, the expected improvements will be mainly for Z couplings to muons and W couplings to leptons for the same reason as for higher-energy stages of e^+e^- . The aTGCs can benefit a lot from higher-energy stages of e^+e^- or muon colliders, in particular for λ_Z , which is sensitive to the transverse modes of W bosons. In general, circular e^+e^- can deliver the best precision for electroweak effective couplings, while linear e^+e^- can bring comparable improvements in particular when the Giga-Z option with beam polarization is included.

For results on Higgs effective couplings, HL-LHC will push the constraints to 2-5% for many couplings while future e^+e^- or muon colliders will improve further to 1% or below. In addition, future e^+e^- can bring a qualitative difference when the Higgs total width is a free parameter in the global fit. High-energy muon colliders can also bring this advantage when a dedicated scan at the Higgs pole is included. There is a potential at the HL-LHC to determine the Higgs total width using off-shell Higgs measurements [171,172] with an uncertainty of 0.75 MeV [173,174]**. This piece of input has not been included in the global fit since the full EFT treatment for this measurement is not yet available [175].

It is worth noting the interplay between Higgs couplings and EWPOs as shown in Fig. 9: For circurlar e^+e^- the achievable precision for Higgs couplings and aTGCs improves by a factor of around 2 when including measurements at the Z-pole and WW threshold. The possibility for similar improvements at muon colliders depends on their ability to measure EWPOs with high precision, which likely would require a dedicated Z-pole run. At linear e^+e^- colliders, the availability of beam polarization helps to break degeneracies in the SMEFT parameter space already from measurements at 250/380 GeV alone, and thus the impact of EW precision data on the other couplings is less significant. It is also worth to note the synergies on Higgs rare decays $(H \to \gamma \gamma, Z \gamma, \mu \mu)$ between HL-LHC and future lepton colliders which play an important role in the global fits.

The results of Fit-2 are shown in Fig. 10-12 for present measurement results and fu-

^{||}Alternatively speaking, the Higgs total width can be indirectly determined at future e^+e^- without any assumption on possible decay modes.

^{**}This uncertainty is likely to be improved once the WW channel is employed in addition to the current ZZ analyses.

ture e^+e^- projections including the following: 1σ relative uncertainties for electroweak effective couplings^{††}; 1σ absolute uncertainties for the Wilson Coefficients of 4-fermion operators. The conclusions for electroweak couplings are consistent with that from Fit-1. For 4-fermion operators the higher-energy stages of future e^+e^- will bring much more profound improvements since the sensitivity to 4-fermion interactions grows quadratically as energy increases. Even with the same energy, linear e^+e^- can deliver much higher sensitivity than circular ones since there are crucial degenericies among 4-fermion operators that the beam polarization can help lift.

The results of Fit-3 are shown in Fig. 13–15 for 95% C.L. bounds on the Wilson Coefficients of various top-quark operators, for LHC and future e^+e^- . LHC will bring invaluable constraints on many top-quark operators while those related to top-quark electroweak couplings will be improved by future e^+e^- in particular when the collision energies above 500 GeV are envisaged. There are a range of eett 4-fermion operators whose degeneracties can not be lifted unless there are measurements with at least two distinct energies well above the $t\bar{t}$ threshold. The uncertainty of top-Yukawa coupling is encapsulated in the bound on $C_{t\phi}$, for which synergies between LHC and future e^+e^- play an important role. There will be no direct constraint on the top Yukawa coupling from circular e^+e^- below 500 GeV; thus the projection from HL-LHC will provide the best constraints in such a scenario. The converted 1σ relative uncertainties on the top Yukawa coupling from the global and indivual fits are shown in Tab. 8. It is worth noting that ILC running at 550 GeV would improve top-Yukawa coupling significantly compared to running at nominal 500 GeV.

The result on the triple Higgs coupling (λ_{hhh}) from a global fit performed by the ESG can be found in Fig. 11 of Ref. [46]. A fit of λ_{hhh} has currently not been included in Fit-1, but is not expected to be much different.

Theory uncertainties may play a significant role in either electroweak couplings or Higgs couplings in the global fits. The impact on Higgs couplings was discussed by the ESG in Tab. 10 and 11 in Ref. [46]. One major challenge will come from the intrinsic error of the SM prediction for $e^+e^- \to ZH$ cross section, which would be around 0.5% with NNLO EW correction and will be significant enough to affect the precision of HZZ and HWWcouplings. In addition, an important parametric error is due to the bottom-quark mass uncertainty ($\sim 13 \text{ MeV}$) which would affect the bottom-Yukawa coupling precision. The Higgs mass uncertainty is another source of parametric error, which will become subdominant if a precision of about 10 MeV can be reached. The detailed impact of theory errors on various effective Higgs and EW couplings is quantified in the context of Fit-1 and summarized in Fig. 16. The parametric errors for all EWPOs and Higgs production and decay rates are included, in particular the projected uncertainties of the bottom-quark and Higgs masses mentioned above, the strong coupling constant ($\Delta \alpha_s = 0.0002$), charm quark mass $(\Delta m_c = 7 \text{ MeV})$ and top quark mass $(\Delta m_t = 50 \text{ MeV})$. The intrinsic theory errors for EWPOs are as introduced in Tab. 4, while the intrinsic errors for Higgs production and decay rates can be found in Ref. [46]. It is worth noting that only the theory errors for SM predictions are included while the theory errors related to SMEFT (e.g. the uncertainty due to truncation up to linear effects from dimension-6 operators) are not evaluated here. More

 $^{^{\}dagger\dagger}$ With the flavor assumption relaxed, Z couplings to u,d,c,s are treated separately.




Figure 10: Precision reach on the 4-fermion operators and electroweak effective couplings from a SMEFT global fit at various future lepton colliders. "LEP+SLC+SLD" represents current measurements which are always combined in the future collider scenarios. The horizontal white line for ILC illustrates the global fit results when the pole observables from its Giga-Z option are included.

details can be found in Ref. [128]. We can clearly see the need to improve in particular the intrinsic theory errors in order to make use of the precision that future experiments could deliver.

The results from global fits can be also interpreted in terms of constraints on simple BSM benchmark models with a small set of parameters (i.e. only a small set of SMEFT operators are generated in each model). Three examples are studied for the Fit-2 global fit results. The first example considers a flavor-universal 4-fermion contact interaction, which can be described by the O_{2B} operator mentioned in section 4.1. Figure 17 shows the bounds on the scale of this operator one can draw for different future colliders. The same bound on O_{2B} can also be interpreted in the Y-universal Z' model [176] as a bound on the gauge coupling $g_{Z'}$ versus Z' mass, as shown in Fig. 18. The third benchmark model extends the SM by two leptoquark multiplets, one being an SU(2) singlet, while the other is a SU(2) triplet [177,178]. This model can generate various 2-lepton-2-quark contact interactions by integrating out the heavy leptoquark fields in the t-channel. The bounds on the ratios of Yukawa couplings λ_i over leptoquark mass M_i are shown in Fig. 19, where i = 1 (3) refers to the singlet (triplet) leptoquark.




Figure 11: Figure 10 continued.




Figure 12: Figure 11 continued.




Figure 13: The 95% probability bounds on the Wilson coefficients for dimension-six operators that affect the top-quark production and decay measurements after Run 2 of the LHC (in dark red) and prospects for the bounds expected after completion of the complete LHC program, including the high-luminosity stage (in light red). The individual bounds obtained from a single-parameter fit are shown as solid bars, while the global or marginalised bounds obtained fitting all Wilson coefficients at once are indicated by the full bars (pale shaded region in each bar).




Figure 14: Comparison of current LHC constraints on various top-sector Wilson coefficients with HL-LHC ones, and those derived from ILC runs at 250, 500 and 1000 GeV. The limits on the $q\bar{q}t\bar{t}$ and C_{tG} coefficients are not shown, since the e^+e^- collider measurements considered are not sensitive to them, but all operators are included in the global fit. The improvement expected from HL-LHC on these coefficients is shown in Fig. 13. The additional bar included for $C_{t\varphi}$ in light green shows the effect on this operator of ILC working at 550 GeV. The solid bars provide the individual limits of the single-parameter fit and the shaded ones the marginalised limits of the global fit.

| Values in % units | | LHC | HL-LHC | ILC500 | ILC550 | ILC1000 | CLIC |
|-------------------|------------|------|--------|--------|--------|---------|------|
| δy_t | Global fit | 12.2 | 5.06 | 3.14 | 2.60 | 1.48 | 2.96 |
| | Indiv. fit | 10.2 | 3.70 | 2.82 | 2.34 | 1.41 | 2.52 |

Table 8: Uncertainties for the top-quark Yukawa coupling at 68% probability for different scenarios, in percentage. The ILC500, ILC550 and CLIC scenarios also include the HL-LHC. The ILC1000 scenario includes also ILC500 and HL-LHC.




Figure 15: Constraints expected on top-sector Wilson coefficients from a combination of HL-LHC and lepton collider data. The limits on the $q\bar{q}t\bar{t}$ and C_{tG} coefficients are not shown, since the e^+e^- collider measurements considered are not sensitive to them, but all operators are included in the global fit. The improvement expected from HL-LHC on these coefficients is shown in Fig. 13. The solid bars provide the individual limits of the single-parameter fit and the pale shaded ones the marginalised limits of the global fit. The results for ILC and CLIC are based on a combination of both low- and high-energy run scenarios.




Figure 16: The impact of parametric and intrinsic theory errors on effective EW and Higgs couplings. The impact is plotted in terms of the ratio of uncertainties when theory errors are included in the global fit over the case when they are not included. The dark bars are when only parametric errors are included; the light bars are when both parametric and intrinsic errors as projected in future are included; the thin lines are when current full theory errors are included.




Figure 17: 95% C.L. exclusion reach of different colliders on four-fermion contact interactions from the operator O_{2B} (numbers for ESG are taken from Ref. [179]).




Figure 18: 95% C.L. exclusion reach of different colliders on the Y-Universal Z' model parameters.




Figure 19: 95% C.L. exclusion reach of different colliders on the leptoquark model parameters. Only future e^+e^- scenarios with energies below the $t\bar{t}$ threshold have been considered since the analysis did not include any top-quark observables.

5 Conclusions

- For "canonical" electroweak precision measurements (Z-pole, WW threshold), circular e^+e^- colliders (FCC-ee, CEPC) have in general a higher sensitivity than linear colliders (ILC, CLIC) due to the high luminosity at center-of-mass energies below 200 GeV. Beam polarization at the linear colliders improves their sensitivity and can help to control systematics. In particular, for a linear collider run on the Z pole, beam polarization would enable measurements of the asymmetry parameters A_f with a precision that is only a factor of a few worse than for circular colliders, in spite of several orders of magnitude larger statistics for Z-pole physics at circular colliders.
- At center-of-mass energies $\sqrt{s} \lesssim 160$ GeV, the beam energy can be precisely calibrated using resonant depolarization at circular e^+e^- colliders, thus enabling very precise determinations of Z and W masses and widths. Linear colliders need a physical mass for energy calibration, which could be the Z mass (with 25 ppm precision from LEP) or possibly hadron (kaon and Λ) masses. Using the latter may put an energy calibration with 2ppm precision within reach, but requires further investigation.
- For many of the most precisely measurable precision observables at linear colliders, the most significant source of experimental systematics stems from the polarization calibration. For the circular colliders, on the other hand, modeling uncertainties for hadronic final states appear to be the dominant systematic error source.
- All e^+e^- Higgs factory colliders are similarly affected by a class of systematic uncertainties due to QCD and hadronization modeling, in particular for heavy-flavor final states.
- At any proposed e^+e^- collider it will be possible to measure the W mass with a precision of a few MeV or even better, thus conclusively resolving the recent discrepancy among different W mass determinations at hadron colliders [13].
- Experiments at lower-energy e^+e^- colliders, lepton-proton or lepton-ion colliders, or neutrino scattering facilities can deliver complementary information about electroweak quantities, such as the running electroweak mixing angle at low scales, or the separate determination of up- and down-quark electroweak couplings.
- High-energy lepton colliders (e^+e^- or $\mu^+\mu^-$ with $\sqrt{s} > 1\,\text{TeV}$) are effectively boson colliders. The total cross-section for many production processes is dominated by VBF/VBS-type contributions. However, for studies of BSM effects at very high invariant masses, non-VBF processes become typically more dominant.
- At multi-TeV lepton colliders, multiple electroweak gauge-boson production is ubiquitous, and new theoretical tools are needed for calculating and simulating these effects.
- Hadron colliders and lepton colliders offer complementary information about potential new BSM physics: Measurements of EWPOs at future Higgs factories offer indirect sensitivity to heavy new physics at scales of several TeV, which in many cases substantially exceeds the reach of LHC / HL-LHC. Furthermore, they have unique sensitivity

to very weakly coupled new particles with smaller masses. A hadron collider with 100 TeV center-of-mass energy, on the other hand, is able to directly produce new particles in the parameter space covered by indirect EWPO tests for most BSM scenarios (see EF BSM report [180] for more information). However, multi-TeV e^+e^- or $\mu^+\mu^-$ colliders, while having lower statistical power than a $\mathcal{O}(100)$ -TeV pp collider, can have an advantage for certain multi-boson studies due to the well-defined initial state and clean event signatures. The specific benefits of hadron and lepton colliders depend on the type of BSM physics, and thus a combination of both collider types is needed for the broadest coverage of new physics scenarios.

- Assuming that any new particles are heavy, a model-independent parametrization of the new-physics reach of different colliders is given by the SMEFT framework, where the SM is extended by higher-dimensional operators, with the leading contribution for most processes entering at dimension 6. Several subsets of such dimension-6 operators have been investigated in a set of global fits across a large number of observables: (a) operators contributing to electroweak gauge-boson interactions; (b) operators contributing to Higgs interactions; (c) operators contributing to top-quark interactions; and (d) operators contributing to four-fermion contact interactions.
- Generally, future lepton colliders have a better reach for many of the aforementioned operators than the HL-LHC. Circular e^+e^- colliders have the best sensitivity to electroweak operators, due to the large statistical precision of Z pole and WW threshold measurements. All lepton colliders are comparable in their reach for Higgs operators, although a multi-TeV muon collider cannot constrain exotic Higgs decays in a model-independent way^{‡‡}. Top-quark and four-fermion operators are best constrained at machines with $\sqrt{s} \geq 500$ GeV, and measurements at two or more values of \sqrt{s} are crucial for breaking degeneracies. Many constraints on top-quark operators are improved by combining e^+e^- and (HL-)LHC inputs and exploiting synergies between them.
- Some of the same SMEFT operators contribute to EW precision quantities and to Higgs observables or anomalous gauge-boson coupings (aGCs). At circular e^+e^- colliders, measurements from a high-luminosity Z-pole run can improve the constraints on Higgs couplings and aGCs by a factor of up to 2. At linear e^+e^- colliders, beam polarization provides additional information for Higgs/aGC measurements, and inputs from Z-pole data are less important.
- Low-energy measurements (below the W/Z mass scales) are needed to close the fit for four-fermion operators, when allowing non-universality among the three fermion generations.
- At this point, not enough information was available to include pp colliders beyond the LHC (such as HE-LHC or a $\mathcal{O}(100)$ -TeV collider) in the global fit. It is likely that these machines have superior sensitivity to many energy-dependent operators, such as

 $^{^{\}ddagger \ddagger}$ At future e^+e^- Higgs factories or a 125-GeV muon collider run, a model-independent study of exotic Higgs decays is possible.

4-fermion operators involving quarks and several operators that mediate multi-boson interactions.

Acknowledgments

The conveners are grateful to C. Grojean and M. Shapiro for detailed reviews of the draft report, and to S. Eno for many helpful comments.

References

- [1] UA1 collaboration, Experimental Observation of Isolated Large Transverse Energy Electrons with Associated Missing Energy at $\sqrt{s} = 540$ GeV, Phys. Lett. B 122 (1983) 103.
- [2] UA2 collaboration, Observation of Single Isolated Electrons of High Transverse Momentum in Events with Missing Transverse Energy at the CERN anti-p p Collider, Phys. Lett. B 122 (1983) 476.
- [3] UA1 collaboration, Experimental Observation of Lepton Pairs of Invariant Mass Around 95-GeV/c**2 at the CERN SPS Collider, Phys. Lett. B 126 (1983) 398.
- [4] UA2 collaboration, Evidence for $Z^0 \to e^+e^-$ at the CERN $\overline{p}p$ Collider, Phys. Lett. B 129 (1983) 130.
- [5] T. Appelquist and J. Carazzone, Infrared Singularities and Massive Fields, Phys. Rev. D 11 (1975) 2856.
- [6] Particle Data Group collaboration, Review of Particle Physics, PTEP 2020 (2020) 083C01.
- [7] C. Sturm, Leptonic contributions to the effective electromagnetic coupling at four-loop order in QED, Nucl. Phys. B 874 (2013) 698 [1305.0581].
- [8] A. Blondel et al., Theory report on the 11th FCC-ee workshop, 1905.05078.
- [9] M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, A new evaluation of the hadronic vacuum polarisation contributions to the muon anomalous magnetic moment and to α(m²_Z), Eur. Phys. J. C 80 (2020) 241 [1908.00921].
- [10] A. Keshavarzi, D. Nomura and T. Teubner, g-2 of charged leptons, $\alpha(M_Z^2)$, and the hyperfine splitting of muonium, Phys. Rev. D 101 (2020) 014029 [1911.00367].
- [11] F. Burger, K. Jansen, M. Petschlies and G. Pientka, Leading hadronic contributions to the running of the electroweak coupling constants from lattice QCD, JHEP 11 (2015) 215 [1505.03283].

- [12] M. Cè, A. Gérardin, G. von Hippel, H.B. Meyer, K. Miura, K. Ottnad et al., The hadronic running of the electromagnetic coupling and the electroweak mixing angle from lattice QCD, 2203.08676.
- [13] CDF collaboration, High-precision measurement of the W boson mass with the CDF II detector, Science 376 (2022) 170.
- [14] R. Schwienhorst et al., Report of the Topical Group on Top quark physics and heavy flavor production for Snowmass 2021, in 2022 Snowmass Summer Study, 2022 [2209.11267].
- [15] ATLAS, CMS collaboration, Snowmass White Paper Contribution: Physics with the Phase-2 ATLAS and CMS Detectors, in Snowmass 2021 proceedings (2022).
- [16] S. Yang, Y. Fu, M. Liu, R. Zhang, T.-J. Hou, C. Wang et al., Reduction of the electroweak correlation in the PDF updating by using the forward-backward asymmetry of Drell-Yan process, 2108.06550.
- [17] L. Buonocore, M. Grazzini, S. Kallweit, C. Savoini and F. Tramontano, *Mixed QCD-EW corrections to* $pp \rightarrow \ell\nu_{\ell} + X$ *at the LHC*, *Phys. Rev. D* **103** (2021) 114012 [2102.12539].
- [18] R. Bonciani, L. Buonocore, M. Grazzini, S. Kallweit, N. Rana, F. Tramontano et al., Mixed Strong-Electroweak Corrections to the Drell-Yan Process, Phys. Rev. Lett. 128 (2022) 012002 [2106.11953].
- [19] F. Buccioni, F. Caola, H.A. Chawdhry, F. Devoto, M. Heller, A. von Manteuffel et al., *Mixed QCD-electroweak corrections to dilepton production at the LHC in the high invariant mass region*, 2203.11237.
- [20] S. Camarda, L. Cieri and G. Ferrera, Drell-Yan lepton-pair production: qT resummation at N3LL accuracy and fiducial cross sections at N3LO, Phys. Rev. D 104 (2021) L111503 [2103.04974].
- [21] H. Baer et al., eds., The International Linear Collider Technical Design Report -Volume 2: Physics, 1306.6352.
- [22] P. Bambade et al., The International Linear Collider: A Global Project, 1903.01629.
- [23] ILC International Development Team collaboration, *The International Linear Collider: Report to Snowmass 2021*, in *Snowmass 2021 proceedings* (2022) [2203.07622].
- [24] L. Linssen, A. Miyamoto, M. Stanitzki and H. Weerts, eds., Physics and Detectors at CLIC: CLIC Conceptual Design Report, 1202.5940.
- [25] CLICDP, CLIC collaboration, The Compact Linear Collider (CLIC) 2018 Summary Report, 1812.06018.
- [26] FCC collaboration, FCC-ee: The Lepton Collider: Future Circular Collider Conceptual Design Report Volume 2, Eur. Phys. J. ST 228 (2019) 261.

- [27] G. Bernardi et al., The Future Circular Collider: a Summary for the US 2021 Snowmass Process, in Snowmass 2021 proceedings (2022) [2203.06520].
- [28] CEPC Study Group collaboration, CEPC Conceptual Design Report: Volume 2 Physics & Detector, 1811.10545.
- [29] CEPC ACCELERATOR STUDY GROUP collaboration, Snowmass2021 White Paper AF3-CEPC, 2203.09451.
- [30] H. Cheng et al., The Physics potential of the CEPC. Prepared for the US Snowmass Community Planning Exercise (Snowmass 2021), in Snowmass 2021 proceedings (2022) [2205.08553].
- [31] M. Bai et al., C^8 : A "Cool" Route to the Higgs Boson and Beyond, in Snowmass 2021 proceedings, 2021 [2110.15800].
- [32] S. Dasu et al., Strategy for Understanding the Higgs Physics: The Cool Copper Collider, in Snowmass 2021 proceedings, 2022 [2203.07646].
- [33] M. Narain et al., The Future of US Particle Physics The Snowmass 2021 Energy Frontier Report, in 2022 Snowmass Summer Study, 2022 [2211.11084].
- [34] C. Gohil, A. Latina, D. Schulte and S. Stapnes, "High-Luminosity CLIC Studies." https://cds.cern.ch/record/2687090, 2020.
- [35] LCC Physics Working Group collaboration, Tests of the Standard Model at the International Linear Collider, 1908.11299.
- [36] A. Blondel and P. Janot, FCC-ee overview: new opportunities create new challenges, Eur. Phys. J. Plus 137 (2022) 92 [2106.13885].
- [37] P. Roloff, "Precision measurements at CLIC." https://indico.fnal.gov/event/43576/contributions/189845/attachments/130766/159625/snowmass_ef04_meeting_roloff_02_07_2020.pdf.
- [38] ALEPH, DELPHI, L3, OPAL, SLD, LEP ELECTROWEAK WORKING GROUP, SLD ELECTROWEAK GROUP, SLD HEAVY FLAVOUR GROUP collaboration, Precision electroweak measurements on the Z resonance, Phys. Rept. 427 (2006) 257 [hep-ex/0509008].
- [39] J. Alcaraz Maestre, Revisiting QCD corrections to the forward-backward charge asymmetry of heavy quarks in electron-positron collisions at the Z pole: really a problem?, 2010.08604.
- [40] D. d'Enterria and C. Yan, Revised QCD effects on the $Z \to b\bar{b}$ forward-backward asymmetry, 2011.00530.
- [41] ALEPH collaboration, Determination of the LEP center-of-mass energy from Z gamma events: Preliminary, in 29th International Conference on High-Energy Physics, 10, 1998 [hep-ex/9810047].

- [42] G.W. Wilson, "Exploring Precision EW Measurement Potential of e^+e^- Colliders." https://indico.fnal.gov/event/50480/contributions/225096/attachments/148222/190348/GWWTalk_Oct8_2021.pdf.
- [43] J. Beyer and J. List, Interplay of Beam Polarisation and Systematic Uncertainties in Electroweak Precision Measurements at Future e⁺e⁻ Colliders, in 24th International Symposium on Spin Physics, 2, 2022 [2202.07385].
- [44] T. Mizuno, K. Fujii and J. Tian, Measurement of A_{LR} using radiative return at ILC 250, in Snowmass 2021 proceedings (2022) [2203.07944].
- [45] M. Baak et al., Working Group Report: Precision Study of Electroweak Interactions, in Community Summer Study 2013: Snowmass on the Mississippi, 2013 [1310.6708].
- [46] J. de Blas et al., Higgs Boson Studies at Future Particle Colliders, JHEP **01** (2020) 139 [1905.03764].
- [47] J. De Blas, G. Durieux, C. Grojean, J. Gu and A. Paul, On the future of Higgs, electroweak and diboson measurements at lepton colliders, JHEP 12 (2019) 117 [1907.04311].
- [48] D. d'Enterria et al., The strong coupling constant: State of the art and the decade ahead, in Snowmass 2021 proceedings (2022) [2203.08271].
- [49] P. Janot, Direct measurement of $\alpha_{QED}(m_Z^2)$ at the FCC-ee, JHEP **02** (2016) 053 [1512.05544].
- [50] M. Awramik, M. Czakon, A. Freitas and G. Weiglein, Precise prediction for the W boson mass in the standard model, Phys. Rev. D 69 (2004) 053006 [hep-ph/0311148].
- [51] I. Dubovyk, A. Freitas, J. Gluza, T. Riemann and J. Usovitsch, Electroweak pseudo-observables and Z-boson form factors at two-loop accuracy, JHEP 08 (2019) 113 [1906.08815].
- [52] K. Yumino and D. Jeans, Measuring the tau polarization at ILC, in Snowmass 2021 proceedings (2022) [2203.07668].
- [53] D.M. Asner et al., Upgrading SuperKEKB with a Polarized Electron Beam: Discovery Potential and Proposed Implementation, in Snowmass 2021 proceedings (2022).
- [54] J. Erler, Global Vision of Precision Measurements, in 55th Rencontres de Moriond on Electroweak Interactions and Unified Theories, 5, 2021 [2105.00217].
- [55] Y.X. Zhao, A. Deshpande, J. Huang, K.S. Kumar and S. Riordan, Neutral-Current Weak Interactions at an EIC, Eur. Phys. J. A 53 (2017) 55 [1612.06927].
- [56] R. Boughezal, A. Emmert, T. Kutz, S. Mantry, M. Nycz, F. Petriello et al., Neutral-Current Electroweak Physics and SMEFT Studies at the EIC, 2204.07557.

- [57] LHEC, FCC-HE STUDY GROUP collaboration, The Large Hadron-Electron Collider at the HL-LHC, J. Phys. G 48 (2021) 110501 [2007.14491].
- [58] FCC collaboration, FCC Physics Opportunities: Future Circular Collider Conceptual Design Report Volume 1, Eur. Phys. J. C 79 (2019) 474.
- [59] D. Britzger, M. Klein and H. Spiesberger, Electroweak physics in inclusive deep inelastic scattering at the LHeC, Eur. Phys. J. C 80 (2020) 831 [2007.11799].
- [60] D. Britzger, M. Klein and H. Spiesberger, Precision electroweak measurements at the LHeC and the FCC-eh, PoS EPS-HEP2021 (2022) 485 [2203.06237].
- [61] A. Blondel, A muon collider as Z factory, AIP Conf. Proc. 435 (1998) 597.
- [62] A. Freitas, *Precision Tests of the Standard Model*, *PoS* **TASI2020** (2021) 005 [2012.11642].
- [63] S. Jadach and M. Skrzypek, QED challenges at FCC-ee precision measurements, Eur. Phys. J. C 79 (2019) 756 [1903.09895].
- [64] S. Jadach, W. Płaczek and M. Skrzypek, QED exponentiation for quasi-stable charged particles: the $e^-e^+ \rightarrow W^-W^+$ process, Eur. Phys. J. C 80 (2020) 499 [1906.09071].
- [65] S. Frixione et al., Initial state QED radiation aspects for future e⁺e⁻ colliders, in Snowmass 2021 proceedings (2022) [2203.12557].
- [66] A. Freitas et al., Theoretical uncertainties for electroweak and Higgs-boson precision measurements at FCC-ee, 1906.05379.
- [67] X. Liu and Y.-Q. Ma, Multiloop corrections for collider processes using auxiliary mass flow, Phys. Rev. D 105 (2022) L051503 [2107.01864].
- [68] X. Liu and Y.-Q. Ma, AMFlow: a Mathematica package for Feynman integrals computation via Auxiliary Mass Flow, 2201.11669.
- [69] Q. Song and A. Freitas, On the evaluation of two-loop electroweak box diagrams for $e^+e^- \to HZ$ production, JHEP **04** (2021) 179 [2101.00308].
- [70] I. Dubovyk, A. Freitas, J. Gluza, K. Grzanka, M. Hidding and J. Usovitsch, Evaluation of multi-loop multi-scale Feynman integrals for precision physics, 2201.02576.
- [71] S. Dawson et al., Report of the Topical Group on Higgs Physics for Snowmass 2021: The Case for Precision Higgs Physics, in 2022 Snowmass Summer Study, 2022 [2209.07510].
- [72] ATLAS collaboration, Measurement of ZZ production in the $\ell\ell\nu\nu$ final state with the ATLAS detector in pp collisions at $\sqrt{s} = 13$ TeV, JHEP 10 (2019) 127 [1905.07163].

- [73] ATLAS collaboration, Measurement of fiducial and differential W^+W^- production cross-sections at $\sqrt{s} = 13$ TeV with the ATLAS detector, Eur. Phys. J. C **79** (2019) 884 [1905.04242].
- [74] ATLAS collaboration, Measurement of $W^{\pm}Z$ production cross sections and gauge boson polarisation in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, Eur. Phys. J. C 79 (2019) 535 [1902.05759].
- [75] ATLAS collaboration, Measurement of the $Z\gamma \to \nu \overline{\nu} \gamma$ production cross section in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector and limits on anomalous triple gauge-boson couplings, JHEP 12 (2018) 010 [1810.04995].
- [76] ATLAS collaboration, $ZZ \to \ell^+\ell^-\ell'^+\ell'^-$ cross-section measurements and search for anomalous triple gauge couplings in 13 TeV pp collisions with the ATLAS detector, Phys. Rev. D **97** (2018) 032005 [1709.07703].
- [77] CMS collaboration, Search for anomalous triple gauge couplings in WW and WZ production in lepton + jet events in proton-proton collisions at $\sqrt{s} = 13$ TeV, JHEP 12 (2019) 062 [1907.08354].
- [78] CMS collaboration, Measurement of electroweak WZ boson production and search for new physics in WZ + two jets events in pp collisions at $\sqrt{s} = 13$ TeV, Phys. Lett. B 795 (2019) 281 [1901.04060].
- [79] CMS collaboration, Measurements of the $pp \to WZ$ inclusive and differential production cross section and constraints on charged anomalous triple gauge couplings at $\sqrt{s} = 13$ TeV, JHEP **04** (2019) 122 [1901.03428].
- [80] CMS collaboration, Measurements of the pp \rightarrow ZZ production cross section and the Z \rightarrow 4 ℓ branching fraction, and constraints on anomalous triple gauge couplings at $\sqrt{s} = 13 \text{ TeV}$, Eur. Phys. J. C 78 (2018) 165 [1709.08601].
- [81] ATLAS collaboration, Evidence for the production of three massive vector bosons with the ATLAS detector, Phys. Lett. B 798 (2019) 134913 [1903.10415].
- [82] ATLAS collaboration, Study of WW γ and WZ γ production in pp collisions at $\sqrt{s} = 8$ TeV and search for anomalous quartic gauge couplings with the ATLAS experiment, Eur. Phys. J. C 77 (2017) 646 [1707.05597].
- [83] CMS collaboration, Observation of the production of three massive gauge bosons at $\sqrt{s} = 13$ TeV, 2006.11191.
- [84] CMS collaboration, Measurements of the $pp \to W\gamma\gamma$ and $pp \to Z\gamma\gamma$ cross sections and limits on anomalous quartic gauge couplings at $\sqrt{s} = 8$ TeV, JHEP 10 (2017) 072 [1704.00366].
- [85] ATLAS collaboration, Observation of electroweak production of two jets and a Z-boson pair with the ATLAS detector at the LHC, 2004.10612.

- [86] ATLAS collaboration, Evidence for electroweak production of two jets in association with a $Z\gamma$ pair in pp collisions at $\sqrt{s}=13$ TeV with the ATLAS detector, Phys. Lett. B 803 (2020) 135341 [1910.09503].
- [87] ATLAS collaboration, Search for the electroweak diboson production in association with a high-mass dijet system in semileptonic final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, Phys. Rev. D 100 (2019) 032007 [1905.07714].
- [88] ATLAS collaboration, Observation of electroweak production of a same-sign W boson pair in association with two jets in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, Phys. Rev. Lett. 123 (2019) 161801 [1906.03203].
- [89] ATLAS collaboration, Observation of electroweak $W^{\pm}Z$ boson pair production in association with two jets in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, Phys. Lett. B **793** (2019) 469 [1812.09740].
- [90] CMS collaboration, Evidence for vector boson scattering in events with four leptons and two jets in proton-proton collisions at sqrt(s) = 13 TeV,.
- [91] CMS collaboration, Measurements of production cross sections of WZ and same-sign WW boson pairs in association with two jets in proton-proton collisions at $\sqrt{s} = 13$ TeV, 2005.01173.
- [92] CMS collaboration, Measurement of the cross section for electroweak production of a Z boson, a photon and two jets in proton-proton collisions at $\sqrt{s} = 13$ TeV and constraints on anomalous quartic couplings, JHEP **06** (2020) 076 [2002.09902].
- [93] CMS collaboration, Search for anomalous electroweak production of vector boson pairs in association with two jets in proton-proton collisions at 13 TeV, Phys. Lett. B 798 (2019) 134985 [1905.07445].
- [94] CMS collaboration, Measurement of vector boson scattering and constraints on anomalous quartic couplings from events with four leptons and two jets in proton-proton collisions at √s = 13 TeV, Phys. Lett. B 774 (2017) 682 [1708.02812].
- [95] CMS collaboration, Measurement of differential cross sections for Z boson pair production in association with jets at $\sqrt{s} = 8$ and 13 TeV, Phys. Lett. B **789** (2019) 19 [1806.11073].
- [96] "CMS summary of limits on anomalous triple and quartic gauge couplings." https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResultsSMPaTGC.
- [97] "ATLAS experiment public results standard model physics." https://twiki.cern.ch/twiki/bin/view/AtlasPublic/StandardModelPublicResults.
- [98] P. Azzi et al., Report from Working Group 1: Standard Model Physics at the HL-LHC and HE-LHC, in Report on the Physics at the HL-LHC, and Perspectives for the HE-LHC, A. Dainese, M. Mangano, A.B. Meyer, A. Nisati, G. Salam and M.A. Vesterinen, eds., vol. 7, pp. 1–220 (2019), DOI [1902.04070].

- [99] C. Grojean, M. Montull and M. Riembau, *Diboson at the LHC vs LEP*, *JHEP* **03** (2019) 020 [1810.05149].
- [100] C. Garcia-Garcia, M. Herrero and R.A. Morales, Unitarization effects in EFT predictions of WZ scattering at the LHC, Phys. Rev. D 100 (2019) 096003 [1907.06668].
- [101] X. Li, H. Xu, C. Yang, C. Zhang and S.-Y. Zhou, Positivity in Multifield Effective Field Theories, Phys. Rev. Lett. 127 (2021) 121601 [2101.01191].
- [102] C. de Rham, S. Kundu, M. Reece, A.J. Tolley and S.-Y. Zhou, Snowmass White Paper: UV Constraints on IR Physics, in Snowmass 2021 proceedings, 2022 [2203.06805].
- [103] W. Shepherd, SMEFT at the LHC and Beyond: A Snowmass White Paper, in 2022 Snowmass Summer Study, 2022 [2203.07406].
- [104] J. Marshall, A. Münnich and M. Thomson, Performance of Particle Flow Calorimetry at CLIC, Nucl. Instrum. Meth. A 700 (2013) 153 [1209.4039].
- [105] C. Benedetti et al., Linear collider based on laser-plasma accelerators, in 2022 Snowmass Summer Study, 3, 2022 [2203.08366].
- [106] V. Shiltsev and F. Zimmermann, *Modern and Future Colliders*, *Rev. Mod. Phys.* **93** (2021) 015006 [2003.09084].
- [107] A. Costantini, F. De Lillo, F. Maltoni, L. Mantani, O. Mattelaer, R. Ruiz et al., Vector boson fusion at multi-TeV muon colliders, JHEP 09 (2020) 080 [2005.10289].
- [108] T. Han, Y. Ma and K. Xie, *High Energy Leptonic Collisions and Electroweak Parton Distribution Functions*, 2007.14300.
- [109] M. Beyer, W. Kilian, P. Krstonosic, K. Monig, J. Reuter, E. Schmidt et al., Determination of New Electroweak Parameters at the ILC - Sensitivity to New Physics, Eur. Phys. J. C 48 (2006) 353 [hep-ph/0604048].
- [110] C. Fleper, W. Kilian, J. Reuter and M. Sekulla, Scattering of W and Z Bosons at High-Energy Lepton Colliders, Eur. Phys. J. C 77 (2017) 120 [1607.03030].
- [111] S. Green, Calorimetry at a Future Linear Collider, Ph.D. thesis, Cambridge U., 2017. 10.17863/CAM.15859.
- [112] D. Buarque Franzosi et al., Vector boson scattering processes: Status and prospects, Rev. Phys. 8 (2022) 100071 [2106.01393].
- [113] R. Ruiz, A. Costantini, F. Maltoni and O. Mattelaer, *The Effective Vector Boson Approximation in High-Energy Muon Collisions*, 2111.02442.
- [114] T. Han, Y. Ma and K. Xie, Electroweak fragmentation at high energies: A Snowmass White Paper, in 2022 Snowmass Summer Study, 2022 [2203.11129].

- [115] V. Bertone, M. Cacciari, S. Frixione and G. Stagnitto, The partonic structure of the electron at the next-to-leading logarithmic accuracy in QED, JHEP 03 (2020) 135 [1911.12040].
- [116] C. Paranjape, D. Stolarski and Y. Wu, Tree-level Interference in VBF production of Vh, in 2022 Snowmass Summer Study, 2022 [2203.05729].
- [117] B. Abbott et al., Anomalous quartic gauge couplings at a muon collider, in 2022 Snowmass Summer Study, 2022 [2203.08135].
- [118] T. Yang, S. Qian, Z. Guan, C. Li, F. Meng, J. Xiao et al., Longitudinally polarized ZZ scattering at a muon collider, Phys. Rev. D 104 (2021) 093003 [2107.13581].
- [119] L. Huang, S.D. Lane, I.M. Lewis and Z. Liu, Electroweak Restoration at the LHC and Beyond: The Vh Channel, Phys. Rev. D 103 (2021) 053007 [2012.00774].
- [120] A. Apyan, C. Mwewa, L. Nedic, M.-A. Pleier and K. Potamianos, Sensitivity to longitudinal vector boson scattering in W[±]W[±]jj at future hadron colliders, in 2022 Snowmass Summer Study, 2022 [2203.07994].
- [121] W. Buchmuller and D. Wyler, Effective Lagrangian Analysis of New Interactions and Flavor Conservation, Nucl. Phys. B 268 (1986) 621.
- [122] B. Grzadkowski, M. Iskrzynski, M. Misiak and J. Rosiek, *Dimension-Six Terms in the Standard Model Lagrangian*, *JHEP* **10** (2010) 085 [1008.4884].
- [123] G. Buchalla, O. Catà and C. Krause, Complete Electroweak Chiral Lagrangian with a Light Higgs at NLO, Nucl. Phys. B 880 (2014) 552 [1307.5017].
- [124] G. Buchalla, O. Cata, A. Celis and C. Krause, Note on Anomalous Higgs-Boson Couplings in Effective Field Theory, Phys. Lett. B 750 (2015) 298 [1504.01707].
- [125] I. Brivio and M. Trott, The Standard Model as an Effective Field Theory, Phys. Rept. 793 (2019) 1 [1706.08945].
- [126] M.E. Peskin and T. Takeuchi, A New constraint on a strongly interacting Higgs sector, Phys. Rev. Lett. 65 (1990) 964.
- [127] G. Altarelli and R. Barbieri, Vacuum polarization effects of new physics on electroweak processes, Phys. Lett. B 253 (1991) 161.
- [128] J. de Blas, Y. Du, C. Grojean, J. Gu, V. Miralles, M.E. Peskin et al., Global SMEFT Fits at Future Colliders, in 2022 Snowmass Summer Study, 6, 2022 [2206.08326].
- [129] S. Weinberg, Baryon and Lepton Nonconserving Processes, Phys. Rev. Lett. 43 (1979) 1566.
- [130] F. Wilczek and A. Zee, Operator Analysis of Nucleon Decay, Phys. Rev. Lett. 43 (1979) 1571.

- [131] L.F. Abbott and M.B. Wise, *The Effective Hamiltonian for Nucleon Decay*, *Phys. Rev. D* **22** (1980) 2208.
- [132] L. Lehman, Extending the Standard Model Effective Field Theory with the Complete Set of Dimension-7 Operators, Phys. Rev. D 90 (2014) 125023 [1410.4193].
- [133] L. Lehman and A. Martin, Low-derivative operators of the Standard Model effective field theory via Hilbert series methods, JHEP **02** (2016) 081 [1510.00372].
- [134] B. Henning, X. Lu, T. Melia and H. Murayama, 2, 84, 30, 993, 560, 15456, 11962, 261485, ...: Higher dimension operators in the SM EFT, JHEP 08 (2017) 016 [1512.03433].
- [135] T. Barklow, K. Fujii, S. Jung, R. Karl, J. List, T. Ogawa et al., Improved Formalism for Precision Higgs Coupling Fits, Phys. Rev. D 97 (2018) 053003 [1708.08912].
- [136] A. Falkowski, M. González-Alonso and K. Mimouni, Compilation of low-energy constraints on 4-fermion operators in the SMEFT, JHEP 08 (2017) 123 [1706.03783].
- [137] A. Efrati, A. Falkowski and Y. Soreq, *Electroweak constraints on flavorful effective theories*, *JHEP* **07** (2015) 018 [1503.07872].
- [138] G.F. Giudice, C. Grojean, A. Pomarol and R. Rattazzi, *The Strongly-Interacting Light Higgs*, *JHEP* **06** (2007) 045 [hep-ph/0703164].
- [139] R. Barbieri, A. Pomarol, R. Rattazzi and A. Strumia, *Electroweak symmetry breaking after LEP-1 and LEP-2*, *Nucl. Phys. B* **703** (2004) 127 [hep-ph/0405040].
- [140] J. Ellis, M. Madigan, K. Mimasu, V. Sanz and T. You, Top, Higgs, Diboson and Electroweak Fit to the Standard Model Effective Field Theory, JHEP 04 (2021) 279 [2012.02779].
- [141] SMEFIT collaboration, Combined SMEFT interpretation of Higgs, diboson, and top quark data from the LHC, JHEP 11 (2021) 089 [2105.00006].
- [142] E. Vryonidou and C. Zhang, Dimension-six electroweak top-loop effects in Higgs production and decay, JHEP 08 (2018) 036 [1804.09766].
- [143] G. Durieux, J. Gu, E. Vryonidou and C. Zhang, *Probing top-quark couplings indirectly at Higgs factories*, *Chin. Phys. C* **42** (2018) 123107 [1809.03520].
- [144] S. Jung, J. Lee, M. Perelló, J. Tian and M. Vos, *Higgs*, top and electro-weak precision measurements at future e⁺e⁻ colliders; a combined effective field theory analysis with renormalization mixing, 2006.14631.
- [145] J. De Blas et al., HEPfit: a code for the combination of indirect and direct constraints on high energy physics models, Eur. Phys. J. C 80 (2020) 456 [1910.14012].

- [146] S. Chen, A. Glioti, R. Rattazzi, L. Ricci and A. Wulzer, *Learning from radiation at a very high energy lepton collider*, *JHEP* **05** (2022) 180 [2202.10509].
- [147] C. Aimè et al., Muon Collider Physics Summary, 2203.07256.
- [148] MUON COLLIDER collaboration, The physics case of a 3 TeV muon collider stage, in 2022 Snowmass Summer Study, 2022 [2203.07261].
- [149] J. de Blas, J. Gu and Z. Liu, Higgs Precision at a 125 GeV Muon Collider, 2203.04324.
- [150] M. Diehl and O. Nachtmann, Optimal observables for the measurement of three gauge boson couplings in e+ e- -> W+ W-, Z. Phys. C 62 (1994) 397.
- [151] G. Durieux, M. Perelló, M. Vos and C. Zhang, Global and optimal probes for the top-quark effective field theory at future lepton colliders, JHEP 10 (2018) 168 [1807.02121].
- [152] CHARM-II collaboration, Precision measurement of electroweak parameters from the scattering of muon-neutrinos on electrons, Phys. Lett. B 335 (1994) 246.
- [153] J. Erler and S. Su, The Weak Neutral Current, Prog. Part. Nucl. Phys. 71 (2013) 119 [1303.5522].
- [154] Particle Data Group collaboration, Review of Particle Physics, Chin. Phys. C 38 (2014) 090001.
- [155] CHARM collaboration, A Precise Determination of the Electroweak Mixing Angle from Semileptonic Neutrino Scattering, Z. Phys. C 36 (1987) 611.
- [156] A. Blondel et al., Electroweak Parameters From a High Statistics Neutrino Nucleon Scattering Experiment, Z. Phys. C 45 (1990) 361.
- [157] CCFR, E744, E770 collaboration, A Precision measurement of electroweak parameters in neutrino nucleon scattering, Eur. Phys. J. C 1 (1998) 509 [hep-ex/9701010].
- [158] CHARM collaboration, Experimental Verification of the Universality of ν_e and ν_{μ} Coupling to the Neutral Weak Current, Phys. Lett. B 180 (1986) 303.
- [159] Particle Data Group collaboration, Review of Particle Physics, Chin. Phys. C 40 (2016) 100001.
- [160] SLAC E158 collaboration, Precision measurement of the weak mixing angle in Moller scattering, Phys. Rev. Lett. 95 (2005) 081601 [hep-ex/0504049].
- [161] A. Czarnecki and W.J. Marciano, Electroweak radiative corrections to polarized Moller scattering asymmetries, Phys. Rev. D 53 (1996) 1066 [hep-ph/9507420].
- [162] QWEAK collaboration, First Determination of the Weak Charge of the Proton, Phys. Rev. Lett. 111 (2013) 141803 [1307.5275].

- [163] PVDIS collaboration, Measurement of parity violation in electron-quark scattering, Nature **506** (2014) 67.
- [164] E.J. Beise, M.L. Pitt and D.T. Spayde, The SAMPLE experiment and weak nucleon structure, Prog. Part. Nucl. Phys. **54** (2005) 289 [nucl-ex/0412054].
- [165] A. Argento et al., Electroweak Asymmetry in Deep Inelastic Muon Nucleon Scattering, Phys. Lett. B 120 (1983) 245.
- [166] VENUS collaboration, Measurement of tau polarization in e+e- annihilation at $s^{**}(1/2) = 58$ -GeV, Phys. Lett. B **403** (1997) 155 [hep-ex/9703003].
- [167] CHARM-II collaboration, First observation of neutrino trident production, Phys. Lett. B 245 (1990) 271.
- [168] CCFR collaboration, Neutrino tridents and W Z interference, Phys. Rev. Lett. 66 (1991) 3117.
- [169] W. Altmannshofer, S. Gori, M. Pospelov and I. Yavin, Neutrino Trident Production: A Powerful Probe of New Physics with Neutrino Beams, Phys. Rev. Lett. 113 (2014) 091801 [1406.2332].
- [170] M. González-Alonso and J. Martin Camalich, Global Effective-Field-Theory analysis of New-Physics effects in (semi)leptonic kaon decays, JHEP 12 (2016) 052 [1605.07114].
- [171] F. Caola and K. Melnikov, Constraining the Higgs boson width with ZZ production at the LHC, Phys. Rev. D 88 (2013) 054024 [1307.4935].
- [172] J.M. Campbell, R.K. Ellis and C. Williams, Bounding the Higgs Width at the LHC Using Full Analytic Results for $gg > e^-e^+\mu^-\mu^+$, JHEP **04** (2014) 060 [1311.3589].
- [173] ATLAS collaboration, Snowmass White Paper Contribution: Physics with the Phase-2 ATLAS and CMS Detectors, .
- [174] CMS collaboration, First evidence for off-shell production of the Higgs boson and measurement of its width, 2202.06923.
- [175] A. Azatov et al., Off-shell Higgs Interpretations Task Force: Models and Effective Field Theories Subgroup Report, 2203.02418.
- [176] T. Appelquist, B.A. Dobrescu and A.R. Hopper, *Nonexotic Neutral Gauge Bosons*, *Phys. Rev. D* **68** (2003) 035012 [hep-ph/0212073].
- [177] V. Gherardi, D. Marzocca and E. Venturini, Matching scalar leptoquarks to the SMEFT at one loop, JHEP 07 (2020) 225 [2003.12525].
- [178] J. Aebischer, W. Dekens, E.E. Jenkins, A.V. Manohar, D. Sengupta and P. Stoffer, Effective field theory interpretation of lepton magnetic and electric dipole moments, JHEP 07 (2021) 107 [2102.08954].

- [179] R.K. Ellis et al., Physics Briefing Book: Input for the European Strategy for Particle Physics Update 2020, 1910.11775.
- [180] T. Bose et al., Report of the Topical Group on Physics Beyond the Standard Model at Energy Frontier for Snowmass 2021, in 2022 Snowmass Summer Study, 2022 [2209.13128].