

# On Some Exact Solutions of Nonlinear Wave Equations

Anatoly BARANNYK † and Ivan YURYK ‡

† *Institute of Mathematics, Pedagogical University,  
22b Arciszewskiego Str., 76-200 Slupsk, Poland  
E-mail: office@bicom.slupsk.pl*

‡ *Ukrainian State University of Food Technologies,  
68 Volodymyrs'ka Str., Kyiv 33, Ukraine*

## Abstract

A new simple method for constructing solutions of multidimensional nonlinear wave equations is proposed

## 1 Introduction

The method of the symmetry reduction of an equation to equations with fewer variables, in particular, to ordinary differential equations [1–3] is among efficient methods for constructing solutions of nonlinear equations of mathematical physics. This method is based on investigation of the subgroup structure of an invariance group of a given differential equation. Solutions being obtained in this way are invariant with respect to a subgroup of the invariance group of the equation. It is worth to note that the invariance imposes very severe constraints on solutions. For this reason, the symmetry reduction doesn't allow to obtain in many cases sufficiently wide classes of solutions.

At last time, the idea of the conditional invariance of differential equations, proposed in [3–6], draws intent attention to itself. By conditional symmetry of an equation, one means the symmetry of some solution set. For a lot of important nonlinear equations of mathematical physics, there exist solution subsets, the symmetry of which is essentially different from that of the whole solution set. One chooses such solution subsets, as a rule, with the help of additional conditions representing partial differential equations. The description of these additional conditions in the explicit form is a difficult problem and unfortunately there are no efficient methods to solve it.

In this paper, we propose a constructive and simple method for constructing some classes of exact solutions to nonlinear equations of mathematical physics. The essence of the method is the following. Let we have a partial differential equation

$$F\left(x, u, u_1, u_2, \dots, u_m\right) = 0, \tag{1}$$

where  $u = u(x)$ ,  $x = (x_0, x_1, \dots, x_n) \in \mathbb{R}_{1,n}$ ,  $u_m$  is a collection of all possible derivatives of order  $m$ , and let equation (1) have a nontrivial symmetry algebra. To construct solutions

of equation (1), we use the symmetry (or conditional symmetry) ansatz [3]. Suppose that it is of the form

$$u = f(x)\varphi(\omega_1, \dots, \omega_k) + g(x), \quad (2)$$

where  $\omega_1 = \omega_1(x_0, x_1, \dots, x_n), \dots, \omega_k = \omega_k(x_0, x_1, \dots, x_n)$  are new independent variables. Ansatz (2) singles out some subset  $S$  from the whole solution set of equation (1). Construct (if it is possible) a new ansatz

$$u = f(x)\varphi(\omega_1, \dots, \omega_k, \omega_{k+1}, \dots, \omega_l) + g(x), \quad (3)$$

being a generalization of ansatz (2). Here  $\omega_{k+1}, \dots, \omega_l$  are new variables that should be determined. We choose the variables  $\omega_{k+1}, \dots, \omega_l$  from the condition that the reduced equation corresponding to ansatz (3) coincides with the reduced equation corresponding to ansatz (2). Ansatz (3) singles out a subset  $S_1$  of solutions to equation (1), being an extension of the subset  $S$ . If solutions of the subset  $S$  are known, then one also can construct solutions of the subset  $S_1$ . These solutions are constructed in the following way. Let  $u = u(x, C_1, \dots, C_t)$  be a multiparameter solution set of the form (2) of equation (1), where  $C_1, \dots, C_t$  are arbitrary constants. We shall obtain a more general solution set of equation (1) if we take constants  $C_i$  in the solution  $u = u(x, C_1, \dots, C_t)$  to be arbitrary smooth functions of  $\omega_{k+1}, \dots, \omega_l$ .

Basic aspects of our approach are presented by the examples of d'Alembert, Liouville and eikonal equations.

## 2 Nonlinear d'Alembert equations

Let us consider a nonlinear Poincaré-invariant d'Alembert equation

$$\square u + F(u) = 0, \quad (4)$$

where

$$\square u = \frac{\partial^2 u}{\partial x_0^2} - \frac{\partial^2 u}{\partial x_1^2} - \dots - \frac{\partial^2 u}{\partial x_n^2},$$

$F(u)$  is an arbitrary smooth function. Papers [3, 7–9] are devoted to the construction of exact solutions to equation (4) for different restrictions on the function  $F(x)$ . Majority of these solutions is invariant with respect to a subgroup of the invariance group of equation (4), i.e., they are Lie solutions. One of the methods for constructing solutions is the method of symmetry reduction of equation (4) to ordinary differential equations. The essence of this method for equation (4) consists in the following.

Equation (4) is invariant under the Poincaré algebra  $AP(1, n)$  with the basis elements

$$\begin{aligned} J_{0a} &= x_0 \partial_a + x_a \partial_0, & J_{ab} &= x_b \partial_a - x_a \partial_b, \\ P_0 &= \partial_0, & P_a &= \partial_a \quad (a, b = 1, 2, \dots, n). \end{aligned}$$

Let  $L$  be an arbitrary rank  $n$  subalgebra of the algebra  $AP(1, n)$ . The subalgebra  $L$  has two main invariants  $u, \omega = \omega(x_0, x_1, \dots, x_n)$ . The ansatz  $u = \varphi(\omega)$  corresponding to the subalgebra  $L$  reduces equation (4) to the ordinary differential equation

$$\ddot{\varphi}(\nabla\omega)^2 + \dot{\varphi}\square\omega + F(\varphi) = 0, \quad (5)$$

where

$$(\nabla\omega)^2 \equiv \left(\frac{\partial\omega}{\partial x_0}\right)^2 - \left(\frac{\partial\omega}{\partial x_1}\right)^2 - \dots - \left(\frac{\partial\omega}{\partial x_n}\right)^2.$$

Such a reduction is called the *symmetry reduction*, and the ansatz is called the *symmetry ansatz*. There exist eight types of nonequivalent rank  $n$  subalgebras of the algebra  $AP(1, n)$  [7]. In Table 1, we write out these subalgebras, their invariants and values of  $(\nabla\omega)^2$ ,  $\square\omega$  for each invariant.

**Table 1.**

N	Algebra	Invariant $\omega$	$(\nabla\omega)^2$	$\square\omega$
1.	$P_1, \dots, P_n$	$x_0$	1	0
2.	$P_0, P_1, \dots, P_{n-1}$	$x_n$	-1	0
3.	$P_1, \dots, P_{n-1}, J_{0n}$	$(x_0^2 - x_n^2)^2$	1	$\frac{1}{\omega}$
4.	$J_{ab}$ ( $a, b = 1, \dots, k$ ), $P_{k+1}, \dots, P_n, P_0$ ( $k \geq 2$ )	$(x_1^2 + \dots + x_k^2)^{1/2}$	-1	$-\frac{k-1}{\omega}$
5.	$G_a = J_{0a} - J_{ak}, J_{ab}$ ( $a, b = 1, \dots, k-1$ ) $J_{0k}, P_{k+1}, \dots, P_n$ ( $k \geq 1$ )	$(x_0^2 - x_1^2 - \dots - x_k^2)^{1/2}$	1	$\frac{k}{\omega}$
6.	$P_1, \dots, P_{n-2}, P_0 + P_n$ $J_{0n} + \alpha P_{n-1}$	$\alpha \ln(x_0 - x_n) + x_{n-1}$	-1	0
7.	$P_0 + P_n, P_1, \dots, P_{n-1}$	$x_0 - x_n$	0	0
8.	$P_a$ ( $a = 1, \dots, n-2$ ), $G_{n-1} + P_0 - P_n, P_0 + P_n$	$(x_0 - x_n)^2 - 4x_{n-1}$	-1	0

The method proposed in [11] of reduction of equation (4) to ODE is a generalization of the symmetry reduction method. Equation (4) is reduced to ODE with the help of the ansatz  $u = \varphi(\omega)$ , where  $\omega = \omega(x)$  is a new variable, if  $\omega(x)$  satisfies the equations

$$\square\omega = F_1(\omega), \quad (\nabla\omega)^2 = F_2(\omega). \quad (6)$$

Here  $F_1, F_2$  are arbitrary smooth functions depending only on  $\omega$ .

Thus, if we construct all solutions to system (6), hence we get the set of all values of the variable  $\omega$ , for which the ansatz  $u = \varphi(\omega)$  reduces equation (4) to ODE in the variable  $\omega$ . Papers [10–11] are devoted to the investigation of system (6).

Note, however, that ansatzes obtained by solving system (6), don't exhaust the set of all ansatzes reducing equation (4) to ordinary differential equations. For this purpose, let us consider the process of finding generalized ansatzes (3) on the known symmetry ansatzes (2) of equation (4).

a) Consider the symmetry ansatz  $u = \varphi(\omega_1)$  for equation (4), where  $\omega_1 = (x_0^2 - x_1^2 - \dots - x_k^2)$ ,  $k \geq 2$ . The ansatz reduces equation (4) to the equation

$$\varphi_{11} + \frac{k}{\omega_1} \varphi_1 + F(\omega_1) = 0, \tag{7}$$

where  $\varphi_{11} = \frac{d^2\varphi}{d\omega_1^2}$ ,  $\varphi_1 = \frac{d\varphi}{d\omega_1}$ . This ansatz should be regarded as a partial case of the more general ansatz  $u = \varphi(\omega_1, \omega_2)$ , where  $\omega_2$  is an unknown variable. The ansatz  $u = \varphi(\omega_1, \omega_2)$  reduces equation (4) to the equation

$$\varphi_{11} + \frac{k}{\omega_1} \varphi_1 + 2\varphi_{12}(\nabla\omega_1 \cdot \nabla\omega_2) + \varphi_2 \square\omega_2 + \varphi_{22}(\nabla\omega_2)^2 + F(\varphi) = 0, \tag{8}$$

where

$$\nabla\omega_1 \cdot \nabla\omega_2 = \frac{\partial\omega_1}{\partial x_0} \cdot \frac{\partial\omega_2}{\partial x_0} - \frac{\partial\omega_1}{\partial x_1} \cdot \frac{\partial\omega_2}{\partial x_1} - \dots - \frac{\partial\omega_1}{\partial x_n} \cdot \frac{\partial\omega_2}{\partial x_n}.$$

Let us impose the condition on equation (8), under which equation (8) coincides with the reduced equation (7). Under such assumption, equation (8) decomposes into two equations

$$\varphi_{11} + \frac{k}{\omega_1} \varphi_1 + F(\varphi) = 0, \tag{9}$$

$$2\varphi_{12}(\nabla\omega_1 \cdot \nabla\omega_2) + \varphi_{22}(\nabla\omega_2)^2 + \varphi_{12} \square\omega_2 = 0. \tag{10}$$

Equation (10) will be fulfilled for an arbitrary function  $\varphi$  if we impose the conditions

$$\square\omega_2 = 0, \quad (\nabla\omega_2)^2 = 0, \tag{11}$$

$$\nabla\omega_1 \cdot \nabla\omega_2 = 0 \tag{12}$$

on the variable  $\omega_2$ . Therefore, if we choose the variable  $\omega_2$  such that conditions (11), (12) are satisfied, then the multidimensional equation (4) is reduced to the ordinary differential equation (7) and solutions of the latter equation give us solutions of equation (4). So, the problem of reduction is reduced to the construction of general or partial solutions to system (11), (12).

The overdetermined system (11) is studied in detail in papers [12–13]. A wide class of solutions to system (11) is constructed in papers [12–13]. These solutions are constructed in the following way. Let us consider a linear algebraic equation in variables  $x_0, x_1, \dots, x_n$  with coefficients depending on the unknown  $\omega_2$ :

$$a_0(\omega_2)x_0 - a_1(\omega_2)x_1 - \dots - a_n(\omega_2)x_n - b(\omega_2) = 0. \tag{13}$$

Let the coefficients of this equation represent analytic functions of  $\omega_2$  satisfying the condition

$$[a_0(\omega_2)]^2 - [a_1(\omega_2)]^2 - \dots - [a_n(\omega_2)]^2 = 0.$$

Suppose that equation (13) is solvable for  $\omega_2$  and let a solution of this equation represent some real or complex function

$$\omega_2(x_0, x_1, \dots, x_n). \tag{14}$$

Then function (14) is a solution to system (11). Single out those solutions (14), that possess the additional property  $\nabla\omega_1 \cdot \nabla\omega_2 = 0$ . It is obvious that

$$\frac{\partial\omega_2}{\partial x_0} = -\frac{a_0}{\delta'}, \quad \frac{\partial\omega_2}{\partial x_1} = \frac{a_1}{\delta'}, \quad \dots, \quad \frac{\partial\omega_2}{\partial x_n} = \frac{a_n}{\delta'},$$

where

$$\delta(\omega_2) \equiv a_0(\omega_2)x_0 - a_1(\omega_2)x_1 - \dots - a_n(\omega_2)x_n - b(\omega_2)$$

and  $\delta'$  is the derivative of  $\delta$  with respect to  $\omega_2$ . Since

$$\frac{\partial\omega_1}{\partial x_0} = \frac{x_0}{\omega_1}, \quad \frac{\partial\omega_1}{\partial x_1} = -\frac{x_1}{\omega_1}, \quad \dots, \quad \frac{\partial\omega_1}{\partial x_n} = -\frac{x_n}{\omega_1},$$

we have

$$\nabla\omega_1 \cdot \nabla\omega_2 = -\frac{1}{\omega_1\delta'}(a_0x_0 - a_1x_1 - \dots - a_nx_n).$$

Hence, with regard for (13), the equality  $\nabla\omega_1 \cdot \nabla\omega_2 = 0$  is fulfilled if and only if  $b(\omega_2) = 0$ . Therefore, we have constructed the wide class of ansatzes reducing the d'Alembert equation to ordinary differential equations. The arbitrariness in choosing the function  $\omega_2$  may be used to satisfy some additional conditions (initial, boundary and so on).

**b)** The symmetry ansatz  $u = \varphi(\omega_1)$ ,  $\omega_1 = (x_1^2 + \dots + x_l^2)^{1/2}$ ,  $1 \leq l < n-1$ , is generalized in the following way. Let  $\omega_2$  be an arbitrary solution to the system of equations

$$\begin{aligned} \frac{\partial^2\omega}{\partial x_0^2} - \frac{\partial^2\omega}{\partial x_{l+1}^2} - \dots - \frac{\partial^2\omega}{\partial x_n^2} &= 0, \\ \left(\frac{\partial\omega}{\partial x_0}\right)^2 - \left(\frac{\partial\omega}{\partial x_{l+1}}\right)^2 - \dots - \left(\frac{\partial\omega}{\partial x_n}\right)^2 &= 0. \end{aligned} \tag{15}$$

The ansatz  $u = \varphi(\omega_1, \omega_2)$  reduces equation (4) to the equation

$$-\frac{d^2\varphi}{d\omega_1^2} - \frac{k-1}{\omega_1} \frac{d\varphi}{d\omega_1} + F(\varphi) = 0.$$

If  $l = n-1$ , then the ansatz  $u = \varphi(\omega_1, \omega_2)$ ,  $\omega_2 = x_0 - x_n$  is a generalization of the symmetry ansatz  $u = \varphi(\omega_1)$ .

Ansatzes corresponding to subalgebras 2, 6 and 8 in Table 1, are particular cases of the ansatz constructed above. Doing in a similar way, one can obtain wide classes of ansatzes reducing equation (4) to two-dimensional, three-dimensional and so on equations. Let us present some of them.

**c)** The ansatz  $u = \varphi(\omega_1, \dots, \omega_l, \omega_{l+1})$ , where  $\omega_1 = x_1, \dots, \omega_l = x_l$ ,  $\omega_{l+1}$  is an arbitrary solution of system (15),  $l \leq n-1$ , is a generalization of the symmetry ansatz  $u = \varphi(\omega_1, \dots, \omega_l)$  and reduces equation (4) to the equation

$$-\frac{\partial^2\varphi}{\partial\omega_1^2} - \frac{\partial^2\varphi}{\partial\omega_2^2} - \dots - \frac{\partial^2\varphi}{\partial\omega_l^2} + F(\varphi) = 0.$$

**d)** The ansatz  $u = \varphi(\omega_1, \dots, \omega_s, \omega_{s+1})$ , where  $\omega_1 = (x_0^2 - x_1^2 - \dots - x_l^2)^{1/2}$ ,  $\omega_2 = x_{l+1}, \dots, \omega_s = x_{l+s-1}$ ,  $l \geq 2$ ,  $l + s - 1 \leq n$ ,  $\omega_{s+1}$  is an arbitrary solution of the system

$$\square\omega_{s+1} = 0, \quad (\nabla\omega_{s+1})^2 = 0, \quad \nabla\omega_i \cdot \nabla\omega_{s+1} = 0, \quad i = 1, 2, \dots, s, \quad (16)$$

is a generalization of the symmetry ansatz  $u = \varphi(\omega_1, \dots, \omega_s)$  and reduces equation (4) to the equation

$$\varphi_{11} - \frac{l}{\omega_1}\varphi_1 - \varphi_{22} - \dots - \varphi_{ss} + F(\varphi) = 0.$$

Let us construct in the way described above some classes of exact solutions of the equation

$$\square u + \lambda u^k = 0, \quad k \neq 1. \quad (17)$$

The following solution of equation (17) is obtained in paper [9]:

$$u^{1-k} = \sigma(k, l)(x_1^2 + \dots + x_l^2), \quad (18)$$

where

$$\sigma(k, l) = \frac{\lambda(1-k)^2}{2(l-lk+2k)}, \quad l = 1, 2, \dots, n.$$

Solution (18) defines a multiparameter solution set

$$u^{1-k} = \sigma(k, l) \left[ (x_1 + C_1)^2 + \dots + (x_l + C_l)^2 \right],$$

where  $C_1, \dots, C_l$  are arbitrary constants. Hence, according to c), we obtain the following set of solutions to equation (17) for  $l \leq n - 1$ :

$$u^{1-k} = \sigma(k, l) \left[ (x_1 + h_1(\omega))^2 + \dots + (x_l + h_l(\omega))^2 \right], \quad k \neq \frac{l}{l-2},$$

where  $\omega$  is an arbitrary solution of system (15) and  $h_1(\omega), \dots, h_l(\omega)$  are arbitrary twice differentiable functions of  $\omega$ . In particular, if  $n = 3$  and  $l = 1$ , then equation (17) possesses in the space  $\mathbb{R}_{1,3}$  the solution set

$$u^{1-k} = \frac{\lambda(1-k)^2}{2(1+k)} [x_1 + h_1(\omega)]^2, \quad k \neq -1.$$

Next, let us consider the following solution of equation (4) [9]:

$$u^{1-k} = \sigma(k, s)(x_0^2 - x_1^2 - \dots - x_s^2), \quad s = 2, \dots, n, \quad (19)$$

where

$$\sigma(k, s) = -\frac{\lambda(1-k)^2}{2(s-ks+k+1)}, \quad k \neq \frac{s+1}{s-1}.$$

Solution (19) defines the multiparameter solution set

$$u^{1-k} = \sigma(k, s) \left[ x_0^2 - x_1^2 - \dots - x_l^2 - (x_{l+1} + C_{l+1})^2 - \dots - (x_s + C_s)^2 \right],$$

where  $C_{l+1}, \dots, C_s$  are arbitrary constants. According to d) we obtain the following solution set for  $l \geq 2$

$$u^{1-k} = \sigma(k, s) \left[ x_0^2 - x_1^2 - \dots - x_l^2 - (x_{l+1} + h_{l+1}(\omega))^2 - \dots - (x_s + h_s(\omega))^2 \right],$$

where  $\omega$  is an arbitrary solution of system (16), and  $h_{l+1}(\omega), \dots, h_s(\omega)$  are arbitrary twice differentiable functions. In particular, if  $l = 2$  and  $s = 3$ , then equation (4) possesses in the space  $\mathbb{R}_{1,3}$  the following solution set

$$u^{1-k} = \frac{\lambda(1-k)^2}{4(k-2)} \left[ x_0^2 - x_1^2 - x_2^2 - (x_3 - h_3(\omega))^2 \right], \quad k \neq 2.$$

The equation

$$\square u + 6u^2 = 0 \tag{20}$$

possesses the solution  $u = \mathcal{P}(x_3 + C_2)$ , where  $\mathcal{P}(x_3 + C_2)$  is an elliptic Weierstrass function with the invariants  $g_2 = 0$  and  $g_3 = C_1$ . Therefore, according to c) we get the following set of solutions of equation (20):

$$u = \mathcal{P}(x_3 + h(\omega)),$$

where  $\omega$  is an arbitrary solution to system (15) and  $h(\omega)$  is an arbitrary twice differentiable function of  $\omega$ .

Next consider the Liouville equation

$$\square u + \lambda \exp u = 0. \tag{21}$$

The symmetry ansatz  $u = \varphi(\omega_1)$ ,  $\omega_1 = x_3$ , reduces equation (21) to the equation

$$\frac{d^2 \varphi}{d\omega_1^2} = \lambda \exp \varphi(\omega_1).$$

Integrating this equation, we obtain that  $\varphi$  coincides with one of the following functions:

$$\ln \left\{ \left( -\frac{C_1}{2\lambda} \sec^2 \left[ \frac{\sqrt{-C_1}}{2} (\omega_1 + C_2) \right] \right) \right\} \quad (C_1 < 0, \lambda > 0, C_2 \in \mathbb{R});$$

$$\ln \left\{ \frac{2C_1 C_2 \exp(\sqrt{C_1} \omega_1)}{\lambda [1 - C_2 \exp(\sqrt{C_1} \omega_1)]^2} \right\} \quad (C_1 > 0, \lambda C_2 > 0);$$

$$-\ln \left( \sqrt{\frac{\lambda}{2}} \omega_1 + C \right)^2.$$

Hence, according to c) we get the following solutions set for equation (21):

$$u = \ln \left\{ \left( -\frac{h_1(\omega)}{2\lambda} \sec^2 \left[ \frac{\sqrt{-h_1(\omega)}}{2} (\omega_1 + h_2(\omega)) \right] \right) \right\} \quad (h_1(\omega) < 0, \lambda > 0);$$

$$u = \ln \left\{ \frac{2h_1(\omega)h_2(\omega) \exp(\sqrt{h_1(\omega)} \omega_1)}{\lambda [1 - h_2(\omega) \exp(\sqrt{h_1(\omega)} \omega_1)]^2} \right\} \quad (h_1(\omega) > 0, \lambda h_2(\omega) > 0);$$

$$u = -\ln \left( \sqrt{\frac{\lambda}{2}} \omega_1 + h(\omega) \right)^2,$$

where  $h_1(\omega)$ ,  $h_2(\omega)$ ,  $h(\omega)$  are arbitrary twice differentiable functions;  $\omega$  is an arbitrary solution to system (15).

Using, for example, the solution to the Liouville equation (21) [9]

$$u = \ln \frac{2(s-2)}{\lambda[x_0^2 - x_1^2 - \dots - x_s^2]}, \quad s \neq 2,$$

we obtain the wide class of solutions to the Liouville equation

$$u = \ln \frac{2(s-2)}{\lambda[x_0^2 - x_1^2 - \dots - x_l^2 - (x_{l+1} + h_{l+1}(\omega))^2 - \dots - (x_s + h_s(\omega))^2]},$$

where  $\omega$  is an arbitrary solution to system (16), and  $h_{l+1}(\omega), \dots, h_s(\omega)$  are arbitrary twice differentiable functions. If  $s = 3$ , then equation (21) possesses in the space  $\mathbb{R}_{1,3}$  the following solution set

$$u = \ln \frac{2}{\lambda[x_0^2 - x_1^2 - x_2^2 - (x_3 + h_3(\omega))^2]}.$$

Let us consider now the sine-Gordon equation

$$\square u + \sin u = 0.$$

Doing in an analogous way, we get the following solutions:

$$u = 4 \arctan h_1(\omega) \exp(\varepsilon_0 x_3) - \frac{1}{2}(1 - \varepsilon)\pi, \quad \varepsilon_0 = \pm 1, \quad \varepsilon = \pm 1;$$

$$u = 2 \arccos[\operatorname{dn}(x_3 + h_1(\omega)), m] + \frac{1}{2}(1 + \varepsilon)\pi, \quad 0 < m < 1;$$

$$u = 2 \arccos \left[ \operatorname{cn} \left( \frac{x_3 + h_1(\omega)}{m} \right), m \right] + \frac{1}{2}(1 + \varepsilon)\pi, \quad 0 < m < 1,$$

where  $h_1(\omega)$  is an arbitrary twice differentiable function,  $\omega$  is an arbitrary solution to system (15).

### 3 Eikonal equation

Consider the eikonal equation

$$\left( \frac{\partial u}{\partial x_0} \right)^2 - \left( \frac{\partial u}{\partial x_1} \right)^2 - \left( \frac{\partial u}{\partial x_2} \right)^2 - \left( \frac{\partial u}{\partial x_3} \right)^2 = 1. \tag{22}$$

The symmetry ansatz  $u = \varphi(\omega_1)$ ,  $\omega_1 = x_0^2 - x_1^2 - x_2^2 - x_3^2$ , reduces equation (22) to the equation

$$4\omega_1 \left( \frac{\partial \varphi}{\partial \omega_1} \right)^2 - 1 = 0. \tag{23}$$



We shall look for a generalized ansatz in the form  $u = \varphi(\omega_1, \omega_2)$ . This ansatz reduces equation (22) to the equation

$$4\omega_1 \left( \frac{\partial \varphi}{\partial \omega_1} \right)^2 + 2(\nabla \omega_1 \cdot \nabla \omega_2) \frac{\partial \varphi}{\partial \omega_1} + (\nabla \omega_2)^2 \left( \frac{\partial \varphi}{\partial \omega_2} \right)^2 = 1. \quad (24)$$

Impose the condition on equation (24), under which equation (24) coincides with equation (23). It is obvious that this condition will be fulfilled if we impose the conditions

$$(\nabla \omega_2)^2 = 0, \quad \nabla \omega_1 \cdot \nabla \omega_2 = 0 \quad (25)$$

on the variable  $\omega_2$ . Having solved system (25), we get the explicit form of the variable  $\omega_2$ . It is easy to see that an arbitrary function of a solution to system (25) is also a solution to this system.

Having integrated equation (23), we obtain  $(u + C)^2 = x_0^2 - x_1^2 - x_2^2 - x_3^2$ , where  $C$  is an arbitrary constant. We shall obtain a more general solution set for the eikonal equation if we take  $C$  to be an arbitrary solution to system (25).

The symmetry ansatz  $u = \varphi(\omega_1, \omega_2)$ ,  $\omega_1 = x_0^2 - x_1^2 - x_2^2$ ,  $\omega_2 = x_3$  can be generalized in the following way. Let  $\omega_3$  be an arbitrary solution to the system of equations

$$\begin{aligned} \left( \frac{\partial \omega_3}{\partial x_0} \right)^2 - \left( \frac{\partial \omega_3}{\partial x_1} \right)^2 - \left( \frac{\partial \omega_3}{\partial x_2} \right)^2 &= 0, \\ x_0 \frac{\partial \omega_3}{\partial x_0} + x_1 \frac{\partial \omega_3}{\partial x_1} + x_3 \frac{\partial \omega_3}{\partial x_2} &= 0. \end{aligned} \quad (26)$$

Then the ansatz  $u = \varphi(\omega_1, \omega_2, \omega_3)$  reduces the eikonal equation to the equation

$$4\omega_1 \left( \frac{\partial \varphi}{\partial \omega_1} \right)^2 - \left( \frac{\partial \varphi}{\partial \omega_2} \right)^2 - 1 = 0. \quad (27)$$

Equation (27) possesses the solution [9]

$$\varphi = \frac{C_1^2 + 1}{2C_1} (x_0^2 - x_1^2 - x_2^2)^{1/2} + \frac{C_1^2 - 1}{2C_1} x_3 + C_2,$$

$$(\varphi + C_2)^2 = x_0^2 - x_1^2 - x_2^2 - (x_3 + C_1)^2,$$

that can be easily found by using the symmetry reduction method of equation (27) to ordinary differential equation. Having replaced arbitrary constants  $C_1$  and  $C_2$  by arbitrary functions  $h_1(\omega)$  and  $h_2(\omega)$ , we get the more wide classes of exact solutions to the eikonal equation:

$$u = \frac{h_1(\omega_3)^2 + 1}{2h_1(\omega_3)} (x_0^2 - x_1^2 - x_2^2)^{1/2} + \frac{h_1(\omega_3)^2 - 1}{2h_1(\omega_3)} x_3 + h_2(\omega_3),$$

$$(u + h_2(\omega_3))^2 = x_0^2 - x_1^2 - x_2^2 - (x_3 + h_1(\omega_3))^2.$$

Let us note, since the Born–Infeld equation is a differential consequence of the eikonal equation [3], hence we also constructed wide classes of exact solutions of the Born–Infeld equation.

## References

- [1] Ovsyannikov L.V., Group Analysis of Differential Equations, Academic Press, 1982.
- [2] Olver P., Applications of Lie Groups to Differential Equations, Springer, New York, 1986.
- [3] Fushchych W.I., Shtelen V.M. and Serov N.I., Symmetry Analysis and Exact Solutions of Equations of Nonlinear Mathematical Physics, Kluwer, Dordrecht, 1993.
- [4] Fushchych W.I., Nikitin A.G., Symmetries of Maxwell's Equations, Dordrecht, Reidel, 1987.
- [5] Fushchych W.I. and Tsyfra I.M., On a reduction and solutions of nonlinear wave equations with broken symmetry, *J. Phys. A: Math. Gen.*, 1987, V.20, N 2, L45–L48.
- [6] Levi D. and Winternitz P., *J. Phys. A*, 1989. V.22, 2915–2924.
- [7] Grundland A.M., Harnad J. and Winternitz P., Symmetry reduction for nonlinear relativistically invariant equations, *J. Math. Phys.*, 1984, V.25, N 4, 791–806.
- [8] Cieciora G. and Grundland A., A certain class of solutions of the nonlinear wave equations, *J. Math. Phys.*, 1984, V.25, N 12, 3460–3469.
- [9] Fushchych W.I., Barannyk L.F. and Barannyk A.F., Subgroup Analysis of Galilei and Poincaré Groups, and Reduction of Nonlinear Equations, Naukova Dumka, Kyiv, 1991 (in Russian).
- [10] Collins C.B., Complex potential equations. I. A technique for solutions, *Proc. Cambridge Phil. Soc.*, 1976, N 9, 225–236.
- [11] Fushchych W.I. and Zhdanov R.Z., Nonlinear Spinor Equations: Symmetry and Exact Solutions, Kyiv, Naukova Dumka, 1992 (in Russian).
- [12] Smirnov V.I. and Sobolev S.L., New method for solving a plane problem of elastic oscillations, *Proc. of Seismological Institute of Acad. Sci. USSR*, 1932, V.20, 37–42.
- [13] Smirnov V.I. and Sobolev S.L., On application of a new method to the study of elastic oscillations in a space with the axial symmetry, *Proc. of Seismological Institute of Acad. Sci. USSR*, 1933, V.29, 43–51.