Renormalization Scheme References

References

[1] H.J. Lu and S.J. Brodsky, Phys. Rev. D**48** 3310 (1993).
Relating Physical Observables in QCD without Scale-Scheme ambiguities

[2] S.J. Brodsky, G.T. Gabadadze, A.L. Kataev, H.J. Lu, SLAC-PUB-95-7030, Dec. 1995, HEP-PH/9512367.
The generalized Crewther relation in QCD and its experimental consequences.

[3] G. Grunberg and A.L. Kataev, Phys. Lett. **B279** 352 (1992).
On some possible extensions of the Brodsky-Lepage-Mackenzie approach beyond next to-leading order.

[4] S. J. Brodsky and H. J. Lu, Preprint SLAC-PUB-6369, November 1993.
Commensurate Scale Relations: Relating Observables in QCD without Renormalization Scale or Scheme Ambiguity.

[5] S. J. Brodsky and H. J. Lu, Preprint SLAC-PUB-6481, April 1994, Phys. Rev. **D51,** 3652, 1995.
Commensurate Scale Relations in Quantum Chromodynamics.

[6] S. J. Brodky and H. J. Lu, Preprint SLAC-PUB-6683, October 1994.
Commensurate Scale Relations: precise test of Quantum Chromodynamics without Scale or scheme ambiguity.

[7] P.A. Raczka and A. Szymacha, Preprint IFT-13-94, September 1994, Warsaw, HEP-PH-9412236.
Improved analysis of the renormalization scheme ambiguities in the QCD corrections to the semileptonic decay of the tau lepton.

[8] P. A. Raczka. Z. Phys. **C65** 481, 1995.
Renormalization scheme dependence and the problem of the uncertainties next-to-leading order QCD corrections.

[9] R. Ball *et al.*, Phys. Lett. **B347** 80, 1995.
Scheme independence and the exact renormalization group.

[10] Wing Kai Wong, SLAC-PUB-95-7063, Dec. 1995, HEP-PH/9601215.
Renormalization scale setting for evolution equation of nonsinglet structure functions and their moments.

[11] D. de Florian *et al.,* Phys. Rev. **D53,** 73, 1996.
Scale dependence of polarized DIS asymmetries.

[12] K. Jansen, *et al.,* Phys. Lett. **B372,** 275, 1996, HEP-LAT/9512009.
Non pertubative renormalization of lattice QCD at all scales.

[13] S.J. Brodsky, Hung Jung Lu, SLAC-PUB-96-7098, Aug. 1995. HEP-PH/9601301.
Optimal renormalization scales and commensurate scale relations.

[14] G. Grunberg, Phys. Lett. **B372,** 121, 1996.
Renormalons at fixed points.

[15] F. T. Hawes, A. G. Williams, C. D. Roberts, HEP-PH/9604402, April 1996.
Renormalization and chiral symmetry breaking in quenched QED in arbitrary covariant gauge.

[16] J. Ellis, Phys. Rev. **D54,** 6986, 1996, HEP-PH/9607404.
Renormalization scheme dependence of PADE summation in QCD.

[17] P. A. Raczka, ICHEP96, Warsaw ,Poland, 25-31 July 1996, HEP-PH/9611257.
Analysis of the renormalization scheme ambiguities in the QCD corrections to the hadronic decays of the tau lepton.

[18] R. S. Thorne, RAL-TR-96-088, Phys. Lett. **B392,** 463, 1997, HEP-PH/9610334.
Renormalization scheme consistent, including leading LN(1/X) terms structure function.

[19] S. Catani, *et al.,* Phys. Rev. **D55,** 6819, 1997, HEP-PH/9610553, Oct.1996.
Regularization scheme independence and unitarity in QCD cross-sections.

[20] F. Litim, Phys. Lett. **B393,** 103, 1997, HEP-TH/9609040, Sept. 1996.
Scheme independence at first order phase transitions and the renormalization group.

[21] S. J. Brodsky , Chueng-Ryong Ji, A. Pang, D.G, Robertson, SLAC-PUB-7473, HEP-PH/9705221, May 1997. Phys. Rev. **D57,** 245, 1998.
Optimal renormalization and scheme for exclusive processes.

[22] S. Brodsky *et al.,* SLAC-PUB-7566, HEP-PH/9706467, June, 1997. Phys. Rev. **D56,** 6980, 1997.
Pade approximants, optimal renormalization scales , and momentum flow in Feynman diagrams.

[23] J. Ellis, *et al.,* CERN-TH-97-267, HEP-PH/9710302, Oct. 199. Phys. Rev. **D57,** 2665, 1998.
Asymptotic pade predictions: up to five loops in QCD and SQCD.

[24] R. Sommer, 36 Int. Universitat Wochen fuer kernphysik and teilchenphysik, Schladming, Austria, 1-8 Mar. 1997. HEP-PH/9711243.
Nonpertubative remormalization of QCD.

[25] S. Capitani, *et al.,* HEP-PH/9711007, Nov. 1997.
Local bilinear operators on the lattice and their pertubative renormalization including O(A) effects.

[26] F. del Aguila, M. Perez-Victoria, HEP-PH/9710442, USTRON 97, Ustron, Poland 19-24 Sept. 1997.
Constrained differential renormalization.

[27] R. S. Thorne, HEP-PH/9710541, Jul. 1997, Nucl. Phys. **B512,** 323, 1998.
A complete leading order, renormalization scheme consistent calculation of structure functions.

[28] S. Brodsky, *et al.,* SLAC-PUB-7157, HEP-PH/9801330, Jan. 1998.
An Analytic extension of the MS-BAR renormalization scheme.

[29] C.R. Stephens, *et al.,* Phys. Rev. **B414,** 333, 1997,HEP-PH/9706084.
Regge behavior from an environmentally friendly renormalization group.

[30] P.A. Raczka, Phys. Rev. **D57,** 6862, 1998, HEP-PH/9707366.
Renormalization scheme dependence and the problem of determination of α_s and the condensates from the semi-leptonic τ decays.

[31] E. Gardi, Acta Phys. Polon. **B28,** 3067, 1997, HEP-PH/9709272.
Resummation of pertubative QCD by Pade Approximants.

[32] Y. Koike, *et al.,* HEP-PH/9805460, Phys. Lett. **B437,** 153, 1998.
Renormalization of chiral-even twist-3 light cone wave functions for vector mesons in QCD.

[33] S. J. Brodsky, HEP-PH/9806445, SLAC-PUB-7861, Fun. 1998.
Commensurate scale relations and the abelian correspondance.

[34] G. Grunberg, HEP-PH/9807494, Jul. 1998. JEP 9811:006, 1998.
On power corrections in the dispersive approach.

[35] M. Gockeler, *et al.,* HEP-LAT/9809001, Sept. 1998.
Composite operators in lattice QCD: nonpertubative renormalization.

[36] M. Crisafulli *et al.,* Eur. Phys. J. **C4,** 75, 1998, HEP-LAT/9707025. CERN-TH-97-015.

Improved renormalization of lattice operators: a critical reappraisal.

[37] G. Cvetic, HEP-PH/9808273, Jul. 1998, submitted to Nucl. Phys. Proc. Suppl.
Global approximants with renormalization invariance in PQCD.

[38] E. Leader, *et al.,* HEP-PH/9808248, Aug. 1998, Phys. Lett. **B445,** 232, 1998.
Scheme dependence in polarized deep inelastic scattering.

[39] G.M. de Divitiis, *et al.,* Phys. Lett. **B419,** 311, 1998. HEP-LAT/9710071.
Nonpertubative renormalization constants on the lattice from flavor nonsinglet ward identities.

[40] A.D. Martin, *et al.,* HEP-PH/9808371. Aug. 1998.
Scheme dependence, leading order and higher twist studies of MRST.

[41] D.V. Shirkov, Int. J. Mod. Phys. B12, 1247, 1998.
The role of renormalization group in fundamental theoretical physics.

[42] Y. Taniguchi, *et al.,* HEP-LAT/9809070, Jul. 1998.
One loop renormalization factors and mixing coefficients of bilinear quark operators for improved gluon and quark actions.

[43] S. Brodsky and J. Rathsman, SLAC-PUB-8176, June 1999. HEP-PH/9906339
Conformal symmetry as a template : commensurate scale relations and physical renormalization schemes.

[44] S. Brodsky, M. Melles, J. Rathsman, SLAC-PUB-8019, HEP-PH/9906324, Jun. 1999.
The two loop scale dependence of the static QCD potential including quark masses.

[45] S. Brodsky , SLAC-PUB-8240, HEP-PH/9909234.
QCD technology: light cone quantization and commensurate scale relation.

[46] G.P. Salam, G. Zanderighi, HEP-PH/9909324, Jul. 1999.
Power corrections and the interplay between pertubative and non pertubative phenomena.

[47] M. Binger *et al.,* HEP-PH/9912241, Dec. 1999.
Renormalization-scale-invariant PQCD predictions for $R_(e+e-)$ and the Bjorkensum rule at next-to-leading order.

[48] G. Cvetic, HEP-PH/0003123, Mar. 2000.
Achieving renormalization-scale and scheme-independence in Pade-related resummation in QCD.