

COMMENT OF THE 1S_0 NUCLEON-NUCLEON EFFECTIVE
RANGE EXPANSION PARAMETERS*

H. Pierre Noyes

Stanford Linear Accelerator Center
Stanford University, Stanford, California 94305

ABSTRACT

The most accurate nucleon-nucleon scattering experiments below 10 MeV are now consistent with the following values and uncertainties for the 1S_0 scattering lengths and effective ranges:

$$\begin{array}{ll} a_{pp}^c = -7.823 \pm 0.01 \text{ FM} & r_{pp}^c = 2.794 \pm 0.015 \text{ FM} \\ a_{nn} = -17 \pm 1 \text{ FM} & r_{nn} = 2.84 \pm 0.03 \text{ FM} \\ a_{np} = -23.715 \pm 0.015 \text{ FM} & r_{np} = 2.73 \pm 0.03 \text{ FM} \end{array}$$

Unless strong theoretical arguments can be advanced that all charge-dependent corrections to the parameters can be calculated accurately enough to prove that the values of a_{nn} , r_{nn} , or r_{np} must be moved outside these limits, or strong experimental reasons given to believe that experiments leading to these results were significantly in error, it is argued that new calculations or experiments aimed at changing these values are likely to fail. This argument is supported by an analysis of the latest p-p experiments of Jarmie, Jett, Detch and Hutson at 9.918 MeV, which reconfirm the one-pion-exchange shape effect and give $G_{\pi^0 p}^2 = 15.3 \pm 2.4$ for the pion-nucleon coupling constant.

*Work performed under the auspices of the U. S. Atomic Energy Commission.

The corrections to the shape-independent effective range approximation for nucleon-nucleon scattering in the 1S_0 state due to one-pion exchange (OPE) can be unambiguously predicted.^{1,2,3} This prediction was confirmed using p-p scattering data below about 3 MeV in 1964.⁴ If this shape effect is accepted, it removes a large systematic ambiguity in the projection of the effective range plot to zero energy to determine a_{pp} , and the data then allow the very accurate determination of two parameters referring to the short-range nucleon-nucleon interaction in any model which has OPE as the longest-range component. The simplest interpretation of the charge-independence hypothesis is that this same nuclear model should predict both n-n and n-p scattering in the 1S_0 state, once the e^2/r Coulomb interaction is removed. This prediction fails, since it gives about -17 FM for the nuclear scattering length, while it has been known since the early 1950's that the n-p scattering length is close to -23.7 FM. Electromagnetic corrections corresponding to the extended charge and magnetic moment distributions of the nucleons measured by electron scattering change this prediction by less than a fermi. In contrast to the highly sensitive parameters a_{nn} and a_{np} , the effective ranges are insensitive to small corrections; the p-p data below 3 MeV plus the simplest version of the charge-independence hypothesis requires⁵ both r_{nn} and r_{np} to be about 2.84 FM. If the $\pi^\pm - \pi^0$ mass difference is included in OPE, and some parameter (e.g., a charge-dependent splitting of the pion-nucleon coupling constants or a phenomenological parameter of the short-range interaction) is adjusted to fit the observed value of a_{np} , then the prediction for r_{np} falls to about 2.73 FM with an estimated uncertainty⁵ of only about 0.03 FM.

The prediction of $r_{np} = 2.73 \pm 0.03$ FM used to be in conflict with n-p total cross sections at 0.4926 and 3.205 MeV measured by Engelke, Benenson, Melkonian, and Lebowitz,⁶ assuming all other experiments in the analysis correct. The other data⁷ used determine a_{np}^t , ϵ_d , and a_{np}^s , and combined with EBML gave⁵

$r_{np} = 2.44 \pm 0.11$ FM. Remeasurement of the total neutron-hydrogen cross section for epithermal neutrons did not change this situation according to Houk and Wilson.⁸ However, the discovery of an error in the evaluation of their experiment,⁹ and a new measurement of the coherent neutron-hydrogen scattering length by Koester,¹⁰ change a_{np}^t and a_{np}^s sufficiently to raise the values of r_{np} calculated¹¹ from the Columbia experiments to 2.646 ± 0.072 FM. The situation has been still further improved by Davis and Barschall,¹² who have shown that the energy scale to which many of the neutron measurements have been referred is in error. Their revision does not change the (new) value of 2.66 ± 0.09 FM obtained from the Columbia experiment at 0.4926 MeV, but their revision of the energy of 3.205 MeV down to 3.186 MeV raises the value of r_{np} calculated from that experiment to 2.77 ± 0.14 FM. The case for an r_{np} close to 2.73 FM is still further strengthened by the preliminary results of a new n-p total cross section measurement at 0.525 MeV by Simmons, Cramer, and Cranberg.¹³ Thus there currently remains no significant discrepancy between this prediction from charge-independence and experiment. Any quantitative theoretical explanation for the discrepancy between the predicted value of -17 FM and the observed value of -23.7 FM for a_{np} remains as elusive as ever.

Much less is known quantitatively about the n-n parameters. While some of the relevant experiments would probably appear anomalous if the n-n effective range were less than 2 or more than 4 FM, none of them, even potentially, are within an order of magnitude of the accuracy needed to check the prediction for r_{nn} . Several attempts to measure a_{nn} have been pushed to the level of statistical uncertainty of one or two fermis, and while some measurements are consistent with -17 ± 1 FM, some values fall more than a standard deviation away. However, in the opinion of this author, in no case is the three-particle theory used for the evaluation of the final state sufficiently under control to make any of these discrepancies

troublesome. Until this theoretical situation is improved, it is probably best to consider these experiments as tests of certain approximations in final state interaction theories using $a_{nn} = -17$ FM as a calibration, rather than as "measurements" of a_{nn} .

Until recently, the situation with regard to p-p scattering has been more confusing. It proved possible to validate¹⁴ the assumptions needed about the P-waves in the previous analysis⁴ of the data below 3 MeV by combining the value of A_{yy}/A_{xx} at 11.4 MeV measured by Catillon, Chapellier, and Garreta¹⁵ with the accurate differential cross section measurement obtained by Johnston and Young¹⁶ at 9.69 MeV; even the use of this spin-dependent information leaves the analysis ambiguous if L · S effects are not assumed to be small. There is strong theoretical reason to believe that the L · S interaction is of such short range that this must be so, but direct proof by spin-dependent experiments below 10 MeV appears hopeless with current techniques.¹⁴ Slobodrian^{17, 18} distrusted the large amount of theoretical input needed to extract a_{pp} and r_{pp} from the data below 3 MeV, and also had reason to suspect that the method used to separate elastic scattering events from background in the 9.69 MeV experiment might be a source of systematic error.¹⁸ His group therefore undertook accurate differential cross section measurements at 6.141, 8.097, and 9.918 MeV. Unfortunately, these experiments¹⁹ failed to yield convincing results, as was demonstrated by MacGregor, Arndt, and Wright,²⁰ this author,²¹ and on somewhat different grounds by Sher, Signell, and Heller.²² Comparison with the corrected²³ predictions from the 1964 analysis shown in Fig. 1 makes it clear that either these experiments, or the entire theory of the OPE shape correction, is wrong. This figure also makes clear the fact that the main problem with p-p experiments is systematic rather than statistical error, and that large variations in the p-p parameters can be achieved by injudicious data selection and analysis.

Experiments at 9.69 MeV and 9.918 MeV undertaken at Los Alamos by Jarmie, Brown, Hutson, and Detch²⁴ with the specific objective of providing an alternative to the Berkeley data produced the theoretically expected shape for the differential cross section as a function of angle, and hence the expected value for the central-force P-wave parameter Δ_c , but again failed to obtain a believable value for the absolute value of the cross section, and hence for δ_0^E : This situation was pointed out at the time of publication in an accompanying letter by Holdeman, Signell, and Sher.²⁵ This new discrepancy has finally been resolved by the discovery²⁶ that there had been an error in the measurement of one of the slit widths. Reevaluation²⁶ of the experiments at 9.69 and 9.918 MeV and a new experiment²⁶ at 13.6 MeV by Jarmie, Jett, Detch, and Huston give results in accord with theory, as can be seen from Fig. 1. There is now every reason to believe that the value of $55.23 \pm 0.13^\circ$ for δ_0^E obtained by Signell and Holdeman²⁷ from the 9.918 MeV experiment plus five previously published values from experiments at 3 MeV and below are the most accurate 1S_0 phase shifts available below 10 MeV; these values are collected in Table I. A preliminary analysis by the author²⁸ of the newly evaluated data from Los Alamos at 9.918 MeV gives a value of δ_0^E agreeing with that quoted to 0.01° , but with a somewhat larger error. Further checks will be made, but it appears unlikely that these values could be changed by as much as a standard deviation without invoking some bizzare assumptions.

The phase shifts in Table I still contain the long range vacuum polarization effect. Model independent values of a_{pp}^E and r_{pp}^E can be extracted from them by using the modified effective range expansion derived by Heller,²⁹ but we believe it more instructive to apply the Foldy³⁰ correction in order to obtain the phase shifts which would be produced by the nuclear and nonrelativistic electromagnetic interactions in the absence of vacuum polarization. These corrections have been

computed for three models by H. M. Lipinski and are given in Table I. Since the effective range expansion about $k^2 = 0$ diverges beyond 9.71 MeV due to the OPE branch cut, it would be a serious error to represent the shape correction phenomenologically by the terms $-Pr^3k^4 + Qr^5k^6$. Instead we use the Cini-Fubini-Stanghellini² approximation for the OPE cut, with Wong-Noyes Coulomb correction,³¹ which extends the radius of convergence to 38.8 MeV; explicitly:

$$\mathcal{E}^2 k \operatorname{ctn} \delta_0^c + 2k\eta h(\eta) = -1/a_{pp}^c + \frac{1}{2} r_{pp}^c k^2 - \frac{pk^4}{1+qk^2}$$

$$\eta = \frac{\alpha (m_p c^2 + T)}{[\Gamma(2m_p c^2 + T)]^{1/2}} \quad k^2 = \frac{m_p c^2 T}{\hbar^2 c^2} \quad \mathcal{E}^2 = \frac{2\pi n}{e^{2\pi n} - 1}$$

$$h(\eta) = \sum_{s=1}^{\infty} \frac{\eta^2}{s(s+\eta)^2} - \gamma - \ln \eta$$

$$\hbar c = 197.327891 \text{ MeV FM} \quad \alpha = 1/137.03602$$

$$m_p c^2 = 938.2592 \text{ MeV} \quad T = \text{Laboratory energy in MeV} \quad (1)$$

The parameters p and q are not phenomenological constants, but are computed from the pion-nucleon coupling constant and pion mass, with Coulomb corrections,³¹ according to the formulae

$$q = \frac{2 - \Gamma \left(\sqrt{2} \left(2f - \frac{1}{2} g \right) - 4/\mu a_{pp}^c + \frac{1}{2} \mu r_{pp}^c \right)}{\mu^2 \left(1 - \Gamma \left(\frac{1}{2} \sqrt{2} \left(f - \frac{1}{2} g \right) - 1/\mu a_{pp}^c \right) \right)}$$

$$p = \left(1 - \frac{1}{2} q\mu^2 \right) \left(2\sqrt{2} f - 4/\mu a_{pp}^c - \mu r_{pp}^c \right)$$

$$\mu = \frac{m_{\pi^0} c^2}{\hbar c} \quad m_{\pi^0} c^2 = 134.975 \text{ MeV} \quad y = \frac{\alpha m_p}{m_{\pi^0} \sqrt{2}}$$

$$f = 2y \left(\frac{\pi}{2} \operatorname{ctn} \pi y + \gamma + y^2 \sum_s \frac{1}{s(s^2 - y^2)} + \ln y \right)$$

$$g = \frac{m_{\pi^0}}{m_p} \alpha \left[\frac{\pi^2}{\sqrt{2}} \operatorname{csc}^2 \pi y - \sqrt{2} - 2\sqrt{2} \sum_s \frac{1}{(s^2 - y^2)^2} \right]$$

$$\Gamma = \frac{m_{\pi^0}}{4m_p} G^2 (\sqrt{2} - 1)^{-2y} \quad (2)$$

A more refined treatment using solutions of the Schroedinger equation for specific nuclear models including the long range OPE interaction such as that carried out by Sher, Signell, and Heller²² might appear desirable, but we doubt³² that it would produce values of a_{pp}^c and r_{pp}^c in meaningful disagreement with the values quoted in the abstract. The result of this analysis using either the hard core Hamada-Johnston potential or the stiff core Yukawa potential with the ω -meson mass for the Foldy correction (a or b in Table I) is given in Table II. If instead we took the extreme view that the nuclear interaction is purely attractive at low energy and that the repulsion seen in the 1S_0 state above 250 MeV is due to a velocity-dependent effect (Foldy correction c in Table I), the scattering length decreases by only .0024 FM and the effective range increases by only .0006 FM; we believe this represents an upper limit for the model dependence of the Foldy correction.

If we assume that the pion-nucleon coupling constant is determined by other experiments, the results of this analysis can be summarized by

$$\begin{aligned} a_{pp}^c &= -7.8243 \pm 0.0054 - 0.0014 (G^2 - 14.0) \pm 0.0024 \text{ FM} \\ r_{pp}^c &= 2.7958 \pm 0.0080 + 0.0056 (G^2 - 14.0) \pm 0.0006 \text{ FM} \end{aligned} \quad (3)$$

where the first error is statistical and the last is the upper limit for the model dependence due to the Foldy correction. If we use the same OPE shape correction in the Heller expansion,²⁹ we find $a_{pp}^E = -7.8146$, $r_{pp}^E = 2.7950$. It is hard to think of applications of these numbers, outside of the prediction of values of δ_0^E for direct comparison with experimental data, for which this much precision in the nuclear parameters a and r is of any use. If, as originally proposed by Cini, Fubini, and Stanghellini,² we use these experiments to measure the pion-nucleon coupling constant, Table II yields the value $G_{\pi^0 p}^2 = 15.29 \pm 2.38$. A decade of work has finally allowed the determination of this constant from p-p scattering data using the OPE singularity in the k^2 complex plane (in S waves) to an accuracy comparable

with that obtained earlier from the corresponding singularity in $\cos \theta$ (in high partial waves).

The values of a_{pp}^c and r_{pp}^c obtained still agree with the values previously obtained from the data below 3 MeV²³ (-7.8275 ± 0.0049 , 2.7937 ± 0.0065) and even with the preliminary values obtained in 1964⁴ ($-7.8259 \pm .0048$, 2.786 ± 0.014). Therefore the conclusions reached in 1965⁵ as to the values of the n-n and n-p parameters required by charge independence still hold. The effects of the charge and current distributions can be included by fitting a nuclear model to the values of a and r derived from the data,³³ which will change the interaction parameters in the nuclear model compared to those which would be obtained under the assumption of point charges. Schneider and Thayer³⁴ found that these effects change the prediction for a_{nn} by -0.02 FM; using charge and current distribution derived from more recent electron scattering results, including a portion of the electromagnetic interaction omitted by ST, and a different nuclear model, SSH²² find the prediction for a_{nn} changed by $+0.31$ FM compared to that for point charges. The effect on r_{nn} is only a few percent of the statistical uncertainty in r_{pp} , so can be ignored. Clearly the 6.7 FM discrepancy between the predicted and observed values of a_{np} cannot be explained in this way. These calculations assume that the charge-current distribution follows the matter distribution given by the nonrelativistic wave function, whereas we know^{5, 35} that in the case of the reaction $n + p \rightarrow \gamma + d$ the meson exchange currents which are ignored in such a calculation contribute 10% of the observed cross section at threshold. The calculation becomes even more ambiguous if the strong interaction itself is highly nonlocal, which may very well be the case.³⁶ Thus, while we believe that it is still worth while to show that these effects are indeed small for conventional models, the quantitative significance of such calculations is highly uncertain. A much more detailed discussion of these problems which reaches much the same conclusion has been given by Breit et al.³⁷

The obvious conclusion to be drawn from this survey of a decade of work on the determination of the effective range expansion parameters is that further calculations or experiments are unlikely to increase our knowledge of them without some dramatic change in both the theoretical and the experimental situation. We have omitted the new 9.69 MeV data because it does not cover a wide enough angular range to be useful for a single-energy analysis, and the 13.6 MeV data because it might be beyond the range of quantitative reliability of the CFS formula; a repeat of the detailed analysis of all data below 30 MeV like that given by SSH²² might be useful, but we doubt it will change the final results for a and r outside our errors. A purist might still like to see a highly accurate measurement of the absolute value of $\sigma_{pp}(90^\circ)$ as a function of energy between 3 and 10 MeV in order to exhibit the shape effect in more detail. This approach would avoid the necessity of measuring Δ_C (which does not contribute at 90°), but still would necessarily require that the tensor force parameter for the P waves Δ_T be taken from theory; the prospect that any spin-dependent experiment could determine this tensor parameter to anything like the accuracy to which it is already known theoretically is extremely dim.¹⁴ Such a 90° cross section experiment might well produce marginal discrepancies with Eq. 2, which in turn would give marginally useful restrictions on the shape of the two-nucleon interaction, but until the theory of strong interactions is under more control, this hardly seems worth the effort.

It remains to ask whether new p-p experiments below 10 MeV might give useful information not included in the 1S_0 parameters we have been discussing up to now. Existing experiments do determine very precise values for the P-wave central force combination of phase shifts Δ_C thanks to interference with the triplet Coulomb amplitude. SSH have already shown that this information can be put together with the experiments around 25 MeV to determine 3P wave scattering lengths and

effective ranges, but the uncertain experimental situation made these six parameters subject to considerable systematic uncertainty. This could now presumably be reduced, and could be reduced still further by constructing the analog of the Cini-Fubini-Stanghellini formula for the P-waves. Since putting $1/a$ equal to zero in the usual N/D approach gives the correct threshold behavior for P waves, this should reduce the number of free parameters from six to three, but extensive checks against model calculations, similar to those already made by SSH to justify their particular effective range formulae, would be needed before one could trust such three-parameter formulae. Only if it could be shown that additional values of Δ_C would improve the determination of these P-wave parameters would additional p-p differential cross section measurements as a function of angle be worth considering. Even if that were the case, the author of this comment would like to see some specific nuclear or elementary particle problem where this increased precision in the triplet-odd P-waves is needed before he could encourage such experiments.

If no new nuclear information is likely to be forthcoming from p-p experiments below 10 MeV, the only other reason of which the author is aware for undertaking them is as a test of vacuum polarization; a specific proposal for new experiments near the Coulomb-nuclear interference minimum has been proposed for this purpose by Brolley.³⁸ A recent review of the tests of quantum electrodynamics at both low and high energy³⁹ shows complete agreement between theory and experiment to fantastically high accuracy, so it would be surprising indeed if this test failed. Expected modifications of the photon propagator due to muon and hadron pairs have, by the usual uncertainty principle argument, the same range as nuclear effects, so could not be disentangled without a complete theory of the strong interactions; expected modifications from a convergence factor, if indefinite metric theories of

finite quantum electrodynamics are followed, are necessarily of opposite sign, so would tend to cancel even these modifications, and in any case are of still shorter range. It is still true that vacuum polarization in p-p scattering is about the only place that the vacuum polarization correction to the photon propagator can be tested in a system containing only hadrons as physical particles,⁴⁰ and an attempt was made to check this test some years ago using the Wisconsin data.⁴¹ The result is that the apparent strength of the vacuum polarization correction when treated as a free parameter deviates significantly from the theoretical prediction. Since the effect varies systematically with energy, it is probably an indication of an unknown systematic error in the Wisconsin data, and very unlikely to be evidence for a breakdown of quantum electrodynamics. It scarcely seems worth while to spend another ten years on those, or new, differential cross section measurements chasing that will-of-the-wisp. The experiment proposed by Brolley³⁸ would provide a more sensitive test, but also is many orders of magnitude too gross to be expected to show up a discrepancy with theory.

The earlier discussion of n-p experiments should make it clear that there may well still be systematic errors lurking in the low energy neutron-proton scattering data. Continued attention to that problem is obviously desirable, but it seems unlikely that systematic error in these experiments can be driven down to the level where a test of the prediction for r_{np} to an accuracy of ± 0.03 FM would become believable. Any novel ideas here would be welcome, since three-nucleon calculations are quite sensitive to the value used for this parameter⁴²; the same applies, of course, to r_{nn} , but experimental precision there looks close to impossible.

We conclude that the era when new information about nuclear force parameters could be derived directly from nucleon-nucleon scattering experiments below 10 MeV is drawing to a close, and that in the absence of novel experimental or theoretical

ideas, the strenuous efforts which would be needed to improve the precision of the knowledge already obtained might better be directed to other objectives.

Extensive use has been made in this paper of unpublished calculations of the Foldy correction and charge-current distribution corrections using various potential models made by H. M. Lipinski in an earlier attempt to analyze the low-energy p-p data; this attempt was frustrated by the systematic errors discussed above. Extensive help with the more recent work was supplied by J. Post and W. Ross; a check on the assumption that charge-current distribution corrections affect only the nuclear parameters and not the values of a and r_0 derived from the data was made by E. Zeiger and V. N. Athavale. The author is indebted to Nelson Jarmie for informing him of the essential result of Ref. 26 in a private discussion; he is sorry that this acknowledgement must⁴³ now take so impersonal a form.

REFERENCES

1. H. P. Noyes and D. Y. Wong, Phys. Rev. Letters 3, 191 (1959).
2. M. Cini, S. Fubini, and A. Stanghellini, Phys. Rev. 114, 1633 (1959).
3. D. Y. Wong and H. P. Noyes, Phys. Rev. 126, 1988 (1962); in the numerical work, a wrong sign leading to a 12% numerical error in the Coulomb correction resulted in a value of a_{nn} of -28 FM rather than -19 FM; see discussion in Ref. 5 for details.
4. H. P. Noyes, Phys. Rev. Letters 12, 171 (1964).
5. H. P. Noyes, Nucl. Phys. 74, 508 (1965).
6. C. E. Engelke, R. E. Benenson, E. Melkoniam, and J. M. Lebowitz, Phys. Rev. 129, 324 (1963), hereinafter referred to as EBML.
7. H. P. Noyes, Phys. Rev. 130, 2025 (1963), and references therein.
8. T. L. Houk and R. L. Wilson, Rev. Mod. Phys. 39, 546 (1967) [Erratum: 40, 672 (1968)].
9. Erratum to Ref. 8.
10. L. Koester, Z. Physik 198, 187 (1967) [Erratum: 203, 515 (1967)].
11. H. P. Noyes and H. Fiedelney in Three Particle Scattering in Quantum Mechanics, J. Nuttall and J. Gillespie, eds. (W. A. Benjamin, Inc., New York, 1968); p. 195.
12. J. C. Davis and H. H. Barschall, Phys. Letters 27B, 636 (1968); I am indebted to Prof. Barschall for informing me of this result prior to publication.
13. D. F. Simmons, D. F. Cramer, and L. Cranberg, Bull. Am. Phys. Soc. 15, 475 (AE5) (1970).
14. H. P. Noyes and H. M. Lipinski, Phys. Rev. 162, 884 (1967).
15. P. Catillon, M. Chapellier, and D. Garreta, Nucl. Phys. B2, 93 (1967).
16. L. H. Johnston and D. E. Young, Phys. Rev. 116, 989 (1959); L. H. Johnston and Y. S. Tsai, Phys. Rev. 115, 1793 (1959).

17. R. J. Slobodrian, Phys. Rev. Letters 21, 438 (1968); Nuovo Cimento 40B, 443 (1965); Nucl. Phys. 85, 33 (1966).
18. R. J. Slobodrian, private communication and discussion.
19. R. J. Slobodrian, H. E. Conzett, E. Shield, and W. F. Tivol, Phys. Rev. 174, 1122 (1968).
20. M. H. MacGregor, R. A. Arndt, and R. M. Wright, Phys. Rev. 179, 1624 (1969).
21. Reference 20, footnote 13; the basis for this conclusion was the peculiar values and energy dependence of δ_0^E , which is obvious from Fig. 1 of this paper.
22. M. S. Sher, P. Signell, and L. Heller, Ann. Phys. 58, 1 (1970), hereinafter referred to as SSH; I am indebted to these authors for a preprint of this careful and detailed analysis.
23. The analysis of Ref. 4 used a phase shift determined by the author from the preliminary position of the interference minimum (M. L. Gursky and L. Heller, Bull. Am. Phys. Soc. 8, 605 (1963)) rather than the final value (Ref. GH in Table I) and preliminary values of the Wisconsin cross sections analyzed by the author rather than the final values (Ref. KDM of Table I). A more serious error was that the exponent in the Coulomb correction was of the wrong sign, as explained in Ref. 5. These deficiencies were corrected in the analysis presented by H. P. Noyes in Few Body Problems, Light Nuclei, and Nuclear Interactions, G. Paic and I. Slaus, eds., (Gordon and Breach, New York, 1968); p. 9; none of these corrections have much significance as is shown in the text.
24. N. Jarmie, R. E. Brown, R. L. Hutson, and J. L. Detch, Jr., Phys. Rev. Letters 24, 240 (1970).
25. J. Holdeman, P. Signell, and M. Sher, Phys. Rev. Letters 24, 243 (1970).
26. N. Jarmie, J. L. Lett, J. L. Detch, Jr., and R. L. Hutson, Phys. Rev. Letters 25, 34 (1970).

27. P. Signell and J. Holdeman, value from unpublished analysis quoted in Ref. 26.
28. Assistance by J. Post and W. A. Ross is gratefully acknowledged.
29. L. Heller, Phys. Rev. 120, 677 (1960).
30. L. L. Foldy and E. Eriksen, Phys. Rev. 98, 775 (1955).
31. The method for obtaining these approximations to the integral equation of Ref. 3 is discussed in Comptes Rendues du Congres International de Physique Nucleaire (Centre National de la Recherche Scientifique, Paris 1964), Vol. II, Paper i/C 104, p. 172. Comparison with an exact treatment using the Bargmann potential is made in Ref. 5, where it is noted that they are quantitatively inadequate for a model-independent calculation of a_{nn} , since they give -19.76 FM rather than the exact result of -17.84 FM; the reason for this is made clear by the work of Heller and Rich (Phys. Rev. 144, 1324 (1966)) who show that the pole in the amplitude calculated from the Bargmann potential becomes a branch cut when the e^2/r interaction is added. Thus calculations of the n-n scattering length from the p-p data to better than 2 FM require explicit assumptions about the two-nucleon wave function inside the range of forces. We still believe that for representing the energy variation of p-p data of currently available accuracy, Eq. (1) is adequate below 10 MeV. (see below).
32. Extensive unpublished numerical investigation of the effect of using this formula rather than the Hamada-Johnston potential for the energy variation below 3 MeV conducted by Noyes, Osborn, and Lipinski show that it might distort the results by close to one standard deviation in that energy region using available data. Now that the lever arm in the $k \cot \delta$ plot has been extended by more than a factor of 3 thanks to the results near 10 MeV, this sensitivity disappears. The purist will probably still prefer to fit his nuclear model directly to the phase shifts given in Table I rather than to a_{pp}^c and r_{pp}^c derived in this way, but I have yet to discover any experimental way to invalidate the simpler treatment.

33. This assumes that the charge-current distributions only change the interaction inside the range of nuclear forces, which is a priori probable because they are believed to arise from meson currents. There is a small effect due to the long-range part of the nonnuclear $1/r^3$ static magnetic dipole interaction, but this is detectable, if at all, only in the higher partial waves (cf. G. Breit and M. M. Ruppel, Phys. Rev. 127, 2123 (1967) and also Ref. 22). That the effect of these corrections is approximately linear in energy over this energy range, and therefore only affects the interpretation of a and r as derived from the data rather than their numerical values has been shown for the Hamada-Johnston potential by H. M. Lipinski, by the authors of Ref. 22 for a quite different nuclear model, and by E. Zeiger and V. N. Athavale for two uniformly charged spheres of appropriate radius; cf., also numerical tables in the Nuovo Cimento reference of footnote 17.
34. R. E. Schneider and R. M. Thaler, Phys. Rev. 137, B874 (1965), hereinafter referred to as ST.
35. R. J. Adler, B. T. Chertok, and H. C. Miller, Phys. Rev. C2, 69 (1970).
36. H. P. Noyes, in The Three Body Problem in Nuclear and Particle Physics, J.S.C. McKee and P. M. Rolph, eds. (North Holland Publishing Co., London 1970); p. 2.
37. G. Breit, K. A. Friedman, J. M. Holt, and R. E. Seamon, Phys. Rev. 170, 1424 (1968); I am indebted to Prof. Breit for sending me a copy of this very interesting survey prior to publication. A recent preprint by Breit and Rustgi notes that the $n+p \rightarrow \gamma+d$ discrepancy might be due to the neglect of $^3S_1 \rightarrow ^3S_1$ transitions.
38. J. E. Brolley, Aust. J. Phys. 22, 327 (1969).
39. S. D. Drell and S. J. Brodsky, Ann. Rev. Nucl. Sci. 20 (in press).
40. S. D. Drell first pointed this out to the author; this was the motivation for the analysis of Ref. 41.

41. H. P. Noyes, unpublished, but reported by L. Heller, Bull. Am. Phys. Soc. 9, 154 (T) (1964); the numerical results are quoted in Ref. 38, and have been reconfirmed in Ref. 22.
42. A. C. Phillips, Nucl. Phys. A107, 209 (1968); see also Ref. 11 and 35.
43. H. P. Noyes, Bull. Am. Phys. Soc. 15, 617 (IA2) (1970), and J. Shapiro, Ibid. (IA3).

TABLE I

Accurate proton-proton electric 1S_0 phase shifts below 10 MeV, and the corresponding vacuum polarization phases and Foldy corrections; the phase shift used in the conventional effective range expansion is obtained by $a_0^c = \delta_0^E + \tau_0 - \Delta_0$. The Foldy correction Δ_0 was computed by H. M. Lipinski for (a) the Hamada-Johnston potential, (b) a sum of three Yukawa potentials: OPE, intermediate range attraction with mass and coupling constant adjusted to fit a and r and short range repulsion with the ω -meson mass and coupling constant adjusted to make δ_0 go negative at about 250 MeV, and (c) OPE plus a purely attractive Bargmann potential: $m_p V(r)/\hbar^2 = -2\beta^2(\beta^2 - \alpha^2)/[\beta \cosh \beta r + \alpha \sinh \beta r]^2$. Model (c) gives a correction close to that for a single attractive Yukawa

potential as quoted in Ref. 30. References are given in the caption to Fig. 1.

Lab. Energy (MeV)	δ_0^E ($^\circ$)	Reference		$-\tau_0$ ($^\circ$)		$-\Delta_0$	
		Data	Analysis	(a)	(b)	(c)	
0.38243	14.6110 ± 0.0115	BSB	GH	0.1925	0.1926	0.1980	
1.397	39.3213 ± 0.028	KDM	(NH)	0.1935	0.1935	0.2056	
1.855	44.3292 ± 0.023	KDM	(NH)	0.1745	0.1745	0.1870	
2.425	48.3553 ± 0.026	KDM	(NH)	0.1544	0.1544	0.1667	
3.037	51.0233 ± 0.040	KDM	(NH)	0.1369	0.1368	0.1488	
9.918	55.23 ± 0.13	JJDH	SH	0.0683	0.0680	0.0774	

TABLE II

Analysis of the six accurate p-p phase shifts below 10 MeV given in Table I, using the Coulomb-corrected Cini-Fubini-Stanghellini shape dependence for fixed values of G^2 ; the Foldy correction used corresponds to a hard or stiff (Yukawa) repulsive core potential.

G^2	a_{pp}^c (FM)	r_{pp}^c (FM)	Chi-squared	p (FM ³)	q (FM ²)
10.0	-7.8192	2.7772	5.1609	0.4377	3.7339
12.0	-7.8216	2.7862	3.0732	0.5400	3.6037
14.0	-7.8243	2.7958	1.7932	0.6481	3.4642
16.0	-7.8272	2.8062	1.5863	0.7626	3.3146
18.0	-7.8304	2.8175	2.7952	0.8838	3.1538
20.0	-7.8337	2.8296	5.8800	1.0124	2.9803

FIGURE CAPTION

1. Comparison of the predictions of Eq. (1) fitted to the data of KDM and BSB as reported in Ref. 23 ($a_{pp}^c = -7.8275$, $r_{pp}^c = 2.7937$, $p = .64788$, $q = 3.4619$, $G^2 = 14$) with the shape-independent result ($p = 0$) subtracted. References noted in the figure are:

BSB	J. E. Brolley, J. P. Seagrave, and J. G. Beery, Phys. Rev. <u>135</u> , B1119 (1964).
GH	M. Gursky and L. Heller, Phys. Rev. <u>136</u> , B1693 (1964).
KDM	D. J. Knecht, P. F. Dahl, and S. Messelt, Phys. Rev. <u>148</u> , 1031 (1966).
(NH)	The OPE corrections for the higher partial waves used by KDM in the analysis of their experiment were supplied by H. P. Noyes, and the vacuum polarization correction was computed by him from the formulae of GH.
JJDH	N. Jarmie, J. L. Jett, J. L. Detch, Jr., and R. L. Hutson, Phys. Rev. Letters <u>25</u> , 34 (1970).
SH	P. Signell and J. Holdeman, as quoted in JJDH.
WMF	H. R. Worthington, J. M. McGruer, and D. E. Findley, Phys. Rev. <u>90</u> , 899 (1953).
S	R. J. Slobodrian, Phys. Rev. Letters <u>21</u> , 438 (1968).
SCST	R. J. Slobodrian, H. E. Conzett, E. Schield, and W. F. Tivol, Phys. Rev. <u>174</u> , 1122 (1968); — BGS background subtracted data, —D discriminator data; clearly the two interpretations of the data are mutually incompatible, and at most one set should be used.
JY	L. H. Johnston and D. E. Young, Phys. Rev. <u>116</u> , 989 (1959).
NL	H. P. Noyes and H. M. Lipinski, Phys. Rev. <u>162</u> , 884 (1967).
N	Preliminary analysis by this author.




Fig. 1