

Measurement of the branching fractions and CP -asymmetry of $B^- \rightarrow D_{(CP)}^0 K^-$ decays with the *BABAR* detector

B. Aubert,¹ R. Barate,¹ D. Boutigny,¹ J.-M. Gaillard,¹ A. Hicheur,¹ Y. Karyotakis,¹ J. P. Lees,¹ P. Robbe,¹ V. Tisserand,¹ A. Zghiche,¹ A. Palano,² A. Pompili,² J. C. Chen,³ N. D. Qi,³ G. Rong,³ P. Wang,³ Y. S. Zhu,³ G. Eigen,⁴ I. Ofte,⁴ B. Stugu,⁴ G. S. Abrams,⁵ A. W. Borgland,⁵ A. B. Breon,⁵ D. N. Brown,⁵ J. Button-Shafer,⁵ R. N. Cahn,⁵ E. Charles,⁵ C. T. Day,⁵ M. S. Gill,⁵ A. V. Gritsan,⁵ Y. Groysman,⁵ R. G. Jacobsen,⁵ R. W. Kadel,⁵ J. Kadyk,⁵ L. T. Kerth,⁵ Yu. G. Kolomensky,⁵ G. Kukartsev,⁵ C. LeClerc,⁵ M. E. Levi,⁵ G. Lynch,⁵ L. M. Mir,⁵ P. J. Oddone,⁵ T. J. Orimoto,⁵ M. Pripstein,⁵ N. A. Roe,⁵ A. Romosan,⁵ M. T. Ronan,⁵ V. G. Shelkov,⁵ A. V. Telnov,⁵ W. A. Wenzel,⁵ K. Ford,⁶ T. J. Harrison,⁶ C. M. Hawkes,⁶ D. J. Knowles,⁶ S. E. Morgan,⁶ R. C. Penny,⁶ A. T. Watson,⁶ N. K. Watson,⁶ K. Goetzen,⁷ T. Held,⁷ H. Koch,⁷ B. Lewandowski,⁷ M. Pelizaeus,⁷ K. Peters,⁷ H. Schmuecker,⁷ M. Steinke,⁷ J. T. Boyd,⁸ N. Chevalier,⁸ W. N. Cottingham,⁸ M. P. Kelly,⁸ T. E. Latham,⁸ C. Mackay,⁸ F. F. Wilson,⁸ K. Abe,⁹ T. Cuhadar-Donszelmann,⁹ C. Hearty,⁹ T. S. Mattison,⁹ J. A. McKenna,⁹ D. Thiessen,⁹ P. Kyberd,¹⁰ A. K. McKemey,¹⁰ L. Teodorescu,¹⁰ V. E. Blinov,¹¹ A. D. Bukin,¹¹ V. B. Golubev,¹¹ V. N. Ivanchenko,¹¹ E. A. Kravchenko,¹¹ A. P. Onuchin,¹¹ S. I. Serednyakov,¹¹ Yu. I. Skovpen,¹¹ E. P. Solodov,¹¹ A. N. Yushkov,¹¹ D. Best,¹² M. Bruinsma,¹² M. Chao,¹² I. Eschrich,¹² D. Kirkby,¹² A. J. Lankford,¹² M. Mandelkern,¹² R. K. Mommsen,¹² W. Roethel,¹² D. P. Stoker,¹² C. Buchanan,¹³ B. L. Hartfiel,¹³ J. W. Gary,¹⁴ B. C. Shen,¹⁴ K. Wang,¹⁴ D. del Re,¹⁵ H. K. Hadavand,¹⁵ E. J. Hill,¹⁵ D. B. MacFarlane,¹⁵ H. P. Paar,¹⁵ Sh. Rahatlou,¹⁵ V. Sharma,¹⁵ J. W. Berryhill,¹⁶ C. Campagnari,¹⁶ B. Dahmes,¹⁶ N. Kuznetsova,¹⁶ S. L. Levy,¹⁶ O. Long,¹⁶ A. Lu,¹⁶ M. A. Mazur,¹⁶ J. D. Richman,¹⁶ W. Verkerke,¹⁶ T. W. Beck,¹⁷ J. Beringer,¹⁷ A. M. Eisner,¹⁷ C. A. Heusch,¹⁷ W. S. Lockman,¹⁷ T. Schalk,¹⁷ R. E. Schmitz,¹⁷ B. A. Schumm,¹⁷ A. Seiden,¹⁷ M. Turri,¹⁷ W. Walkowiak,¹⁷ D. C. Williams,¹⁷ M. G. Wilson,¹⁷ J. Albert,¹⁸ E. Chen,¹⁸ G. P. Dubois-Felsmann,¹⁸ A. Dvoretzskii,¹⁸ D. G. Hitlin,¹⁸ I. Narsky,¹⁸ T. Piatenko,¹⁸ F. C. Porter,¹⁸ A. Ryd,¹⁸ A. Samuel,¹⁸ S. Yang,¹⁸ S. Jayatilake,¹⁹ G. Mancinelli,¹⁹ B. T. Meadows,¹⁹ M. D. Sokoloff,¹⁹ T. Abe,²⁰ F. Blanc,²⁰ P. Bloom,²⁰ S. Chen,²⁰ P. J. Clark,²⁰ W. T. Ford,²⁰ U. Nauenberg,²⁰ A. Olivas,²⁰ P. Rankin,²⁰ J. Roy,²⁰ J. G. Smith,²⁰ W. C. van Hoek,²⁰ L. Zhang,²⁰ J. L. Harton,²¹ T. Hu,²¹ A. Soffer,²¹ W. H. Toki,²¹ R. J. Wilson,²¹ J. Zhang,²¹ R. Aleksan,²² S. Emery,²² A. Gaidot,²² S. F. Ganzhur,²² P.-F. Giraud,²² G. Hamel de Monchenault,²² W. Kozanecki,²² M. Langer,²² M. Legendre,²² G. W. London,²² B. Mayer,²² G. Schott,²² G. Vasseur,²² Ch. Yeche,²² M. Zito,²² D. Altenburg,²³ T. Brandt,²³ J. Brose,²³ T. Colberg,²³ M. Dickopp,²³ A. Hauke,²³ H. M. Lacker,²³ E. Maly,²³ R. Müller-Pfefferkorn,²³ R. Nogowski,²³ S. Otto,²³ J. Schubert,²³ K. R. Schubert,²³ R. Schwierz,²³ B. Spaan,²³ D. Bernard,²⁴ G. R. Bonneaud,²⁴ F. Brochard,²⁴ J. Cohen-Tanugi,²⁴ P. Grenier,²⁴ Ch. Thiebaux,²⁴ G. Vasileiadis,²⁴ M. Verderi,²⁴ A. Khan,²⁵ D. Lavin,²⁵ F. Muheim,²⁵ S. Playfer,²⁵ J. E. Swain,²⁵ M. Andreotti,²⁶ V. Azzolini,²⁶ D. Bettoni,²⁶ C. Bozzi,²⁶ R. Calabrese,²⁶ G. Cibinetto,²⁶ E. Luppi,²⁶ M. Negrini,²⁶ L. Piemontese,²⁶ A. Sarti,²⁶ E. Treadwell,²⁷ F. Anulli,²⁸ * R. Baldini-Ferrolì,²⁸ A. Calcaterra,²⁸ R. de Sangro,²⁸ D. Falciari,²⁸ G. Finocchiaro,²⁸ P. Patteri,²⁸ I. M. Peruzzi,²⁸ * M. Piccolo,²⁸ A. Zallo,²⁸ A. Buzzo,²⁹ R. Capra,²⁹ R. Contri,²⁹ G. Crosetti,²⁹ M. Lo Vetere,²⁹ M. Macri,²⁹ M. R. Monge,²⁹ S. Passaggio,²⁹ C. Patrignani,²⁹ E. Robutti,²⁹ A. Santroni,²⁹ S. Tosi,²⁹ S. Bailey,³⁰ M. Morii,³⁰ E. Won,³⁰ R. S. Dubitzky,³¹ U. Langenegger,³¹ W. Bhimji,³² D. A. Bowerman,³² P. D. Dauncey,³² U. Egede,³² J. R. Gaillard,³² G. W. Morton,³² J. A. Nash,³² P. Sanders,³² G. P. Taylor,³² G. J. Grenier,³³ S.-J. Lee,³³ U. Mallik,³³ J. Cochran,³⁴ H. B. Crawley,³⁴ J. Lamsa,³⁴ W. T. Meyer,³⁴ S. Prell,³⁴ E. I. Rosenberg,³⁴ J. Yi,³⁴ M. Biasini,³⁵ M. Pioppi,³⁵ M. Davier,³⁶ G. Grosdidier,³⁶ A. Höcker,³⁶ S. Laplace,³⁶ F. Le Diberder,³⁶ V. Lepeltier,³⁶ A. M. Lutz,³⁶ T. C. Petersen,³⁶ S. Plaszczynski,³⁶ M. H. Schune,³⁶ L. Tantot,³⁶ G. Wormser,³⁶ V. Brigljević,³⁷ C. H. Cheng,³⁷ D. J. Lange,³⁷ D. M. Wright,³⁷ A. J. Bevan,³⁸ J. P. Coleman,³⁸ J. R. Fry,³⁸ E. Gabathuler,³⁸ R. Gamet,³⁸ M. Kay,³⁸ R. J. Parry,³⁸ D. J. Payne,³⁸ R. J. Sloane,³⁸ C. Touramanis,³⁸ J. J. Back,³⁹ C. M. Cormack,³⁹ P. F. Harrison,³⁹ H. W. Shorthouse,³⁹ P. B. Vidal,³⁹ C. L. Brown,⁴⁰ G. Cowan,⁴⁰ R. L. Flack,⁴⁰ H. U. Flaecher,⁴⁰ S. George,⁴⁰ M. G. Green,⁴⁰ A. Kurup,⁴⁰ C. E. Marker,⁴⁰ T. R. McMahon,⁴⁰ S. Ricciardi,⁴⁰ F. Salvatore,⁴⁰ G. Vaitsas,⁴⁰ M. A. Winter,⁴⁰

D. Brown,⁴¹ C. L. Davis,⁴¹ J. Allison,⁴² N. R. Barlow,⁴² R. J. Barlow,⁴² P. A. Hart,⁴² M. C. Hodgkinson,⁴² F. Jackson,⁴² G. D. Lafferty,⁴² A. J. Lyon,⁴² J. H. Weatherall,⁴² J. C. Williams,⁴² A. Farbin,⁴³ A. Jawahery,⁴³ D. Kovalskyi,⁴³ C. K. Lae,⁴³ V. Lillard,⁴³ D. A. Roberts,⁴³ G. Blaylock,⁴⁴ C. Dallapiccola,⁴⁴ K. T. Flood,⁴⁴ S. S. Hertzbach,⁴⁴ R. Kofler,⁴⁴ V. B. Koptchev,⁴⁴ T. B. Moore,⁴⁴ S. Saremi,⁴⁴ H. Staengle,⁴⁴ S. Willocq,⁴⁴ R. Cowan,⁴⁵ G. Sciolla,⁴⁵ F. Taylor,⁴⁵ R. K. Yamamoto,⁴⁵ D. J. J. Mangeol,⁴⁶ P. M. Patel,⁴⁶ S. H. Robertson,⁴⁶ A. Lazzaro,⁴⁷ F. Palombo,⁴⁷ J. M. Bauer,⁴⁸ L. Cremaldi,⁴⁸ V. Eschenburg,⁴⁸ R. Godang,⁴⁸ R. Kroeger,⁴⁸ J. Reidy,⁴⁸ D. A. Sanders,⁴⁸ D. J. Summers,⁴⁸ H. W. Zhao,⁴⁸ S. Brunet,⁴⁹ D. Cote-Ahern,⁴⁹ P. Taras,⁴⁹ H. Nicholson,⁵⁰ G. Raven,⁵¹ L. Wilden,⁵¹ C. Cartaro,⁵² N. Cavallo,⁵² G. De Nardo,⁵² F. Fabozzi,⁵²,[†] C. Gatto,⁵² L. Lista,⁵² P. Paolucci,⁵² D. Piccolo,⁵² C. Sciacca,⁵² C. P. Jessop,⁵³ J. M. LoSecco,⁵³ T. A. Gabriel,⁵⁴ B. Brau,⁵⁵ K. K. Gan,⁵⁵ K. Honscheid,⁵⁵ D. Hufnagel,⁵⁵ H. Kagan,⁵⁵ R. Kass,⁵⁵ T. Pulliam,⁵⁵ Q. K. Wong,⁵⁵ J. Brau,⁵⁶ R. Frey,⁵⁶ C. T. Potter,⁵⁶ N. B. Sinev,⁵⁶ D. Strom,⁵⁶ E. Torrence,⁵⁶ F. Colecchia,⁵⁷ A. Dorigo,⁵⁷ F. Galeazzi,⁵⁷ M. Margoni,⁵⁷ M. Morandin,⁵⁷ M. Posocco,⁵⁷ M. Rotondo,⁵⁷ F. Simonetto,⁵⁷ R. Stroili,⁵⁷ G. Tiozzo,⁵⁷ C. Voci,⁵⁷ M. Benayoun,⁵⁸ H. Briand,⁵⁸ J. Chauveau,⁵⁸ P. David,⁵⁸ Ch. de la Vaissière,⁵⁸ L. Del Buono,⁵⁸ O. Hamon,⁵⁸ M. J. J. John,⁵⁸ Ph. Leruste,⁵⁸ J. Ocariz,⁵⁸ M. Pivk,⁵⁸ L. Roos,⁵⁸ J. Stark,⁵⁸ S. T’Jampens,⁵⁸ G. Therin,⁵⁸ P. F. Manfredi,⁵⁹ V. Re,⁵⁹ P. K. Behera,⁶⁰ L. Gladney,⁶⁰ Q. H. Guo,⁶⁰ J. Panetta,⁶⁰ C. Angelini,⁶¹ G. Batignani,⁶¹ S. Bettarini,⁶¹ M. Bondioli,⁶¹ F. Bucci,⁶¹ G. Calderini,⁶¹ M. Carpinelli,⁶¹ V. Del Gamba,⁶¹ F. Forti,⁶¹ M. A. Giorgi,⁶¹ A. Lusiani,⁶¹ G. Marchiori,⁶¹ F. Martinez-Vidal,⁶¹ M. Morganti,⁶¹ N. Neri,⁶¹ E. Paoloni,⁶¹ M. Rama,⁶¹ G. Rizzo,⁶¹ F. Sandrelli,⁶¹ J. Walsh,⁶¹ M. Haire,⁶² D. Judd,⁶² K. Paick,⁶² D. E. Wagoner,⁶² G. Cavoto,⁶³,[‡] N. Danielson,⁶³ P. Elmer,⁶³ C. Lu,⁶³ V. Miftakov,⁶³ J. Olsen,⁶³ A. J. S. Smith,⁶³ H. A. Tanaka,⁶³ F. Bellini,⁶⁴ R. Faccini,⁶⁴,[§] F. Ferrarotto,⁶⁴ F. Ferroni,⁶⁴ M. Gaspero,⁶⁴ M. A. Mazzoni,⁶⁴ S. Morganti,⁶⁴ M. Pierini,⁶⁴ G. Piredda,⁶⁴ F. Safai Tehrani,⁶⁴ C. Voena,⁶⁴ S. Christ,⁶⁵ G. Wagner,⁶⁵ R. Waldi,⁶⁵ T. Adye,⁶⁶ N. De Groot,⁶⁶ B. Franek,⁶⁶ N. I. Geddes,⁶⁶ G. P. Gopal,⁶⁶ E. O. Olaiya,⁶⁶ S. M. Xella,⁶⁶ M. V. Purohit,⁶⁷ A. W. Weidemann,⁶⁷ F. X. Yumiceva,⁶⁷ D. Aston,⁶⁸ R. Bartoldus,⁶⁸ N. Berger,⁶⁸ A. M. Boyarski,⁶⁸ O. L. Buchmueller,⁶⁸ M. R. Convery,⁶⁸ D. P. Coupal,⁶⁸ D. Dong,⁶⁸ J. Dorfan,⁶⁸ D. Dujmic,⁶⁸ W. Dunwoodie,⁶⁸ R. C. Field,⁶⁸ T. Glanzman,⁶⁸ S. J. Gowdy,⁶⁸ E. Grauges-Pous,⁶⁸ T. Hadig,⁶⁸ V. Halyo,⁶⁸ T. Hryn’ova,⁶⁸ W. R. Innes,⁶⁸ M. H. Kelsey,⁶⁸ P. Kim,⁶⁸ M. L. Kocian,⁶⁸ D. W. G. S. Leith,⁶⁸ J. Libby,⁶⁸ S. Luitz,⁶⁸ V. Luth,⁶⁸ H. L. Lynch,⁶⁸ H. Marsiske,⁶⁸ R. Messner,⁶⁸ D. R. Muller,⁶⁸ C. P. O’Grady,⁶⁸ V. E. Ozcan,⁶⁸ A. Perazzo,⁶⁸ M. Perl,⁶⁸ S. Petrak,⁶⁸ B. N. Ratcliff,⁶⁸ A. Roodman,⁶⁸ A. A. Salnikov,⁶⁸ R. H. Schindler,⁶⁸ J. Schwiening,⁶⁸ G. Simi,⁶⁸ A. Snyder,⁶⁸ A. Soha,⁶⁸ J. Stelzer,⁶⁸ D. Su,⁶⁸ M. K. Sullivan,⁶⁸ J. Va’vra,⁶⁸ S. R. Wagner,⁶⁸ M. Weaver,⁶⁸ A. J. R. Weinstein,⁶⁸ W. J. Wisniewski,⁶⁸ D. H. Wright,⁶⁸ C. C. Young,⁶⁸ P. R. Burchat,⁶⁹ A. J. Edwards,⁶⁹ T. I. Meyer,⁶⁹ B. A. Petersen,⁶⁹ C. Roat,⁶⁹ M. Ahmed,⁷⁰ S. Ahmed,⁷⁰ M. S. Alam,⁷⁰ J. A. Ernst,⁷⁰ M. A. Saeed,⁷⁰ M. Saleem,⁷⁰ F. R. Wappler,⁷⁰ W. Bugg,⁷¹ M. Krishnamurthy,⁷¹ S. M. Spanier,⁷¹ R. Eckmann,⁷² H. Kim,⁷² J. L. Ritchie,⁷² R. F. Schwitters,⁷² J. M. Izen,⁷³ I. Kitayama,⁷³ X. C. Lou,⁷³ S. Ye,⁷³ F. Bianchi,⁷⁴ M. Bona,⁷⁴ F. Gallo,⁷⁴ D. Gamba,⁷⁴ C. Borean,⁷⁵ L. Bosisio,⁷⁵ G. Della Ricca,⁷⁵ S. Dittongo,⁷⁵ S. Grancagnolo,⁷⁵ L. Lanceri,⁷⁵ P. Poropat,⁷⁵ L. Vitale,⁷⁵ G. Vuagnin,⁷⁵ R. S. Panvini,⁷⁶ Sw. Banerjee,⁷⁷ C. M. Brown,⁷⁷ D. Fortin,⁷⁷ P. D. Jackson,⁷⁷ R. Kowalewski,⁷⁷ J. M. Roney,⁷⁷ H. R. Band,⁷⁸ S. Dasu,⁷⁸ M. Datta,⁷⁸ A. M. Eichenbaum,⁷⁸ J. R. Johnson,⁷⁸ P. E. Kutter,⁷⁸ H. Li,⁷⁸ R. Liu,⁷⁸ F. Di Lodovico,⁷⁸ A. Mihalyi,⁷⁸ A. K. Mohapatra,⁷⁸ Y. Pan,⁷⁸ R. Prepost,⁷⁸ S. J. Sekula,⁷⁸ J. H. von Wimmersperg-Toeller,⁷⁸ J. Wu,⁷⁸ S. L. Wu,⁷⁸ Z. Yu,⁷⁸ and H. Neal⁷⁹

(The BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

²Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

³Institute of High Energy Physics, Beijing 100039, China

⁴University of Bergen, Inst. of Physics, N-5007 Bergen, Norway

⁵Lawrence Berkeley National Laboratory and University of California, Berkeley, CA 94720, USA

⁶University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁷Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁸University of Bristol, Bristol BS8 1TL, United Kingdom

⁹University of British Columbia, Vancouver, BC, Canada V6T 1Z1

¹⁰Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹¹Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹²University of California at Irvine, Irvine, CA 92697, USA

¹³University of California at Los Angeles, Los Angeles, CA 90024, USA

¹⁴Univ. of California, Riverside, CA 92521

¹⁵University of California at San Diego, La Jolla, CA 92093, USA

- ¹⁶University of California at Santa Barbara, Santa Barbara, CA 93106, USA
- ¹⁷University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, CA 95064, USA
- ¹⁸California Institute of Technology, Pasadena, CA 91125, USA
- ¹⁹University of Cincinnati, Cincinnati, OH 45221, USA
- ²⁰University of Colorado, Boulder, CO 80309, USA
- ²¹Colorado State University, Fort Collins, CO 80523, USA
- ²²DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France
- ²³Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany
- ²⁴Ecole Polytechnique, LLR, F-91128 Palaiseau, France
- ²⁵University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom
- ²⁶Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy
- ²⁷Florida A&M University, Tallahassee, FL 32307, USA
- ²⁸Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy
- ²⁹Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy
- ³⁰Harvard University, Cambridge, MA 02138, USA
- ³¹Univ. Heidelberg, Philosophenweg 12, D-69120 Heidelberg, Germany
- ³²Imperial College London, London, SW7 2AZ, United Kingdom
- ³³University of Iowa, Iowa City, IA 52242, USA
- ³⁴Iowa State University, Ames, IA 50011-3160, USA
- ³⁵Istituto Naz. Fis. Nucleare, I-06100 Perugia, Italy
- ³⁶Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France
- ³⁷Lawrence Livermore National Laboratory, Livermore, CA 94550, USA
- ³⁸University of Liverpool, Liverpool L69 3BX, United Kingdom
- ³⁹Queen Mary, University of London, E1 4NS, United Kingdom
- ⁴⁰University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom
- ⁴¹University of Louisville, Louisville, KY 40292, USA
- ⁴²University of Manchester, Manchester M13 9PL, United Kingdom
- ⁴³University of Maryland, College Park, MD 20742, USA
- ⁴⁴University of Massachusetts, Amherst, MA 01003, USA
- ⁴⁵Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, MA 02139, USA
- ⁴⁶McGill University, Montréal, QC, Canada H3A 2T8
- ⁴⁷Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy
- ⁴⁸University of Mississippi, University, MS 38677, USA
- ⁴⁹Université de Montréal, Laboratoire René J. A. Lévesque, Montréal, QC, Canada H3C 3J7
- ⁵⁰Mount Holyoke College, South Hadley, MA 01075, USA
- ⁵¹NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands
- ⁵²Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy
- ⁵³University of Notre Dame, Notre Dame, IN 46556, USA
- ⁵⁴Oak Ridge National Laboratory, Oak Ridge, TN 37831, USA
- ⁵⁵Ohio State University, Columbus, OH 43210, USA
- ⁵⁶University of Oregon, Eugene, OR 97403, USA
- ⁵⁷Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy
- ⁵⁸Universités Paris VI et VII, Lab de Physique Nucléaire H. E., F-75252 Paris, France
- ⁵⁹Università di Pavia, Dipartimento di Elettronica and INFN, I-27100 Pavia, Italy
- ⁶⁰University of Pennsylvania, Philadelphia, PA 19104, USA
- ⁶¹Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy
- ⁶²Prairie View A&M University, Prairie View, TX 77446, USA
- ⁶³Princeton University, Princeton, NJ 08544, USA
- ⁶⁴Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy
- ⁶⁵Universität Rostock, D-18051 Rostock, Germany
- ⁶⁶Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom
- ⁶⁷University of South Carolina, Columbia, SC 29208, USA
- ⁶⁸Stanford Linear Accelerator Center, Stanford, CA 94309, USA
- ⁶⁹Stanford University, Stanford, CA 94305-4060, USA
- ⁷⁰State Univ. of New York, Albany, NY 12222, USA
- ⁷¹University of Tennessee, Knoxville, TN 37996, USA
- ⁷²University of Texas at Austin, Austin, TX 78712, USA
- ⁷³University of Texas at Dallas, Richardson, TX 75083, USA
- ⁷⁴Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy
- ⁷⁵Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy
- ⁷⁶Vanderbilt University, Nashville, TN 37235, USA
- ⁷⁷University of Victoria, Victoria, BC, Canada V8W 3P6
- ⁷⁸University of Wisconsin, Madison, WI 53706, USA
- ⁷⁹Yale University, New Haven, CT 06511, USA

We present a study of $B^- \rightarrow D_{CP}^0 K^-$ decays, where D_{CP}^0 is reconstructed in CP -even channels, based on a sample of 88.8 million $\Upsilon(4S) \rightarrow B\bar{B}$ decays collected with the *BABAR* detector at the PEP-II e^+e^- storage ring. We measure the ratio of Cabibbo-suppressed to Cabibbo-favored branching fractions $\mathcal{B}(B^- \rightarrow D_{CP}^0 K^-)/\mathcal{B}(B^- \rightarrow D_{CP}^0 \pi^-) = (8.8 \pm 1.6(\text{stat}) \pm 0.5(\text{syst})) \times 10^{-2}$ and the CP asymmetry $A_{CP} = 0.07 \pm 0.17(\text{stat}) \pm 0.06(\text{syst})$. We also measure $\mathcal{B}(B^- \rightarrow D^0 K^-)/\mathcal{B}(B^- \rightarrow D^0 \pi^-) = (8.31 \pm 0.35(\text{stat}) \pm 0.20(\text{syst})) \times 10^{-2}$ using a sample of 61.0 million $B\bar{B}$ pairs.

PACS numbers: 14.40.Nd, 13.25.Hw

The recent observation of CP violation in the B meson system [1] has provided a clean measurement of the angle β of the unitarity triangle. Although this measurement is in good agreement with the expectations of the Standard Model derived from other measurements of weak interactions, further measurements of CP violation in B decays are needed to overconstrain the unitarity triangle and confirm the CKM mechanism or observe deviations from it. A theoretically clean measurement of the angle γ can be obtained from the study of $B^- \rightarrow D^{(*)0} K^{(*)-}$ decays by reconstructing the D^0 meson in Cabibbo-allowed CP eigenstates and doubly Cabibbo-suppressed decays [2].

In this Letter we present the measurement of the ratios of Cabibbo-suppressed to Cabibbo-favored branching fractions

$$R_{(CP)} = \frac{\mathcal{B}(B^- \rightarrow D^0(D_{CP}^0)K^-) + \mathcal{B}(B^+ \rightarrow \bar{D}^0(D_{CP}^0)K^+)}{\mathcal{B}(B^- \rightarrow D^0(D_{CP}^0)\pi^-) + \mathcal{B}(B^+ \rightarrow \bar{D}^0(D_{CP}^0)\pi^+)} \quad (1)$$

with D^0 reconstructed in Cabibbo-allowed or CP -even (D_{CP}^0) channels. The direct CP asymmetry

$$A_{CP} = \frac{\mathcal{B}(B^- \rightarrow D_{CP}^0 K^-) - \mathcal{B}(B^+ \rightarrow D_{CP}^0 K^+)}{\mathcal{B}(B^- \rightarrow D_{CP}^0 K^-) + \mathcal{B}(B^+ \rightarrow D_{CP}^0 K^+)} \quad (2)$$

is also measured. The measurement of R uses a sample of 61.0 million $\Upsilon(4S)$ decays in $B\bar{B}$ pairs collected with the *BABAR* detector at the PEP-II asymmetric-energy B factory. The analysis of $B^\pm \rightarrow D_{CP}^0 K^\pm$ decays uses a sample of 88.8 million $B\bar{B}$ pairs. Since the *BABAR* detector is described in detail elsewhere [3], only the components that are crucial to this analysis are summarized here. Charged-particle tracking is provided by a five-layer silicon vertex tracker (SVT) and a 40-layer drift chamber (DCH). For charged-particle identification, ionization energy loss in the DCH and SVT, and Cherenkov radiation detected in a ring-imaging device (DIRC) are used. Photons are identified by the electromagnetic calorimeter (EMC), which comprises 6580 thallium-doped CsI crystals. These systems are mounted inside a 1.5-T solenoidal superconducting magnet. We use the GEANT [4] software to simulate interactions of particles traversing the detector, taking into account the varying accelerator and detector conditions.

We reconstruct $B^- \rightarrow D^0 h^-$ decays, where the prompt track h^- is a kaon or a pion (reference to the charge-conjugate state is implied here and throughout the text

unless otherwise stated). Candidates for D^0 are reconstructed in the non- CP flavor eigenstates $K^-\pi^+$, $K^-\pi^+\pi^+\pi^-$, $K^-\pi^+\pi^0$ (non- CP modes) and in the CP -even eigenstates $\pi^-\pi^+$ and K^-K^+ (CP modes).

To reduce the combinatorial background, only charged tracks with momenta greater than 150 MeV/ c are used to reconstruct $D^0 \rightarrow K^-\pi^+\pi^+\pi^-$ and $D^0 \rightarrow K^-\pi^+\pi^0$; the prompt particle h is required to have momentum greater than 1.4 GeV/ c . Particle ID information from the drift chamber and, when available, from the DIRC must be consistent with the kaon hypothesis for the K meson candidate in all D^0 modes and with the pion hypothesis for the π^\pm meson candidates in the $D^0 \rightarrow \pi^-\pi^+$ mode. For the prompt track to be identified as a pion or a kaon, we require that its Cherenkov angle be reconstructed with at least five photons. We reject a candidate track if its Cherenkov angle is consistent with that of a proton or if it is identified as an electron by the DCH and the EMC.

Photon candidates are required to have energies greater than 70 MeV. Photon pairs with invariant mass within the range 124–144 MeV/ c^2 and total energy greater than 200 MeV are considered π^0 candidates. To improve the momentum resolution, the π^0 candidates are kinematically fit with their mass constrained to the nominal π^0 mass [5].

The invariant mass of a D^0 candidate, $M(D^0)$, must be within 3σ of the mean fitted mass for the channels $K^-\pi^+$, $K^-\pi^+\pi^+\pi^-$ and K^-K^+ , and within 2σ for the $K^-\pi^+\pi^0$ channel. Candidates for $D^0 \rightarrow \pi^-\pi^+$ are selected in the range $1.80 < M(D^0) < 1.93$ GeV/ c^2 and the invariant mass of the $(h^-\pi^+)$ system, where π^+ is the pion from D^0 and h^- is the prompt track taken with the kaon mass hypothesis, must be greater than 1.9 GeV/ c^2 to reject background from $B^- \rightarrow D^0[\rightarrow K^-\pi^+]\pi^-$ and $B^- \rightarrow K^*[\rightarrow K^-\pi^+]\pi^-$ decays. For all the D^0 decay channels except the $\pi^-\pi^+$ mode a kinematical fit to the nominal D^0 mass [5] is applied. The $D^0 \rightarrow \pi^-\pi^+$ selection differs because of its particular background, as described later.

We reconstruct B meson candidates by combining a D^0 candidate with a track h . For the non- CP modes, the charge of the track h must match that of the kaon from the D^0 meson decay. We select B meson candidates by using the beam-energy-substituted mass $m_{ES} = \sqrt{(E_i^{*2}/2 + \mathbf{p}_i \cdot \mathbf{p}_B)^2/E_i^2 - p_B^2}$ and the energy difference $\Delta E = E_B^* - E_i^*/2$, where the subscripts i and B refer to

the initial e^+e^- system and the B candidate respectively, and the asterisk denotes the CM frame. The m_{ES} distributions for $B^- \rightarrow D^0 h^-$ signals are Gaussian distributions centered at the B mass with a resolution of $2.6 \text{ MeV}/c^2$, which does not depend on the decay mode or on the nature of the prompt track. In contrast, the ΔE distributions depend on the mass assigned to the prompt track and on the D^0 momentum resolution. We evaluate ΔE with the kaon mass hypothesis so that the distributions are centered near zero for $B^- \rightarrow D^0 K^-$ events and shifted by approximately 50 MeV for $B^- \rightarrow D^0 \pi^-$ events. The ΔE resolution is about 20 MeV for the $D^0 \rightarrow \pi^- \pi^+$ mode, and typically 17 MeV for the other D^0 decay modes. We select B mesons in the range $5.2 < m_{\text{ES}} < 5.3 \text{ GeV}/c^2$ with the exception of the $D^0 \rightarrow \pi^- \pi^+$ mode, for which m_{ES} is required to be within 3σ of the mean value. All B candidates are selected in the range $-0.10 < \Delta E < 0.13 \text{ GeV}$. For events with multiple $B^- \rightarrow D^0 h^-$ candidates, the best candidate is chosen based on the values of $M(D^0)$ and m_{ES} ; this happens in fewer than 1% of the selected events for two-body D^0 decays and in $\approx 4\%$ of the events for the other D^0 decays.

To reduce backgrounds from continuum production of light quarks, we make use of two quantities that exploit the different topologies of $e^+e^- \rightarrow q\bar{q}$ ($q = u, d, s, c$) and $B\bar{B}$ events. The first quantity is the normalized second Fox-Wolfram moment [6], $R_2 \equiv H_2/H_0$, where H_l is the l -order Fox-Wolfram moment of all the charged tracks and neutral clusters in the event. Only events with $R_2 < 0.5$ are selected. The second quantity is the angle θ_T between the thrust axes of the B candidate and of the remaining charged tracks and neutral clusters, evaluated in the CM. We require $|\cos\theta_T| < 0.9$ for the $D^0 \rightarrow K^- \pi^+$ mode, and $|\cos\theta_T| < 0.7$ for the $D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$ and $D^0 \rightarrow K^- \pi^+ \pi^0$ modes. For the $D^0 \rightarrow K^- K^+$ and $D^0 \rightarrow \pi^- \pi^+$ modes an additional quantity is used to suppress further the continuum background: the angle θ_{Dh} between the direction of one of the decay products of the D^0 and the direction of flight of the B , in the D^0 rest frame. The quantities $\cos\theta_T$ and $\cos\theta_{Dh}$ are uncorrelated for the signal but *not* for the continuum background. This correlation is exploited to make a more efficient cut in the $\cos\theta_T - \cos\theta_{Dh}$ plane.

The total reconstruction efficiencies, based on simulated signal events, are 42% ($K^- \pi^+$), 14% ($K^- \pi^+ \pi^+ \pi^-$), 8% ($K^- \pi^+ \pi^0$), 34% ($K^- K^+$) and 36% ($\pi^- \pi^+$).

The main contributions to the $B\bar{B}$ background for the non- CP modes come from the processes $B \rightarrow D^* h$ ($h = \pi, K$), $B^- \rightarrow D^0 \rho^-$ and mis-reconstructed $B^- \rightarrow D^0 h^-$. For D_{CP}^0 decays, the backgrounds $B^- \rightarrow K^- K^+ K^-$ and $B^- \rightarrow K^- \pi^+ \pi^-$ [7] must also be considered, since they have the same ΔE and m_{ES} distribution as the $D^0 K^-$ signal. The resonant component of these decays is negligible after the selection requirements for the CP modes.

For each D^0 decay mode an extended unbinned maximum-likelihood fit to the selected data events deter-

mines the signal and background yields n_i ($i = 1$ to M , where M is the total number of signal and background channels). Two kinds of signal events, $B^- \rightarrow D^0 \pi^-$ and $B^- \rightarrow D^0 K^-$, are considered, while the number of background sources depends on the D^0 channel. For non- CP modes we consider four kinds of backgrounds: candidates selected either from continuum or from $B\bar{B}$ events, in which the prompt track is either a pion or a kaon. In the case of $D^0 \rightarrow K^- K^+$ we consider two kinds of background depending on the nature of the prompt track. Finally, in the case of $D^0 \rightarrow \pi^- \pi^+$ we consider four contributions: the $B^- \rightarrow K^- \pi^+ \pi^-$ and $B^- \rightarrow \pi^- \pi^+ \pi^-$ decays and two kinds of generic background depending on the nature of the prompt track.

The input variables to the fit for the non- CP and the $D^0 \rightarrow K^- K^+$ modes are m_{ES} , ΔE , and a particle identification probability for the prompt track based on the Cherenkov angle θ_C , the momentum p and the polar angle θ of the track. For the $D^0 \rightarrow \pi^- \pi^+$ mode, m_{ES} is replaced by $M(D^0)$. This allows us to separate the $B^- \rightarrow D^0 K^-$ from the non-resonant $B^- \rightarrow K^- \pi^+ \pi^-$ contributions since the $(\pi^- \pi^+)$ -invariant-mass distribution peaks at the D^0 mass for signal while it is featureless for background. The extended likelihood function \mathcal{L} is defined as

$$\mathcal{L} = \exp \left(- \sum_{i=1}^M n_i \right) \prod_{j=1}^N \left[\sum_{i=1}^M n_i \mathcal{P}_i(\vec{x}_j; \vec{\alpha}_i) \right], \quad (3)$$

where N is the total number of observed events. The M functions $\mathcal{P}_i(\vec{x}_j; \vec{\alpha}_i)$ are the probability density functions (PDFs) for the variables \vec{x}_j , given the set of parameters $\vec{\alpha}_i$. They are evaluated as a product $\mathcal{P}_i = \mathcal{P}_i(\Delta E, x) \times \mathcal{P}_i(\theta_C)$ where $x = m_{\text{ES}}$ or $M(D^0)$ depending on the D^0 channel.

The Gaussian shape of the m_{ES} PDF for signal events is determined from a pure sample of $B^- \rightarrow D^0 \pi^-$, $D^0 \rightarrow K^- \pi^+$ decays selected from on-resonance data. The ΔE distribution for $B^- \rightarrow D^0 K^-$ signal events is parameterized with a Gaussian distribution whose parameters are determined from a pure sample of $B^- \rightarrow D^0 \pi^-$ events selected after the pion mass is assigned to the prompt track. The displaced ΔE distribution for $B^- \rightarrow D^0 \pi^-$ is parameterized with a sum of two Gaussian distributions. The shape of $M(D^0)$ is also described by a Gaussian distribution whose parameters are determined from data.

The parameters for the ΔE and m_{ES} distributions for the continuum background of the non- CP and $D^0 \rightarrow K^- K^+$ modes are determined from off-resonance data. The background shape in ΔE is parameterized with a linear function, while that of the m_{ES} is parameterized with an ARGUS threshold function [8] $f(m_{\text{ES}}) \propto m_{\text{ES}} \sqrt{1 - y^2} \exp[-\xi(1 - y^2)]$, where $y = m_{\text{ES}}/m_0$ and m_0 is the mean CM energy of the beams. The correlation between m_{ES} and ΔE for the generic $B\bar{B}$ background is taken into account with a two-dimensional PDF

determined from simulated events through a method based on the Kernel Estimation [9] technique. For the $D^0 \rightarrow K^- K^+$ mode the contribution from the non-resonant $B^- \rightarrow K^- K^- K^+$ decays is estimated [7] and added to the continuum and the generic $B\bar{B}$ background.

The ΔE and $M(D^0)$ distributions for the continuum and generic $B\bar{B}$ background of the $D^0 \rightarrow \pi^- \pi^+$ mode are determined from off-resonance data and simulated events, respectively. The ΔE distribution is described by a linear function while $M(D^0)$ is parameterized with the sum of a linear function (combinatorial background) and a Gaussian distribution (real $D^0 \rightarrow \pi^- \pi^+$). The $M(D^0)$ PDFs of the non-resonant $B^- \rightarrow h^- \pi^- \pi^+$ decays are described by linear functions, while the ΔE distributions are parameterized with one or two Gaussian distributions, as for the $B^- \rightarrow D^0 h^-$ signals.

Finally, the parameterization of the particle identification PDF is performed by fitting with a Gaussian distribution the background-subtracted distribution of the difference between the reconstructed and expected Cherenkov angles of the kaons and pions from D^0 decays, in a pure $D^{*+} \rightarrow D^0 \pi^+$ ($D^0 \rightarrow K^- \pi^+$) control sample.

The results of the fit are summarized in Table I. Figure 1 shows the distributions of ΔE for the combined non- CP and CP modes after enhancing the $B \rightarrow D^0 K$ purity by requiring that the prompt track be consistent with the kaon hypothesis and that $|m_{ES} - \langle m_{ES} \rangle| < 3\sigma$ ($|M(\pi^- \pi^+) - \langle M(\pi^- \pi^+) \rangle| < 3\sigma$ for $D^0 \rightarrow \pi^- \pi^+$). The projection of a likelihood fit, modified to take into account the tighter selection criteria, is overlaid in the Figure.

TABLE I: Results from the maximum-likelihood fit. For the $D^0 \rightarrow K^- K^+$ and $D^0 \rightarrow \pi^- \pi^+$ modes we quote the results for the fits performed on the whole sample and on the B^+ and B^- subsamples.

D^0 mode	$N(B \rightarrow D^0 \pi)$	$N(B \rightarrow D^0 K)$	$N(\Upsilon(4S))$
$K^- \pi^+$	4440 ± 69	360 ± 21	61.0×10^6
$K^- \pi^+ \pi^+ \pi^-$	2914 ± 56	242 ± 18	61.0×10^6
$K^- \pi^+ \pi^0$	2650 ± 56	208 ± 18	61.0×10^6
$K^- K^+$	565 ± 25	44.3 ± 9.0	88.8×10^6
$K^- K^+ [B^+]$	286 ± 18	16.7 ± 5.8	88.8×10^6
$K^- K^+ [B^-]$	280 ± 18	27.8 ± 6.8	88.8×10^6
$\pi^- \pi^+$	195 ± 17	24.2 ± 7.2	88.8×10^6
$\pi^- \pi^+ [B^+]$	99 ± 12	$16.8^{+5.6}_{-4.9}$	88.8×10^6
$\pi^- \pi^+ [B^-]$	96 ± 12	$6.5^{+5.1}_{-4.3}$	88.8×10^6

The ratios R and R_{CP} are computed by scaling the ratios of the numbers of $B^- \rightarrow D^0 K^-$ and $B^- \rightarrow D^0 \pi^-$ mesons by correction factors that account for small differences in the efficiency between $B^- \rightarrow D^0 K^-$ and $B^- \rightarrow D^0 \pi^-$ selection, estimated with simulated signal samples. The results are listed in Table II.

The direct CP asymmetry A_{CP} for the $B^\pm \rightarrow D_{CP}^0 K^\pm$ decays is calculated from the measured yields of positive




FIG. 1: Distributions of ΔE for events enhanced in $B \rightarrow D^0 K$ signal. Top: $D^0 \rightarrow K^- \pi^+, K^- \pi^+ \pi^+ \pi^-, K^- \pi^+ \pi^0$; bottom: $D^0 \rightarrow K^- K^+, \pi^- \pi^+$. Solid curves represent projections of the maximum likelihood fit; dashed, dashed-dotted and dotted curves represent the $B \rightarrow D^0 K$, $B \rightarrow D^0 \pi$ and background contributions.

and negative charged meson decays reported in Table I. We measure $A_{CP} = 0.07 \pm 0.17(\text{stat}) \pm 0.06(\text{syst})$.

Systematic uncertainties in the ratios R , R_{CP} and in the CP asymmetry A_{CP} arise primarily from uncertainties in signal yields due to imperfect knowledge of the PDF shapes. The parameters of the analytical PDFs are varied by $\pm 1\sigma$ and the difference in the signal yields is taken as a systematic uncertainty. When a $B\bar{B}$ PDF is parameterized through the kernel estimation, we repeat the fit using several statistically independent simulated $B\bar{B}$ samples to define the PDF. The width of the distribution of the difference between the new yields and the original yield is taken as the systematic uncertainty.

The uncertainties in the branching fractions of the channels contributing to the $B\bar{B}$ background have been taken into account. The correlations between the different sources of systematic errors, when non-negligible, are considered. An upper limit on intrinsic detector charge bias due to acceptance, tracking, and particle identification efficiency has been obtained from the measured asymmetries in the processes $B^- \rightarrow D^0 [\rightarrow K^- \pi^+] h^-$ and $B^- \rightarrow D_{CP}^0 \pi^-$, where CP violation is expected to be negligible. This limit (0.04) has been added in quadrature to the total systematic uncertainty on the CP asymmetry.

In conclusion, we have reconstructed $B^- \rightarrow D^0 K^-$ decays with D^0 mesons decaying to non- CP and CP -even eigenstates. The ratios $R_{(CP)}$ of the branching fractions $\mathcal{B}(B^- \rightarrow D_{(CP)}^0 K^-)$ and $\mathcal{B}(B^- \rightarrow D_{(CP)}^0 \pi^-)$ and the direct CP asymmetry A_{CP} have been measured. The

TABLE II: Measured ratios R and R_{CP} for different D^0 decay modes. The first error is statistical, the second is systematic.

$B^- \rightarrow D^0 h^-$ decay mode	$\mathcal{B}(B \rightarrow DK)/\mathcal{B}(B \rightarrow D\pi)$ (%)
$D^0 \rightarrow K^- \pi^+$	$8.4 \pm 0.5 \pm 0.2$
$D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$	$8.7 \pm 0.7 \pm 0.2$
$D^0 \rightarrow K^- \pi^+ \pi^0$	$7.7 \pm 0.7 \pm 0.3$
weighted mean	$8.31 \pm 0.35 \pm 0.20$
$D^0 \rightarrow K^- K^+$	$8.0 \pm 1.7 \pm 0.6$
$D^0 \rightarrow \pi^- \pi^+$	$12.9 \pm 4.0^{+1.1}_{-1.5}$
weighted mean	$8.8 \pm 1.6 \pm 0.5$

measured ratio R is consistent with Standard Model expectation ($\approx 7.5\%$) assuming factorization [10]. In the Standard Model $R_{CP}/R = 1 + r^2 + 2r \cos \delta \cos \gamma$ and $A_{CP} = 2r \sin \delta \sin \gamma / (1 + r^2 + 2r \cos \delta \cos \gamma)$, where $r \approx 0.1 - 0.2$ is the magnitude of the ratio of the amplitudes for the processes $B^- \rightarrow \bar{D}^0 K^-$ and $B^- \rightarrow D^0 K^-$, and δ is the (unknown) relative strong phase between these two amplitudes [2]. The measured values of R and R_{CP} are equal within errors, and A_{CP} is consistent with zero. These results, together with the ones obtained by CLEO and Belle [11], represent a first step towards the measurement of the angle γ and of direct CP violation in the B system using the $B^- \rightarrow D^0 K^-$ decays.

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support BABAR. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (the Netherlands), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Indi-

viduals have received support from the A. P. Sloan Foundation, Research Corporation, and Alexander von Humboldt Foundation.

- * Also with Università di Perugia, I-06100 Perugia, Italy
 - † Also with Università della Basilicata, I-85100 Potenza, Italy
 - ‡ Also with Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy
 - § Also with University of California at San Diego, La Jolla, CA 92093, USA
- [1] BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **87**, 091801 (2001); BELLE Collaboration, K. Abe *et al.*, Phys. Rev. Lett. **87**, 091802 (2001).
 - [2] M. Gronau and D. Wyler, Phys. Lett. **B265**, 172 (1991); M. Gronau and D. London, Phys. Lett. **B253** 483 (1991); D. Atwood, I. Dunietz and A. Soni, Phys. Rev. Lett. **78**, 3257 (1997); M. Gronau, Phys. Lett. **B557** 198-206 (2003).
 - [3] BABAR Collaboration, B. Aubert *et al.*, Nucl. Instr. and Methods **A479** 1 (2002).
 - [4] GEANT4 Collaboration, S. Agostinelli *et al.*, Nucl. Instr. and Methods **A506** 250 (2003).
 - [5] Particle Data Group, K. Hagiwara *et al.*, Phys. Rev. **D66** 010001 (2002).
 - [6] G.C. Fox and S. Wolfram, Phys. Rev. Lett. **41** 1581 (1978).
 - [7] Belle Collaboration, K. Abe *et al.*, Phys. Rev. **D65** 092005 (2002); BABAR Collaboration, B. Aubert *et al.*, hep-ex/0308065, submitted to Phys. Rev. Lett. .
 - [8] ARGUS Collaboration, H. Albrecht *et al.*, Z. Phys. **C48**, 543 (1990).
 - [9] K. S. Cranmer, Comp. Phys. Commun. **136**, 198-207 (2001).
 - [10] M Gronau *et al.*, Phys. Rev. **D52** 6356-6373 (1995).
 - [11] CLEO Collaboration, A. Bornheim *et al.*, Phys. Rev. **D68** 052002 (2003); Belle Collaboration, S. K. Swain *et al.*, Phys. Rev. **D68** 051101 (2003).