

Search for decays of B^0 mesons into pairs of charged leptons:
 $B^0 \rightarrow e^+e^-$, $B^0 \rightarrow \mu^+\mu^-$, $B^0 \rightarrow e^\pm\mu^\mp$

B. Aubert,¹ R. Barate,¹ D. Boutigny,¹ F. Couderc,¹ J.-M. Gaillard,¹ A. Hicheur,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Tisserand,¹ A. Zghiche,¹ A. Palano,² A. Pompili,² J. C. Chen,³ N. D. Qi,³ G. Rong,³ P. Wang,³ Y. S. Zhu,³ G. Eigen,⁴ I. Ofte,⁴ B. Stugu,⁴ G. S. Abrams,⁵ A. W. Borgland,⁵ A. B. Breon,⁵ D. N. Brown,⁵ J. Button-Shafer,⁵ R. N. Cahn,⁵ E. Charles,⁵ C. T. Day,⁵ M. S. Gill,⁵ A. V. Gritsan,⁵ Y. Groyzman,⁵ R. G. Jacobsen,⁵ R. W. Kadel,⁵ J. Kadyk,⁵ L. T. Kerth,⁵ Yu. G. Kolomensky,⁵ G. Kukartsev,⁵ G. Lynch,⁵ L. M. Mir,⁵ P. J. Oddone,⁵ T. J. Orimoto,⁵ M. Pripstein,⁵ N. A. Roe,⁵ M. T. Ronan,⁵ V. G. Shelkov,⁵ W. A. Wenzel,⁵ M. Barrett,⁶ K. E. Ford,⁶ T. J. Harrison,⁶ A. J. Hart,⁶ C. M. Hawkes,⁶ S. E. Morgan,⁶ A. T. Watson,⁶ M. Fritsch,⁷ K. Goetzen,⁷ T. Held,⁷ H. Koch,⁷ B. Lewandowski,⁷ M. Pelizaeus,⁷ M. Steinke,⁷ J. T. Boyd,⁸ N. Chevalier,⁸ W. N. Cottingham,⁸ M. P. Kelly,⁸ T. E. Latham,⁸ F. F. Wilson,⁸ T. Cuhadar-Donszelmann,⁹ C. Hearty,⁹ N. S. Knecht,⁹ T. S. Mattison,⁹ J. A. McKenna,⁹ D. Thiessen,⁹ A. Khan,¹⁰ P. Kyberd,¹⁰ L. Teodorescu,¹⁰ A. E. Blinov,¹¹ V. E. Blinov,¹¹ V. P. Druzhinin,¹¹ V. B. Golubev,¹¹ V. N. Ivanchenko,¹¹ E. A. Kravchenko,¹¹ A. P. Onuchin,¹¹ S. I. Serednyakov,¹¹ Yu. I. Skovpen,¹¹ E. P. Solodov,¹¹ A. N. Yushkov,¹¹ D. Best,¹² M. Bruinsma,¹² M. Chao,¹² I. Eschrich,¹² D. Kirkby,¹² A. J. Lankford,¹² M. Mandelkern,¹² R. K. Mommsen,¹² W. Roethel,¹² D. P. Stoker,¹² C. Buchanan,¹³ B. L. Hartfiel,¹³ S. D. Foulkes,¹⁴ J. W. Gary,¹⁴ B. C. Shen,¹⁴ K. Wang,¹⁴ D. del Re,¹⁵ H. K. Hadavand,¹⁵ E. J. Hill,¹⁵ D. B. MacFarlane,¹⁵ H. P. Paar,¹⁵ Sh. Rahatlou,¹⁵ V. Sharma,¹⁵ J. W. Berryhill,¹⁶ C. Campagnari,¹⁶ B. Dahmes,¹⁶ S. L. Levy,¹⁶ O. Long,¹⁶ A. Lu,¹⁶ M. A. Mazur,¹⁶ J. D. Richman,¹⁶ W. Verkerke,¹⁶ T. W. Beck,¹⁷ A. M. Eisner,¹⁷ C. A. Heusch,¹⁷ W. S. Lockman,¹⁷ G. Nesom,¹⁷ T. Schalk,¹⁷ R. E. Schmitz,¹⁷ B. A. Schumm,¹⁷ A. Seiden,¹⁷ P. Spradlin,¹⁷ D. C. Williams,¹⁷ M. G. Wilson,¹⁷ J. Albert,¹⁸ E. Chen,¹⁸ G. P. Dubois-Felsmann,¹⁸ A. Dvoretzkii,¹⁸ D. G. Hitlin,¹⁸ I. Narsky,¹⁸ T. Piatenko,¹⁸ F. C. Porter,¹⁸ A. Ryd,¹⁸ A. Samuel,¹⁸ S. Yang,¹⁸ S. Jayatilake,¹⁹ G. Mancinelli,¹⁹ B. T. Meadows,¹⁹ M. D. Sokoloff,¹⁹ T. Abe,²⁰ F. Blanc,²⁰ P. Bloom,²⁰ S. Chen,²⁰ W. T. Ford,²⁰ U. Nauenberg,²⁰ A. Olivas,²⁰ P. Rankin,²⁰ J. G. Smith,²⁰ J. Zhang,²⁰ L. Zhang,²⁰ A. Chen,²¹ J. L. Harton,²¹ A. Soffer,²¹ W. H. Toki,²¹ R. J. Wilson,²¹ Q. L. Zeng,²¹ D. Altenburg,²² T. Brandt,²² J. Brose,²² M. Dickopp,²² E. Feltresi,²² A. Hauke,²² H. M. Lacker,²² R. Müller-Pfefferkorn,²² R. Nogowski,²² S. Otto,²² A. Petzold,²² J. Schubert,²² K. R. Schubert,²² R. Schwierz,²² B. Spaan,²² J. E. Sundermann,²² D. Bernard,²³ G. R. Bonneaud,²³ F. Brochard,²³ P. Grenier,²³ S. Schrenk,²³ Ch. Thiebaut,²³ G. Vasileiadis,²³ M. Verderi,²³ D. J. Bard,²⁴ P. J. Clark,²⁴ D. Lavin,²⁴ F. Muheim,²⁴ S. Playfer,²⁴ Y. Xie,²⁴ M. Andreotti,²⁵ V. Azzolini,²⁵ D. Bettoni,²⁵ C. Bozzi,²⁵ R. Calabrese,²⁵ G. Cibinetto,²⁵ E. Luppi,²⁵ M. Negrini,²⁵ L. Piemontese,²⁵ A. Sarti,²⁵ E. Treadwell,²⁶ R. Baldini-Ferrolli,²⁷ A. Calcaterra,²⁷ R. de Sangro,²⁷ G. Finocchiaro,²⁷ P. Patteri,²⁷ M. Piccolo,²⁷ A. Zallo,²⁷ A. Buzzo,²⁸ R. Capra,²⁸ R. Contri,²⁸ G. Crosetti,²⁸ M. Lo Vetere,²⁸ M. Macri,²⁸ M. R. Monge,²⁸ S. Passaggio,²⁸ C. Patrignani,²⁸ E. Robutti,²⁸ A. Santroni,²⁸ S. Tosi,²⁸ S. Bailey,²⁹ G. Brandenburg,²⁹ M. Morii,²⁹ E. Won,²⁹ R. S. Dubitzky,³⁰ U. Langenegger,³⁰ W. Bhimji,³¹ D. A. Bowerman,³¹ P. D. Dauncey,³¹ U. Egede,³¹ J. R. Gaillard,³¹ G. W. Morton,³¹ J. A. Nash,³¹ M. B. Nikolich,³¹ G. P. Taylor,³¹ M. J. Charles,³² G. J. Grenier,³² U. Mallik,³² J. Cochran,³³ H. B. Crawley,³³ J. Lamsa,³³ W. T. Meyer,³³ S. Prell,³³ E. I. Rosenberg,³³ J. Yi,³³ M. Davier,³⁴ G. Grosdidier,³⁴ A. Höcker,³⁴ S. Laplace,³⁴ F. Le Diberder,³⁴ V. Lepeltier,³⁴ A. M. Lutz,³⁴ T. C. Petersen,³⁴ S. Plaszczynski,³⁴ M. H. Schune,³⁴ L. Tantot,³⁴ G. Wormser,³⁴ C. H. Cheng,³⁵ D. J. Lange,³⁵ M. C. Simani,³⁵ D. M. Wright,³⁵ A. J. Bevan,³⁶ C. A. Chavez,³⁶ J. P. Coleman,³⁶ I. J. Forster,³⁶ J. R. Fry,³⁶ E. Gabathuler,³⁶ R. Gamet,³⁶ R. J. Parry,³⁶ D. J. Payne,³⁶ R. J. Sloane,³⁶ C. Touramanis,³⁶ J. J. Back,^{37,*} C. M. Cormack,^{37,*} P. F. Harrison,^{37,*} F. Di Lodovico,³⁷ G. B. Mohanty,^{37,*} C. L. Brown,³⁸ G. Cowan,³⁸ R. L. Flack,³⁸ H. U. Flaecher,³⁸ M. G. Green,³⁸ P. S. Jackson,³⁸ T. R. McMahon,³⁸ S. Ricciardi,³⁸ F. Salvatore,³⁸ M. A. Winter,³⁸ D. Brown,³⁹ C. L. Davis,³⁹ J. Allison,⁴⁰ N. R. Barlow,⁴⁰ R. J. Barlow,⁴⁰ M. C. Hodgkinson,⁴⁰ G. D. Lafferty,⁴⁰ A. J. Lyon,⁴⁰ J. C. Williams,⁴⁰ A. Farbin,⁴¹ W. D. Hulsbergen,⁴¹ A. Jawahery,⁴¹ D. Kovalskyi,⁴¹ C. K. Lae,⁴¹ V. Lillard,⁴¹ D. A. Roberts,⁴¹ G. Blaylock,⁴² C. Dallapiccola,⁴² K. T. Flood,⁴² S. S. Hertzbach,⁴²

R. Kofler,⁴² V. B. Koptchev,⁴² T. B. Moore,⁴² S. Saremi,⁴² H. Staengle,⁴² S. Willocq,⁴² R. Cowan,⁴³ G. Sciolla,⁴³ F. Taylor,⁴³ R. K. Yamamoto,⁴³ D. J. J. Mangeol,⁴⁴ P. M. Patel,⁴⁴ S. H. Robertson,⁴⁴ A. Lazzaro,⁴⁵ F. Palombo,⁴⁵ J. M. Bauer,⁴⁶ L. Cremaldi,⁴⁶ V. Eschenburg,⁴⁶ R. Godang,⁴⁶ R. Kroeger,⁴⁶ J. Reidy,⁴⁶ D. A. Sanders,⁴⁶ D. J. Summers,⁴⁶ H. W. Zhao,⁴⁶ S. Brunet,⁴⁷ D. Côté,⁴⁷ P. Taras,⁴⁷ H. Nicholson,⁴⁸ F. Fabozzi,⁴⁹ † C. Gatto,⁴⁹ L. Lista,⁴⁹ D. Monorchio,⁴⁹ P. Paolucci,⁴⁹ D. Piccolo,⁴⁹ C. Sciacca,⁴⁹ M. Baak,⁵⁰ H. Bulten,⁵⁰ G. Raven,⁵⁰ H. L. Snoek,⁵⁰ L. Wilden,⁵⁰ C. P. Jessop,⁵¹ J. M. LoSecco,⁵¹ T. A. Gabriel,⁵² T. Allmendinger,⁵³ B. Brau,⁵³ K. K. Gan,⁵³ K. Honscheid,⁵³ D. Hufnagel,⁵³ H. Kagan,⁵³ R. Kass,⁵³ T. Pulliam,⁵³ A. M. Rahimi,⁵³ R. Ter-Antonyan,⁵³ Q. K. Wong,⁵³ J. Brau,⁵⁴ R. Frey,⁵⁴ O. Igonkina,⁵⁴ C. T. Potter,⁵⁴ N. B. Sinev,⁵⁴ D. Strom,⁵⁴ E. Torrence,⁵⁴ F. Colecchia,⁵⁵ A. Dorigo,⁵⁵ F. Galeazzi,⁵⁵ M. Margoni,⁵⁵ M. Morandin,⁵⁵ M. Posocco,⁵⁵ M. Rotondo,⁵⁵ F. Simonetto,⁵⁵ R. Stroili,⁵⁵ G. Tiozzo,⁵⁵ C. Voci,⁵⁵ M. Benayoun,⁵⁶ H. Briand,⁵⁶ J. Chauveau,⁵⁶ P. David,⁵⁶ Ch. de la Vaissière,⁵⁶ L. Del Buono,⁵⁶ O. Hamon,⁵⁶ M. J. J. John,⁵⁶ Ph. Leruste,⁵⁶ J. Malcles,⁵⁶ J. Ocariz,⁵⁶ M. Pivk,⁵⁶ L. Roos,⁵⁶ S. T'Jampens,⁵⁶ G. Therin,⁵⁶ P. F. Manfredi,⁵⁷ V. Re,⁵⁷ P. K. Behera,⁵⁸ L. Gladney,⁵⁸ Q. H. Guo,⁵⁸ J. Panetta,⁵⁸ F. Anulli,^{27,59} M. Biasini,⁵⁹ I. M. Peruzzi,^{27,59} M. Pioppi,⁵⁹ C. Angelini,⁶⁰ G. Batignani,⁶⁰ S. Bettarini,⁶⁰ M. Bondioli,⁶⁰ F. Bucci,⁶⁰ G. Calderini,⁶⁰ M. Carpinelli,⁶⁰ F. Forti,⁶⁰ M. A. Giorgi,⁶⁰ A. Lusiani,⁶⁰ G. Marchiori,⁶⁰ F. Martinez-Vidal,^{60,‡} M. Morganti,⁶⁰ N. Neri,⁶⁰ E. Paoloni,⁶⁰ M. Rama,⁶⁰ G. Rizzo,⁶⁰ F. Sandrelli,⁶⁰ J. Walsh,⁶⁰ M. Haire,⁶¹ D. Judd,⁶¹ K. Paick,⁶¹ D. E. Wagoner,⁶¹ N. Danielson,⁶² P. Elmer,⁶² Y. P. Lau,⁶² C. Lu,⁶² V. Miftakov,⁶² J. Olsen,⁶² A. J. S. Smith,⁶² A. V. Telnov,⁶² F. Bellini,⁶³ G. Cavoto,^{62,63} R. Faccini,⁶³ F. Ferrarotto,⁶³ F. Ferroni,⁶³ M. Gaspero,⁶³ L. Li Gioi,⁶³ M. A. Mazzoni,⁶³ S. Morganti,⁶³ M. Pierini,⁶³ G. Piredda,⁶³ F. Safai Tehrani,⁶³ C. Voena,⁶³ S. Christ,⁶⁴ G. Wagner,⁶⁴ R. Waldi,⁶⁴ T. Adye,⁶⁵ N. De Groot,⁶⁵ B. Franek,⁶⁵ N. I. Geddes,⁶⁵ G. P. Gopal,⁶⁵ E. O. Olaiya,⁶⁵ R. Aleksan,⁶⁶ S. Emery,⁶⁶ A. Gaidot,⁶⁶ S. F. Ganzhur,⁶⁶ P.-F. Giraud,⁶⁶ G. Hamel de Monchenault,⁶⁶ W. Kozanecki,⁶⁶ M. Langer,⁶⁶ M. Legendre,⁶⁶ G. W. London,⁶⁶ B. Mayer,⁶⁶ G. Schott,⁶⁶ G. Vasseur,⁶⁶ Ch. Yèche,⁶⁶ M. Zito,⁶⁶ M. V. Purohit,⁶⁷ A. W. Weidemann,⁶⁷ J. R. Wilson,⁶⁷ F. X. Yumiceva,⁶⁷ D. Aston,⁶⁸ R. Bartoldus,⁶⁸ N. Berger,⁶⁸ A. M. Boyarski,⁶⁸ O. L. Buchmueller,⁶⁸ R. Claus,⁶⁸ M. R. Convery,⁶⁸ M. Cristinziani,⁶⁸ G. De Nardo,⁶⁸ D. Dong,⁶⁸ J. Dorfan,⁶⁸ D. Dujmic,⁶⁸ W. Dunwoodie,⁶⁸ E. E. Elsen,⁶⁸ S. Fan,⁶⁸ R. C. Field,⁶⁸ T. Glanzman,⁶⁸ S. J. Gowdy,⁶⁸ T. Hadig,⁶⁸ V. Halyo,⁶⁸ C. Hast,⁶⁸ T. Hryn'ova,⁶⁸ W. R. Innes,⁶⁸ M. H. Kelsey,⁶⁸ P. Kim,⁶⁸ M. L. Kocian,⁶⁸ D. W. G. S. Leith,⁶⁸ J. Libby,⁶⁸ S. Luitz,⁶⁸ V. Luth,⁶⁸ H. L. Lynch,⁶⁸ H. Marsiske,⁶⁸ R. Messner,⁶⁸ D. R. Muller,⁶⁸ C. P. O'Grady,⁶⁸ V. E. Ozcan,⁶⁸ A. Perazzo,⁶⁸ M. Perl,⁶⁸ S. Petrak,⁶⁸ B. N. Ratcliff,⁶⁸ A. Roodman,⁶⁸ A. A. Salnikov,⁶⁸ R. H. Schindler,⁶⁸ J. Schwiening,⁶⁸ G. Simi,⁶⁸ A. Snyder,⁶⁸ A. Soha,⁶⁸ J. Stelzer,⁶⁸ D. Su,⁶⁸ M. K. Sullivan,⁶⁸ J. Va'vra,⁶⁸ S. R. Wagner,⁶⁸ M. Weaver,⁶⁸ A. J. R. Weinstein,⁶⁸ W. J. Wisniewski,⁶⁸ M. Wittgen,⁶⁸ D. H. Wright,⁶⁸ A. K. Yarritu,⁶⁸ C. C. Young,⁶⁸ P. R. Burchat,⁶⁹ A. J. Edwards,⁶⁹ T. I. Meyer,⁶⁹ B. A. Petersen,⁶⁹ C. Roat,⁶⁹ S. Ahmed,⁷⁰ M. S. Alam,⁷⁰ J. A. Ernst,⁷⁰ M. A. Saeed,⁷⁰ M. Saleem,⁷⁰ F. R. Wappler,⁷⁰ W. Bugg,⁷¹ M. Krishnamurthy,⁷¹ S. M. Spanier,⁷¹ R. Eckmann,⁷² H. Kim,⁷² J. L. Ritchie,⁷² A. Satpathy,⁷² R. F. Schwitters,⁷² J. M. Izen,⁷³ I. Kitayama,⁷³ X. C. Lou,⁷³ S. Ye,⁷³ F. Bianchi,⁷⁴ M. Bona,⁷⁴ F. Gallo,⁷⁴ D. Gamba,⁷⁴ C. Borean,⁷⁵ L. Bosisio,⁷⁵ C. Cartaro,⁷⁵ F. Cossutti,⁷⁵ G. Della Ricca,⁷⁵ S. Dittongo,⁷⁵ S. Grancagnolo,⁷⁵ L. Lanceri,⁷⁵ P. Poropat,^{75,§} L. Vitale,⁷⁵ G. Vuagnin,⁷⁵ R. S. Panvini,⁷⁶ Sw. Banerjee,⁷⁷ C. M. Brown,⁷⁷ D. Fortin,⁷⁷ P. D. Jackson,⁷⁷ R. Kowalewski,⁷⁷ J. M. Roney,⁷⁷ R. J. Sobie,⁷⁷ H. R. Band,⁷⁸ S. Dasu,⁷⁸ M. Datta,⁷⁸ A. M. Eichenbaum,⁷⁸ M. Graham,⁷⁸ J. J. Hollar,⁷⁸ J. R. Johnson,⁷⁸ P. E. Kutter,⁷⁸ H. Li,⁷⁸ R. Liu,⁷⁸ A. Mihalyi,⁷⁸ A. K. Mohapatra,⁷⁸ Y. Pan,⁷⁸ R. Prepost,⁷⁸ A. E. Rubin,⁷⁸ S. J. Sekula,⁷⁸ P. Tan,⁷⁸ J. H. von Wimmersperg-Toeller,⁷⁸ J. Wu,⁷⁸ S. L. Wu,⁷⁸ Z. Yu,⁷⁸ M. G. Greene,⁷⁹ and H. Neal⁷⁹

(The BABAR Collaboration)

BABAR Collaboration

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

²Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

³Institute of High Energy Physics, Beijing 100039, China

⁴University of Bergen, Inst. of Physics, N-5007 Bergen, Norway

⁵Lawrence Berkeley National Laboratory and University of California, Berkeley, CA 94720, USA

⁶University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁷Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁸University of Bristol, Bristol BS8 1TL, United Kingdom

⁹University of British Columbia, Vancouver, BC, Canada V6T 1Z1

¹⁰Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

- ¹¹*Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia*
- ¹²*University of California at Irvine, Irvine, CA 92697, USA*
- ¹³*University of California at Los Angeles, Los Angeles, CA 90024, USA*
- ¹⁴*University of California at Riverside, Riverside, CA 92521, USA*
- ¹⁵*University of California at San Diego, La Jolla, CA 92093, USA*
- ¹⁶*University of California at Santa Barbara, Santa Barbara, CA 93106, USA*
- ¹⁷*University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, CA 95064, USA*
- ¹⁸*California Institute of Technology, Pasadena, CA 91125, USA*
- ¹⁹*University of Cincinnati, Cincinnati, OH 45221, USA*
- ²⁰*University of Colorado, Boulder, CO 80309, USA*
- ²¹*Colorado State University, Fort Collins, CO 80523, USA*
- ²²*Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany*
- ²³*Ecole Polytechnique, LLR, F-91128 Palaiseau, France*
- ²⁴*University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom*
- ²⁵*Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy*
- ²⁶*Florida A&M University, Tallahassee, FL 32307, USA*
- ²⁷*Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy*
- ²⁸*Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy*
- ²⁹*Harvard University, Cambridge, MA 02138, USA*
- ³⁰*Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany*
- ³¹*Imperial College London, London, SW7 2AZ, United Kingdom*
- ³²*University of Iowa, Iowa City, IA 52242, USA*
- ³³*Iowa State University, Ames, IA 50011-3160, USA*
- ³⁴*Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France*
- ³⁵*Lawrence Livermore National Laboratory, Livermore, CA 94550, USA*
- ³⁶*University of Liverpool, Liverpool L69 7ZE, United Kingdom*
- ³⁷*Queen Mary, University of London, E1 4NS, United Kingdom*
- ³⁸*University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom*
- ³⁹*University of Louisville, Louisville, KY 40292, USA*
- ⁴⁰*University of Manchester, Manchester M13 9PL, United Kingdom*
- ⁴¹*University of Maryland, College Park, MD 20742, USA*
- ⁴²*University of Massachusetts, Amherst, MA 01003, USA*
- ⁴³*Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, MA 02139, USA*
- ⁴⁴*McGill University, Montréal, QC, Canada H3A 2T8*
- ⁴⁵*Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy*
- ⁴⁶*University of Mississippi, University, MS 38677, USA*
- ⁴⁷*Université de Montréal, Laboratoire René J. A. Lévesque, Montréal, QC, Canada H3C 3J7*
- ⁴⁸*Mount Holyoke College, South Hadley, MA 01075, USA*
- ⁴⁹*Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy*
- ⁵⁰*NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands*
- ⁵¹*University of Notre Dame, Notre Dame, IN 46556, USA*
- ⁵²*Oak Ridge National Laboratory, Oak Ridge, TN 37831, USA*
- ⁵³*Ohio State University, Columbus, OH 43210, USA*
- ⁵⁴*University of Oregon, Eugene, OR 97403, USA*
- ⁵⁵*Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy*
- ⁵⁶*Universités Paris VI et VII, Laboratoire de Physique Nucléaire et de Hautes Energies, F-75252 Paris, France*
- ⁵⁷*Università di Pavia, Dipartimento di Elettronica and INFN, I-27100 Pavia, Italy*
- ⁵⁸*University of Pennsylvania, Philadelphia, PA 19104, USA*
- ⁵⁹*Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy*
- ⁶⁰*Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy*
- ⁶¹*Prairie View A&M University, Prairie View, TX 77446, USA*
- ⁶²*Princeton University, Princeton, NJ 08544, USA*
- ⁶³*Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy*
- ⁶⁴*Universität Rostock, D-18051 Rostock, Germany*
- ⁶⁵*Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom*
- ⁶⁶*DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France*
- ⁶⁷*University of South Carolina, Columbia, SC 29208, USA*
- ⁶⁸*Stanford Linear Accelerator Center, Stanford, CA 94309, USA*
- ⁶⁹*Stanford University, Stanford, CA 94305-4060, USA*
- ⁷⁰*State Univ. of New York, Albany, NY 12222, USA*
- ⁷¹*University of Tennessee, Knoxville, TN 37996, USA*
- ⁷²*University of Texas at Austin, Austin, TX 78712, USA*
- ⁷³*University of Texas at Dallas, Richardson, TX 75083, USA*
- ⁷⁴*Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy*

⁷⁵Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy

⁷⁶Vanderbilt University, Nashville, TN 37235, USA

⁷⁷University of Victoria, Victoria, BC, Canada V8W 3P6

⁷⁸University of Wisconsin, Madison, WI 53706, USA

⁷⁹Yale University, New Haven, CT 06511, USA

(Dated: August 19, 2004)

We present a search for the decays $B^0 \rightarrow e^+e^-$, $B^0 \rightarrow \mu^+\mu^-$, and $B^0 \rightarrow e^\pm\mu^\mp$ in data collected at the $\Upsilon(4S)$ resonance with the *BABAR* detector at the SLAC *B* Factory. Using a data set of 111 fb^{-1} , we find no evidence for a signal in any of the three channels investigated and set the following branching fraction upper limits at the 90% confidence level: $\mathcal{B}(B^0 \rightarrow e^+e^-) < 6.1 \times 10^{-8}$, $\mathcal{B}(B^0 \rightarrow \mu^+\mu^-) < 8.3 \times 10^{-8}$, and $\mathcal{B}(B^0 \rightarrow e^\pm\mu^\mp) < 18 \times 10^{-8}$.

PACS numbers: 13.20.He,14.40.Nd

In the Standard Model (SM), rare B decays such as $B^0 \rightarrow \ell^+\ell^-$, where ℓ refers to e or μ , are expected to proceed through diagrams such as those shown in Fig. 1 (charge conjugate processes are included implicitly throughout). These decays are highly suppressed since they involve a $b \rightarrow d$ transition and require an internal quark annihilation within the B meson. In addition, the decays are helicity suppressed by factors of $(m_\ell/m_B)^2$ where m_ℓ is the mass of the lepton ℓ and m_B is the mass of the B meson. B^0 decays to leptons of two different flavors, violate lepton flavor conservation and are therefore forbidden in the SM, although permitted in extensions to the SM with non-zero neutrino mass [1]. The SM expectations are given in Table I.

Since these processes are highly suppressed in the SM, they are potentially sensitive probes of physics beyond the SM. In the minimally supersymmetric Standard Model (MSSM) the branching fraction for these decays can be enhanced by orders of magnitude [2]. In particular, for MSSM models with modified minimal flavor violation ($\overline{\text{MFV}}$) and large $\tan\beta$ [3], the branching fraction can be increased by up to four orders of magnitude. Experimental bounds can restrict allowed regions of parameter space, specifically the mass of the charged Higgs boson. In non-MSSM models with two Higgs doublets and natural flavor conservation at large $\tan\beta$, an increase in the branching fraction of several orders of magnitude is expected [4]. $B^0 \rightarrow \ell^+\ell^-$ decays are also allowed in specific models containing leptoquarks [5] and supersymmetric (SUSY) models without R-parity [6]. The branching fractions for the flavor violating channels $B^0 \rightarrow \ell_i^+\ell_j^-$ ($i \neq j$) are expected to be exceedingly small but can be enhanced by leptoquarks or R -parity violating operators in SUSY models.

To date, $B^0 \rightarrow \ell^+\ell^-$ decays have not been observed. As shown in Table I, experimental limits are approaching a level of sensitivity that will restrict the allowed parameter space of models that produce $B^0 \rightarrow \ell^+\ell^-$ branching fraction enhancements of a few orders of magnitude with respect to the SM rates.

The data used in these analyses were collected with the *BABAR* detector at the PEP-II e^+e^- storage ring




FIG. 1: Representative Feynman diagrams for $B^0 \rightarrow \ell^+\ell^-$ in the Standard Model.

TABLE I: The expected branching fractions in the Standard Model [7] and the current best upper limits (U.L.) at the 90% C.L.

Decay Mode	SM Expectation	CLEO [8] 9.1 fb^{-1}	Belle [9] 78 fb^{-1}	CDF [10] 0.17 fb^{-1}
e^+e^-	1.9×10^{-15}	8.3×10^{-7}	1.9×10^{-7}	–
$\mu^+\mu^-$	8.0×10^{-11}	6.1×10^{-7}	1.6×10^{-7}	1.5×10^{-7}
$e^\pm\mu^\mp$	–	15×10^{-7}	1.7×10^{-7}	–

and correspond to an integrated luminosity of 111 fb^{-1} accumulated at the $\Upsilon(4S)$ resonance (“on-resonance”) and 11.9 fb^{-1} accumulated at a center-of-mass (c.m.) energy about 40 MeV below the $\Upsilon(4S)$ resonance (“off-resonance”). The latter sample is used for non-resonant $q\bar{q}$ ($q = u, d, s$, and c) background studies. The collider is operated with asymmetric beam energies, producing a boost ($\beta\gamma = 0.55$) of the $\Upsilon(4S)$ along the collision axis.

The *BABAR* detector is optimized for the asymmetric beam configuration at PEP-II and is described in detail in [11]. The 1.5-T superconducting solenoidal magnet, whose cylindrical volume is 1.4 m in radius and 3 m long, contains a charged-particle tracking system, a Cherenkov detector dedicated to charged-particle identification, and central and forward electromagnetic CsI calorimeters (EMC). The segmented flux return, includ-

ing endcaps, is instrumented with resistive plate chambers for muon and K_L^0 identification.

The presence of two charged high-momentum leptons provides a very clean signature for the three decay modes under consideration. We require two oppositely-charged high-momentum leptons (*i.e.* $|p_\ell^*| \sim m_B/2$ where p_ℓ^* is the c.m. momentum of lepton ℓ from a common vertex consistent with the decay of a B^0 meson. Since the signal events contain two B^0 mesons and no additional particles, the total energy of each B^0 in the c.m. must be equal to half of the total beam energy. We define

$$m_{\text{ES}} = \sqrt{(E_{\text{beam}}^*)^2 - \left(\sum_i \mathbf{p}_i^*\right)^2} \quad (1)$$

$$\Delta E = \sum_i \sqrt{m_i^2 + (\mathbf{p}_i^*)^2} - E_{\text{beam}}^*, \quad (2)$$

where E_{beam}^* is the (e^+ or e^-) beam energy in the c.m. frame, \mathbf{p}_i^* is the momentum of lepton i in the c.m. frame, and m_i is the mass of lepton i . In Eq. (1), E_{beam}^* is used as opposed to E_B^* because E_{beam}^* is known with much greater precision. For correctly reconstructed B^0 mesons, m_{ES} peaks at the mass of the B^0 meson with a resolution of about 2.5 MeV/ c^2 and ΔE peaks near zero.

To reduce background from lepton misidentification, we require the leptons to satisfy stringent electron and muon identification criteria [12]. The electron identification efficiency is greater than 93% with a mis-identification rate of less than 0.3%. The muon identification efficiency ranges from (55 – 70)% (depending on run period) with a mis-identification rate of 3%. Electron energy lost through bremsstrahlung is partially recovered by adding the energy of photons that lie within a 3 degree cone about the electron direction.

Suppression of background from non-resonant $q\bar{q}$ production is provided by a series of topological requirements. In particular, we require $|\cos\theta_T| < 0.8$, where θ_T is the angle in the c.m. frame between the thrust axis of the particles that form the reconstructed B^0 candidate and the thrust axis of the remaining tracks and neutral clusters in the event. In addition we employ cuts on the invariant mass of the “Rest Of the Event” (all tracks not associated with the B^0 candidate where all non-leptonic tracks are assumed to be pions) of $m_{\text{ROE}} > 0.5$ GeV and on the second normalized Fox-Wolfram moment of $R_2 < 0.8$ [13]. We also cut on the total multiplicity of both charged tracks and neutral particles by means of the variable N_{mult} defined as $N_{\text{mult}} = N_{\text{trk}} + N_\gamma/2$, where N_{trk} is the total number of tracks in the event and N_γ is the number of photons found with an energy $E_\gamma > 80$ MeV. We require $N_{\text{mult}} \geq 5.5$ for the ee and $e\mu$ channels and $N_{\text{mult}} \geq 5.0$ for the $\mu\mu$ channel. This variable is especially useful in the rejection of radiative Bhabha events. We also require that the total energy in the EMC (E_{EMC}) be less than 11 GeV. This cut is ef-




FIG. 2: Distributions of signal MC (hatched) and sideband data (points) for the e^+e^- channel after the initial selection cuts for (a) $|\cos\theta_T|$, (b) m_{ROE} , (c) R_2 , and (d) N_{mult} . Arrows indicate final cut values. All distributions are normalized to unity.

fective in reducing background from QED e^+e^- events, including radiative Bhabhas with many conversions.

Four of the selection criteria given above ($|\cos\theta_T|$, m_{ROE} , N_{mult} , and E_{EMC}) were simultaneously optimized for the best upper limit on $\mathcal{B}(B^0 \rightarrow \ell^+\ell^-)$ where the assumed number of observed events is determined from a Poisson distribution with the mean equal to the expected background. Sideband data are compared with signal Monte Carlo (MC) for the e^+e^- channel for four of these variables, $|\cos\theta_T|$, m_{ROE} , R_2 , and N_{mult} in Fig. 2.

The $B^0 \rightarrow \ell^+\ell^-$ candidates are selected by simultaneous requirements on the energy difference ΔE and the energy-substituted mass m_{ES} . For the $B^0 \rightarrow \mu^+\mu^-$ decay mode, the size of this “signal box” is chosen to be $[+2, -2]\sigma$ of the m_{ES} distribution and $[+2, -2]\sigma$ for the ΔE . In the cases of the $B^0 \rightarrow e^+e^-$ and $B^0 \rightarrow e^\pm\mu^\mp$ decay modes, the signal box sizes in m_{ES} are also $[+2, -2]\sigma$ but in ΔE are relaxed to $[+2, -3]\sigma$ and $[+2, -2.5]\sigma$, re-

TABLE II: Definition of the three different sideband boxes used for all three decay modes.

Sideband Box	span in ΔE [GeV]	span in m_{ES} [GeV/ c^2]
Grand Sideband	(-0.40, 0.40)	(5.20, 5.26)
Upper m_{ES}	(0.20, 0.50)	(5.20, 5.29)
Lower m_{ES}	(-0.50, -0.20)	(5.20, 5.29)

spectively, to accommodate the tail in the distribution resulting from uncorrected bremsstrahlung and final state radiation. The resolution in m_{ES} is obtained from a fit to a Gaussian distribution, whereas the resolution in ΔE is obtained from a fit to an empirical function [14] that gives a good description of this tail.

We estimate the background level in the signal box from the data sidebands assuming that it is described by the ARGUS function [15] in m_{ES} and an exponential function in ΔE . We use these parameterizations to extrapolate the background level found in the sidebands into the signal box. As indicated in Table II, three different sideband boxes are used. The grand sideband box is used to estimate the functional form of the ΔE distribution. The upper and lower m_{ES} sideband boxes are used to estimate the functional form of the m_{ES} distribution. Peaking backgrounds from misidentified two-body B decay modes were estimated using an MC sample equivalent to more than 20 times the data luminosity and found to be negligible. The total background expectations and signal efficiencies are given in Table III.

The systematic uncertainties on the signal efficiency ε , the number of B^0 mesons produced in the data, and the background estimate are incorporated into the determination of the upper limit on $\mathcal{B}(B^0 \rightarrow \ell^+\ell^-)$. Since the signal efficiency is determined from MC simulation only, differences between data and the simulation would result in an error in our normalization. To estimate this uncertainty we perform comparisons of data and MC using high statistics control samples that have similar characteristics to our $B^0 \rightarrow \ell^+\ell^-$ signal. The optimal control samples are $B^0 \rightarrow J/\psi K_S^0$, with $J/\psi \rightarrow e^+e^-$ for $B^0 \rightarrow e^+e^-$ and $J/\psi \rightarrow \mu^+\mu^-$ for $B^0 \rightarrow \mu^+\mu^-$, respectively. Since there exists no appropriate control sample for the $e^\pm\mu^\mp$ mode, we use the larger of the systematic errors derived from either the ee or $\mu\mu$ modes. In performing these comparisons we found a substantial uncertainty on the signal efficiency to be due to differences between data and the MC simulation in the mean and resolutions of various quantities, depending on the channel. For the electron channels the dominant quantities are ΔE and m_{ROE} whereas for the muon channels they are $|\cos\theta_T|$, N_{mult} , and m_{ROE} . When combined with the uncertainties on tracking efficiency of 2.6% and that for particle identification (1.0% per electron, 3.0% per muon), the total systematic uncertainty on the efficiency is estimated to be 5.7%, 7.1%, and 6.8% for the ee , $\mu\mu$, and $e\mu$ modes respectively.

The background estimate is obtained from a fit to sideband data, so the primary uncertainty is due to fluctuations in the fit procedure as events fall in or out of the sideband box. We have studied the stability of the fit and the background estimate when adding or removing events from the m_{ES} and ΔE histograms. We find that the fit is unbiased and stable to a level significantly less than the statistical uncertainty on the background estimate.

TABLE III: Summary of the analyses where N_{obs} and $N_{\text{exp}}^{\text{bg}}$ are the observed and expected number of events in the signal box, ε is the efficiency, and $\mathcal{B}_{\text{UL}}(B^0 \rightarrow \ell^+\ell^-)$ is the upper limit on the branching fraction at the 90% C.L. Systematic uncertainties on $N_{\text{exp}}^{\text{bg}}$ and ε are given.

Decay Mode	N_{obs}	$N_{\text{exp}}^{\text{bg}}$	$\varepsilon[\%]$	$\mathcal{B}_{\text{UL}}(B^0 \rightarrow \ell^+\ell^-)$
e^+e^-	0	0.71 ± 0.31	21.8 ± 1.2	6.1×10^{-8}
$\mu^+\mu^-$	0	0.72 ± 0.26	15.9 ± 1.1	8.3×10^{-8}
$e^\pm\mu^\mp$	2	1.29 ± 0.44	18.1 ± 1.2	18×10^{-8}

As shown in Fig. 3 and Table III, when the contents of the signal box were revealed, 0, 0, and 2 events were found in the ee , $\mu\mu$, and $e\mu$ channels respectively. As can be seen in Table III, the numbers of events found in the signal boxes are compatible with the expected background for each mode.

An upper limit on the branching fraction is computed using

$$\mathcal{B}(B^0 \rightarrow \ell^+\ell^-) = \frac{N_{\text{UL}}(N_{\text{obs}})}{(N_{B^0} + N_{\bar{B}^0}) \cdot \varepsilon}, \quad (3)$$

where $N_{\text{UL}}(N_{\text{obs}})$ is the Poisson 90% U.L. on the number of events assuming N_{obs} events have been observed, $N_{B^0}(N_{\bar{B}^0})$ is the number of $B^0(\bar{B}^0)$ mesons produced in the data, and ε is the signal efficiency. We have $N_{B^0} + N_{\bar{B}^0} = N_{B\bar{B}}$ under the assumption of equal production of $B^0\bar{B}^0$ and B^+B^- in $\Upsilon(4S)$ decays. For our data set $N_{B\bar{B}} = (122.5 \pm 1.2) \times 10^6$.

We follow the technique of [16] in order to account for the presence of background and to include our systematic




FIG. 3: Distribution of events in m_{ES} and ΔE for $B^0 \rightarrow e^+e^-$ (top left), $B^0 \rightarrow \mu^+\mu^-$ (top right), and $B^0 \rightarrow e^\pm\mu^\mp$ (bottom).

uncertainties in the determination of the upper limit. As summarized in Table III, the resulting upper limits at the 90% C.L. for $\mathcal{B}(B^0 \rightarrow e^+e^-)$, $\mathcal{B}(B^0 \rightarrow \mu^+\mu^-)$, and $\mathcal{B}(B^0 \rightarrow e^\pm\mu^\mp)$ are 6.1×10^{-8} , 8.3×10^{-8} , and 18×10^{-8} respectively.

These bounds are stringent enough to place interesting constraints on popular models. For example, for the MSSM (MFV) models, the relation between $\mathcal{B}(B^0 \rightarrow \mu^+\mu^-)$ and the mass of the charged Higgs boson m_H is given as a function of $\tan\beta$ in [3]. We find that for $\tan\beta = 60$, $m_H > 138$ GeV (90% C.L.).

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support BABAR. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from CONACyT (Mexico), A. P. Sloan Foundation, Research Corporation, and Alexander von Humboldt Foundation.

* Now at Department of Physics, University of Warwick, Coventry, United Kingdom

† Also with Università della Basilicata, Potenza, Italy

‡ Also with IFIC, Instituto de Física Corpuscular, CSIC-Universidad de Valencia, Valencia, Spain

§ Deceased

[1] M. Maki, M. Nakagawa, and S. Sakata, Prog. Theor.

Phys. **28**, 870 (1962); V.N. Gribov and B. Pontecorvo, Phys. Lett. **B28**, 493 (1969).

- [2] A. Dedes, H. K. Dreiner and U. Nierste, Phys. Rev Lett. **87** (2001) 251804.
- [3] C. Bobeth *et al.*, Phys. Rev. D **66**, 074021 (2002), J. Urban, Proceedings of the 10th International Conference on Supersymmetry and Unification of Fundamental Interactions (SUSY02), hep-ph/0210286.
- [4] G. L. Kane, C. Kolda and J. E. Lennon, hep-ph/0310042.
- [5] S. Davidson, D. C. Bailey and B. A. Campbell, Z. Phys. C **61**, 613 (1994).
- [6] D. Roy, Phys. Lett. **B283** (1992) 270.
- [7] A. Ali, C. Greub, and T. Mannel, in *Proceedings of the ECFA Workshop on the Physics of the European B Meson Factory*, edited by R. Aleksan and A. Ali, p. 155 (ECFA-93-151,C93/03/26), DESY-93-016.
- [8] CLEO Collaboration, T. Bergfeld *et al.*, Phys. Rev. D **62**, 091102(R) (2000).
- [9] Belle Collaboration, M.-C. Chang *et al.*, Phys. Rev. D **68**, 111101(R) (2003).
- [10] CDF Collaboration, D. Acosta *et al.*, hep-ex/0403032.
- [11] BABAR Collaboration, B. Aubert *et al.*, Nucl. Instr. and Methods **A479**, 1 (2002).
- [12] BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. D **67**, 032002 (2003).
- [13] G.C. Fox and S. Wolfram, Phys. Rev. Lett. **41** (1978) 1581.
- [14] BABAR defines this empirical function to be
- $$f(E) = A \cdot \exp \left[-\frac{1}{2} \left(\frac{\log(1+\tau(E-\nu) \cdot \frac{\sinh(\tau\sqrt{\log 4})}{\sigma\tau\sqrt{\log 4}})}{\tau} \right)^2 + \tau^2 \right],$$
- where τ is the “tail parameter” (describing how much is contained in the tail), σ is the width, and ν is the peak position.
- [15] ARGUS Collaboration, H. Albrecht *et al.*, Phys. Lett. B **241**, 278 (1990);
- [16] R. Barlow, Comput. Phys. Commun. **149**, 97 (2002).