

Production and decay of Ξ_c^0 at **BABAR**

- B. Aubert,¹ R. Barate,¹ D. Boutigny,¹ F. Couderc,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Poireau,¹ V. Tisserand,¹
A. Zghiche,¹ E. Grauges,² A. Palano,³ M. Pappagallo,³ A. Pompili,³ J. C. Chen,⁴ N. D. Qi,⁴ G. Rong,⁴ P. Wang,⁴
Y. S. Zhu,⁴ G. Eigen,⁵ I. Ofte,⁵ B. Stugu,⁵ G. S. Abrams,⁶ A. W. Borgland,⁶ A. B. Breon,⁶ D. N. Brown,⁶
J. Button-Shafer,⁶ R. N. Cahn,⁶ E. Charles,⁶ C. T. Day,⁶ M. S. Gill,⁶ A. V. Gritsan,⁶ Y. Groysman,⁶
R. G. Jacobsen,⁶ R. W. Kadel,⁶ J. Kadyk,⁶ L. T. Kerth,⁶ Yu. G. Kolomensky,⁶ G. Kukartsev,⁶ G. Lynch,⁶
L. M. Mir,⁶ P. J. Oddone,⁶ T. J. Orimoto,⁶ M. Pripstein,⁶ N. A. Roe,⁶ M. T. Ronan,⁶ W. A. Wenzel,⁶ M. Barrett,⁷
K. E. Ford,⁷ T. J. Harrison,⁷ A. J. Hart,⁷ C. M. Hawkes,⁷ S. E. Morgan,⁷ A. T. Watson,⁷ M. Fritsch,⁸ K. Goetzen,⁸
T. Held,⁸ H. Koch,⁸ B. Lewandowski,⁸ M. Pelizaeus,⁸ K. Peters,⁸ T. Schroeder,⁸ M. Steinke,⁸ J. T. Boyd,⁹
J. P. Burke,⁹ N. Chevalier,⁹ W. N. Cottingham,⁹ M. P. Kelly,⁹ T. Cuhadar-Donszelmann,¹⁰ C. Hearty,¹⁰
N. S. Knecht,¹⁰ T. S. Mattison,¹⁰ J. A. McKenna,¹⁰ D. Thiessen,¹⁰ A. Khan,¹¹ P. Kyberd,¹¹ L. Teodorescu,¹¹
A. E. Blinov,¹² V. E. Blinov,¹² A. D. Bukan,¹² V. P. Druzhinin,¹² V. B. Golubev,¹² V. N. Ivanchenko,¹²
E. A. Kravchenko,¹² A. P. Onuchin,¹² S. I. Serednyakov,¹² Yu. I. Skovpen,¹² E. P. Solodov,¹² A. N. Yushkov,¹²
D. Best,¹³ M. Bondioli,¹³ M. Bruinsma,¹³ M. Chao,¹³ I. Eschrich,¹³ D. Kirkby,¹³ A. J. Lankford,¹³ M. Mandelkern,¹³
R. K. Mommsen,¹³ W. Roethel,¹³ D. P. Stoker,¹³ C. Buchanan,¹⁴ B. L. Hartfiel,¹⁴ A. J. R. Weinstein,¹⁴
S. D. Foulkes,¹⁵ J. W. Gary,¹⁵ O. Long,¹⁵ B. C. Shen,¹⁵ K. Wang,¹⁵ L. Zhang,¹⁵ D. del Re,¹⁶ H. K. Hadavand,¹⁶
E. J. Hill,¹⁶ D. B. MacFarlane,¹⁶ H. P. Paar,¹⁶ S. Rahatlou,¹⁶ V. Sharma,¹⁶ J. W. Berryhill,¹⁷ C. Campagnari,¹⁷
A. Cunha,¹⁷ B. Dahmes,¹⁷ T. M. Hong,¹⁷ A. Lu,¹⁷ M. A. Mazur,¹⁷ J. D. Richman,¹⁷ W. Verkerke,¹⁷ T. W. Beck,¹⁸
A. M. Eisner,¹⁸ C. J. Flacco,¹⁸ C. A. Heusch,¹⁸ J. Kroseberg,¹⁸ W. S. Lockman,¹⁸ G. Nesom,¹⁸ T. Schalk,¹⁸
B. A. Schumm,¹⁸ A. Seiden,¹⁸ P. Spradlin,¹⁸ D. C. Williams,¹⁸ M. G. Wilson,¹⁸ J. Albert,¹⁹ E. Chen,¹⁹
G. P. Dubois-Felsmann,¹⁹ A. Dvoretskii,¹⁹ D. G. Hitlin,¹⁹ I. Narsky,¹⁹ T. Piatenko,¹⁹ F. C. Porter,¹⁹ A. Ryd,¹⁹
A. Samuel,¹⁹ S. Yang,¹⁹ R. Andreassen,²⁰ S. Jayatilleke,²⁰ G. Mancinelli,²⁰ B. T. Meadows,²⁰ M. D. Sokoloff,²⁰
F. Blanc,²¹ P. Bloom,²¹ S. Chen,²¹ W. T. Ford,²¹ U. Nauenberg,²¹ A. Olivas,²¹ P. Rankin,²¹ W. O. Ruddick,²¹
J. G. Smith,²¹ K. A. Ulmer,²¹ J. Zhang,²¹ A. Chen,²² E. A. Eckhart,²² J. L. Harton,²² A. Soffer,²² W. H. Toki,²²
R. J. Wilson,²² Q. Zeng,²² B. Spaan,²³ D. Altenburg,²⁴ T. Brandt,²⁴ J. Brose,²⁴ M. Dickopp,²⁴ E. Feltresi,²⁴
A. Hauke,²⁴ V. Klose,²⁴ H. M. Lacker,²⁴ E. Maly,²⁴ R. Nogowski,²⁴ S. Otto,²⁴ A. Petzold,²⁴ G. Schott,²⁴
J. Schubert,²⁴ K. R. Schubert,²⁴ R. Schwierz,²⁴ J. E. Sundermann,²⁴ D. Bernard,²⁵ G. R. Bonneau,²⁵ P. Grenier,²⁵
S. Schrenk,²⁵ Ch. Thiebaut,²⁵ G. Vasileiadis,²⁵ M. Verderi,²⁵ D. J. Bard,²⁶ P. J. Clark,²⁶ W. Gradl,²⁶ F. Muheim,²⁶
S. Playfer,²⁶ Y. Xie,²⁶ M. Andreotti,²⁷ V. Azzolini,²⁷ D. Bettoni,²⁷ C. Bozzi,²⁷ R. Calabrese,²⁷ G. Cibinetto,²⁷
E. Luppi,²⁷ M. Negrini,²⁷ L. Piemontese,²⁷ A. Sarti,²⁷ F. Anulli,²⁸ R. Baldini-Ferroli,²⁸ A. Calcaterra,²⁸ R. de
Sangro,²⁸ G. Finocchiaro,²⁸ P. Patteri,²⁸ I. M. Peruzzi,²⁸ M. Piccolo,²⁸ A. Zallo,²⁸ A. Buzzo,²⁹ R. Capra,²⁹
R. Contri,²⁹ M. Lo Vetere,²⁹ M. Macri,²⁹ M. R. Monge,²⁹ S. Passaggio,²⁹ C. Patrignani,²⁹ E. Robutti,²⁹
A. Santroni,²⁹ S. Tosi,²⁹ S. Bailey,³⁰ G. Brandenburg,³⁰ K. S. Chaisanguanthum,³⁰ M. Morii,³⁰ E. Won,³⁰
R. S. Dubitzky,³¹ U. Langenegger,³¹ J. Marks,³¹ S. Schenk,³¹ U. Uwer,³¹ W. Bhimji,³² D. A. Bowerman,³²
P. D. Dauncey,³² U. Egede,³² J. R. Gaillard,³² G. W. Morton,³² J. A. Nash,³² M. B. Nikolic,³² G. P. Taylor,³²
X. Chai,³³ M. J. Charles,³³ G. J. Grenier,³³ U. Mallik,³³ A. K. Mohapatra,³³ V. Ziegler,³³ J. Cochran,³⁴
H. B. Crawley,³⁴ V. Eyges,³⁴ W. T. Meyer,³⁴ S. Prell,³⁴ E. I. Rosenberg,³⁴ A. E. Rubin,³⁴ J. Yi,³⁴ N. Arnaud,³⁵
M. Davier,³⁵ X. Giroux,³⁵ G. Grosdidier,³⁵ A. Höcker,³⁵ F. Le Diberder,³⁵ V. Lepeltier,³⁵ A. M. Lutz,³⁵
T. C. Petersen,³⁵ M. Pierini,³⁵ S. Plaszczynski,³⁵ S. Rodier,³⁵ P. Roudeau,³⁵ M. H. Schune,³⁵ A. Stocchi,³⁵
G. Wormser,³⁵ C. H. Cheng,³⁶ D. J. Lange,³⁶ M. C. Simani,³⁶ D. M. Wright,³⁶ A. J. Bevan,³⁷ C. A. Chavez,³⁷
J. P. Coleman,³⁷ I. J. Forster,³⁷ J. R. Fry,³⁷ E. Gabathuler,³⁷ R. Gamet,³⁷ K. A. George,³⁷ D. E. Hutchcroft,³⁷
R. J. Parry,³⁷ D. J. Payne,³⁷ C. Touramanis,³⁷ C. M. Cormack,³⁸ F. Di Lodovico,³⁸ C. L. Brown,³⁹ G. Cowan,³⁹
R. L. Flack,³⁹ H. U. Flaecher,³⁹ M. G. Green,³⁹ P. S. Jackson,³⁹ T. R. McMahon,³⁹ S. Ricciardi,³⁹ F. Salvatore,³⁹
D. Brown,⁴⁰ C. L. Davis,⁴⁰ J. Allison,⁴¹ N. R. Barlow,⁴¹ R. J. Barlow,⁴¹ M. C. Hodgkinson,⁴¹ G. D. Lafferty,⁴¹
M. T. Naisbit,⁴¹ J. C. Williams,⁴¹ C. Chen,⁴² A. Farbin,⁴² W. D. Hulsbergen,⁴² A. Jawahery,⁴² D. Kovalskyi,⁴²
C. K. Lae,⁴² V. Lillard,⁴² D. A. Roberts,⁴² G. Blaylock,⁴³ C. Dallapiccola,⁴³ S. S. Hertzbach,⁴³ R. Kofler,⁴³
V. B. Koptchev,⁴³ T. B. Moore,⁴³ S. Saremi,⁴³ H. Staengle,⁴³ S. Willocq,⁴³ R. Cowan,⁴⁴ K. Koeneke,⁴⁴ G. Sciolla,⁴⁴
S. J. Sekula,⁴⁴ F. Taylor,⁴⁴ R. K. Yamamoto,⁴⁴ H. Kim,⁴⁵ P. M. Patel,⁴⁵ S. H. Robertson,⁴⁵ A. Lazzaro,⁴⁶
V. Lombardo,⁴⁶ F. Palombo,⁴⁶ J. M. Bauer,⁴⁷ L. Cremaldi,⁴⁷ V. Eschenburg,⁴⁷ R. Godang,⁴⁷ R. Kroeger,⁴⁷
J. Reidy,⁴⁷ D. A. Sanders,⁴⁷ D. J. Summers,⁴⁷ H. W. Zhao,⁴⁷ S. Brunet,⁴⁸ D. Côté,⁴⁸ P. Taras,⁴⁸ B. Viaud,⁴⁸
H. Nicholson,⁴⁹ N. Cavallo,^{50,*} G. De Nardo,^{50,*} C. Gatto,⁵⁰ L. Lista,⁵⁰ D. Monorchio,⁵⁰

Submitted to Physical Review Letters

Work supported in part by the Department of Energy contract DE-AC02-76SF00515

SLAC, Stanford University, Stanford, CA 94309

P. Paolucci,⁵⁰ D. Piccolo,⁵⁰ C. Sciacca,⁵⁰ M. Baak,⁵¹ H. Bulten,⁵¹ G. Raven,⁵¹ H. L. Snoek,⁵¹ L. Wilden,⁵¹ C. P. Jessop,⁵² J. M. LoSecco,⁵² T. Allmendinger,⁵³ G. Benelli,⁵³ K. K. Gan,⁵³ K. Honscheid,⁵³ D. Hufnagel,⁵³ P. D. Jackson,⁵³ H. Kagan,⁵³ R. Kass,⁵³ T. Pulliam,⁵³ A. M. Rahimi,⁵³ R. Ter-Antonyan,⁵³ Q. K. Wong,⁵³ J. Brau,⁵⁴ R. Frey,⁵⁴ O. Igonkina,⁵⁴ M. Lu,⁵⁴ C. T. Potter,⁵⁴ N. B. Sinev,⁵⁴ D. Strom,⁵⁴ E. Torrence,⁵⁴ F. Colecchia,⁵⁵ A. Dorigo,⁵⁵ F. Galeazzi,⁵⁵ M. Margoni,⁵⁵ M. Morandin,⁵⁵ M. Posocco,⁵⁵ M. Rotondo,⁵⁵ F. Simonetto,⁵⁵ R. Stroili,⁵⁵ C. Voci,⁵⁵ M. Benayoun,⁵⁶ H. Briand,⁵⁶ J. Chauveau,⁵⁶ P. David,⁵⁶ L. Del Buono,⁵⁶ Ch. de la Vaissière,⁵⁶ O. Hamon,⁵⁶ M. J. J. John,⁵⁶ Ph. Leruste,⁵⁶ J. Malclès,⁵⁶ J. Ocariz,⁵⁶ L. Roos,⁵⁶ G. Therin,⁵⁶ P. K. Behera,⁵⁷ L. Gladney,⁵⁷ Q. H. Guo,⁵⁷ J. Panetta,⁵⁷ M. Biasini,⁵⁸ R. Covarelli,⁵⁸ M. Pioppi,⁵⁸ C. Angelini,⁵⁹ G. Batignani,⁵⁹ S. Bettarini,⁵⁹ F. Bucci,⁵⁹ G. Calderini,⁵⁹ M. Carpinelli,⁵⁹ F. Forti,⁵⁹ M. A. Giorgi,⁵⁹ A. Lusiani,⁵⁹ G. Marchiori,⁵⁹ M. Morganti,⁵⁹ N. Neri,⁵⁹ E. Paoloni,⁵⁹ M. Rama,⁵⁹ G. Rizzo,⁵⁹ G. Simi,⁵⁹ J. Walsh,⁵⁹ M. Haire,⁶⁰ D. Judd,⁶⁰ K. Paick,⁶⁰ D. E. Wagoner,⁶⁰ J. Biesiada,⁶¹ N. Danielson,⁶¹ P. Elmer,⁶¹ Y. P. Lau,⁶¹ C. Lu,⁶¹ J. Olsen,⁶¹ A. J. S. Smith,⁶¹ A. V. Telnov,⁶¹ F. Bellini,⁶² G. Cavoto,⁶² A. D'Orazio,⁶² E. Di Marco,⁶² R. Faccini,⁶² F. Ferrarotto,⁶² F. Ferroni,⁶² M. Gaspero,⁶² L. Li Gioi,⁶² M. A. Mazzoni,⁶² S. Morganti,⁶² G. Piredda,⁶² F. Polci,⁶² F. Safai Tehrani,⁶² C. Voena,⁶² S. Christ,⁶³ H. Schröder,⁶³ G. Wagner,⁶³ R. Waldi,⁶³ T. Adye,⁶⁴ N. De Groot,⁶⁴ B. Franek,⁶⁴ G. P. Gopal,⁶⁴ E. O. Olaiya,⁶⁴ F. F. Wilson,⁶⁴ R. Aleksan,⁶⁵ S. Emery,⁶⁵ A. Gaidot,⁶⁵ S. F. Ganzhur,⁶⁵ P.-F. Giraud,⁶⁵ G. Graziani,⁶⁵ G. Hamel de Monchenault,⁶⁵ W. Kozanecki,⁶⁵ M. Legendre,⁶⁵ G. W. London,⁶⁵ B. Mayer,⁶⁵ G. Vasseur,⁶⁵ Ch. Yèche,⁶⁵ M. Zito,⁶⁵ M. V. Purohit,⁶⁶ A. W. Weidemann,⁶⁶ J. R. Wilson,⁶⁶ F. X. Yumiceva,⁶⁶ T. Abe,⁶⁷ M. T. Allen,⁶⁷ D. Aston,⁶⁷ R. Bartoldus,⁶⁷ N. Berger,⁶⁷ A. M. Boyarski,⁶⁷ O. L. Buchmueller,⁶⁷ R. Claus,⁶⁷ M. R. Convery,⁶⁷ M. Cristinziani,⁶⁷ J. C. Dingfelder,⁶⁷ D. Dong,⁶⁷ J. Dorfan,⁶⁷ D. Dujmic,⁶⁷ W. Dunwoodie,⁶⁷ S. Fan,⁶⁷ R. C. Field,⁶⁷ T. Glanzman,⁶⁷ S. J. Gowdy,⁶⁷ T. Hadig,⁶⁷ V. Halyo,⁶⁷ C. Hast,⁶⁷ T. Hrynn'ova,⁶⁷ W. R. Innes,⁶⁷ M. H. Kelsey,⁶⁷ P. Kim,⁶⁷ M. L. Kocian,⁶⁷ D. W. G. S. Leith,⁶⁷ J. Libby,⁶⁷ S. Luitz,⁶⁷ V. Luth,⁶⁷ H. L. Lynch,⁶⁷ H. Marsiske,⁶⁷ R. Messner,⁶⁷ D. R. Muller,⁶⁷ C. P. O'Grady,⁶⁷ V. E. Ozcan,⁶⁷ A. Perazzo,⁶⁷ M. Perl,⁶⁷ B. N. Ratcliff,⁶⁷ A. Roodman,⁶⁷ A. A. Salnikov,⁶⁷ R. H. Schindler,⁶⁷ J. Schwiening,⁶⁷ A. Snyder,⁶⁷ A. Soha,⁶⁷ J. Stelzer,⁶⁷ J. Strube,^{54, 67} D. Su,⁶⁷ M. K. Sullivan,⁶⁷ K. Suzuki,⁶⁷ J. M. Thompson,⁶⁷ J. Va'vra,⁶⁷ S. R. Wagner,⁶⁷ M. Weaver,⁶⁷ W. J. Wisniewski,⁶⁷ M. Wittgen,⁶⁷ D. H. Wright,⁶⁷ A. K. Yarritu,⁶⁷ C. C. Young,⁶⁷ P. R. Burchat,⁶⁸ A. J. Edwards,⁶⁸ S. A. Majewski,⁶⁸ B. A. Petersen,⁶⁸ C. Roat,⁶⁸ M. Ahmed,⁶⁹ S. Ahmed,⁶⁹ M. S. Alam,⁶⁹ J. A. Ernst,⁶⁹ M. A. Saeed,⁶⁹ M. Saleem,⁶⁹ F. R. Wappler,⁶⁹ W. Bugg,⁷⁰ M. Krishnamurthy,⁷⁰ S. M. Spanier,⁷⁰ R. Eckmann,⁷¹ J. L. Ritchie,⁷¹ A. Satpathy,⁷¹ R. F. Schwitters,⁷¹ J. M. Izen,⁷² I. Kitayama,⁷² X. C. Lou,⁷² S. Ye,⁷² F. Bianchi,⁷³ M. Bona,⁷³ F. Gallo,⁷³ D. Gamba,⁷³ M. Bomben,⁷⁴ L. Bosisio,⁷⁴ C. Cartaro,⁷⁴ F. Cossutti,⁷⁴ G. Della Ricca,⁷⁴ S. Dittongo,⁷⁴ S. Grancagnolo,⁷⁴ L. Lanceri,⁷⁴ P. Poropat,^{74, †} L. Vitale,⁷⁴ G. Vuagnin,⁷⁴ F. Martinez-Vidal,⁷⁵ R. S. Panvini,^{76, †} Sw. Banerjee,⁷⁷ B. Bhuyan,⁷⁷ C. M. Brown,⁷⁷ D. Fortin,⁷⁷ K. Hamano,⁷⁷ R. Kowalewski,⁷⁷ J. M. Roney,⁷⁷ R. J. Sobie,⁷⁷ J. J. Back,⁷⁸ P. F. Harrison,⁷⁸ T. E. Latham,⁷⁸ G. B. Mohanty,⁷⁸ H. R. Band,⁷⁹ X. Chen,⁷⁹ B. Cheng,⁷⁹ S. Dasu,⁷⁹ M. Datta,⁷⁹ A. M. Eichenbaum,⁷⁹ K. T. Flood,⁷⁹ M. Graham,⁷⁹ J. J. Hollar,⁷⁹ J. R. Johnson,⁷⁹ P. E. Kutter,⁷⁹ H. Li,⁷⁹ R. Liu,⁷⁹ B. Mellado,⁷⁹ A. Mihalyi,⁷⁹ Y. Pan,⁷⁹ R. Prepost,⁷⁹ P. Tan,⁷⁹ J. H. von Wimmersperg-Toeller,⁷⁹ J. Wu,⁷⁹ S. L. Wu,⁷⁹ Z. Yu,⁷⁹ M. G. Greene,⁸⁰ and H. Neal⁸⁰

(The BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

²IFAE, Universitat Autònoma de Barcelona, E-08193 Bellaterra, Barcelona, Spain

³Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

⁴Institute of High Energy Physics, Beijing 100039, China

⁵University of Bergen, Inst. of Physics, N-5007 Bergen, Norway

⁶Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

⁷University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁸Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁹University of Bristol, Bristol BS8 1TL, United Kingdom

¹⁰University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

¹¹Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹²Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹³University of California at Irvine, Irvine, California 92697, USA

¹⁴University of California at Los Angeles, Los Angeles, California 90024, USA

¹⁵University of California at Riverside, Riverside, California 92521, USA

¹⁶University of California at San Diego, La Jolla, California 92093, USA

¹⁷University of California at Santa Barbara, Santa Barbara, California 93106, USA

¹⁸University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

- ¹⁹California Institute of Technology, Pasadena, California 91125, USA
²⁰University of Cincinnati, Cincinnati, Ohio 45221, USA
²¹University of Colorado, Boulder, Colorado 80309, USA
²²Colorado State University, Fort Collins, Colorado 80523, USA
²³Universität Dortmund, Institut für Physik, D-44221 Dortmund, Germany
²⁴Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany
²⁵Ecole Polytechnique, LLR, F-91128 Palaiseau, France
²⁶University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom
²⁷Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy
²⁸Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy
²⁹Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy
³⁰Harvard University, Cambridge, Massachusetts 02138, USA
³¹Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany
³²Imperial College London, London, SW7 2AZ, United Kingdom
³³University of Iowa, Iowa City, Iowa 52242, USA
³⁴Iowa State University, Ames, Iowa 50011-3160, USA
³⁵Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France
³⁶Lawrence Livermore National Laboratory, Livermore, California 94550, USA
³⁷University of Liverpool, Liverpool L69 7ZE, United Kingdom
³⁸Queen Mary, University of London, E1 4NS, United Kingdom
³⁹University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom
⁴⁰University of Louisville, Louisville, Kentucky 40292, USA
⁴¹University of Manchester, Manchester M13 9PL, United Kingdom
⁴²University of Maryland, College Park, Maryland 20742, USA
⁴³University of Massachusetts, Amherst, Massachusetts 01003, USA
⁴⁴Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA
⁴⁵McGill University, Montréal, Québec, Canada H3A 2T8
⁴⁶Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy
⁴⁷University of Mississippi, University, Mississippi 38677, USA
⁴⁸Université de Montréal, Laboratoire René J. A. Lévesque, Montréal, Québec, Canada H3C 3J7
⁴⁹Mount Holyoke College, South Hadley, Massachusetts 01075, USA
⁵⁰Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy
⁵¹NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands
⁵²University of Notre Dame, Notre Dame, Indiana 46556, USA
⁵³Ohio State University, Columbus, Ohio 43210, USA
⁵⁴University of Oregon, Eugene, Oregon 97403, USA
⁵⁵Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy
⁵⁶Universités Paris VI et VII, Laboratoire de Physique Nucléaire et de Hautes Energies, F-75252 Paris, France
⁵⁷University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
⁵⁸Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy
⁵⁹Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy
⁶⁰Prairie View A&M University, Prairie View, Texas 77446, USA
⁶¹Princeton University, Princeton, New Jersey 08544, USA
⁶²Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy
⁶³Universität Rostock, D-18051 Rostock, Germany
⁶⁴Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom
⁶⁵DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France
⁶⁶University of South Carolina, Columbia, South Carolina 29208, USA
⁶⁷Stanford Linear Accelerator Center, Stanford, California 94309, USA
⁶⁸Stanford University, Stanford, California 94305-4060, USA
⁶⁹State University of New York, Albany, New York 12222, USA
⁷⁰University of Tennessee, Knoxville, Tennessee 37996, USA
⁷¹University of Texas at Austin, Austin, Texas 78712, USA
⁷²University of Texas at Dallas, Richardson, Texas 75083, USA
⁷³Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy
⁷⁴Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy
⁷⁵IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain
⁷⁶Vanderbilt University, Nashville, Tennessee 37235, USA
⁷⁷University of Victoria, Victoria, British Columbia, Canada V8W 3P6
⁷⁸Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom
⁷⁹University of Wisconsin, Madison, Wisconsin 53706, USA
⁸⁰Yale University, New Haven, Connecticut 06511, USA

(Dated: April 8, 2005)

Using 116.1 fb^{-1} of data collected by the *BABAR* detector, we present an analysis of Ξ_c^0 production in B decays and from the $c\bar{c}$ continuum, with the Ξ_c^0 decaying into $\Omega^- K^+$ and $\Xi^- \pi^+$ final states. We measure the ratio of branching fractions $\mathcal{B}(\Xi_c^0 \rightarrow \Omega^- K^+)/\mathcal{B}(\Xi_c^0 \rightarrow \Xi^- \pi^+)$ to be $0.294 \pm 0.018 \pm 0.016$, where the first uncertainty is statistical and the second is systematic. The Ξ_c^0 momentum spectrum is measured on and 40 MeV below the $\Upsilon(4S)$ resonance. From these spectra the branching fraction product $\mathcal{B}(B \rightarrow \Xi_c^0 X) \times \mathcal{B}(\Xi_c^0 \rightarrow \Xi^- \pi^+)$ is measured to be $(2.11 \pm 0.19 \pm 0.25) \times 10^{-4}$, and the cross-section product $\sigma(e^+ e^- \rightarrow \Xi_c^0 X) \times \mathcal{B}(\Xi_c^0 \rightarrow \Xi^- \pi^+)$ from the continuum is measured to be $(388 \pm 39 \pm 41) \text{ fb}$ at a center-of-mass energy of 10.58 GeV.

PACS numbers: 14.20.Lq, 14.40.Nd

Considerable progress in the charmed baryon sector [1] has been made over the past two decades with the advent of high-luminosity colliders and flavor factory experiments such as CLEO, FOCUS and SELEX. Today the high-luminosity B -factory experiments BELLE and *BABAR* present excellent opportunities to build upon this knowledge and study the production and decay of charmed baryons with high precision.

In this Letter we present a study of the $\Xi_c^0(csd)$ [2] charmed baryon through two decay modes: $\Xi_c^0 \rightarrow \Omega^- K^+$ and $\Xi_c^0 \rightarrow \Xi^- \pi^+$. We measure the ratio of branching fractions of these decay modes, which has been predicted to be 0.32 with a spectator quark model calculation [3]. A previous measurement was consistent with this prediction but had a large (40%) statistical uncertainty [4].

We also study Ξ_c^0 production by measuring the spectrum of the Ξ_c^0 momentum in the $e^+ e^-$ center-of-mass frame (p^*). A number of theoretical predictions of the rate of Ξ_c production in B decays have been made [5–8]. Insight into the contributing processes can be gained by studying the shape of the p^* spectrum. Evidence for Ξ_c^0 production in B decays was presented previously by the CLEO collaboration, with a statistical significance of $\sim 3\sigma$ in the $\Xi_c^0 \rightarrow \Xi^- \pi^+$ decay mode and $\sim 4\sigma$ in the $\Xi_c^+ \rightarrow \Xi^- \pi^+ \pi^+$ decay mode [9].

The data for this analysis were collected with the *BABAR* detector at the SLAC PEP-II asymmetric energy $e^+ e^-$ collider; the detector is described in detail elsewhere [10]. A total integrated luminosity of 116.1 fb^{-1} is used, of which 105.4 fb^{-1} was collected at the $\Upsilon(4S)$ resonance [1] (corresponding to 116 million $B\bar{B}$ pairs) and 10.7 fb^{-1} was collected at a center-of-mass energy of 10.54 GeV, which is below the $B\bar{B}$ production threshold. These are referred to as the on-resonance and off-resonance data samples, respectively.

The reconstruction of Ξ_c^0 candidates takes place as follows. A Λ candidate is reconstructed by identifying a proton and combining it with an oppositely charged track interpreted as a π^- , fitting the tracks to a common vertex. The Λ candidate is then combined with a negatively charged track interpreted as a $\pi^- (K^-)$ to form a $\Xi^- (\Omega^-)$ candidate. For each intermediate hyperon, the invariant mass is required to be within 3σ of the central value, where σ is the fitted mass resolution. The invariant mass is then constrained to the nominal value [1].

Each resulting $\Xi^- (\Omega^-)$ candidate passing the selection criteria is then combined with a positively charged track interpreted as a $\pi^+ (K^+)$ to form a Ξ_c^0 candidate. For the $\Omega^- K^+$ final state, the two K^\pm tracks must be identified as kaons. Particle identification is performed with dE/dx and Cherenkov angle measurements [10].

Additional selection criteria, described below, are used to improve the signal-to-background ratio. As a precaution against selection bias, these are optimized with sub-samples of the data: 20 fb^{-1} and 40 fb^{-1} for the $\Xi^- \pi^+$ and $\Omega^- K^+$ final states, respectively. A minimum decay distance of 2.5 mm (1.5 mm) between the event primary vertex and the $\Xi^- (\Omega^-)$ decay vertex in the plane perpendicular to the beam direction is required. The distance between the Ω^- and Λ decay vertices is required to be at least 3 mm. In addition, the relative positioning of vertices is required to be causally connected: we reject candidates in which the Ξ^- decays further from the primary vertex than its daughter Λ does, or where the displacement vector from the Ω^- decay point to the Λ decay point is anti-parallel to the Λ momentum vector. The invariant mass distributions for the Ξ_c^0 candidates satisfying these criteria are shown in Fig. 1 (a) and (b) for $\Xi^- \pi^+$ and $\Omega^- K^+$ combinations, respectively.

Simulated events with the Ξ_c^0 decaying into the two desired final states are generated for the processes $e^+ e^- \rightarrow c\bar{c} \rightarrow \Xi_c^0 X$ and $e^+ e^- \rightarrow \Upsilon(4S) \rightarrow B\bar{B} \rightarrow \Xi_c^0 X$, where X represents the rest of the event. The PYTHIA simulation package [11] is used for the $c\bar{c}$ fragmentation and for B decays to Ξ_c^0 , and GEANT4 [12] is used to simulate the detector response. For $c\bar{c}$ production, samples of 90,000 events for the $\Xi^- \pi^+$ final state and 60,000 for the $\Omega^- K^+$ final state are generated. For $B\bar{B}$ production, samples of 255,000 and 120,000 events are used, respectively.

Additional generic Monte Carlo events are used to investigate possible background contributions. The sample sizes are equivalent to 245 fb^{-1} , 64 fb^{-1} , and 33 fb^{-1} for $e^+ e^- \rightarrow B\bar{B}$, $c\bar{c}$, and $q\bar{q}$, respectively, where $q = u, d, s$. Excluding signal contributions, the mass distribution varies smoothly throughout the region near the Ξ_c^0 mass.

To measure the ratio of branching fractions, a further requirement that $p^* > 1.8 \text{ GeV}/c$ is imposed on the Ξ_c^0 candidates, improving the signal purity. Additionally, the candidates are required to be within the region of high Ξ_c^0 reconstruction efficiency $-0.8 \leq \cos \theta^* \leq 0.6$,




FIG. 1: Invariant mass distributions for Ξ_c^0 candidates in 116.1 fb^{-1} of data, for (a) $\Xi^- \pi^+$, and (b) $\Omega^- K^+$.

where θ^* is the polar angle of the Ξ_c^0 candidate with respect to the collision axis in the center-of-mass frame. The efficiency is calculated from signal Monte Carlo events as a function of p^* and $\cos \theta^*$ for each of the decay modes. For each mode, a fifteen-parameter fit gives a smooth parameterization of the efficiency with small statistical uncertainty. The efficiency is then corrected by weighting each candidate by the inverse of its efficiency, and the efficiency-corrected mass spectrum is fitted with a double Gaussian for signal plus a linear background function. Including efficiency loss due to the Ω^- and Λ branching fractions, we obtain 25889 ± 516 events in the $\Xi^- \pi^+$ mode and 7615 ± 443 events in the $\Omega^- K^+$ mode.

We evaluate several sources of systematic uncertainty in the ratio of branching fractions: the fits to the mass spectra (3.4%), the efficiency parameterization (3.1%), particle identification (2.0%), finite Monte Carlo statistics (1.4%), multiple candidates in the same event (1.0%), charge asymmetries in detection efficiency (1.0%), the $\cos \theta^*$ distribution (1.0%), and the Ω^- branching fraction (1.0%). No baryon polarization is considered and any systematic uncertainty due to this is neglected. Adding all of the uncertainties in quadrature, we obtain:

$$\frac{\mathcal{B}(\Xi_c^0 \rightarrow \Omega^- K^+)}{\mathcal{B}(\Xi_c^0 \rightarrow \Xi^- \pi^+)} = 0.294 \pm 0.018 \pm 0.016.$$

Throughout this Letter, the first uncertainty is statistical and the second is systematic.

After obtaining the ratio of branching fractions, we next measure the p^* spectrum of the Ξ_c^0 baryons in order

to study the production mechanisms in both $c\bar{c}$ and $B\bar{B}$ events. The same selection criteria and data samples described above are used, except that no requirement on p^* or $\cos \theta^*$ is made. Instead, the Ξ_c^0 candidates are divided into intervals of p^* . The yield is then measured in each interval with two different methods: first with a fitting method, where the mass spectrum is fitted with a single Gaussian for signal plus a linear background function and the integral of the Gaussian is taken as the yield; second with a counting method, where the background is estimated from mass sidebands and the signal yield is then taken as the statistical excess above this background in a mass window around the peak. The use of two different methods serves as a cross-check.

The efficiency in each p^* interval is estimated with signal Monte Carlo events from that p^* range. For both methods, the simulated events are reconstructed and the yield is measured, then divided by the number of events generated to obtain the efficiency. Due to the different angular distributions, the efficiencies for Ξ_c^0 produced from $c\bar{c}$ ($\varepsilon_{c\bar{c}}$) and from $B\bar{B}$ ($\varepsilon_{B\bar{B}}$) differ slightly. In the region $1.2 < p^* < 2.0 \text{ GeV}/c$ where both production mechanisms are significant and the difference is approximately 8% (relative), the efficiency is taken to be $(\varepsilon_{c\bar{c}} + \varepsilon_{B\bar{B}})/2$. The systematic uncertainty on the efficiency is then $|\varepsilon_{c\bar{c}} - \varepsilon_{B\bar{B}}|/\sqrt{12}$. The angular distributions produced in PYTHIA fragmentation are assumed to be correct when calculating the efficiency; the data are fully consistent with these distributions within available statistics. The efficiency-corrected yield in each p^* interval is then calculated, including loss of efficiency due to the Λ and Ω^- branching fractions. The spectra obtained with the two methods are in good agreement; we use the counting method for the quoted results since it is more stable for low statistics.

A number of systematic uncertainties are considered, the most important of which are the uncertainties associated with the track-finding and particle identification efficiencies (5.8% and 3.5%, respectively). Uncertainties from the simulated Ξ_c^0 mass resolution (1%), the mass resolutions of the intermediate hyperon states (0.5%), the p^* resolution ($\mathcal{O}(1\%)$), the effect of finite interval width ($\mathcal{O}(2\%)$), multiple candidates (0%), non-linearity of the background ($\mathcal{O}(1\%)$), the signal measurement method used (2%), the finite Monte Carlo statistics available ($\mathcal{O}(3\%)$), and uncertainties in the Λ and Ω^- branching fractions (0.8%, 1.0%) are all considered individually; the notation $\mathcal{O}(x\%)$ indicates the typical value when the exact uncertainty varies among p^* intervals. The total systematic uncertainty for each p^* interval is obtained by adding the individual contributions in quadrature. In addition, a systematic correction of 1.0% is applied to account for a known data-Monte Carlo discrepancy in the track-finding efficiency, and small corrections are applied to each interval to account for the broadening effect of the p^* experimental resolution on the spectrum. The fi-




FIG. 2: The p^* spectrum measurements. In (a), the p^* spectrum of Ξ_c^0 decaying via $\Xi^- \pi^+$ is shown for the on-resonance data sample. In (b), the Ξ_c^0 spectra obtained with the $\Xi^- \pi^+$ and $\Omega^- K^+$ modes are compared after scaling the $\Omega^- K^+$ normalization by the ratio of branching fractions presented in this Letter. In (c), the on-resonance and off-resonance data samples are shown together, with the off-resonance normalization scaled to account for the difference in integrated luminosity and cross-section. In each plot, the inner error bars give the statistical uncertainty and the outer error bars give the sum in quadrature of the statistical and systematic uncertainties. The vertical line at 2.15 GeV/ c in (c) shows the kinematic cutoff for Ξ_c^0 produced in B decays at *BABAR*. Note that the vertical axes show events per unit p^* , not events per p^* interval as given in Table I.

nal p^* spectrum for the on-resonance data set, obtained with the counting method in the $\Xi^- \pi^+$ mode, is shown in Fig. 2(a). Table I shows the corresponding values.

A further check is performed by comparing the two decay modes. The $\Omega^- K^+$ yields are scaled by a factor of $(1/0.294)$, the ratio of branching fractions previously presented in this Letter. The spectra are shown in Fig. 2(b), with yields measured with the counting method. Because the $\Omega^- K^+$ signal has fewer events, wider p^* intervals are used. The spectra of the two modes show good agreement in both shape and normalization; this serves as a cross-check both of the p^* spectrum measurement and of

TABLE I: Efficiency-corrected yield and cross-section product including B production $\sigma(e^+e^- \rightarrow \Xi_c^0 X) \times \mathcal{B}(\Xi_c^0 \rightarrow \Xi^- \pi^+)$.

p^* range (GeV/ c)	Corrected yield	Cross-section product (fb)
0.0–0.2	$1046 \pm 201 \pm 128$	$10 \pm 2 \pm 1$
0.2–0.4	$5889 \pm 446 \pm 483$	$56 \pm 4 \pm 5$
0.4–0.6	$10681 \pm 631 \pm 801$	$101 \pm 6 \pm 8$
0.6–0.8	$10709 \pm 660 \pm 817$	$102 \pm 6 \pm 8$
0.8–1.0	$8811 \pm 647 \pm 679$	$84 \pm 6 \pm 7$
1.0–1.2	$6834 \pm 573 \pm 530$	$65 \pm 5 \pm 5$
1.2–1.4	$2954 \pm 501 \pm 252$	$28 \pm 5 \pm 2$
1.4–1.6	$2429 \pm 470 \pm 212$	$23 \pm 4 \pm 2$
1.6–1.8	$2252 \pm 424 \pm 202$	$21 \pm 4 \pm 2$
1.8–2.0	$2159 \pm 350 \pm 217$	$20 \pm 3 \pm 2$
2.0–2.2	$2375 \pm 347 \pm 205$	$23 \pm 3 \pm 2$
2.2–2.4	$2743 \pm 340 \pm 227$	$26 \pm 3 \pm 2$
2.4–2.6	$3537 \pm 315 \pm 285$	$34 \pm 3 \pm 3$
2.6–2.8	$3920 \pm 282 \pm 306$	$37 \pm 3 \pm 3$
2.8–3.0	$4595 \pm 294 \pm 359$	$44 \pm 3 \pm 3$
3.0–3.2	$4873 \pm 263 \pm 401$	$46 \pm 2 \pm 4$
3.2–3.4	$4442 \pm 244 \pm 348$	$42 \pm 2 \pm 3$
3.4–3.6	$4084 \pm 223 \pm 355$	$39 \pm 2 \pm 3$
3.6–3.8	$2282 \pm 171 \pm 189$	$22 \pm 2 \pm 2$
3.8–4.0	$2095 \pm 155 \pm 166$	$20 \pm 1 \pm 2$
4.0–4.2	$1168 \pm 123 \pm 177$	$11 \pm 1 \pm 2$
4.2–4.4	$233 \pm 53 \pm 32$	$2.2 \pm 0.5 \pm 0.3$
4.4–4.6	$88 \pm 37 \pm 21$	$0.8 \pm 0.3 \pm 0.2$
4.6–4.8	$5 \pm 13 \pm 7$	$0.0 \pm 0.1 \pm 0.1$
4.8–5.0	$24 \pm 17 \pm 16$	$0.2 \pm 0.2 \pm 0.1$

the ratio of branching fractions.

The double-peak structure seen in the p^* spectrum is due to two production mechanisms: the peak at lower p^* is due to Ξ_c^0 production in B meson decays and the peak at higher p^* is due to Ξ_c^0 production from the $c\bar{c}$ continuum. This is evident from Fig. 2(c), where the p^* spectra for the on-resonance and off-resonance data are shown separately (with the off-resonance spectrum scaled to the on-resonance integrated luminosity and corrected for the change in $c\bar{c}$ cross-section). The $c\bar{c}$ peak is present in both samples, but the $B\bar{B}$ peak is only present in the on-resonance sample. Assuming baryon number conservation, the kinematic limit for Ξ_c^0 produced in the decays of B mesons at *BABAR* is $p^* = 2.135$ GeV/ c . We compare the on-resonance and scaled off-resonance samples for $p^* \leq 2.15$ GeV/ c to obtain the yield of Ξ_c^0 produced in B decays. This is scaled by the number of B mesons in the data sample (introducing a further 1.1% systematic

uncertainty) to obtain:

$$\begin{aligned}\mathcal{B}(B \rightarrow \Xi_c^0 X) \times \mathcal{B}(\Xi_c^0 \rightarrow \Xi^- \pi^+) \\ = (2.11 \pm 0.19 \pm 0.25) \times 10^{-4}.\end{aligned}$$

The yield of Ξ_c^0 produced in $c\bar{c}$ events at an energy of 10.58 GeV is calculated from the scaled off-resonance data set (for $p^* \leq 2.15$ GeV/c) and the on-resonance data set (for $p^* > 2.15$ GeV/c). The yield is then divided by the integrated luminosity (introducing a further 1.5% systematic uncertainty) to obtain the cross-section from the continuum:

$$\begin{aligned}\sigma(e^+ e^- \rightarrow \Xi_c^0 X) \times \mathcal{B}(\Xi_c^0 \rightarrow \Xi^- \pi^+) \\ = (388 \pm 39 \pm 41) \text{ fb},\end{aligned}$$

where both Ξ_c^0 and Ξ_c^0 are included in the cross-section. The effect of initial state radiation is not isolated.

In summary, we have studied the Ξ_c^0 charmed baryon at *BABAR* through its decays to the $\Omega^- K^+$ and $\Xi^- \pi^+$ final states using 116.1 fb^{-1} of data. The ratio of branching fractions of these decay modes was measured to be $0.294 \pm 0.018 \pm 0.016$. This represents a substantial improvement on the previous measurement [4] and is consistent with a spectator quark model prediction [3]. We have also measured the p^* spectrum for Ξ_c^0 produced at the $\Upsilon(4S)$ resonance. The high rate of Ξ_c^0 production at low p^* in B decays (below 1.2 GeV/c) is particularly intriguing, implying that the invariant mass of the recoiling antibaryon system is typically above 2.0 GeV/c². This can be explained naturally by a substantial rate of charmed baryon pair production through the $b \rightarrow c\bar{c}s$ weak decay process [5–8] which was observed indirectly in a previous *BABAR* analysis [13]. In this Letter we measured the branching fraction product $\mathcal{B}(B \rightarrow \Xi_c^0 X) \times \mathcal{B}(\Xi_c^0 \rightarrow \Xi^- \pi^+)$ to be $(2.11 \pm 0.19 \pm 0.25) \times 10^{-4}$; the precision is significantly improved over the previous measurement [9]. We have also measured the cross-section product $\sigma(e^+ e^- \rightarrow \Xi_c^0 X) \times \mathcal{B}(\Xi_c^0 \rightarrow \Xi^- \pi^+)$ from the continuum to be $(388 \pm 39 \pm 41)$ fb.

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from CONACyT (Mexico), A. P. Sloan Foundation, Research Corporation, and Alexander von Humboldt Foundation.

* Also with Università della Basilicata, Potenza, Italy

† Deceased

- [1] Particle Data Group, S. Eidelman *et al.*, Phys. Lett. B **592**, 1 (2004).
- [2] Charge conjugate reactions are implied throughout.
- [3] J. G. Körner and M. Krämer, Z. Phys. C **55**, 659 (1992).
- [4] CLEO Collaboration, S. Henderson *et al.*, Phys. Lett. B **283**, 161 (1992).
- [5] P. Ball and H. G. Dosch, Z. Phys. C **51**, 445 (1991).
- [6] V. L. Chernyak and I. R. Zhitnitsky, Nucl. Phys. **B345**, 137 (1990).
- [7] S. M. Sheikholeslami and M. P. Khanna, Phys. Rev. D **44**, 770 (1991).
- [8] I. Dunietz, P. S. Cooper, A. F. Falk and M. B. Wise, Phys. Rev. Lett. **73**, 1075 (1994).
- [9] CLEO Collaboration, B. Barish *et al.*, Phys. Rev. Lett. **79**, 3599 (1997).
- [10] *BABAR* Collaboration, B. Aubert *et al.*, Nucl. Instrum. Methods A **479**, 1 (2002).
- [11] T. Sjostrand *et al.* Comput. Phys. Commun. **135**, 238 (2001).
- [12] S. Agostinelli *et al.*, Nucl. Instrum. Methods A **506**, 250 (2003).
- [13] B. Aubert *et al.*, Phys. Rev. D **70**, 091106 (2004).