

Light-Front Dynamics and AdS/QCD: The Pion Form Factor in the Space- and Time-Like Regions

Stanley J. Brodsky¹ and Guy F. de Téramond^{2,1}

¹*Stanford Linear Accelerator Center, Stanford University, Stanford, California 94309, USA*

²*Universidad de Costa Rica, San José, Costa Rica*

(Dated: July 26, 2007)

The AdS/CFT correspondence between string theory in AdS space and conformal field theories in physical space-time leads to an analytic, semi-classical model for strongly-coupled QCD which has scale invariance and dimensional counting at short distances and color confinement at large distances. The AdS/CFT correspondence also provides insights into the inherently non-perturbative aspects of QCD such as the orbital and radial spectra of hadrons and the form of hadronic wavefunctions. In particular, we show that there is an exact correspondence between the fifth-dimensional coordinate of AdS space z and a specific impact variable ζ which measures the separation of the quark and gluonic constituents within the hadron in ordinary space-time. This connection allows one to compute the analytic form of the frame-independent light-front wavefunctions of mesons and baryons, the fundamental entities which encode hadron properties and which allow the computation of decay constants, form factors and other exclusive scattering amplitudes. Relativistic light-front equations in ordinary space-time are found which reproduce the results obtained using the fifth-dimensional theory. As specific examples we compute the pion coupling constant f_π and examine the propagation of the electromagnetic current in AdS space, which determines the space and time-like behavior of the pion form factor and the value of M_ρ .

PACS numbers: 11.15.Tk, 11.25.Tq, 12.38Aw, 12.40Yx

I. INTRODUCTION

Quantum Chromodynamics, the Yang-Mills local gauge field theory of $SU(3)_C$ color symmetry provides a fundamental description of hadron and nuclear physics in terms of quark and gluon degrees of freedom. However, because of its strong coupling nature, it is difficult to find analytic solutions to QCD or to make precise predictions outside of its perturbative domain. An important goal is thus to find an initial approximation to QCD which is both analytically tractable and which can be systematically improved. For example, in quantum electrodynamics, the Coulombic Schrödinger and Dirac equations provide quite accurate first approximations to atomic bound state problems, which can then be systematically improved using the Bethe-Salpeter formalism and correcting for quantum fluctuations, such as the Lamb Shift and vacuum polarization.

One of the most significant theoretical advances in recent years has been the application of the AdS/CFT correspondence [1] between string states defined on the 5-dimensional Anti-de Sitter (AdS) space-time and conformal field theories in physical space-time. The essential principle underlying the AdS/CFT approach to conformal gauge theories is the isomorphism of the group of Poincaré and conformal transformations $SO(4,2)$ to the group of isometries of Anti-de Sitter space. The AdS metric is

$$ds^2 = \frac{R^2}{z^2} (\eta_{\mu\nu} dx^\mu dx^\nu - dz^2), \quad (1)$$

which is invariant under scale changes of the coordinate in the fifth dimension $z \rightarrow \lambda z$ and $x_\mu \rightarrow \lambda x_\mu$. Thus one

can match scale transformations of the theory in $3+1$ physical space-time to scale transformations in the fifth dimension z .

QCD is not itself a conformal theory; however as we shall discuss, there are indications, both from theory and phenomenology, that the QCD coupling is slowly varying at small momentum transfer. Thus in the domain where the QCD coupling is approximately constant and quark masses can be neglected, QCD resembles a conformal theory. As shown by Polchinski and Strassler [2], one can simulate confinement by imposing boundary conditions in the holographic variable at $z = z_0 = 1/\Lambda_{\text{QCD}}$. Confinement can also be introduced by modifying the AdS metric to mimic a confining potential. The resulting models, although *ad hoc*, provide a simple semi-classical approximation to QCD which has both constituent counting rule behavior at short distances and confinement at large distances. This simple approach, which has been described as a “bottom-up” approach, has been successful in obtaining general properties of scattering amplitudes of hadronic bound states at strong coupling [2–5], the low-lying hadron spectra [6–15], hadron couplings and chiral symmetry breaking [9, 16–18], quark potentials in confining backgrounds [19, 20] and a description of weak hadron decays [21]. Geometry back-reaction in AdS may also be relevant to the infrared physics [22]. The gauge theory/gravity duality also provides a convenient framework for the description of deep inelastic scattering structure functions at small x [23], a unified description of hard and soft pomeron physics [24] and gluon scattering amplitudes at strong coupling [25]. Recent applications to describe chiral symmetry breaking [26] and other meson

and baryon properties, have also been carried out within the framework of a top-bottom approach to AdS/CFT using higher dimensional branes [27–31].

The AdS/QCD correspondence is particularly relevant for the description of hadronic form factors, since it incorporates the connection between the twist of the hadron to the fall-off of its current matrix elements, as well as essential aspects of vector meson dominance. It also provides a convenient framework for analytically continuing the space-like results to the time-like region. Recent applications to the form factors of mesons and nucleons [11, 32–37] have followed from the original work described in [23, 38]

A physical hadron in four-dimensional Minkowski space has four-momentum P_μ and invariant mass given by $P_\mu P^\mu = \mathcal{M}^2$. The physical states in AdS₅ space are represented by normalizable “string” modes

$$\Phi(x, z) \sim e^{-iP \cdot x} \Phi^P(z), \quad (2)$$

with plane waves along the Poincaré coordinates and a profile function $\Phi^P(z)$ along the holographic coordinate z . For small- z , Φ scales as $\Phi \sim z^\Delta$, where Δ is the conformal dimension of the string state, the same dimension of the interpolating operator \mathcal{O} which creates a hadron out of the vacuum [2], $\langle P|\mathcal{O}|0\rangle \neq 0$. The scale dependence of each string mode $\Phi(x, z)$ is thus determined by matching its behavior at $z \rightarrow 0$ with the dimension of the corresponding hadronic state at short distances $x^2 \rightarrow 0$. Changes in length scale are mapped to evolution in the holographic variable z . The string mode in z thus represents the extension of the hadron wave function into the fifth dimension. There are also non-normalizable modes which are related to external currents: they propagate into the interior of AdS space.

As we shall discuss, there is a remarkable mapping between the AdS description of hadrons and the Hamiltonian formulation of QCD in physical space-time quantized on the light front. The light-front wavefunctions of bound states in QCD are relativistic and frame-independent generalizations of the familiar Schrödinger wavefunctions of atomic physics, but they are determined at fixed light-cone time $\tau = t + z/c$ —the “front form” advocated by Dirac—rather than at fixed ordinary time t . An important feature of light-front quantization is the fact that it provides exact formulas to write down matrix elements as a sum of bilinear forms, which can be mapped into their AdS/CFT counterparts in the semi-classical approximation. One can thus obtain not only an accurate description of the hadron spectrum for light quarks, but also a remarkably simple but realistic model of the valence wavefunctions of mesons, baryons, and glueballs. In fact, we find an exact correspondence between the fifth-dimensional coordinate of anti-de Sitter space z and a specific impact variable ζ in the light-front formalism which measures the separation of the constituents within the hadron in ordinary space-time. Since $\tau = 0$ in the front form, it corresponds to the transverse separation $x^\mu x_\mu = -\mathbf{x}_\perp^2$. The amplitude $\Phi(z)$ describing the

hadronic state in AdS₅ can then be precisely mapped to the light-front wavefunctions $\psi_{n/h}$ of hadrons in physical space-time [39], thus providing a relativistic description of hadrons in QCD at the amplitude level.

This paper is organized as follows. We first review in section II recent evidence which indicates that QCD has near-conformal behavior in the infrared region. The QCD light-front Fock representation is then briefly reviewed in section III. In section IV, we derive the exact form of current matrix elements in the light-front formalism which sets the framework for the identification with the corresponding amplitudes in AdS/CFT, and constitutes the basis of the partonic interpretation of the AdS/CFT correspondence. The actual mapping to AdS matrix elements is carried out in section V for the hard wall model and in section VI for the soft wall model, where the truncated space boundary conditions are replaced by a smooth cutoff. Results for the pion form factor in the space and time-like regions are presented in section VII for both models. In section VIII we compute the pion decay constant in light-front QCD and find its precise expression in AdS, following the mapping discussed in sections V and VI. Some final remarks are given in the conclusions in section IX. Other technical aspects useful for the discussion of the article are given in the appendices. In particular we relate our results for the soft wall model to the analytical results found recently by Grigoryan and Radyushkin [36].

II. CONFORMAL QCD WINDOW

It was originally believed that the AdS/CFT mathematical correspondence could only be applied to strictly conformal theories, such as $\mathcal{N} = 4$ supersymmetric Yang-Mills gauge theory. In our approach, we will apply the gauge theory/gravity duality to the strong coupling regime of QCD where the coupling appears to be approximately constant. Theoretical studies based on Dyson-Schwinger equation [40–43] and phenomenological [44, 45] evidence is in fact accumulating that the QCD couplings defined from physical observables such as τ decay [46] become constant at small virtuality; *i.e.*, effective charges develop an infrared fixed point in contradiction to the usual assumption of singular growth in the infrared. Recent lattice gauge theory simulations [47] also indicate an infrared fixed point for QCD. The near-constant behavior of effective couplings thus suggests that QCD can be approximated as a conformal theory at relatively small momentum transfer.

It is clear from a physical perspective that in a confining theory where gluons and quarks have an effective mass or maximal wavelength, all vacuum polarization corrections to the gluon self-energy must decouple at long wavelength; thus an infrared fixed point appears to be a natural consequence of confinement. Furthermore, if one considers a semi-classical approximation to QCD with massless quarks and without particle creation or absorp-

tion, then the resulting β function is zero, the coupling is constant, and the approximate theory is scale and conformal invariant.

In the case of hard exclusive reactions [48], the virtuality of the gluons exchanged in the underlying QCD process is typically much less than the momentum transfer scale Q , since typically several gluons share the total momentum transfer. Since the coupling is probed in the conformal window, this kinematic feature can explain why the measured proton Dirac form factor scales as $Q^4 F_1(Q^2) \simeq \text{const}$ up to $Q^2 < 35 \text{ GeV}^2$ [49] with little sign of the logarithmic running of the QCD coupling.

Thus conformal symmetry can be a useful first approximant even for physical QCD. Conformal symmetry is broken in physical QCD by quantum effects and quark masses; nevertheless, one can use conformal symmetry as a *template*, systematically correcting for its nonzero β function as well as higher-twist effects. For example, “commensurate scale relations” [50] which relate QCD observables to each other, such as the generalized Crewther relation [51], have no renormalization scale or scheme ambiguity and retain a convergent perturbative structure which reflects the underlying conformal symmetry of the classical theory. In general, the scale is set such that one has the correct analytic behavior at the heavy particle thresholds [52]. Analytic effective charges [53] such as the pinch scheme [54, 55] also provide an important perspective for unifying the electroweak and strong couplings [56].

III. THE LIGHT-FRONT FOCK REPRESENTATION

The light-front expansion of any hadronic system is constructed by quantizing quantum chromodynamics at fixed light-cone time [57] $\tau = t + z/c$. In terms of the hadron four-momentum $P = (P^+, P^-, \mathbf{P}_\perp)$, $P^\pm = P^0 \pm P^3$, the light-cone Lorentz invariant Hamiltonian for the composite system, $H_{LC}^{QCD} = P^- P^+ - \mathbf{P}_\perp^2$, has eigenvalues given in terms of the eigenmass \mathcal{M} squared corresponding to the mass spectrum of the color-singlet states in QCD [58]. The momentum generators P^+ and \mathbf{P}_\perp are kinematical; *i.e.*, they are independent of the interactions. The LC time evolution operator $P^- = i \frac{d}{d\tau}$ can be derived directly from the QCD Lagrangian in the light-cone gauge $A^+ = 0$. In principle, the complete set of bound state and scattering eigensolutions of H_{LC} can be obtained by solving the light-front Heisenberg equation

$$H_{LC} |\psi_h\rangle = \mathcal{M}_h^2 |\psi_h\rangle, \quad (3)$$

where $|\psi_h\rangle$ is an expansion in multi-particle Fock eigenstates $\{|n\rangle\}$ of the free light-front Hamiltonian: $|\psi_h\rangle = \sum_n \psi_{n/h} |n\rangle$. The LF Heisenberg equation has in fact been solved for QCD(1+1) and a number of other theories using the discretized light-cone quantization method [59]. The light-cone gauge has the advantage that all gluon

degrees of freedom have physical polarization and positive metric. In addition, orbital angular momentum has a simple physical interpretation in this representation. The light-front wavefunctions (LFWFs) $\psi_{n/h}$ provide a frame independent representation of a hadron, and relate the quark and gluon degrees of freedom with their asymptotic hadronic state.

Given the light-front wavefunctions $\psi_{n/h}$ one can compute a large range of hadron observables. For example, the valence and sea quark and gluon distributions which are measured in deep inelastic lepton scattering are defined from the squares of the LFWFs summed over all Fock states n . Form factors, exclusive weak transition amplitudes [60] such as $B \rightarrow \ell\nu\pi$, and the generalized parton distributions [61] measured in deeply virtual Compton scattering are (assuming the “handbag” approximation) overlaps of the initial and final LFWFs with $n = n'$ and $n = n' + 2$. The gauge-invariant distribution amplitude $\phi_H(x_i, Q)$ defined from the integral over the transverse momenta $\mathbf{k}_{\perp i}^2 \leq Q^2$ of the valence (smallest n) Fock state provides a fundamental measure of the hadron at the amplitude level [62, 63]; they are the nonperturbative input to the factorized form of hard exclusive amplitudes and exclusive heavy hadron decays in perturbative QCD. The resulting distributions obey the DGLAP and ERL evolution equations as a function of the maximal invariant mass, thus providing a physical factorization scheme [48]. In each case, the derived quantities satisfy the appropriate operator product expansions, sum rules, and evolution equations. However, at large x where the struck quark is far-off shell, DGLAP evolution is quenched [64], so that the fall-off of the DIS cross sections in Q^2 satisfies inclusive-exclusive duality at fixed W^2 .

The hadron wavefunction is an eigenstate of the total momentum P^+ and \mathbf{P}_\perp and the longitudinal spin projection S_z , and is normalized according to

$$\begin{aligned} \langle \psi_h(P^+, \mathbf{P}_\perp, S_z) | \psi_h(P'^+, \mathbf{P}'_\perp, S'_z) \rangle \\ = 2P^+ (2\pi)^3 \delta_{S_z, S'_z} \delta(P^+ - P'^+) \delta^{(2)}(\mathbf{P}_\perp - \mathbf{P}'_\perp). \end{aligned} \quad (4)$$

Each hadronic eigenstate $|\psi_h\rangle$ is expanded in a Fock-state complete basis of non-interacting n -particle states $|n\rangle$ with an infinite number of components

$$\begin{aligned} |\psi_h(P^+, \mathbf{P}_\perp, S_z)\rangle = \\ \sum_{n, \lambda_i} \int [dx_i] [d^2\mathbf{k}_{\perp i}] \psi_{n/h}(x_i, \mathbf{k}_{\perp i}, \lambda_i) \\ \frac{1}{\sqrt{x_i}} |n : x_i P^+, x_i \mathbf{P}_\perp + \mathbf{k}_{\perp i}, \lambda_i\rangle, \end{aligned} \quad (5)$$

where the sum begins with the valence state; *e.g.*, $n \geq 2$ for mesons. The coefficients of the Fock expansion

$$\psi_{n/h}(x_i, \mathbf{k}_{\perp i}, \lambda_i) = \langle n : x_i, \mathbf{k}_{\perp i}, \lambda_i | \psi_h \rangle, \quad (6)$$

are independent of the total momentum P^+ and \mathbf{P}_\perp of the hadron and depend only on the relative partonic coordinates, the longitudinal momentum fraction

$x_i = k_i^+ / P^+$, the relative transverse momentum $\mathbf{k}_{\perp i}$, and λ_i , the projection of the constituent's spin along the z direction. Thus, given the Fock-projection (6), the wavefunction of a hadron is determined in any frame. The amplitudes $\psi_{n/h}$ represent the probability amplitudes to find on-mass-shell constituents i with longitudinal momentum $x_i P^+$, transverse momentum $x_i \mathbf{P}_{\perp} + \mathbf{k}_{\perp i}$, helicity λ_i and invariant mass

$$\mathcal{M}_n^2 = \sum_{i=1}^n k_i^\mu k_{i\mu} = \sum_{i=1}^n \frac{k_{\perp i}^2 + m_i^2}{x_i}, \quad (7)$$

in the hadron h . Momentum conservation requires $\sum_{i=1}^n x_i = 1$ and $\sum_{i=1}^n \mathbf{k}_{\perp i} = 0$. In addition, each light front wavefunction $\psi_{n/h}(x_i, \vec{k}_{\perp i}, \lambda_i)$ obeys the angular momentum sum rule [65]

$$J^z = \sum_{i=1}^n S_i^z + \sum_{i=1}^{n-1} L_i^z, \quad (8)$$

where $S_i^z = \lambda_i$ and the $n-1$ orbital angular momenta have the operator form

$$L_i^z = -i \left(\frac{\partial}{\partial k_i^x} k_i^y - \frac{\partial}{\partial k_i^y} k_i^x \right). \quad (9)$$

The LFWFs are normalized according to

$$\sum_n \int [dx_i] [d^2 \mathbf{k}_{\perp i}] |\psi_{n/h}(x_i, \mathbf{k}_{\perp i})|^2 = 1, \quad (10)$$

where the measure of the constituents phase-space momentum integration is

$$\int [dx_i] \equiv \prod_{i=1}^n \int dx_i \delta \left(1 - \sum_{j=1}^n x_j \right), \quad (11)$$

$$\int [d^2 \mathbf{k}_{\perp i}] \equiv \prod_{i=1}^n \int \frac{d^2 \mathbf{k}_{\perp i}}{2(2\pi)^3} (16\pi^3) \delta^{(2)} \left(\sum_{j=1}^n \mathbf{k}_{\perp j} \right), \quad (12)$$

for the normalization given by (4). The spin indices have been suppressed.

A. Construction of Partonic States

The complete basis of Fock-states $|n\rangle$ is constructed by applying free-field creation operators to the vacuum state $|0\rangle$ which has no particle content, $P^+|0\rangle = 0$, $\mathbf{P}_{\perp}|0\rangle = 0$. The basic constituents appear from the light-front quantization of the excitations of the dynamical fields, the Dirac field ψ_+ , $\psi_{\pm} = \Lambda_{\pm} \psi$, $\Lambda_{\pm} = \gamma^0 \gamma^{\pm}$, and the transverse field \mathbf{A}_{\perp} in the $A^+ = 0$ gauge, expanded in terms of creation and annihilation operators on the transverse

plane with coordinates $x^- = x^0 - x^3$ and \vec{x}_{\perp} at fixed light-front time $x^+ = x^0 + x^3$ [58]

$$\psi_+(x)_{\alpha} = \sum_{\lambda} \int_{q^+ > 0} \frac{dq^+}{\sqrt{2q^+}} \frac{d^2 \mathbf{q}_{\perp}}{(2\pi)^3} [b_{\lambda}(q) u_{\alpha}(q, \lambda) e^{-iq \cdot x} + d_{\lambda}(q)^{\dagger} v_{\alpha}(q, \lambda) e^{iq \cdot x}], \quad (13)$$

$$\mathbf{A}_{\perp}(x) = \sum_{\lambda} \int_{q^+ > 0} \frac{dq^+}{\sqrt{2q^+}} \frac{d^2 \mathbf{q}_{\perp}}{(2\pi)^3} [a(q) \vec{\epsilon}_{\perp}(q, \lambda) e^{-iq \cdot x} + a(q)^{\dagger} \vec{\epsilon}_{\perp}^*(q, \lambda) e^{iq \cdot x}], \quad (14)$$

with commutation relations

$$[a(q), a^{\dagger}(q')] = \{b(q), b^{\dagger}(q')\} = \{d(q), d^{\dagger}(q')\} = (2\pi)^3 \delta(q^+ - q'^+) \delta^{(2)}(\mathbf{q}_{\perp} - \mathbf{q}'_{\perp}). \quad (15)$$

We shall use the conventions of Appendix A of Ref. [48] for the properties of the light-cone spinors. A one-particle state is defined by

$$|q\rangle = \sqrt{2q^+} b^{\dagger}(q) |0\rangle, \quad (16)$$

so that its normalization has the Lorentz invariant form

$$\langle q|q'\rangle = 2q^+ (2\pi)^3 \delta(q^+ - q'^+) \delta^{(2)}(\mathbf{q}_{\perp} - \mathbf{q}'_{\perp}), \quad (17)$$

and this fixes our normalization. Each n -particle Fock state $|p_i^+, \mathbf{p}_{\perp i}\rangle$ is an eigenstate of P^+ and \mathbf{P}_{\perp} and is normalized according to

$$\langle p_i^+, \mathbf{p}_{\perp i}, \lambda | p_i'^+, \mathbf{p}_{\perp i}', \lambda' \rangle = 2p_i^+ (2\pi)^3 \delta(p_i^+ - p_i'^+) \delta^{(2)}(\mathbf{p}_{\perp i} - \mathbf{p}_{\perp i}') \delta_{\lambda, \lambda'}. \quad (18)$$

B. Impact Space Representation

The holographic mapping of hadronic LFWFs to AdS string modes is most transparent when one uses the impact parameter space representation. The total position coordinate of a hadron or its transverse center of momentum \mathbf{R}_{\perp} , is defined in terms of the energy momentum tensor $T^{\mu\nu}$

$$\mathbf{R}_{\perp} = \frac{1}{P^+} \int dx^- \int d^2 \mathbf{x}_{\perp} T^{++} \mathbf{x}_{\perp}. \quad (19)$$

In terms of partonic transverse coordinates

$$x_i \mathbf{r}_{\perp i} = x_i \mathbf{R}_{\perp} + \mathbf{b}_{\perp i}, \quad (20)$$

where the $\mathbf{r}_{\perp i}$ are the physical transverse position coordinates and $\mathbf{b}_{\perp i}$ frame independent internal coordinates, conjugate to the relative coordinates $\mathbf{k}_{\perp i}$. Thus, $\sum_i \mathbf{b}_{\perp i} = 0$ and $\mathbf{R}_{\perp} = \sum_i x_i \mathbf{r}_{\perp i}$. The LFWFs

$\psi_n(x_j, \mathbf{k}_{\perp j})$ can be expanded in terms of the $n-1$ independent transverse coordinates $\mathbf{b}_{\perp j}$, $j = 1, 2, \dots, n-1$

$$\psi_n(x_j, \mathbf{k}_{\perp j}) = (4\pi)^{(n-1)/2} \prod_{j=1}^{n-1} \int d^2\mathbf{b}_{\perp j} \exp\left(i \sum_{j=1}^{n-1} \mathbf{b}_{\perp j} \cdot \mathbf{k}_{\perp j}\right) \tilde{\psi}_n(x_j, \mathbf{b}_{\perp j}). \quad (21)$$

The normalization is defined by

$$\sum_n \prod_{j=1}^{n-1} \int dx_j d^2\mathbf{b}_{\perp j} \left| \tilde{\psi}_n(x_j, \mathbf{b}_{\perp j}) \right|^2 = 1. \quad (22)$$

IV. THE FORM FACTOR IN QCD

The hadron form factors can be predicted from the overlap integral of string modes propagating in AdS space with the boundary electromagnetic source which probes the AdS interior, or equivalently by using the Drell-Yan-West formula in physical space-time. If both quantities represent the same physical observable for any value of the transfer momentum q^2 , an exact correspondence can be established between the string modes Φ in fifth-dimensional AdS space and the light-front wavefunctions of hadrons $\psi_{n/h}$ in 3+1 spacetime [39].

A. The Drell-Yan-West Form Factor

The light-front formalism provides an exact Lorentz-invariant representation of current matrix elements in terms of the overlap of light-front wave functions. The electromagnetic current has elementary couplings to the charged constituents since the full Heisenberg current can be replaced by the free quark current $J^\mu(0)$, evaluated at fixed light-cone time $x^+ = 0$ in the $q^+ = 0$ frame [66]. In contrast to the covariant Bethe-Salpeter equation, in the light front Fock expansion one does not need to include the contributions to the current from an infinite number of irreducible kernels, or the interactions of the electromagnetic current with vacuum fluctuations [58].

We choose the light-front frame coordinates

$$P = (P^+, P^-, \mathbf{P}_\perp) = \left(P^+, \frac{M^2}{P^+}, \vec{0}_\perp\right), \quad (23)$$

$$q = (q^+, q^-, \mathbf{q}_\perp) = \left(0, \frac{-q^2}{P^+}, \mathbf{q}_\perp\right),$$

where $q^2 = -Q^2 = -2P \cdot q = -\mathbf{q}_\perp^2$ is the space-like four momentum squared transferred to the composite system. The electromagnetic form factor of a meson is defined in terms of the hadronic amplitude of the electromagnetic current evaluated at light-cone time $x^+ = 0$

$$\langle P' | J^+(0) | P \rangle = 2(P + P')^+ F(Q^2), \quad (24)$$

where $P' = P + q$ and $F(0) = 1$.

The expression for the current operator $J^+(x) = \sum_q e_q \bar{\psi}(x) \gamma^+ \psi(x)$ in the particle number representation follows from the momentum expansion of $\psi(x)$ in terms of creation and annihilation operators given by (13). Since γ^+ conserves the spin component of the struck quark (Appendix A), we obtain for $J^+(0)$

$$J^+(0) = \sum_q e_q \int \frac{dq^+ d^2\mathbf{q}_\perp}{(2\pi)^3} \int \frac{dq'^+ d^2\mathbf{q}'_\perp}{(2\pi)^3} \{b_\uparrow^\dagger(q) b_\uparrow(q') + b_\downarrow^\dagger(q) b_\downarrow(q') - d_\uparrow^\dagger(q) d_\uparrow(q') - d_\downarrow^\dagger(q) d_\downarrow(q')\}. \quad (25)$$

The operator J annihilates a quark (antiquark) with with charge e_q ($-e_q$), momentum q' and spin-up (spin-down) along the z direction and creates a quark (antiquark) with the same spin and momentum q .

The matrix element $\langle \psi_{P'} | J^+(0) | \psi_P \rangle$ can be computed by expanding the initial and final hadronic states in terms of its Fock components using (5). The transition amplitude can then be expressed as a sum of overlap integrals with diagonal J^+ -matrix elements in the n -particle Fock-state basis. For each Fock-state, we label with q the struck constituent quark with charge e_q and $j = 1, 2, \dots, n-1$ each spectator. Using the normalization condition (18) for each individual constituent and after integration over the intermediate variables in the $q^+ = 0$ frame, we find the Drell-Yan-West (DYW) expression for the meson form factor [66, 67]

$$F(q^2) = \sum_n \int [dx_i] [d^2\mathbf{k}_{\perp i}] \sum_j e_j \psi_{n/P'}^*(x_i, \mathbf{k}'_{\perp i}, \lambda_i) \psi_{n/P}(x_i, \mathbf{k}_{\perp i}, \lambda_i), \quad (26)$$

where the variables of the light-cone Fock components in the final-state are given by $\mathbf{k}'_{\perp i} = \mathbf{k}_{\perp i} + (1-x_i)\mathbf{q}_\perp$ for a struck constituent quark and $\mathbf{k}'_{\perp i} = \mathbf{k}_{\perp i} - x_i\mathbf{q}_\perp$ for each spectator. The formula is exact if the sum is over all Fock states n . Notice that there is a factor of N_C from a closed quark loop where the photon is attached and a normalization factor of $1/\sqrt{N_C}$ for each meson wave function; thus color factors cancel out from the expression of the form factor.

B. The Form Factor in Impact Space

One of the important advantages of the light-front formalism is that current matrix elements can be represented without approximation as overlaps of light-front wavefunctions. In the case of the elastic space-like form factors, the matrix element of the J^+ current only couples Fock states with the same number of constituents, as expressed by the Drell-Yan-West formula (26). Suppose that the charged parton n is the active constituent

struck by the current, and the quarks $i = 1, 2, \dots, n-1$ are spectators. We substitute (21) in the DYW formula (26). Integration over k_\perp phase space gives us $n-1$ delta functions to integrate over the $n-1$ intermediate transverse variables with the result

$$F(q^2) = \sum_n \prod_{j=1}^{n-1} \int dx_j d^2 \mathbf{b}_{\perp j} \exp \left(i \mathbf{q}_\perp \cdot \sum_{j=1}^{n-1} x_j \mathbf{b}_{\perp j} \right) \left| \tilde{\psi}_n(x_j, \mathbf{b}_{\perp j}) \right|^2, \quad (27)$$

corresponding to a change of transverse momentum $x_j \mathbf{q}_\perp$ for each of the $n-1$ spectators. This is a convenient form for comparison with AdS results, since the form factor may be expressed in terms of the product of light-front wave functions with identical variables.

C. Effective Single-Particle Distribution

The form factor in the light-front formulation has an exact representation in terms of an effective single particle density [68]

$$F(q^2) = \int_0^1 dx \rho(x, \mathbf{q}_\perp), \quad (28)$$

where $\rho(x, \mathbf{q}_\perp)$ is given by

$$\rho(x, \mathbf{q}_\perp) = \sum_n \prod_{j=1}^{n-1} \int dx_j d^2 \mathbf{b}_{\perp j} \delta \left(1 - x - \sum_{j=1}^{n-1} x_j \right) \exp \left(i \mathbf{q}_\perp \cdot \sum_{j=1}^{n-1} x_j \mathbf{b}_{\perp j} \right) \left| \psi_n(x_j, \mathbf{b}_\perp) \right|^2. \quad (29)$$

The integration is over the coordinates of the $n-1$ spectator partons, and $x = x_n$ is the coordinate of the active charged quark. We can also write the form factor in terms of an effective single particle transverse distribution $\tilde{\rho}(x, \vec{\eta}_\perp)$ [68]

$$F(q^2) = \int_0^1 dx \int d^2 \vec{\eta}_\perp e^{i \vec{\eta}_\perp \cdot \mathbf{q}_\perp} \tilde{\rho}(x, \vec{\eta}_\perp), \quad (30)$$

where $\vec{\eta}_\perp = \sum_{j=1}^{n-1} x_j \mathbf{b}_{\perp j}$ is the x -weighted transverse position coordinate of the $n-1$ spectators [69]. The corresponding transverse density is

$$\begin{aligned} \tilde{\rho}(x, \vec{\eta}_\perp) &= \int \frac{d^2 \mathbf{q}_\perp}{(2\pi)^2} e^{-i \vec{\eta}_\perp \cdot \mathbf{q}_\perp} \rho(x, \mathbf{q}_\perp) \\ &= \sum_n \prod_{j=1}^{n-1} \int dx_j d^2 \mathbf{b}_{\perp j} \delta \left(1 - x - \sum_{j=1}^{n-1} x_j \right) \\ &\quad \delta^{(2)} \left(\sum_{j=1}^{n-1} x_j \mathbf{b}_{\perp j} - \vec{\eta}_\perp \right) \left| \tilde{\psi}_n(x_j, \mathbf{b}_{\perp j}) \right|^2. \end{aligned} \quad (31)$$

The procedure is valid for any Fock state n , and thus the results can be summed over n to obtain an exact representation.

V. THE FORM FACTOR IN ADS/CFT AND LIGHT-FRONT MAPPING OF STRING AMPLITUDES

We now derive the corresponding expression for the electromagnetic form factor of a pion in AdS. The derivation can be extended to vector mesons and baryons, although the actual analysis becomes more complex, since the matrix elements of space-like local operators include the dependence for the different spin transitions. For example, in the case of the proton, the Dirac and Pauli form factors correspond to the light-cone spin-conserving and spin-flip matrix elements of the J^+ current [70], and a mapping to the corresponding AdS matrix elements has to be carried out for each case. The electric vector-meson form factor has been discussed recently in the context of the hard wall model in [35, 38] and in the soft-wall model in [36]. A detailed discussion of the form factor of nucleons in AdS/QCD will be given elsewhere.

In AdS/CFT, the hadronic matrix element for the electromagnetic current has the form of a convolution of the string modes for the initial and final hadrons with the external electromagnetic source which propagates inside AdS. We discuss first the truncated space or hard wall [2] holographic model, where quark and gluons as well as the external electromagnetic current propagate freely into the AdS interior according to the AdS metric. We discuss latter the case where the sharp boundary of the hard wall in the infrared region is replaced by the introduction of a soft cutoff.

AdS coordinates are the Minkowski coordinates x^μ and z , the holographic coordinate, which we label collectively $x^\ell = (x^\mu, z)$. The metric of the full space-time is

$$ds^2 = g_{\ell m} dx^\ell dx^m \quad (32)$$

where $g_{\ell m} \rightarrow (R^2/z^2) \eta_{\ell m}$ in the conformal $z \rightarrow 0$ limit, and $\eta_{\ell m}$ has diagonal components $(1, -1, \dots, -1)$. Unless stated otherwise, 5-dimensional fields are represented by capital letters such as Φ . Holographic fields in 4-dimensional Minkowski space are represented by lower case letters such as ϕ . The metric (32) is AdS in the finite interval $0 \leq z \leq z_0 = 1/\Lambda_{QCD}$, and the dual conformal field theory is strongly coupled at all scales up to the confining scale.

Assuming minimal coupling the form factor has the form [23, 38]

$$i g_5 \int d^4 x dz \sqrt{g} A^\ell(x, z) \Phi^*(x, z) \overleftrightarrow{\partial}_\ell \Phi(x, z), \quad (33)$$

where g_5 is a five-dimensional effective coupling constant and $\Phi(x, z)$ is a normalizable mode representing a hadronic state, $\Phi(x, z) \sim e^{-iP \cdot x} \Phi^P(z)$, with hadronic invariant mass given by $P_\mu P^\mu = \mathcal{M}^2$.

We consider the propagation inside AdS space of an electromagnetic probe polarized along Minkowski coordinates ($Q^2 = -q^2 > 0$)

$$A(x, z)_\mu = \epsilon_\mu e^{-iQ \cdot x} J(Q, z), \quad A_z = 0, \quad (34)$$

where $J(Q, z)$ has the value 1 at zero momentum transfer, since we are normalizing the bulk solutions to the total charge operator, and as boundary limit the external current $A_\mu(x, z \rightarrow 0) = \epsilon_\mu e^{-iQ \cdot x}$. Thus

$$J(Q = 0, z) = J(Q, z = 0) = 1. \quad (35)$$

The propagation of the external current inside AdS space is described by the AdS wave equation

$$[z^2 \partial_z^2 - z \partial_z - z^2 Q^2] J(Q, z) = 0, \quad (36)$$

and its solution subject to the boundary conditions (35) is

$$J(Q, z) = zQK_1(zQ). \quad (37)$$

Notice that $J(Q, z)$ can also be obtained from the Green's function of (36), since it is the bulk-to-boundary propagator [9, 38].

Substituting the normalizable mode $\Phi(x, z)$ in (33) and extracting an overall delta function from momentum conservation at the vertex, we find for the elastic form factor of the pion

$$\langle P' | J^\mu(0) | P \rangle = 2(P + P')^\mu F(Q^2), \quad (38)$$

with

$$F(Q^2)_{P \rightarrow P'} = R^3 \int \frac{dz}{z^3} \Phi^{P'}(z) J(Q, z) \Phi^P(z). \quad (39)$$

The form factor in AdS is thus represented as the overlap in the fifth dimension coordinate z of the normalizable modes dual to the incoming and outgoing hadrons, Φ^P and $\Phi^{P'}$, with the non-normalizable mode, $J(Q, z)$, dual to the external source [23]. Since $K_n(x) \sim \sqrt{\pi/2x} e^{-x}$ for large x , it follows that the external electromagnetic field is suppressed inside the AdS cavity for large Q . At small z the string modes scale as $\Phi \sim z^\Delta$. At large enough Q , the important contribution to (39) is from the region near $z \sim 1/Q$

$$F(Q^2) \rightarrow \left[\frac{1}{Q^2} \right]^{\Delta-1}, \quad (40)$$

and the ultraviolet point-like behavior responsible for the power law scaling [71, 72] is recovered. This is a remarkable consequence of truncating AdS space [2], since we are indeed describing the coupling of an electromagnetic current to an extended mode, and we should expect soft collision amplitudes characteristic of strings, instead of hard point-like ultraviolet behavior [2].

A. Light-Front Mapping in the Hard-Wall Model

We can now establish a connection of the AdS/CFT and the LF formulae. It is useful to integrate (30) over angles to obtain

$$F(q^2) = 2\pi \int_0^1 dx \frac{(1-x)}{x} \int \zeta d\zeta J_0 \left(\zeta q \sqrt{\frac{1-x}{x}} \right) \tilde{\rho}(x, \zeta), \quad (41)$$

where we have introduced the variable

$$\zeta = \sqrt{\frac{x}{1-x}} \left| \sum_{j=1}^{n-1} x_j \mathbf{b}_{\perp j} \right|, \quad (42)$$

representing the x -weighted transverse impact coordinate of the spectator system. We substitute the integral representation of the bulk-to boundary propagator $J(Q, z)$ (Appendix B)

$$J(Q, z) = \int_0^1 dx J_0 \left(\zeta Q \sqrt{\frac{1-x}{x}} \right) = \zeta Q K_1(\zeta Q), \quad (43)$$

in the expression for the form factor in AdS space (39) for arbitrary values of Q . Comparing with the light-front expression (41), we can identify the spectator density function appearing in the light-front formalism with the corresponding AdS density

$$\tilde{\rho}(x, \zeta) = \frac{R^3}{2\pi} \frac{x}{1-x} \frac{|\Phi(\zeta)|^2}{\zeta^4}. \quad (44)$$

Equation (44) gives a precise relation between string modes $\Phi(\zeta)$ in AdS₅ and the QCD transverse charge density $\tilde{\rho}(x, \zeta)$. The variable ζ , $0 \leq \zeta \leq \Lambda_{\text{QCD}}^{-1}$, represents a measure of the transverse separation between point-like constituents, and it is also the holographic variable z [39].

In the case of a two-parton constituent bound state the correspondence between the string amplitude $\Phi(z)$ in AdS space and the QCD light-front wave wavefunction $\tilde{\psi}(x, \mathbf{b}_\perp)$ follows immediately from (44). For two partons the transverse density (31) has the simple form

$$\tilde{\rho}_{n=2}(x, \zeta) = \frac{|\tilde{\psi}(x, \zeta)|^2}{(1-x)^2}, \quad (45)$$

and a closed form solution for the two-constituent bound state light-front wave function is obtained

$$|\tilde{\psi}(x, \zeta)|^2 = \frac{R^3}{2\pi} x(1-x) \frac{|\Phi(\zeta)|^2}{\zeta^4}, \quad (46)$$

where $\zeta^2 = x(1-x)\mathbf{b}_\perp^2$. Here \mathbf{b}_\perp is the impact separation and Fourier conjugate to \mathbf{k}_\perp .

B. Holographic Light-Front Representation

The equations of motion in AdS space can be recast in the form of a light-cone Hamiltonian equation (3)

$$H_{LC} | \phi \rangle = \mathcal{M}^2 | \phi \rangle, \quad (47)$$

which allows the discussion of the AdS/CFT solutions in terms of light-front equations in physical 3+1 space-time. To simplify the discussion we consider the propagation of massive scalar modes in AdS space described by the normalized solutions to the wave equation ($d = 4$)

$$[z^2 \partial_z^2 - (d-1)z \partial_z + z^2 \mathcal{M}^2 - (\mu R)^2] \Phi(z) = 0, \quad (48)$$

where μ is the five-dimensional mass. Since the field Φ couples to a local boundary interpolating operator of dimension Δ , \mathcal{O}_Δ , μ is determined by the relation $(\mu R)^2 = \Delta(\Delta - 4)$ between the fifth-dimensional mass of Φ and its scaling dimension. For spin carrying constituents, the dimension of the operator is replaced by its twist τ , dimension minus spin, $\tau = \Delta - \sigma_i$. For example $\tau = 2$ for a meson.

By substituting

$$\phi(\zeta) = \left(\frac{\zeta}{R}\right)^{-3/2} \Phi(\zeta), \quad (49)$$

in (48) we find an effective Schrödinger equation as a function of the weighted impact variable ζ

$$\left[-\frac{d^2}{d\zeta^2} + V(\zeta)\right] \phi(\zeta) = \mathcal{M}^2 \phi(\zeta), \quad (50)$$

with $-\frac{d^2}{d\zeta^2}$ the light-front kinetic energy operator and

$$V(\zeta) = -\frac{1 - 4L^2}{4\zeta^2}, \quad (51)$$

the effective potential in the conformal limit, where we identify ζ with the fifth dimension z of AdS space: $\zeta = z$. We have written above $(\mu R)^2 = -4 + L^2$, for the fifth dimensional mass μ appearing in the AdS wave equation. The Breitenlohner-Freedman stability bound [73]

$$(\mu R)^2 \geq -\frac{d^2}{4}, \quad (52)$$

requires $L^2 \geq 0$, thus the lowest state corresponds to $L = 0$. Higher excitations are matched to the small z asymptotic behavior of each string mode to the corresponding conformal dimension of the boundary operators of each hadronic state. In the semiclassical approximation the solution for mesons with relative orbital angular momentum L corresponds to an effective five dimensional mass on the bulk side. The allowed values of μR are determined by requiring that asymptotically the dimensions become spaced by integers. The solution to (50) is

$$\phi(\zeta) = \zeta^{-3/2} \Phi(\zeta) = C z^{1/2} J_L(\zeta \mathcal{M}). \quad (53)$$

The holographic light-front wavefunction $\phi(\zeta) = \langle \zeta | \phi \rangle$ is normalized according to

$$\langle \phi | \phi \rangle = \int d\zeta |\langle \zeta | \phi \rangle|^2 = 1, \quad (54)$$

and represent the probability amplitude to find n -partons at transverse impact separation $\zeta = z$. Furthermore, its eigenmodes determine the hadronic mass spectrum [39]. For the truncated space model the eigenvalues are determined by the boundary conditions at $\phi(z = 1/\Lambda_{\text{QCD}}) = 0$ and are given in terms of the roots of Bessel functions: $\mathcal{M}_{L,k} = \beta_{L,k} \Lambda_{\text{QCD}}$. [74]

A closed form of the light-front wavefunctions $\tilde{\psi}(x, \mathbf{b}_\perp)$ in the hard wall model follow from (46)

$$\begin{aligned} \tilde{\psi}_{L,k}(x, \mathbf{b}_\perp) &= \frac{\Lambda_{\text{QCD}}}{\sqrt{\pi} J_{1+L}(\beta_{L,k})} \sqrt{x(1-x)} \\ J_L\left(\sqrt{x(1-x)} |\mathbf{b}_\perp| \beta_{L,k} \Lambda_{\text{QCD}}\right) &\theta\left(\mathbf{b}_\perp^2 \leq \frac{\Lambda_{\text{QCD}}^{-2}}{x(1-x)}\right). \end{aligned} \quad (55)$$

The resulting wavefunction depicted in Fig. 1 display confinement at large interquark separation and conformal symmetry at short distances, reproducing dimensional counting rules for hard exclusive amplitudes and the conformal properties of the LFWFs at high relative momenta in agreement with perturbative QCD.




FIG. 1: AdS/QCD Predictions for the ground state light-front wavefunction of a pion in impact space. Left: truncated space model for $\Lambda_{\text{QCD}} = 0.22$ GeV. Right: soft wall model for $\kappa = 0.375$ GeV.

VI. SOFT-WALL HOLOGRAPHIC MODEL

The predicted mass spectrum in the truncated space hard-wall (HW) model is linear $M \propto L$ at high orbital angular momentum L , in contrast to the quadratic dependence $M^2 \propto L$ in the usual Regge parameterization. It has been shown recently that by choosing a specific profile for a non-constant dilaton, the usual Regge dependence can be obtained [10]. The procedure allows one to retain conformal AdS metrics (1) and to introduce a smooth cutoff which depends on the profile of a dilaton background field φ

$$S = \int d^4x dz \sqrt{g} e^{-\varphi(z)} \mathcal{L}, \quad (56)$$

where φ is a function of the holographic coordinate z which vanishes in the ultraviolet limit $z \rightarrow 0$. The IR hard-wall or truncated space holographic model, discussed in the previous section, correspond to a constant dilaton field $\varphi(z) = \varphi_0$ in the confining region, $z \leq 1/\Lambda_{\text{QCD}}$, and to very large values elsewhere: $\varphi(z) \rightarrow \infty$ for $z > 1/\Lambda_{\text{QCD}}$. The introduction of a soft cutoff avoids the ambiguities in the choice of boundary conditions at the infrared wall. A convenient choice [10] for the background field with usual Regge behavior is $\varphi(z) = \kappa^2 z^2$. The resulting wave equations are equivalent to the radial equation of a two-dimensional oscillator, previously found in the context of mode propagation on $AdS_5 \times S^5$, in the light-cone formulation of Type II supergravity [75]. Also, equivalent results follow from the introduction of a gaussian warp factor in the AdS metric for the particular case of massless vector modes propagating in the distorted metric [76].

A. Light-Front Mapping in the Soft-Wall Model

We consider the propagation in AdS space of an electromagnetic probe polarized along Minkowski coordinates. Since the non-normalizable modes also couple to the dilaton field we must study the solutions of the modified wave equation

$$[z^2 \partial_z^2 - z(1 + 2\kappa^2 z^2) \partial_z - Q^2 z^2] J_\kappa(Q, z) = 0, \quad (57)$$

subject to the boundary conditions (35). The solution is

$$J_\kappa(Q, z) = \Gamma\left(1 + \frac{Q^2}{4\kappa^2}\right) U\left(\frac{Q^2}{4\kappa^2}, 0, \kappa^2 z^2\right), \quad (58)$$

where $U(a, b, c)$ is the confluent hypergeometric function.

The form factor in AdS space in presence of the dilaton background $\varphi = \kappa^2 z^2$ has the additional term $e^{-\kappa^2 z^2}$ in the metric

$$F(Q^2) = R^3 \int \frac{dz}{z^3} e^{-\kappa^2 z^2} \Phi(z) J_\kappa(Q, z) \Phi(z), \quad (59)$$

to be normalized to the charge operator at $Q = 0$. In the large Q^2 limit, $Q^2 \gg 4\kappa^2$, we find (Appendix C)

$$J_\kappa(Q, z) \rightarrow zQK_1(zQ) = J(Q, z). \quad (60)$$

Thus, for large transverse momentum the current decouples from the dilaton field, and we recover our previous scaling results for the ultraviolet behavior of matrix elements.

To obtain the corresponding basis set of LFWFs we compare the DYW expression of the form factor (41) with the AdS form factor (59) for large values of Q , where (60) is valid. Thus, in the large Q limit we can identify the light-front spectator density with the corresponding AdS density

$$\tilde{\rho}(x, \zeta) = \frac{R^3}{2\pi} \frac{x}{1-x} e^{-\kappa^2 \zeta^2} \frac{|\Phi(\zeta)|^2}{\zeta^4}. \quad (61)$$

When summed over all Fock states the Drell-Yan-West formula gives an exact result. The formula describes the coupling of the free electromagnetic current to the elementary constituents in the interaction representation. In the presence of a dilaton field in AdS space, or in the case where the electromagnetic probe propagates in modified confining AdS metrics, the electromagnetic AdS mode is no longer dual to a the free quark current, but dual to a dressed current, *i.e.*, a hadronic electromagnetic current including virtual $\bar{q}q$ pairs and thus confined. Consequently, at finite values of the momentum transfer Q^2 our simple identification discussed above has to be reinterpreted since we are comparing states in different representations: the interaction representation in light-cone QCD versus the Heisenberg representation in AdS. However both quantities should represent the same observables. We thus expect that the modified mapping corresponds to the presence of higher Fock states in the hadron.

B. Holographic Light-Front Representation

The equations of motion for scalar modes with smooth boundary conditions can also be recast in the form of a light-cone Hamiltonian form $H_{LC} |\phi\rangle = \mathcal{M}^2 |\phi\rangle$, with effective potential

$$V(\zeta) = -\frac{1 - 4L^2}{4\zeta^2} + \kappa^4 \zeta^2 + 2\kappa^2(L - 1), \quad (62)$$

in the covariant light-front effective Schrödinger equation (50). Its solution is

$$\phi(\zeta) = \kappa^{1+L} \sqrt{\frac{2n!}{(n+L)!}} \zeta^{1/2+L} e^{-\kappa^2 \zeta^2/2} L_n^L(\kappa^2 \zeta^2), \quad (63)$$

with eigenvalues

$$\mathcal{M}^2 = 4\kappa^2(n + L). \quad (64)$$

The lowest stable solution corresponds to $n = L = 0$. With this choice for the potential, the lowest possible state is a zero mode in the spectrum, which can be identified with the pion.

Upon the substitution

$$\Phi(\zeta) = e^{\kappa^2 \zeta^2/2} \left(\frac{\zeta}{R}\right)^{3/2} \phi(\zeta), \quad \zeta \rightarrow z, \quad (65)$$

we express the solution (63) as the scalar mode

$$\Phi(z) = \kappa^{1+L} \sqrt{\frac{2n!}{(n+L)!}} z^{2+L} L_n^L(\kappa^2 z^2), \quad (66)$$

of scaling dimension $\Delta = 2 + L$ propagating in AdS space. A closed form of the light-front wavefunctions $\tilde{\psi}(x, \mathbf{b}_\perp)$

for a two-parton state in the soft wall model then follows from (61)

$$\tilde{\psi}(x, \mathbf{b}_\perp) = \kappa^{L+1} \sqrt{\frac{2n!}{(n+L)!}} [x(1-x)]^{\frac{1}{2}+L} |\mathbf{b}_\perp|^L e^{-\frac{1}{2}\kappa^2 x(1-x)\mathbf{b}_\perp^2} L_n^L(\kappa^2 x(1-x)\mathbf{b}_\perp^2). \quad (67)$$

The resulting wavefunction for the pion is compared in Fig. 1 with the hard-wall result.

VII. PION FORM FACTOR

We compute the pion form factor from the AdS expressions (39) and (59) for the hadronic string modes Φ_π in the hard-wall (HW)

$$\Phi_\pi^{HW}(z) = \frac{\sqrt{2}\Lambda_{QCD}}{R^{3/2}J_1(\beta_{0,1})} z^2 J_0(z\beta_{0,1}\Lambda_{QCD}), \quad (68)$$

and soft-wall (SW) model

$$\Phi_\pi^{SW}(z) = \frac{\sqrt{2}\kappa}{R^{3/2}} z^2, \quad (69)$$

respectively. We have obtained numerical results for both models. It was found recently that in the case of the SW model the results for form factors can be expressed analytically [36]. Thus, we shall extend the analytical results given in [36] to string modes with arbitrary scaling dimension τ , integer or not, allowing us to study the effect of anomalous dimensions in the expression for the form factors. This is described in Appendix D.

Since the pion mode couples to a twist-two boundary interpolating operator which creates a two-component hadronic bound state, the form factor is given in the SW model by the simple monopole form (D11) corresponding to $n = 2$

$$F_\pi(Q^2) = \frac{4\kappa^2}{4\kappa^2 + Q^2}. \quad (70)$$

The hadronic scale is evaluated by fitting the space-like data for the form factor as shown in Figure 2, where we plot the product $Q^2 F_\pi(Q^2)$ for the soft and hard-wall holographic models. Both models would seem to describe the overall behavior of the space-like data, with a better high- Q^2 description of scaling with the SW model. However, when the low energy data is examined in detail the SW model gives a noticeable better description as shown in Figure 3. When the results for the pion form factor are analytically continued to the time-like region, $q^2 \rightarrow -q^2$ we obtain the results shown in Figure 4 for $\log(|F_\pi(q^2)|)$. The monopole form of the SW model reproduces well the ρ peak with $M_\rho = 4\kappa^2 = 750$ MeV and the scaling behavior in the space-like region, but does not give rise to the additional structure found in the time-like region. The unconfined propagator $J(Q, z) = zQK_1(zQ)$ used in our




FIG. 2: Space-like scaling behavior for $Q^2 F_\pi(Q^2)$ as a function of $Q^2 = -q^2$. The continuous line is the prediction of the soft-wall model for $\kappa = 0.375$ GeV. The dashed line is the prediction of the hard-wall model for $\Lambda_{QCD} = 0.22$ GeV. The black triangles is the data compilation from Baldini *et al.* [77], the red boxes are JLAB 1 data [78] and cobalt green diamonds are JLAB 2 data [79].




FIG. 3: Space-like behavior of the pion form factor $F_\pi(q^2)$ as a function of q^2 for $\kappa = 0.375$ GeV and $\Lambda_{QCD} = 0.22$ GeV. Continuous line: soft-wall model, dashed line: hard-wall model. Black (triangle) is the data compilation from Baldini *et al.* [77], red (box) is JLAB 1 [78] and cobalt green diamonds are JLAB 2 [79].

description of the hard wall model do not give rise to a pole structure in the time like region, and thus fails to reproduce the ρ structure. When a small anomalous dimension is introduced, $\tau = 2 \pm \epsilon$, the formula (D9) for the form factor should be used instead. For a small deviation from the canonical value of two, the behavior in the space

like region remains almost identical, but the form-factor in the time like region oscillates with infinite amplitude, and thus gives a non-physical acceptable solution.




FIG. 4: Space and time-like behavior of the pion form factor $\log(|F_\pi(q^2)|)$ as a function of q^2 for $\kappa = 0.375$ GeV in the soft-wall model. Black (triangle) is the data compilation from Baldini *et al.* [77], red (box) is JLAB 1 [78] and cobalt green diamonds are JLAB 2 [79].




FIG. 5: Time-like pion form factor $|F_\pi(q^2)|$ near the ρ peak as a function of q^2 for $\kappa = 0.375$ GeV in the soft-wall model. The upper curve correspond to $\Gamma = 0$ and the lower to $\Gamma = 100$ MeV. The data is from the compilation Baldini *et al.* [77].

In the strongly coupled semiclassical gauge/gravity limit hadrons have zero widths and are stable. One can nonetheless modify the formula (70) to introduce a finite width Γ

$$F_\pi(Q^2) = \frac{4\kappa^2}{4\kappa^2 - q^2 - 2i\kappa\Gamma}, \quad (71)$$

to compare with the pion form factor data near the ρ peak. The result is shown in Figure 5. The corresponding phases for the pion form factor are shown in Figure 6.




FIG. 6: Time-like pion form factor phase $\delta_\pi(q)$ as a function of q^2 for $\kappa = 0.375$ GeV in the soft-wall model. The sharp curve corresponds to $\Gamma = 0$ and the smooth curve to $\Gamma = 100$ MeV.

VIII. THE PION DECAY CONSTANT

The pion decay constant is given in terms of the matrix element of the axial isospin current $J^{\mu 5a}$ between a physical pion and the vacuum state [80]

$$\langle 0 | J_a^{\mu 5}(x) | \pi_b(P) \rangle = -iP^\mu f_\pi \delta_{ab} e^{-iP \cdot x}, \quad (72)$$

Axial and vector currents are

$$J^{\mu 5a} = \bar{Q} \gamma^\mu \gamma^5 t^a Q, \quad J^{\mu a} = \bar{Q} \gamma^\mu t^a Q \quad (73)$$

with Q the quark doublet

$$Q = \begin{pmatrix} \psi_u \\ \psi_d \end{pmatrix}, \quad (74)$$

and $t^a = \tau^a/2$ the generators of isospin rotations.

Consider the amplitude for the weak decay of a charged pion $\langle 0 | J_W^+(0) | \pi^-(P^+, \vec{P}_\perp) \rangle$, where J_W^+ is the flavor changing weak current

$$J_W^+ = \bar{\psi}_u \gamma^+ \frac{1}{2} (1 - \gamma_5) \psi_d. \quad (75)$$

It follows that [81]

$$\langle 0 | \bar{\psi}_u \gamma^+ \frac{1}{2} (1 - \gamma_5) \psi_d | \pi^- \rangle = i \frac{P^+ f_\pi}{\sqrt{2}}. \quad (76)$$

The expression for the current operator (75) in the particle number representation follows from the expansion of the field operator (13) in terms of quark creation and annihilation operators. The only terms which contribute to the annihilation process are the terms containing the product of operators d and b with opposite spin. Using the results listed in Appendix A for the quark spinor projections, we obtain the effective current operator

$$J_W^+ = \int \frac{dq^+ d^2 \vec{q}_\perp}{2\pi^3} \int \frac{dq'^+ d^2 \vec{q}'_\perp}{2\pi^3} d_{u\uparrow}(q) b_{d\downarrow}(q'), \quad (77)$$

in the limit of massless quarks. The flavor changing operator J annihilates a d -quark with momentum q' and spin-down along the z direction and a u -antiquark momentum q and spin-up. Only the valence state

$$|d\bar{u}\rangle = \frac{1}{\sqrt{N_C}} \frac{1}{\sqrt{2}} \sum_{c=1}^{N_C} \left(b_{cd\downarrow}^\dagger d_{cu\uparrow}^\dagger - b_{cd\uparrow}^\dagger d_{cu\downarrow}^\dagger \right) |0\rangle, \quad (78)$$

with $L_z = 0$, $S_z = 0$, contributes to the decay of the π^\pm . Expanding the hadronic initial state in the amplitude (76) into its Fock components we find

$$f_\pi = 2\sqrt{N_C} \int_0^1 dx \int \frac{d^2\vec{k}_\perp}{16\pi^3} \psi_{\bar{q}q/\pi}(x, k_\perp), \quad (79)$$

the light-cone equation which allows the exact computation of the pion decay constant in terms of the valence pion light-front wave function [48].

In terms of the distribution amplitude $\phi(x, Q)$

$$\phi(x, Q) = \int \frac{d^2\vec{k}_\perp}{16\pi^3} \psi(x, k_\perp), \quad (80)$$

it follows that

$$\phi_\pi(x, Q \rightarrow \infty) = \frac{4}{\sqrt{3}\pi} f_\pi \sqrt{x(1-x)}, \quad (81)$$

with

$$f_\pi = \frac{1}{8} \sqrt{\frac{3}{2}} R^{3/2} \lim_{\zeta \rightarrow 0} \frac{\Phi(\zeta)}{\zeta^2}. \quad (82)$$

since $\phi(x, Q \rightarrow \infty) \rightarrow \tilde{\psi}(x, \vec{b}_\perp \rightarrow 0)/\sqrt{4\pi}$ as $\zeta \rightarrow 0$. The pion decay constant depends only on the behavior of the AdS string mode near the asymptotic boundary, $\zeta = z = 0$, and the mode normalization. For the hard-wall truncated-space pion mode (68) we find

$$f_\pi^{HW} = \frac{\sqrt{3}}{8J_1(\beta_{0,k})} \Lambda_{\text{QCD}} = 91.7 \text{ MeV}, \quad (83)$$

for $\Lambda_{\text{QCD}} = 0.22 \text{ MeV}$. The corresponding result for the soft-wall pion mode (69) is

$$f_\pi^{SW} = \frac{\sqrt{3}}{8} \kappa = 81.2 \text{ MeV}, \quad (84)$$

for $\kappa = 0.375 \text{ MeV}$. The values of Λ_{QCD} and κ are determined from the space-like form factor data as discussed above. The experimental result for f_π is extracted from the rate of weak π decay and has the value $f_\pi = 92.4 \text{ MeV}$ [82].

It is interesting to note that the pion distribution amplitude predicted by AdS/QCD (81) has a quite different x -behavior than the asymptotic distribution amplitude predicted from the PQCD evolution [62] of the pion distribution amplitude $\phi_\pi(x, Q \rightarrow \infty) = \sqrt{3}f_\pi x(1-x)$. This observation appears to be consistent with the results

of the Fermilab diffractive dijet experiment [83] which shows a broader x distribution for the dijets at small transverse momentum $k_\perp \leq 1 \text{ GeV}$. The broader shape of the pion distribution increases the magnitude of the leading twist perturbative QCD prediction for the pion form factor by a factor of 16/9 compared to the prediction based on the asymptotic form, bringing the PQCD prediction close to the empirical pion form factor [37].

IX. CONCLUSIONS

One of the key difficulties in studies of quantum chromodynamics has been the absence of an analytic first approximation to the theory which not only can reproduce the hadronic spectrum, but also provides a good description of hadron wavefunctions. The AdS/CFT correspondence provides an elegant semi-classical approximation to QCD, which incorporates both color confinement and the conformal short-distance behavior appropriate for a theory with an infrared fixed point. Since the hadronic solutions are controlled by their twist dimension z^τ at small z , one also reproduces dimensional counting rules for hard exclusive processes. The AdS/CFT approach thus allows one to construct a model of hadrons which has both confinement at large distances and the conformal scaling properties which reproduce dimensional counting rules for hard exclusive reactions. The fundamental equations of AdS/CFT for mesons have the appearance of a radial Schrödinger Coulomb equation, but they are relativistic, covariant, and analytically tractable. The eigenvalues of the AdS/CFT equations provide a good description of the meson and baryon spectra for light quarks [8, 11, 39], and its eigensolutions provide a remarkably simple but realistic model of their valence wavefunctions. One can also derive analogous equations for baryons composed of massless quarks using a Dirac matrix representation for the baryon system [34].

The lowest stable state of the AdS equations are determined by the Breitenlohner-Freedman bound [73]. We can model confinement by imposing Dirichlet boundary conditions at $\phi(z = 1/\Lambda_{\text{QCD}}) = 0$. The eigenvalues are then given in terms of the roots of the Bessel functions: $\mathcal{M}_{L,k} = \beta_{L,k} \Lambda_{\text{QCD}}$. Alternatively, one can introduce a dilaton field [10] which provides a confinement potential $-\kappa^2 \zeta^2$ to the effective potential $V(\zeta)$. The resulting hadron spectra are given by linear Regge trajectories in the square of the hadron masses, \mathcal{M}^2 , characteristic of the Nambu string model. The AdS/CFT equations are integrable and thus the radial and orbital excitations can be obtained from ladder operators [34].

In this work we have shown that the eigensolutions $\phi_H(z)$ of the AdS/CFT equations in the fifth dimension z have a remarkable mapping to the light-front wavefunctions $\psi_H(x_i, \vec{b}_{\perp i})$, the hadronic amplitudes which describe the valence constituents of hadrons in physical space time, but at fixed light-cone time $\tau = t + z/c = 0$. Similarly, the AdS/CFT equations for hadrons can be

mapped to equivalent light-front equations. The correspondence of AdS/CFT amplitudes to the QCD wavefunctions in light-front coordinates in physical space-time, provides an exact holographic mapping at all energy scales between string modes in AdS space and hadronic boundary states. Most important, the eigen-solutions of the AdS/CFT equation can be mapped to light-front equations of the hadrons in physical space-time, thus providing an elegant description of the light hadrons at the amplitude level.

The mapping of AdS/CFT string modes to light-front wave functions thus provides a remarkable analytic first approximation to QCD. Since they are complete and orthonormal, the AdS/CFT model wavefunctions can also be used as a basis for the diagonalization of the full light-front QCD Hamiltonian, thus systematically improving the AdS/CFT approximation.

We have also shown the correspondence between the expressions for current matrix elements in AdS/CFT with the corresponding expressions for form factors as given in the light-front formalism. In first approximation, where one takes the current propagating in a non-confining background, one obtains the Drell-Yan West formula for valence Fock states, corresponding to the interaction picture of the light-front theory. Hadron form factors can thus be directly predicted from the overlap integrals in AdS space or equivalently by using the Drell-Yan-West formula in physical space-time. The form factor at high Q^2 receives its main contributions from small $\zeta \sim 1/Q$, corresponding to small $\vec{b}_\perp = \mathcal{O}(1/Q)$ and $1-x = \mathcal{O}(1/Q)$. We have also shown how to improve this approximation by studying the propagation of non-normalizable solutions representing the electromagnetic current in a modified AdS confining metric, or equivalently in a dilaton background. This improvement in the description of the current corresponds in the light-front to multiple hadronic Fock states. The introduction of the confined current implies that the timelike form factors of hadrons will be mediated by vector mesons, including radial excitations. The wavefunction of the normalizable vector meson states $\mathcal{A}(z)$ appearing in the spectral decomposition of the Green's function described in Appendix C, which is dual to the non-normalizable photon propagation mode in AdS, is twist-3. This is the expected result for even parity axial mesons in QCD, or $L = 1$ odd parity vector mesons composed of a scalar squark and anti-squarks. In the case of quark-antiquark states, one also expects to find $C = -1$, twist-2 meson solutions for the zero helicity component of the ρ with $S = 1$ and $L = 0$, which is supposed to give a dominant contribution to the ρ form factor.

We have applied our formulation to both the spacelike and timelike pion form factor. The description of the pion form factor in the spacelike domain is in good agreement with experiment in both confinement models, the hard and the soft wall holographic models. In the soft wall model the time-like pion form factor exhibits a pole at the ρ mass with zero width since hadrons are stable in

this theory. If one allows a width, the height of the ρ pole is in reasonable agreement with experiment. The space-like Dirac form factor for the proton is also very well reproduced by the double-pole analytic expression shown in Appendix D for the case $N = 3$. This will be discussed elsewhere.

The deeply virtual Compton amplitude in the handbag approximation can be expressed as overlap of light-front wavefunctions [61]. The deeply virtual Compton amplitudes can be Fourier transformed to b_\perp and $\sigma = x^- P^+ / 2$ space providing new insights into QCD distributions [84–87]. The distributions in the light-front direction σ typically display diffraction patterns arising from the interference of the initial and final state LFWFs [86, 88]. All of these processes can provide detailed tests of the AdS/CFT LFWFs predictions.

The phenomenology of the AdS/CFT model is just beginning, but it can be anticipated that it will have many applications to QCD phenomena at the amplitude level. For example, the model LFWFs obtained from AdS/QCD provide a basis for understanding hadron structure functions and fragmentation functions at the amplitude level; the same wavefunctions can describe hadron formation from the coalescence of co-moving quarks. The spin correlations which underly single and double spin correlations are also described by the AdS/CFT eigensolutions. The AdS/CFT hadronic wavefunctions also provide predictions for the generalized parton distributions of hadrons and their weak decay amplitudes from first principles. The amplitudes relevant to diffractive reactions could also be computed. We also anticipate that the extension of the AdS/CFT formalism to heavy quarks will allow a great variety of heavy hadron phenomena to be analyzed from first principles.

Acknowledgments

This research was supported by the Department of Energy contract DE-AC02-76SF00515. The research of GdT is supported in part by an Aportes grant from Florida Ice & Farm. We thank Garth Huber, Cheng-Ryong Ji, Anatoly Radyushkin, Giovanni Salmè and James Vary for helpful comments.

Appendix A: Light-Cone Spinors

In the light-cone (LC) formalism, spinors are written as [48]

$$u^\dagger(p) = \frac{1}{\sqrt{p^+}} (p^+ + \beta m + \vec{\alpha}_\perp \cdot \mathbf{p}_\perp) \chi^\dagger, \quad (\text{A1})$$

$$v^\dagger(p) = \frac{1}{\sqrt{p^+}} (p^+ - \beta m + \vec{\alpha}_\perp \cdot \mathbf{p}_\perp) \chi^\dagger, \quad (\text{A2})$$

with representation matrices

$$\vec{\alpha} = \begin{pmatrix} 0 & \vec{\sigma} \\ \vec{\sigma} & 0 \end{pmatrix}, \quad \beta = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}, \quad \gamma_5 = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}, \quad (\text{A3})$$

and spinors

$$\chi^\uparrow = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \end{pmatrix}, \quad \chi^\downarrow = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ 1 \\ 0 \\ -1 \end{pmatrix}. \quad (\text{A4})$$

The corresponding solutions for LC spinors $u^\downarrow(p)$ and $v^\downarrow(p)$ follow from reversing the spin component. Useful Dirac matrix elements for the helicity spinors are [48]

$$\bar{u}^\uparrow(\ell)\gamma^+u^\uparrow(k) = \bar{u}^\downarrow(\ell)\gamma^+u^\downarrow(k) = 2\sqrt{\ell^+k^+}, \quad (\text{A5})$$

$$\bar{u}^\downarrow(\ell)\gamma^+u^\uparrow(k) = \bar{u}^\uparrow(\ell)\gamma^+u^\downarrow(k) = 0, \quad (\text{A6})$$

with $\bar{v}(\ell)\gamma^\mu v(k) = \bar{u}(k)\gamma^\mu u(\ell)$ and

$$\bar{v}^\downarrow(\ell)\gamma^+u^\uparrow(k) = \bar{v}^\uparrow(\ell)\gamma^+u^\downarrow(k) = 2\sqrt{\ell^+k^+}, \quad (\text{A7})$$

$$\bar{v}^\uparrow(\ell)\gamma^+u^\uparrow(k) = \bar{v}^\downarrow(\ell)\gamma^+u^\downarrow(k) = 0. \quad (\text{A8})$$

It is simple to verify that

$$\frac{1}{2}(1 - \gamma_5)\chi^\uparrow = 0, \quad \frac{1}{2}(1 - \gamma_5)\chi^\downarrow = \chi^\downarrow. \quad (\text{A9})$$

Thus the chiral projection of the boosted LC spinors $u(p)$ and $v(p)$ in the limit of massless quarks

$$\frac{1}{2}(1 - \gamma_5)u^\downarrow(p) = u^\downarrow(p), \quad \frac{1}{2}(1 - \gamma_5)u^\uparrow(p) = 0, \quad (\text{A10})$$

$$\frac{1}{2}(1 - \gamma_5)v^\downarrow(p) = 0, \quad \frac{1}{2}(1 - \gamma_5)v^\uparrow(p) = v^\uparrow(p), \quad (\text{A11})$$

which follows from the fact that γ_5 commutes with $\vec{\alpha}_\perp$, but not with β . Consequently, in the massless limit we find that LC quark spinors with positive spin projection along the z -component also transform as right-handed spinors and LC spinors with negative spin projection are left-handed spinors. Also spinors representing massless antiquarks with positive spin component along the z -direction are left-handed and right-handed if polarized along the negative z -direction.

Appendix B: A Useful Integral

Consider the integral

$$J(Q, z) = \int_0^1 dx J_0 \left(zQ \sqrt{\frac{1-x}{x}} \right) \quad (\text{B1})$$

Changing the variable x according to $x = \frac{Q^2}{t^2 + Q^2}$, we recast (B1) as

$$J(z, Q) = 2Q^2 \int_0^\infty \frac{tJ_0(zt)}{(t^2 + Q^2)^2} dt, \quad (\text{B2})$$

which is a Hankel-Nicholson type integral [89],

$$\int_0^\infty \frac{t^{\nu+1} J_\nu(zt)}{(t^2 + a^2)^{\mu+1}} dt = \frac{z^\mu a^{\nu-\mu}}{z^\mu \Gamma(\mu+1)} K_{\nu-\mu}(az). \quad (\text{B3})$$

Thus

$$J(Q, z) = QzK_1(zQ). \quad (\text{B4})$$

Appendix C: Bulk-to-Boundary Propagator in the Soft Wall Model

The bulk-to-boundary propagator in the soft-wall holographic model follows from the solution of the wave equation

$$[z^2 \partial_z^2 - (1 + 2\kappa^2 z^2) z \partial_z - Q^2 z^2] J_\kappa(Q, z) = 0, \quad (\text{C1})$$

describing the propagation of an electromagnetic probe in AdS space coupled to a dilaton background field $\varphi(z) = \kappa^2 z^2$. The solution to (C1) subject to the boundary conditions

$$J_\kappa(Q = 0, z) = J_\kappa(Q, z = 0) = 1, \quad (\text{C2})$$

is

$$J_\kappa(Q, z) = \Gamma\left(1 + \frac{Q^2}{4\kappa^2}\right) U\left(\frac{Q^2}{4\kappa^2}, 0, \kappa^2 z^2\right). \quad (\text{C3})$$

Using the integral representation of the confluent hypergeometric function $U(a, b, z)$ [90]

$$\Gamma(a)U(a, b, z) = \int_0^\infty e^{-zt} t^{a-1} (1+t)^{b-a-1} dt, \quad (\text{C4})$$

we find

$$J_\kappa(Q, z) = \frac{Q^2}{4\kappa^2} \int_0^\infty e^{-\kappa^2 z^2 t} \left(\frac{1+t}{t}\right)^{-\frac{Q^2}{4\kappa^2}} \frac{dt}{t(1+t)}. \quad (\text{C5})$$

Making the change of variable $t = \frac{Q^2}{4\kappa^2} \mu$ and taking the limit $Q^2 \gg \kappa^2$

$$J_\kappa(Q \rightarrow \infty, z) = \int_0^\infty \frac{d\mu}{\mu^2} \exp\left(-\frac{z^2 Q^2}{4} \mu - \frac{1}{\mu}\right). \quad (\text{C6})$$

Comparing the above expression with the integral representation of the modified Bessel function $K_\nu(z)$

$$K_\nu(z) = \frac{1}{2} \left(\frac{z}{2}\right)^\nu \int_0^\infty \frac{e^{-t - \frac{z^2}{4t}}}{t^{\nu+1}} dt, \quad (\text{C7})$$

we find

$$\lim_{Q \rightarrow \infty} J_\kappa(Q, z) = zQK_1(zQ). \quad (\text{C8})$$

In the large Q limit the electromagnetic probe in AdS space decouples from the dilaton background.

1. Green's Function Formalism

In a finite spatial dimension the Green's function has an spectral decomposition in terms of eigenfunctions. Normalizable spin-one modes propagating on AdS₅ have polarization and plane waves along Minkowski coordinates $\mathcal{A}(x, z)_\mu = e^{-iP \cdot x} \mathcal{A}(z) \epsilon_\mu$, with invariant mass

$P_\mu P^\mu = \mathcal{M}^2$. The eigenfunctions are determined by the eigenvalue equation

$$[-z^2 \partial_z^2 + (1 + 2\kappa^2 z^2) z \partial_z] \mathcal{A}_n(z) = M_n^2 z^2 \mathcal{A}_n(z), \quad (\text{C9})$$

with solutions

$$\mathcal{A}_n(z) = \frac{\kappa^2}{R^{1/2}} \sqrt{\frac{2}{(n+1)}} z^2 L_n^1(\kappa^2 z^2), \quad (\text{C10})$$

and eigenvalues

$$M_n^2 = 4\kappa^2(n+1). \quad (\text{C11})$$

The eigenfunctions \mathcal{A}_n are orthonormal

$$R \int \frac{dz}{z} e^{-\kappa^2 z^2} \mathcal{A}_n(z) \mathcal{A}_m(z) = \delta_{nm}, \quad (\text{C12})$$

and satisfy the completeness relation

$$R \sum_n \frac{e^{-\kappa^2 z'^2}}{z'} \mathcal{A}_n(z') \mathcal{A}_n(z) = \delta(z - z'). \quad (\text{C13})$$

The Green's function is a solution of the homogeneous equation

$$[-z^2 \partial_z^2 + (1 + 2\kappa^2 z^2) z \partial_z - q^2 z^2] G(z, z'; q) = -\delta(z - z'), \quad (\text{C14})$$

where $q^2 = -Q^2 \leq 0$. Since the eigenfunctions \mathcal{A}_n form a complete basis we can perform a spectral expansion of the Green's function

$$G(z, z'; q) = R \sum_{nm} a_{nm}(Q) \frac{e^{-\kappa^2 z'^2}}{z'} \mathcal{A}_n(z') \mathcal{A}_m(z). \quad (\text{C15})$$

Substituting the above expression in (C14) and using (C13) the expansion coefficients a_{nm} are determined. We find

$$G(x, x'; q) = \sum_n \frac{\mathcal{A}_n(x') \mathcal{A}_n(x)}{q^2 - M_n^2 + i\epsilon}. \quad (\text{C16})$$

2. Relation with the Results of Grigoryan and Radyushkin

Performing the change of variable $t = \frac{x}{1-x}$ in the integral representation for the bulk-to-boundary propagator (C5), and integrating by parts the resulting expression, there follows

$$J_\kappa(Q, z) = \kappa^2 z^2 \int_0^1 \frac{dx}{(1-x)^2} x^{\frac{Q^2}{4\kappa^2}} e^{-\kappa^2 z^2 x/(1-x)}, \quad (\text{C17})$$

the result found by Grigoryan and Radyushkin in [36]. It was also pointed out in [36] that the integrand in (C17) contains the generating function of the associated Laguerre Polynomials

$$\frac{e^{-\kappa^2 z^2 x/(1-x)}}{(1-x)^{k+1}} = \sum_{n=0}^{\infty} L_n^k(\kappa^2 z^2) x^n, \quad (\text{C18})$$

and thus $J_\kappa(Q, z)$ can be expressed as a sum of poles [36]

$$J_\kappa(Q, z) = 4\kappa^4 z^2 \sum_{n=0}^{\infty} \frac{L_n^1(\kappa^2 z^2)}{Q^2 + M_n^2}, \quad (\text{C19})$$

with M_n^2 given by (C11). The above result also follows immediately from (C16) using the expression

$$J(Q, z) = J(Q, 0) \lim_{z' \rightarrow 0} \frac{R}{z'} e^{-\kappa^2 z'^2} \partial_{z'} G(z, z'; q), \quad (\text{C20})$$

which defines the bulk-to-boundary propagator in terms of the Green's function [38].

Appendix D: Analytic Solution of Structure Functions and Form Factors for Arbitrary Twist in the Soft Wall Model

A string mode Φ_τ in the soft-wall model for $n = 0$

$$\Phi_\tau(z) = \frac{1}{R^{3/2}} \sqrt{\frac{2}{\Gamma(\tau-1)}} \kappa^{\tau-1} z^\tau. \quad (\text{D1})$$

is normalized according to

$$\langle \Phi_\tau | \Phi_\tau \rangle = R^3 \int \frac{dz}{z^3} e^{-\kappa^2 z^2} \Phi_\tau(z)^2 = 1. \quad (\text{D2})$$

Since the field ϕ_τ couples to a local hadronic interpolating operator of twist τ defined at the asymptotic boundary of AdS space, the scaling dimension of Φ_τ is τ .

1. The Structure Function

To compute the structure function $q(x)$

$$\int_0^1 dx q(x) = 1, \quad (\text{D3})$$

corresponding to the the string mode Φ_τ , we substitute the unit operator in the warped geometry of the soft-wall model which follows from (C17) for $Q = 0$

$$\mathbf{1} = \kappa^2 z^2 \int_0^1 \frac{dx}{(1-x)^2} e^{-\kappa^2 z^2 x/(1-x)}, \quad (\text{D4})$$

in the normalization condition (D2): $\langle \Phi_\tau | \mathbf{1} | \Phi_\tau \rangle$. Upon integration over the coordinate z we find

$$q(x) = (\tau-1) (1-x)^{\tau-2}, \quad (\text{D5})$$

which gives a constant x -dependence for a two-parton hadronic bound state.

2. Hadronic Form Factor

Likewise, we can compute the hadronic form factor in the soft wall holographic model

$$F(Q^2) = R^3 \int \frac{dz}{z^3} e^{-\kappa^2 z^2} \Phi_\tau(z) J_\kappa(Q, z) \Phi_\tau(z), \quad (\text{D6})$$

by substituting the integral representation (C17) for $J_\kappa(q, z)$ in (D6) and integrating over the variable z . We find the result

$$F(Q^2) = \int_0^1 dx \rho(x, Q), \quad (\text{D7})$$

where

$$\rho(x, Q) = (\tau - 1) (1 - x)^{\tau - 2} x^{\frac{Q^2}{4\kappa^2}}. \quad (\text{D8})$$

The integral (D7) can be expressed in terms of Gamma functions. The final result for the form factor is

$$F(Q^2) = \Gamma(\tau) \frac{\Gamma\left(1 + \frac{Q^2}{4\kappa^2}\right)}{\Gamma\left(\tau + \frac{Q^2}{4\kappa^2}\right)}. \quad (\text{D9})$$

In the absence of anomalous dimensions the twist is an integer, $\tau = N$, and we can simplify (D9) by using the recurrence formula

$$\Gamma(N+z) = (N-1+z)(N-2+z) \dots (1+z)\Gamma(1+z). \quad (\text{D10})$$

We find

$$F(Q^2) = \frac{1}{1 + \frac{Q^2}{4\kappa^2}}, \quad N = 2, \quad (\text{D11})$$

$$F(Q^2) = \frac{2}{\left(1 + \frac{Q^2}{4\kappa^2}\right)\left(2 + \frac{Q^2}{4\kappa^2}\right)}, \quad N = 3, \quad (\text{D12})$$

$$F(Q^2) = \frac{(N-1)!}{\left(1 + \frac{Q^2}{4\kappa^2}\right)\left(2 + \frac{Q^2}{4\kappa^2}\right) \dots \left(N-1 + \frac{Q^2}{4\kappa^2}\right)}, \quad N, \quad (\text{D13})$$

which is expressed as a $N-1$ product of poles, corresponding to the first $N-1$ states along the vector meson radial trajectory. For large Q^2 it follows that

$$F(Q^2) \rightarrow (N-1)! \left[\frac{4\kappa^2}{Q^2} \right]^{(N-1)}, \quad (\text{D14})$$

and we recover the power law counting rules for hard scattering.

Appendix E: Contributions to Meson Form Factors and Structure Functions at Large Momentum Transfer in AdS/QCD

The form factor of a hadron at large Q^2 arises from the small z kinematic domain in AdS space. According

to the AdS/CFT duality, this corresponds to small distances $x^\mu x_\mu \sim 1/Q^2$ in physical space-time, the domain where the current matrix elements are controlled by the conformal twist-dimension, Δ , of the hadron's interpolating operator. In the case of the front form, where $x^+ = 0$, this corresponds to small transverse separation $x^\mu x_\mu = -\mathbf{x}_\perp^2$.

As we have shown in [39], one can use holography to map the functional form of the string modes $\Phi(z)$ in AdS space to the light front wavefunctions in physical space time by identifying z with the transverse variable $\zeta = \sqrt{\frac{x}{1-x}} |\vec{\eta}_\perp|$. Here $\vec{\eta}_\perp = \sum_{i=1}^{n-1} x_i \mathbf{b}_{\perp i}$ is the weighted impact separation, summed over the impact separation of the spectator constituents. The leading large- Q^2 behavior of form factors in AdS/QCD arises from small $\zeta \sim 1/Q$, corresponding to small transverse separation.

For the case of a meson with two constituents the form factor can be written in terms of an effective light-front transverse density in impact space. From (30)

$$F(q^2) = \int_0^1 dx \int db^2 \tilde{\rho}(x, b, Q), \quad (\text{E1})$$

with $\tilde{\rho}(x, b, Q) = \pi J_0(bQ(1-x)) |\tilde{\psi}(x, b)|^2$ and $b = |\mathbf{b}_\perp|$. The kinematics are illustrated in figure 7 for the case of a meson with two constituents in the soft-wall model

$$\tilde{\psi}_{q\bar{q}/\pi}(x, \mathbf{b}_\perp) = \frac{\kappa}{\sqrt{\pi}} \sqrt{x(1-x)} e^{-\frac{1}{2}\kappa^2 x(1-x)\mathbf{b}_\perp^2}, \quad (\text{E2})$$

where the Gaussian form of the LFWF favors short distance configurations with small $\zeta^2 = b_\perp^2 x(1-x) \sim 1/Q^2$. Since we are mainly interested in studying the contribution from different regions to the form factor at large Q^2 , we have replaced the modified bulk-to-boundary propagator $J_\kappa(Q, z)$ (58) by its large Q^2 form $zQK_1(zQ)$. One sees a shift of the integrand $\tilde{\rho}(x, b, Q)$ toward small $|\mathbf{b}_\perp|$ and small $1-x$ at high Q^2 . A similar behavior is observed for the LFWF obtained from the hard wall model.




FIG. 7: Effective partonic density $\rho(x, b, Q)$ in terms of the longitudinal momentum fraction x , the transverse relative impact variable $b = |\mathbf{b}_\perp|$ and momentum transfer Q for the soft wall model. As Q increases the distribution becomes increasingly important near $x = 1$ and $\mathbf{b}_\perp = 0$. This is illustrated in (a) for $Q = 1$ GeV/c. At very large Q (figure (b)), the distribution is peaked towards $\mathbf{b}_\perp = 0$. The value of κ is 0.375 GeV.

The physical situation in momentum space is somewhat more complicated. The LFWF in \mathbf{k}_\perp space is the Fourier transform

$$\psi(x, k) = 4\pi^{3/2} \int_0^\infty b db J_0(kb) \tilde{\psi}(x, b), \quad (\text{E3})$$

with $k = |\mathbf{k}_\perp|$. Thus

$$\psi_{\bar{q}q/\pi}(x, \mathbf{k}_\perp) = \frac{4\pi}{\kappa \sqrt{x(1-x)}} e^{-\frac{\mathbf{k}_\perp^2}{2\kappa^2 x(1-x)}}, \quad (\text{E4})$$

with exponential fall off at high relative transverse momentum. The Drell-Yan-West light-front expression (26) for the form factor for a two-parton bound state has the form

$$F(q^2) = \int_0^1 dx \int \frac{d^2 \vec{k}_\perp}{16\pi^3} \sum_q e_q \psi_{P'}^*(x, \vec{k}_\perp + (1-x)\vec{q}_\perp) \psi_P(x, \vec{k}_\perp). \quad (\text{E5})$$

The form factor can be written in terms of a density in x, \mathbf{k}_\perp space:

$$F(Q^2) = \int_0^1 dx \int d^2 k \rho(x, k, Q), \quad (\text{E6})$$

where

$$\rho(x, k, Q) = \frac{1}{\kappa^2} \frac{1}{x(1-x)} I_0\left(\frac{kQ}{\kappa^2 x}\right) e^{-\frac{(1-x)Q^2}{2\kappa^2 x}} e^{-\frac{\mathbf{k}_\perp^2}{\kappa^2 x(1-x)}}. \quad (\text{E7})$$

In this case the large Q^2 behavior comes from the regime $1-x \sim \kappa^2/Q^2$ and $k_\perp^2 \sim x(1-x)\kappa^2 \sim \kappa^4/Q^2$. Note that the $x \rightarrow 1$ domain for finite k_\perp (or nonzero quark

mass) implies $x_j = k_j^+/P^+ \rightarrow 0$ for each spectator, a regime of high relative longitudinal momentum since the limit $k^+ \rightarrow 0$ requires $k^z \simeq -k^0$; i.e., $k_z^2 \gg \mathbf{k}_\perp^2$ for each spectator.

Integration over k in equation (E6) gives

$$F(Q^2) = \int_0^1 dx e^{-\frac{(1-x)Q^2}{4\kappa^2 x}}, \quad (\text{E8})$$

which is identical to the result obtained in the transverse impact representation after integration over \mathbf{b}_\perp . The above expression for the form factor in the soft wall model is valid for relatively high Q^2 since we have approximated the bulk-to-boundary propagator by its asymptotic form. In the limit $Q^2 \gg \kappa^2$ we find

$$F_\pi(Q^2) \rightarrow \frac{4\kappa^2}{Q^2}, \quad (\text{E9})$$

the scaling result for a two-constituent meson at large Q^2 found in Appendix D.

The pion structure function $q_\pi(x, Q^2)$ is computed by integrating the square of the pion LFWF (E4) up to the scale Q^2

$$q_\pi(x, Q^2) = \int \frac{d^2 \vec{k}_\perp}{16\pi^3} \left| \psi_{\bar{q}q/\pi}(x, \vec{k}_\perp) \right|^2. \quad (\text{E10})$$

We find

$$q_\pi(x, Q^2) = 1 - e^{-\frac{Q^2}{\kappa^2 x(1-x)}}. \quad (\text{E11})$$

In the large Q^2 limit $q_\pi(x, Q^2 \rightarrow \infty) \equiv q_\pi(x) = 1$, which is the behavior of a strongly coupled theory found in QCD(1+1) [59]. The result coincides with (D5) for $\tau = 2$. Identical results [32] are obtained for the pion in the hard wall model, $q(x) = 1$.

-
- [1] J. M. Maldacena, "The large N limit of superconformal field theories and supergravity," *Adv. Theor. Math. Phys.* **2**, 231 (1998) [*Int. J. Theor. Phys.* **38**, 1113 (1999)] [arXiv:hep-th/9711200].
- [2] J. Polchinski and M. J. Strassler, "Hard scattering and gauge/string duality," *Phys. Rev. Lett.* **88**, 031601 (2002) [arXiv:hep-th/0109174].
- [3] R. A. Janik and R. Peschanski, "High energy scattering and the AdS/CFT correspondence," *Nucl. Phys. B* **565**, 193 (2000) [arXiv:hep-th/9907177].
- [4] S. J. Brodsky and G. F. de Teramond, "Light-front hadron dynamics and AdS/CFT correspondence," *Phys. Lett. B* **582**, 211 (2004) [arXiv:hep-th/0310227].
- [5] R. C. Brower, M. J. Strassler and C. I. Tan, "On the Eikonal Approximation in AdS Space," arXiv:0707.2408 [hep-th] and references therein.
- [6] H. Boschi-Filho and N. R. F. Braga, "Gauge / string duality and scalar glueball mass ratios," *JHEP* **0305**, 009 (2003) [arXiv:hep-th/0212207].
- [7] G. F. de Teramond and S. J. Brodsky, "Baryonic states in QCD from gauge / string duality at large N(c)," arXiv:hep-th/0409074.
- [8] G. F. de Teramond and S. J. Brodsky, "The hadronic spectrum of a holographic dual of QCD," *Phys. Rev. Lett.* **94**, 201601 (2005) [arXiv:hep-th/0501022].
- [9] J. Erlich, E. Katz, D. T. Son and M. A. Stephanov, "QCD and a holographic model of hadrons," *Phys. Rev. Lett.* **95**, 261602 (2005) [arXiv:hep-ph/0501128].
- [10] A. Karch, E. Katz, D. T. Son and M. A. Stephanov, "Linear confinement and AdS/QCD," *Phys. Rev. D* **74**, 015005 (2006) [arXiv:hep-ph/0602229].
- [11] G. F. de Teramond, "Mapping string states into partons: Form factors and the hadron spectrum in AdS/QCD," arXiv:hep-ph/0606143;
- [12] N. Evans and A. Tedder, "Perfecting the ultra-violet of holographic descriptions of QCD," *Phys. Lett. B* **642**, 546 (2006) [arXiv:hep-ph/0609112].
- [13] D. K. Hong, T. Inami and H. U. Yee, "Baryons in AdS/QCD," *Phys. Lett. B* **646**, 165 (2007) [arXiv:hep-ph/0609270].

- [14] P. Colangelo, F. De Fazio, F. Jugeau and S. Nicotri, “On the light glueball spectrum in a holographic description of QCD,” arXiv:hep-ph/0703316.
- [15] H. Forkel, M. Beyer and T. Frederico, “Linear square-mass trajectories of radially and orbitally excited hadrons in holographic QCD,” arXiv:0705.1857 [hep-ph].
- [16] L. Da Rold and A. Pomarol, “Chiral symmetry breaking from five dimensional spaces,” Nucl. Phys. B **721**, 79 (2005) [arXiv:hep-ph/0501218]; “The scalar and pseudoscalar sector in a five-dimensional approach to chiral symmetry breaking,” JHEP **0601**, 157 (2006) [arXiv:hep-ph/0510268].
- [17] J. Hirn and V. Sanz, “Interpolating between low and high energy QCD via a 5D Yang-Mills model,” JHEP **0512**, 030 (2005) [arXiv:hep-ph/0507049]; J. Hirn, N. Rius and V. Sanz, “Geometric approach to condensates in holographic QCD,” Phys. Rev. D **73**, 085005 (2006) [arXiv:hep-ph/0512240].
- [18] K. Ghoroku, N. Maru, M. Tachibana and M. Yahiro, “Holographic model for hadrons in deformed AdS(5) background,” Phys. Lett. B **633**, 602 (2006) [arXiv:hep-ph/0510334].
- [19] H. Boschi-Filho, N. R. F. Braga and C. N. Ferreira, “Static strings in Randall-Sundrum scenarios and the quark anti-quark potential,” Phys. Rev. D **73**, 106006 (2006) [arXiv:hep-th/0512295].
- [20] O. Andreev and V. I. Zakharov, “Heavy-quark potentials and AdS/QCD,” Phys. Rev. D **74**, 025023 (2006) [arXiv:hep-ph/0604204].
- [21] T. Hambye, B. Hassanain, J. March-Russell and M. Schvellinger, “On the Delta(I) = 1/2 rule in holographic QCD,” Phys. Rev. D **74**, 026003 (2006) [arXiv:hep-ph/0512089]; “Four-point functions and Kaon decays in AdS/QCD,” arXiv:hep-ph/0612010.
- [22] C. Csaki and M. Reece, “Toward a systematic holographic QCD: A braneless approach,” JHEP **0705**, 062 (2007) [arXiv:hep-ph/0608266].
- [23] J. Polchinski and M. J. Strassler, “Deep inelastic scattering and gauge/string duality,” JHEP **0305**, 012 (2003) [arXiv:hep-th/0209211].
- [24] R. C. Brower, J. Polchinski, M. J. Strassler and C. I. Tan, “The pomeron and gauge / string duality,” arXiv:hep-th/0603115.
- [25] L. F. Alday and J. Maldacena, “Gluon scattering amplitudes at strong coupling,” arXiv:0705.0303 [hep-th].
- [26] J. Babington, J. Erdmenger, N. J. Evans, Z. Guralnik and I. Kirsch, “Chiral symmetry breaking and pions in non-supersymmetric gauge / gravity duals,” Phys. Rev. D **69**, 066007 (2004) [arXiv:hep-th/0306018].
- [27] I. Kirsch, “Spectroscopy of fermionic operators in AdS/CFT,” JHEP **0609** (2006) 052 [arXiv:hep-th/0607205].
- [28] R. Apreda, J. Erdmenger, D. Lust and C. Sieg, “Adding flavour to the Polchinski-Strassler background,” JHEP **0701**, 079 (2007) [arXiv:hep-th/0610276].
- [29] K. Nawa, H. Suganuma and T. Kojo, “Baryons in Holographic QCD,” Phys. Rev. D **75**, 086003 (2007) [arXiv:hep-th/0612187].
- [30] D. K. Hong, M. Rho, H. U. Yee and P. Yi, “Chiral Dynamics of Baryons from String Theory,” arXiv:hep-th/0701276; “Dynamics of Baryons from String Theory and Vector Dominance,” arXiv:0705.2632 [hep-th].
- [31] H. Hata, T. Sakai, S. Sugimoto and S. Yamato, “Baryons from instantons in holographic QCD,” arXiv:hep-th/0701280.
- [32] A. V. Radyushkin, “Holographic wave functions, meromorphization and counting rules,” Phys. Lett. B **642**, 459 (2006) [arXiv:hep-ph/0605116].
- [33] S. J. Brodsky, “New perspectives for QCD from AdS/CFT,” arXiv:hep-ph/0608005.
- [34] S. J. Brodsky and G. F. de Teramond, “AdS/CFT and QCD,” arXiv:hep-th/0702205.
- [35] H. R. Grigoryan and A. V. Radyushkin, “Form factors and wave functions of vector mesons in holographic QCD,” arXiv:hep-ph/0703069.
- [36] H. R. Grigoryan and A. V. Radyushkin, “Structure of Vector Mesons in Holographic Model with Linear Confinement,” arXiv:0706.1543 [hep-ph].
- [37] For a different but complementary approach see for example H. M. Choi and C. R. Ji, “Conformal symmetry and pion form factor: Soft and hard contributions,” Phys. Rev. D **74**, 093010 (2006) [arXiv:hep-ph/0608148]; J. P. B. de Melo, T. Frederico, E. Pace and G. Salme, “Space-like and time-like pion electromagnetic form factor and Fock state components within the light-front dynamics,” Phys. Rev. D **73**, 074013 (2006) [arXiv:hep-ph/0508001] and references therein.
- [38] S. Hong, S. Yoon and M. J. Strassler, “On the couplings of vector mesons in AdS/QCD,” JHEP **0604**, 003 (2006) [arXiv:hep-th/0409118]; S. Hong, S. Yoon and M. J. Strassler, “On the couplings of the rho meson in AdS/QCD,” arXiv:hep-ph/0501197.
- [39] S. J. Brodsky and G. F. de Teramond, “Hadronic spectra and light-front wavefunctions in holographic QCD,” Phys. Rev. Lett. **96**, 201601 (2006) [arXiv:hep-ph/0602252].
- [40] L. von Smekal, R. Alkofer and A. Hauck, “The infrared behavior of gluon and ghost propagators in Landau gauge QCD,” Phys. Rev. Lett. **79**, 3591 (1997) [arXiv:hep-ph/9705242].
- [41] D. Zwanziger, “Non-perturbative Faddeev-Popov formula and infrared limit of QCD,” Phys. Rev. D **69**, 016002 (2004) [arXiv:hep-ph/0303028].
- [42] R. Alkofer, C. S. Fischer and F. J. Llanes-Estrada, “Vertex functions and infrared fixed point in Landau gauge SU(N) Yang-Mills theory,” Phys. Lett. B **611**, 279 (2005) [arXiv:hep-th/0412330].
- [43] D. Epple, H. Reinhardt and W. Schleifenbaum, “Confining Solution of the Dyson-Schwinger Equations in Coulomb Gauge,” Phys. Rev. D **75**, 045011 (2007) [arXiv:hep-th/0612241].
- [44] A. Deur, V. Burkert, J. P. Chen and W. Korsch, “Experimental determination of the effective strong coupling constant,” arXiv:hep-ph/0509113.
- [45] S. J. Brodsky, S. Menke, C. Merino and J. Rathsman, “On the behavior of the effective QCD coupling $\alpha(\tau)(s)$ at low scales,” Phys. Rev. D **67**, 055008 (2003) [arXiv:hep-ph/0212078].
- [46] S. J. Brodsky, J. R. Pelaez and N. Toumbas, “Testing QCD with hypothetical tau leptons,” Phys. Rev. D **60**, 037501 (1999) [arXiv:hep-ph/9810424].
- [47] S. Furui and H. Nakajima, “Infrared features of unquenched finite temperature lattice Landau gauge QCD,” arXiv:hep-lat/0612009; “Correlation of the ghost and the quark in the lattice Landau gauge QCD,” arXiv:hep-lat/0609024.
- [48] G. P. Lepage and S. J. Brodsky, “Exclusive Processes In Perturbative Quantum Chromodynamics,” Phys. Rev. D

- 22**, 2157 (1980).
- [49] M. Diehl, T. Feldmann, R. Jakob and P. Kroll, “Generalized parton distributions from nucleon form factor data,” *Eur. Phys. J. C* **39**, 1 (2005) [arXiv:hep-ph/0408173].
- [50] S. J. Brodsky and H. J. Lu, “Commensurate scale relations in quantum chromodynamics,” *Phys. Rev. D* **51**, 3652 (1995) [arXiv:hep-ph/9405218].
- [51] S. J. Brodsky, G. T. Gabadadze, A. L. Kataev and H. J. Lu, “The generalized Crewther relation in QCD and its experimental consequences,” *Phys. Lett. B* **372**, 133 (1996) [arXiv:hep-ph/9512367].
- [52] S. J. Brodsky, G. P. Lepage and P. B. Mackenzie, “On The Elimination Of Scale Ambiguities In Perturbative Quantum Chromodynamics,” *Phys. Rev. D* **28**, 228 (1983).
- [53] S. J. Brodsky, M. S. Gill, M. Melles and J. Rathsmann, “An analytic extension of the MS-bar renormalization scheme,” *Phys. Rev. D* **58**, 116006 (1998) [arXiv:hep-ph/9801330].
- [54] M. Binger and S. J. Brodsky, “The form factors of the gauge-invariant three-gluon vertex,” *Phys. Rev. D* **74**, 054016 (2006) [arXiv:hep-ph/0602199].
- [55] J. M. Cornwall and J. Papavassiliou, “Gauge Invariant Three Gluon Vertex in QCD,” *Phys. Rev. D* **40**, 3474 (1989).
- [56] M. Binger and S. J. Brodsky, “Physical renormalization schemes and grand unification,” *Phys. Rev. D* **69**, 095007 (2004) [arXiv:hep-ph/0310322].
- [57] P. A. M. Dirac, “Forms Of Relativistic Dynamics,” *Rev. Mod. Phys.* **21**, 392 (1949).
- [58] S. J. Brodsky, H. C. Pauli and S. S. Pinsky, “Quantum chromodynamics and other field theories on the light cone,” *Phys. Rept.* **301**, 299 (1998) [arXiv:hep-ph/9705477].
- [59] H. C. Pauli and S. J. Brodsky, “Discretized Light Cone Quantization: Solution To A Field Theory In One Space One Time Dimensions,” *Phys. Rev. D* **32**, 2001 (1985); K. Hornbostel, S. J. Brodsky and H. C. Pauli, “Light Cone Quantized QCD in (1+1)-Dimensions,” *Phys. Rev. D* **41**, 3814 (1990).
- [60] S. J. Brodsky and D. S. Hwang, “Exact light-cone wavefunction representation of matrix elements of electroweak currents,” *Nucl. Phys. B* **543**, 239 (1999) [arXiv:hep-ph/9806358].
- [61] S. J. Brodsky, M. Diehl and D. S. Hwang, “Light-cone wavefunction representation of deeply virtual Compton scattering,” *Nucl. Phys. B* **596**, 99 (2001) [arXiv:hep-ph/0009254].
- [62] G. P. Lepage and S. J. Brodsky, “Exclusive Processes In Quantum Chromodynamics: Evolution Equations For Hadronic Wave Functions And The Form-Factors Of Mesons,” *Phys. Lett. B* **87** (1979) 359.
- [63] A. V. Efremov and A. V. Radyushkin, “Factorization And Asymptotical Behavior Of Pion Form-Factor In QCD,” *Phys. Lett. B* **94**, 245 (1980).
- [64] S. J. Brodsky and G. P. Lepage, “Exclusive Processes And The Exclusive Inclusive Connection In Quantum Chromodynamics,” SLAC-PUB-2294
- [65] S. J. Brodsky, D. S. Hwang, B. Q. Ma and I. Schmidt, “Light-cone representation of the spin and orbital angular momentum of relativistic composite systems,” *Nucl. Phys. B* **593**, 311 (2001) [arXiv:hep-th/0003082].
- [66] S. D. Drell and T. M. Yan, “Connection Of Elastic Electromagnetic Nucleon Form-Factors At Large Q^{*2} And Deep Inelastic Structure Functions Near Threshold,” *Phys. Rev. Lett.* **24**, 181 (1970).
- [67] G. B. West, “Phenomenological Model For The Electromagnetic Structure Of The Proton,” *Phys. Rev. Lett.* **24**, 1206 (1970).
- [68] D. E. Soper, “The Parton Model And The Bethe-Salpeter Wave Function,” *Phys. Rev. D* **15**, 1141 (1977).
- [69] In the study of generalized parton distributions (GPDs) the transverse distance between the active quark and the center of mass momentum, $\vec{\eta}_{\perp} = \sum_{j=1}^{n-1} x_j \mathbf{b}_{\perp j}$, is often referred as \mathbf{b}_{\perp} , whereas the internal transverse coordinates, labeled here $\mathbf{b}_{\perp i}$, conjugate to the relative transverse momentum coordinates $\mathbf{k}_{\perp i}$, are labeled $\mathbf{c}_{\perp i}$. See M. Burkardt and D. S. Hwang, “Sivers asymmetry and generalized parton distributions in impact parameter space,” *Phys. Rev. D* **69**, 074032 (2004) [arXiv:hep-ph/0309072].
- [70] S. J. Brodsky and S. D. Drell, “The Anomalous Magnetic Moment And Limits On Fermion Substructure,” *Phys. Rev. D* **22**, 2236 (1980).
- [71] S. J. Brodsky and G. R. Farrar, “Scaling Laws At Large Transverse Momentum,” *Phys. Rev. Lett.* **31**, 1153 (1973); G. P. Lepage and S. J. Brodsky, “Exclusive Processes In Perturbative Quantum Chromodynamics,” *Phys. Rev. D* **22**, 2157 (1980).
- [72] V. A. Matveev, R. M. Muradian and A. N. Tavkhelidze, “Automodellism In The Large - Angle Elastic Scattering And Structure Of Hadrons,” *Lett. Nuovo Cim.* **7**, 719 (1973).
- [73] P. Breitenlohner and D. Z. Freedman, “Stability In Gauged Extended Supergravity,” *Annals Phys.* **144**, 249 (1982).
- [74] It is not possible to obtain physically reasonable light-front wave functions satisfying Neumann boundary conditions at the infrared boundary, $\partial_n \phi(z)|_{z=z_0} = 0$, since hadronic LFWFs are expected to vanish at the confinement radius $1/\Lambda_{\text{QCD}}$ [35].
- [75] R. R. Metsaev, “IIB supergravity and various aspects of light-cone formalism in AdS space-time,” arXiv:hep-th/0002008.
- [76] O. Andreev, “ $1/q^{*2}$ corrections and gauge / string duality,” *Phys. Rev. D* **73** (2006) 107901 [arXiv:hep-th/0603170].
- [77] R. Baldini, S. Dubnicka, P. Gauzzi, S. Pacetti, E. Pasqualucci and Y. Srivastava, “Nucleon time-like form factors below the N anti-N threshold,” *Eur. Phys. J. C* **11**, 709 (1999).
- [78] V. Tadevosyan *et al.* [Jefferson Lab F(pi) Collaboration], “Determination of the pion charge form factor for $Q^2=0.60-1.60 \text{ GeV}^2$,” *Phys. Rev. C* **75**, 055205 (2007) [arXiv:nucl-ex/0607007].
- [79] T. Horn *et al.* [Fpi2 Collaboration], “Determination of the charged pion form factor at $Q^{*2} = 1.60-(\text{GeV}/c)^{*2}$ and $2.45-(\text{GeV}/c)^{*2}$,” *Phys. Rev. Lett.* **97**, 192001 (2006) [arXiv:nucl-ex/0607005].
- [80] M. E. Peskin and D. V. Schroeder, “An Introduction To Quantum Field Theory,” Addison-Wesley Publishing Company (1995).
- [81] In obtaining (76) we have used the isospin charge eigenstate basis $|\pi^{\pm}\rangle = \frac{1}{\sqrt{2}} (|\pi^1\rangle \pm i|\pi^2\rangle)$, and the the relation [80]: $\bar{u}\gamma^{\mu} (1 - \gamma^5) d = j^{\mu 1} + i j^{\mu 2} - j^{\mu 51} - i j^{\mu 52}$.
- [82] S. Eidelman *et al.* [Particle Data Group], “Review of Particle Physics,” *Phys. Lett. B* **592**, 1 (2004).

- [83] E. M. Aitala *et al.* [E791 Collaboration], “Direct measurement of the pion valence quark momentum distribution, the pion light-cone wave function squared,” *Phys. Rev. Lett.* **86**, 4768 (2001) [arXiv:hep-ex/0010043].
- [84] M. Burkardt, “Hadron tomography,” *Int. J. Mod. Phys. A* **21**, 926 (2006) [arXiv:hep-ph/0509316].
- [85] X. d. Ji, “Viewing the proton through ‘color’-filters,” *Phys. Rev. Lett.* **91**, 062001 (2003) [arXiv:hep-ph/0304037].
- [86] S. J. Brodsky, D. Chakrabarti, A. Harindranath, A. Mukherjee and J. P. Vary, “Hadron optics: Diffraction patterns in deeply virtual Compton scattering,” *Phys. Lett. B* **641**, 440 (2006) [arXiv:hep-ph/0604262].
- [87] P. Hoyer, “Comments on the relativity of shape,” *AIP Conf. Proc.* **904**, 65 (2007) [arXiv:hep-ph/0608295].
- [88] S. J. Brodsky, D. Chakrabarti, A. Harindranath, A. Mukherjee and J. P. Vary, “Hadron optics in three-dimensional invariant coordinate space from deeply virtual Compton scattering,” *Phys. Rev. D* **75**, 014003 (2007) [arXiv:hep-ph/0611159].
- [89] M. Abramowitz and I. Stegun, Eds, *Handbook of Mathematical Functions*, Dover, New York (1970), p. 488.
- [90] M. Abramowitz and I. Stegun, Eds, *Handbook of Mathematical Functions*, Dover, New York (1970), p. 505.