

## Search for Lepton Flavor Violating Decays $\tau^\pm \rightarrow \ell^\pm \omega$ ( $\ell = e, \mu$ )

B. Aubert,<sup>1</sup> M. Bona,<sup>1</sup> Y. Karyotakis,<sup>1</sup> J. P. Lees,<sup>1</sup> V. Poireau,<sup>1</sup> X. Prudent,<sup>1</sup> V. Tisserand,<sup>1</sup> A. Zghiche,<sup>1</sup>  
J. Garra Tico,<sup>2</sup> E. Grauges,<sup>2</sup> L. Lopez,<sup>3</sup> A. Palano,<sup>3</sup> M. Pappagallo,<sup>3</sup> G. Eigen,<sup>4</sup> B. Stugu,<sup>4</sup> L. Sun,<sup>4</sup> G. S. Abrams,<sup>5</sup>  
M. Battaglia,<sup>5</sup> D. N. Brown,<sup>5</sup> J. Button-Shafer,<sup>5</sup> R. N. Cahn,<sup>5</sup> R. G. Jacobsen,<sup>5</sup> J. A. Kadyk,<sup>5</sup> L. T. Kerth,<sup>5</sup>  
Yu. G. Kolomensky,<sup>5</sup> G. Kukartsev,<sup>5</sup> D. Lopes Pegna,<sup>5</sup> G. Lynch,<sup>5</sup> T. J. Orimoto,<sup>5</sup> I. L. Osipenko,<sup>5</sup>  
M. T. Ronan,<sup>5,\*</sup> K. Tackmann,<sup>5</sup> T. Tanabe,<sup>5</sup> W. A. Wenzel,<sup>5</sup> P. del Amo Sanchez,<sup>6</sup> C. M. Hawkes,<sup>6</sup> N. Soni,<sup>6</sup>  
A. T. Watson,<sup>6</sup> H. Koch,<sup>7</sup> T. Schroeder,<sup>7</sup> D. Walker,<sup>8</sup> D. J. Asgeirsson,<sup>9</sup> T. Cuhadar-Donszelmann,<sup>9</sup> B. G. Fulsom,<sup>9</sup>  
C. Hearty,<sup>9</sup> T. S. Mattison,<sup>9</sup> J. A. McKenna,<sup>9</sup> M. Barrett,<sup>10</sup> A. Khan,<sup>10</sup> M. Saleem,<sup>10</sup> L. Teodorescu,<sup>10</sup>  
V. E. Blinov,<sup>11</sup> A. D. Bukin,<sup>11</sup> A. R. Buzykaev,<sup>11</sup> V. P. Druzhinin,<sup>11</sup> V. B. Golubev,<sup>11</sup> A. P. Onuchin,<sup>11</sup>  
S. I. Serebnyakov,<sup>11</sup> Yu. I. Skovpen,<sup>11</sup> E. P. Solodov,<sup>11</sup> K. Yu. Todyshev,<sup>11</sup> M. Bondioli,<sup>12</sup> S. Curry,<sup>12</sup> I. Eschrich,<sup>12</sup>  
D. Kirkby,<sup>12</sup> A. J. Lankford,<sup>12</sup> P. Lund,<sup>12</sup> M. Mandelkern,<sup>12</sup> E. C. Martin,<sup>12</sup> D. P. Stoker,<sup>12</sup> S. Abachi,<sup>13</sup>  
C. Buchanan,<sup>13</sup> J. W. Gary,<sup>14</sup> F. Liu,<sup>14</sup> O. Long,<sup>14</sup> B. C. Shen,<sup>14,\*</sup> G. M. Vitug,<sup>14</sup> L. Zhang,<sup>14</sup> H. P. Paar,<sup>15</sup>  
S. Rahatlou,<sup>15</sup> V. Sharma,<sup>15</sup> J. W. Berryhill,<sup>16</sup> C. Campagnari,<sup>16</sup> A. Cunha,<sup>16</sup> B. Dahmes,<sup>16</sup> T. M. Hong,<sup>16</sup>  
D. Kovalskiy,<sup>16</sup> J. D. Richman,<sup>16</sup> T. W. Beck,<sup>17</sup> A. M. Eisner,<sup>17</sup> C. J. Flacco,<sup>17</sup> C. A. Heusch,<sup>17</sup> J. Kroseberg,<sup>17</sup>  
W. S. Lockman,<sup>17</sup> T. Schalk,<sup>17</sup> B. A. Schumm,<sup>17</sup> A. Seiden,<sup>17</sup> M. G. Wilson,<sup>17</sup> L. O. Winstrom,<sup>17</sup> E. Chen,<sup>18</sup>  
C. H. Cheng,<sup>18</sup> B. Echenard,<sup>18</sup> F. Fang,<sup>18</sup> D. G. Hitlin,<sup>18</sup> I. Narsky,<sup>18</sup> T. Piatenko,<sup>18</sup> F. C. Porter,<sup>18</sup>  
R. Andreassen,<sup>19</sup> G. Mancinelli,<sup>19</sup> B. T. Meadows,<sup>19</sup> K. Mishra,<sup>19</sup> M. D. Sokoloff,<sup>19</sup> F. Blanc,<sup>20</sup> P. C. Bloom,<sup>20</sup>  
W. T. Ford,<sup>20</sup> J. F. Hirschauer,<sup>20</sup> A. Kreisel,<sup>20</sup> M. Nagel,<sup>20</sup> U. Nauenberg,<sup>20</sup> A. Olivas,<sup>20</sup> J. G. Smith,<sup>20</sup>  
K. A. Ulmer,<sup>20</sup> S. R. Wagner,<sup>20</sup> J. Zhang,<sup>20</sup> R. Ayad,<sup>21,†</sup> A. M. Gabareen,<sup>21</sup> A. Soffer,<sup>21,‡</sup> W. H. Toki,<sup>21</sup>  
R. J. Wilson,<sup>21</sup> D. D. Altenburg,<sup>22</sup> E. Feltresi,<sup>22</sup> A. Hauke,<sup>22</sup> H. Jasper,<sup>22</sup> J. Merkel,<sup>22</sup> A. Petzold,<sup>22</sup> B. Spaan,<sup>22</sup>  
K. Wacker,<sup>22</sup> V. Klose,<sup>23</sup> M. J. Kobel,<sup>23</sup> H. M. Lacker,<sup>23</sup> W. F. Mader,<sup>23</sup> R. Nogowski,<sup>23</sup> J. Schubert,<sup>23</sup>  
K. R. Schubert,<sup>23</sup> R. Schwierz,<sup>23</sup> J. E. Sundermann,<sup>23</sup> A. Volk,<sup>23</sup> D. Bernard,<sup>24</sup> G. R. Bonneaud,<sup>24</sup> E. Latour,<sup>24</sup>  
V. Lombardo,<sup>24</sup> Ch. Thiebaux,<sup>24</sup> M. Verderi,<sup>24</sup> P. J. Clark,<sup>25</sup> W. Gradl,<sup>25</sup> F. Muheim,<sup>25</sup> S. Playfer,<sup>25</sup>  
A. I. Robertson,<sup>25</sup> J. E. Watson,<sup>25</sup> Y. Xie,<sup>25</sup> M. Andreotti,<sup>26</sup> D. Bettoni,<sup>26</sup> C. Bozzi,<sup>26</sup> R. Calabrese,<sup>26</sup> A. Cecchi,<sup>26</sup>  
G. Cibinetto,<sup>26</sup> P. Franchini,<sup>26</sup> E. Luppi,<sup>26</sup> M. Negrini,<sup>26</sup> A. Petrella,<sup>26</sup> L. Piemontese,<sup>26</sup> E. Prencipe,<sup>26</sup>  
V. Santoro,<sup>26</sup> F. Anulli,<sup>27</sup> R. Baldini-Ferrolì,<sup>27</sup> A. Calcaterra,<sup>27</sup> R. de Sangro,<sup>27</sup> G. Finocchiaro,<sup>27</sup> S. Pacetti,<sup>27</sup>  
P. Patteri,<sup>27</sup> I. M. Peruzzi,<sup>27,§</sup> M. Piccolo,<sup>27</sup> M. Rama,<sup>27</sup> A. Zallo,<sup>27</sup> A. Buzzo,<sup>28</sup> R. Contri,<sup>28</sup> M. Lo Vetere,<sup>28</sup>  
M. M. Macri,<sup>28</sup> M. R. Monge,<sup>28</sup> S. Passaggio,<sup>28</sup> C. Patrignani,<sup>28</sup> E. Robutti,<sup>28</sup> A. Santroni,<sup>28</sup> S. Tosi,<sup>28</sup>  
K. S. Chaisanguanthum,<sup>29</sup> M. Morii,<sup>29</sup> J. Wu,<sup>29</sup> R. S. Dubitzky,<sup>30</sup> J. Marks,<sup>30</sup> S. Schenk,<sup>30</sup> U. Uwer,<sup>30</sup> D. J. Bard,<sup>31</sup>  
P. D. Dauncey,<sup>31</sup> J. A. Nash,<sup>31</sup> W. Panduro Vazquez,<sup>31</sup> M. Tibbetts,<sup>31</sup> P. K. Behera,<sup>32</sup> X. Chai,<sup>32</sup> M. J. Charles,<sup>32</sup>  
U. Mallik,<sup>32</sup> J. Cochran,<sup>33</sup> H. B. Crawley,<sup>33</sup> L. Dong,<sup>33</sup> V. Eyges,<sup>33</sup> W. T. Meyer,<sup>33</sup> S. Prell,<sup>33</sup> E. I. Rosenberg,<sup>33</sup>  
A. E. Rubin,<sup>33</sup> Y. Y. Gao,<sup>34</sup> A. V. Gritsan,<sup>34</sup> Z. J. Guo,<sup>34</sup> C. K. Lae,<sup>34</sup> A. G. Denig,<sup>35</sup> M. Fritsch,<sup>35</sup> G. Schott,<sup>35</sup>  
N. Arnaud,<sup>36</sup> J. Béquilleux,<sup>36</sup> A. D’Orazio,<sup>36</sup> M. Davier,<sup>36</sup> G. Grosdidier,<sup>36</sup> A. Höcker,<sup>36</sup> V. Lepeltier,<sup>36</sup>  
F. Le Diberder,<sup>36</sup> A. M. Lutz,<sup>36</sup> S. Pruvot,<sup>36</sup> P. Roudeau,<sup>36</sup> M. H. Schune,<sup>36</sup> J. Serrano,<sup>36</sup> V. Sordini,<sup>36</sup>  
A. Stocchi,<sup>36</sup> W. F. Wang,<sup>36</sup> G. Wormser,<sup>36</sup> D. J. Lange,<sup>37</sup> D. M. Wright,<sup>37</sup> I. Bingham,<sup>38</sup> J. P. Burke,<sup>38</sup>  
C. A. Chavez,<sup>38</sup> J. R. Fry,<sup>38</sup> E. Gabathuler,<sup>38</sup> R. Gamet,<sup>38</sup> D. E. Hutchcroft,<sup>38</sup> D. J. Payne,<sup>38</sup> K. C. Schofield,<sup>38</sup>  
C. Touramanis,<sup>38</sup> A. J. Bevan,<sup>39</sup> K. A. George,<sup>39</sup> F. Di Lodovico,<sup>39</sup> R. Sacco,<sup>39</sup> G. Cowan,<sup>40</sup> H. U. Flaecher,<sup>40</sup>  
D. A. Hopkins,<sup>40</sup> S. Paramesvaran,<sup>40</sup> F. Salvatore,<sup>40</sup> A. C. Wren,<sup>40</sup> D. N. Brown,<sup>41</sup> C. L. Davis,<sup>41</sup> N. R. Barlow,<sup>42</sup>  
R. J. Barlow,<sup>42</sup> Y. M. Chia,<sup>42</sup> C. L. Edgar,<sup>42</sup> G. D. Lafferty,<sup>42</sup> T. J. West,<sup>42</sup> J. I. Yi,<sup>42</sup> J. Anderson,<sup>43</sup> C. Chen,<sup>43</sup>  
A. Jawahery,<sup>43</sup> D. A. Roberts,<sup>43</sup> G. Simi,<sup>43</sup> J. M. Tuggle,<sup>43</sup> C. Dallapiccola,<sup>44</sup> S. S. Hertzbach,<sup>44</sup> X. Li,<sup>44</sup>  
T. B. Moore,<sup>44</sup> E. Salvati,<sup>44</sup> S. Saremi,<sup>44</sup> R. Cowan,<sup>45</sup> D. Dujmic,<sup>45</sup> P. H. Fisher,<sup>45</sup> K. Koeneke,<sup>45</sup> G. Sciolla,<sup>45</sup>  
M. Spitznagel,<sup>45</sup> F. Taylor,<sup>45</sup> R. K. Yamamoto,<sup>45</sup> M. Zhao,<sup>45</sup> S. E. Mclachlin,<sup>46,\*</sup> P. M. Patel,<sup>46</sup> S. H. Robertson,<sup>46</sup>  
A. Lazzaro,<sup>47</sup> F. Palombo,<sup>47</sup> J. M. Bauer,<sup>48</sup> L. Cremaldi,<sup>48</sup> V. Eschenburg,<sup>48</sup> R. Godang,<sup>48</sup> R. Kroeger,<sup>48</sup>  
D. A. Sanders,<sup>48</sup> D. J. Summers,<sup>48</sup> H. W. Zhao,<sup>48</sup> S. Brunet,<sup>49</sup> D. Côté,<sup>49</sup> M. Simard,<sup>49</sup> P. Taras,<sup>49</sup> F. B. Viaud,<sup>49</sup>  
H. Nicholson,<sup>50</sup> G. De Nardo,<sup>51</sup> F. Fabozzi,<sup>51,¶</sup> L. Lista,<sup>51</sup> D. Monorchio,<sup>51</sup> C. Sciacca,<sup>51</sup> M. A. Baak,<sup>52</sup> G. Raven,<sup>52</sup>  
H. L. Snoek,<sup>52</sup> C. P. Jessop,<sup>53</sup> K. J. Knoepfel,<sup>53</sup> J. M. LoSecco,<sup>53</sup> G. Benelli,<sup>54</sup> L. A. Corwin,<sup>54</sup> K. Honscheid,<sup>54</sup>  
H. Kagan,<sup>54</sup> R. Kass,<sup>54</sup> J. P. Morris,<sup>54</sup> A. M. Rahimi,<sup>54</sup> J. J. Regensburger,<sup>54</sup> S. J. Sekula,<sup>54</sup> Q. K. Wong,<sup>54</sup>

N. L. Blount,<sup>55</sup> J. Brau,<sup>55</sup> R. Frey,<sup>55</sup> O. Igonkina,<sup>55</sup> J. A. Kolb,<sup>55</sup> M. Lu,<sup>55</sup> R. Rahmat,<sup>55</sup> N. B. Sinev,<sup>55</sup> D. Strom,<sup>55</sup> J. Strube,<sup>55</sup> E. Torrence,<sup>55</sup> N. Gagliardi,<sup>56</sup> A. Gaz,<sup>56</sup> M. Margoni,<sup>56</sup> M. Morandin,<sup>56</sup> A. Pompili,<sup>56</sup> M. Posocco,<sup>56</sup> M. Rotondo,<sup>56</sup> F. Simonetto,<sup>56</sup> R. Stroili,<sup>56</sup> C. Voci,<sup>56</sup> E. Ben-Haim,<sup>57</sup> H. Briand,<sup>57</sup> G. Calderini,<sup>57</sup> J. Chauveau,<sup>57</sup> P. David,<sup>57</sup> L. Del Buono,<sup>57</sup> Ch. de la Vaissière,<sup>57</sup> O. Hamon,<sup>57</sup> Ph. Leruste,<sup>57</sup> J. Malclès,<sup>57</sup> J. Ocariz,<sup>57</sup> A. Perez,<sup>57</sup> J. Prendki,<sup>57</sup> L. Gladney,<sup>58</sup> M. Biasini,<sup>59</sup> R. Covarelli,<sup>59</sup> E. Manoni,<sup>59</sup> C. Angelini,<sup>60</sup> G. Batignani,<sup>60</sup> S. Bettarini,<sup>60</sup> M. Carpinelli,<sup>60,\*\*</sup> R. Cenci,<sup>60</sup> A. Cervelli,<sup>60</sup> F. Forti,<sup>60</sup> M. A. Giorgi,<sup>60</sup> A. Lusiani,<sup>60</sup> G. Marchiori,<sup>60</sup> M. A. Mazur,<sup>60</sup> M. Morganti,<sup>60</sup> N. Neri,<sup>60</sup> E. Paoloni,<sup>60</sup> G. Rizzo,<sup>60</sup> J. J. Walsh,<sup>60</sup> J. Biesiada,<sup>61</sup> Y. P. Lau,<sup>61</sup> C. Lu,<sup>61</sup> J. Olsen,<sup>61</sup> A. J. S. Smith,<sup>61</sup> A. V. Telnov,<sup>61</sup> E. Baracchini,<sup>62</sup> F. Bellini,<sup>62</sup> G. Cavoto,<sup>62</sup> D. del Re,<sup>62</sup> E. Di Marco,<sup>62</sup> R. Faccini,<sup>62</sup> F. Ferrarotto,<sup>62</sup> F. Ferroni,<sup>62</sup> M. Gaspero,<sup>62</sup> P. D. Jackson,<sup>62</sup> M. A. Mazzoni,<sup>62</sup> S. Morganti,<sup>62</sup> G. Piredda,<sup>62</sup> F. Polci,<sup>62</sup> F. Renga,<sup>62</sup> C. Voena,<sup>62</sup> M. Ebert,<sup>63</sup> T. Hartmann,<sup>63</sup> H. Schröder,<sup>63</sup> R. Waldi,<sup>63</sup> T. Adye,<sup>64</sup> G. Castelli,<sup>64</sup> B. Franek,<sup>64</sup> E. O. Olaiya,<sup>64</sup> W. Roethel,<sup>64</sup> F. F. Wilson,<sup>64</sup> S. Emery,<sup>65</sup> M. Escalier,<sup>65</sup> A. Gaidot,<sup>65</sup> S. F. Ganzhur,<sup>65</sup> G. Hamel de Monchenault,<sup>65</sup> W. Kozanecki,<sup>65</sup> G. Vasseur,<sup>65</sup> Ch. Yèche,<sup>65</sup> M. Zito,<sup>65</sup> X. R. Chen,<sup>66</sup> H. Liu,<sup>66</sup> W. Park,<sup>66</sup> M. V. Purohit,<sup>66</sup> R. M. White,<sup>66</sup> J. R. Wilson,<sup>66</sup> M. T. Allen,<sup>67</sup> D. Aston,<sup>67</sup> R. Bartoldus,<sup>67</sup> P. Bechtle,<sup>67</sup> R. Claus,<sup>67</sup> J. P. Coleman,<sup>67</sup> M. R. Convery,<sup>67</sup> J. C. Dingfelder,<sup>67</sup> J. Dorfan,<sup>67</sup> G. P. Dubois-Felsmann,<sup>67</sup> W. Dunwoodie,<sup>67</sup> R. C. Field,<sup>67</sup> T. Glanzman,<sup>67</sup> S. J. Gowdy,<sup>67</sup> M. T. Graham,<sup>67</sup> P. Grenier,<sup>67</sup> C. Hast,<sup>67</sup> W. R. Innes,<sup>67</sup> J. Kaminski,<sup>67</sup> M. H. Kelsey,<sup>67</sup> H. Kim,<sup>67</sup> P. Kim,<sup>67</sup> M. L. Kocian,<sup>67</sup> D. W. G. S. Leith,<sup>67</sup> S. Li,<sup>67</sup> S. Luitz,<sup>67</sup> V. Luth,<sup>67</sup> H. L. Lynch,<sup>67</sup> D. B. MacFarlane,<sup>67</sup> H. Marsiske,<sup>67</sup> R. Messner,<sup>67</sup> D. R. Muller,<sup>67</sup> S. Nelson,<sup>67</sup> C. P. O'Grady,<sup>67</sup> I. Ofte,<sup>67</sup> A. Perazzo,<sup>67</sup> M. Perl,<sup>67</sup> T. Pulliam,<sup>67</sup> B. N. Ratcliff,<sup>67</sup> A. Roodman,<sup>67</sup> A. A. Salnikov,<sup>67</sup> R. H. Schindler,<sup>67</sup> J. Schwiening,<sup>67</sup> A. Snyder,<sup>67</sup> D. Su,<sup>67</sup> M. K. Sullivan,<sup>67</sup> K. Suzuki,<sup>67</sup> S. K. Swain,<sup>67</sup> J. M. Thompson,<sup>67</sup> J. Va'vra,<sup>67</sup> A. P. Wagner,<sup>67</sup> M. Weaver,<sup>67</sup> W. J. Wisniewski,<sup>67</sup> M. Wittgen,<sup>67</sup> D. H. Wright,<sup>67</sup> H. W. Wulsin,<sup>67</sup> A. K. Yarritu,<sup>67</sup> K. Yi,<sup>67</sup> C. C. Young,<sup>67</sup> V. Ziegler,<sup>67</sup> P. R. Burchat,<sup>68</sup> A. J. Edwards,<sup>68</sup> S. A. Majewski,<sup>68</sup> T. S. Miyashita,<sup>68</sup> B. A. Petersen,<sup>68</sup> L. Wilden,<sup>68</sup> S. Ahmed,<sup>69</sup> M. S. Alam,<sup>69</sup> R. Bula,<sup>69</sup> J. A. Ernst,<sup>69</sup> B. Pan,<sup>69</sup> M. A. Saeed,<sup>69</sup> S. B. Zain,<sup>69</sup> S. M. Spanier,<sup>70</sup> B. J. Wogland,<sup>70</sup> R. Eckmann,<sup>71</sup> J. L. Ritchie,<sup>71</sup> A. M. Ruland,<sup>71</sup> C. J. Schilling,<sup>71</sup> R. F. Schwitters,<sup>71</sup> J. M. Izen,<sup>72</sup> X. C. Lou,<sup>72</sup> S. Ye,<sup>72</sup> F. Bianchi,<sup>73</sup> F. Gallo,<sup>73</sup> D. Gamba,<sup>73</sup> M. Pelliccioni,<sup>73</sup> M. Bomben,<sup>74</sup> L. Bosisio,<sup>74</sup> C. Cartaro,<sup>74</sup> F. Cossutti,<sup>74</sup> G. Della Ricca,<sup>74</sup> L. Lanceri,<sup>74</sup> L. Vitale,<sup>74</sup> V. Azzolini,<sup>75</sup> N. Lopez-March,<sup>75</sup> F. Martinez-Vidal,<sup>75</sup> D. A. Milanes,<sup>75</sup> A. Oyanguren,<sup>75</sup> J. Albert,<sup>76</sup> Sw. Banerjee,<sup>76</sup> B. Bhuyan,<sup>76</sup> K. Hamano,<sup>76</sup> R. Kowalewski,<sup>76</sup> I. M. Nugent,<sup>76</sup> J. M. Roney,<sup>76</sup> R. J. Sobie,<sup>76</sup> P. F. Harrison,<sup>77</sup> J. Ilic,<sup>77</sup> T. E. Latham,<sup>77</sup> G. B. Mohanty,<sup>77</sup> H. R. Band,<sup>78</sup> X. Chen,<sup>78</sup> S. Dasu,<sup>78</sup> K. T. Flood,<sup>78</sup> J. J. Hollar,<sup>78</sup> P. E. Kutter,<sup>78</sup> Y. Pan,<sup>78</sup> M. Pierini,<sup>78</sup> R. Prepost,<sup>78</sup> S. L. Wu,<sup>78</sup> and H. Neal<sup>79</sup>

(The BABAR Collaboration)

<sup>1</sup>Laboratoire de Physique des Particules, IN2P3/CNRS et Université de Savoie, F-74941 Annecy-Le-Vieux, France

<sup>2</sup>Universitat de Barcelona, Facultat de Física, Departament ECM, E-08028 Barcelona, Spain

<sup>3</sup>Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

<sup>4</sup>University of Bergen, Institute of Physics, N-5007 Bergen, Norway

<sup>5</sup>Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

<sup>6</sup>University of Birmingham, Birmingham, B15 2TT, United Kingdom

<sup>7</sup>Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

<sup>8</sup>University of Bristol, Bristol BS8 1TL, United Kingdom

<sup>9</sup>University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

<sup>10</sup>Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

<sup>11</sup>Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

<sup>12</sup>University of California at Irvine, Irvine, California 92697, USA

<sup>13</sup>University of California at Los Angeles, Los Angeles, California 90024, USA

<sup>14</sup>University of California at Riverside, Riverside, California 92521, USA

<sup>15</sup>University of California at San Diego, La Jolla, California 92093, USA

<sup>16</sup>University of California at Santa Barbara, Santa Barbara, California 93106, USA

<sup>17</sup>University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

<sup>18</sup>California Institute of Technology, Pasadena, California 91125, USA

<sup>19</sup>University of Cincinnati, Cincinnati, Ohio 45221, USA

<sup>20</sup>University of Colorado, Boulder, Colorado 80309, USA

<sup>21</sup>Colorado State University, Fort Collins, Colorado 80523, USA

<sup>22</sup>Universität Dortmund, Institut für Physik, D-44221 Dortmund, Germany

<sup>23</sup>Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany

<sup>24</sup>Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, F-91128 Palaiseau, France

<sup>25</sup>University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom

- <sup>26</sup> *Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy*  
<sup>27</sup> *Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy*  
<sup>28</sup> *Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy*  
<sup>29</sup> *Harvard University, Cambridge, Massachusetts 02138, USA*  
<sup>30</sup> *Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany*  
<sup>31</sup> *Imperial College London, London, SW7 2AZ, United Kingdom*  
<sup>32</sup> *University of Iowa, Iowa City, Iowa 52242, USA*  
<sup>33</sup> *Iowa State University, Ames, Iowa 50011-3160, USA*  
<sup>34</sup> *Johns Hopkins University, Baltimore, Maryland 21218, USA*  
<sup>35</sup> *Universität Karlsruhe, Institut für Experimentelle Kernphysik, D-76021 Karlsruhe, Germany*  
<sup>36</sup> *Laboratoire de l'Accélérateur Linéaire, IN2P3/CNRS et Université Paris-Sud 11, Centre Scientifique d'Orsay, B. P. 34, F-91898 ORSAY Cedex, France*  
<sup>37</sup> *Lawrence Livermore National Laboratory, Livermore, California 94550, USA*  
<sup>38</sup> *University of Liverpool, Liverpool L69 7ZE, United Kingdom*  
<sup>39</sup> *Queen Mary, University of London, E1 4NS, United Kingdom*  
<sup>40</sup> *University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom*  
<sup>41</sup> *University of Louisville, Louisville, Kentucky 40292, USA*  
<sup>42</sup> *University of Manchester, Manchester M13 9PL, United Kingdom*  
<sup>43</sup> *University of Maryland, College Park, Maryland 20742, USA*  
<sup>44</sup> *University of Massachusetts, Amherst, Massachusetts 01003, USA*  
<sup>45</sup> *Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA*  
<sup>46</sup> *McGill University, Montréal, Québec, Canada H3A 2T8*  
<sup>47</sup> *Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy*  
<sup>48</sup> *University of Mississippi, University, Mississippi 38677, USA*  
<sup>49</sup> *Université de Montréal, Physique des Particules, Montréal, Québec, Canada H3C 3J7*  
<sup>50</sup> *Mount Holyoke College, South Hadley, Massachusetts 01075, USA*  
<sup>51</sup> *Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy*  
<sup>52</sup> *NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands*  
<sup>53</sup> *University of Notre Dame, Notre Dame, Indiana 46556, USA*  
<sup>54</sup> *Ohio State University, Columbus, Ohio 43210, USA*  
<sup>55</sup> *University of Oregon, Eugene, Oregon 97403, USA*  
<sup>56</sup> *Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy*  
<sup>57</sup> *Laboratoire de Physique Nucléaire et de Hautes Energies, IN2P3/CNRS, Université Pierre et Marie Curie-Paris6, Université Denis Diderot-Paris7, F-75252 Paris, France*  
<sup>58</sup> *University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*  
<sup>59</sup> *Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy*  
<sup>60</sup> *Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy*  
<sup>61</sup> *Princeton University, Princeton, New Jersey 08544, USA*  
<sup>62</sup> *Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy*  
<sup>63</sup> *Universität Rostock, D-18051 Rostock, Germany*  
<sup>64</sup> *Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom*  
<sup>65</sup> *DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France*  
<sup>66</sup> *University of South Carolina, Columbia, South Carolina 29208, USA*  
<sup>67</sup> *Stanford Linear Accelerator Center, Stanford, California 94309, USA*  
<sup>68</sup> *Stanford University, Stanford, California 94305-4060, USA*  
<sup>69</sup> *State University of New York, Albany, New York 12222, USA*  
<sup>70</sup> *University of Tennessee, Knoxville, Tennessee 37996, USA*  
<sup>71</sup> *University of Texas at Austin, Austin, Texas 78712, USA*  
<sup>72</sup> *University of Texas at Dallas, Richardson, Texas 75083, USA*  
<sup>73</sup> *Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy*  
<sup>74</sup> *Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy*  
<sup>75</sup> *IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain*  
<sup>76</sup> *University of Victoria, Victoria, British Columbia, Canada V8W 3P6*  
<sup>77</sup> *Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom*  
<sup>78</sup> *University of Wisconsin, Madison, Wisconsin 53706, USA*  
<sup>79</sup> *Yale University, New Haven, Connecticut 06511, USA*

(Dated: November 9, 2007)

A search for lepton flavor violating decays of a  $\tau$  to a lighter-mass charged lepton and an  $\omega$  vector meson is performed using  $384.1 \text{ fb}^{-1}$  of  $e^+e^-$  annihilation data collected with the BABAR detector at the Stanford Linear Accelerator Center PEP-II storage ring. No signal is found, and the upper limits on the branching ratios are determined to be  $\mathcal{B}(\tau^\pm \rightarrow e^\pm \omega) < 1.1 \times 10^{-7}$  and  $\mathcal{B}(\tau^\pm \rightarrow \mu^\pm \omega) < 1.0 \times 10^{-7}$  at 90% confidence level.

In the Standard Model (SM) with massless neutrinos, lepton number is conserved separately for each generation. However the discovery of large neutrino mixing [1] requires that lepton flavor violation (LFV) occur, although decays involving charged LFV have not yet been observed. In minimal extensions of the SM that account for neutrino oscillations by the seesaw mechanism of neutrino mass generation, the expected rates of LFV decays are too small to be observable. Thus the observation of neutrinoless decays like  $\tau^\pm \rightarrow \ell^\pm \omega$ , where  $\ell = e$  or  $\mu$ , would be an unambiguous signature of new physics [2], while limits on this process provide constraints on theoretical models.

The search for  $\tau^\pm \rightarrow \ell^\pm \omega$  decays presented here uses data recorded by the *BABAR* detector at the SLAC PEP-II asymmetric-energy  $e^+e^-$  storage ring. The data sample corresponds to an integrated luminosity  $\mathcal{L} = 347.5 \text{ fb}^{-1}$  recorded at an  $e^+e^-$  center-of-mass (c.m.) energy  $\sqrt{s} = 10.58 \text{ GeV}$ , and  $36.6 \text{ fb}^{-1}$  recorded at  $\sqrt{s} = 10.54 \text{ GeV}$ . With an average cross section of  $\sigma_{e^+e^- \rightarrow \tau^+\tau^-} = (0.919 \pm 0.003) \text{ nb}$  [3], this corresponds to a sample of  $3.53 \times 10^8$   $\tau$ -pair events.

The details of the *BABAR* detector are described elsewhere [4]. Charged particles are reconstructed as tracks with a five-layer silicon vertex tracker (SVT) and a 40-layer drift chamber (DCH) inside a 1.5 T solenoidal magnet. An electromagnetic calorimeter (EMC) consisting of 6580 CsI(Tl) crystals is used to identify electrons and photons. A ring-imaging Cherenkov detector (DIRC) is used to identify charged pions and kaons. The flux return of the solenoid, instrumented with resistive plate chambers and limited streamer tubes, is used to identify muons.

The signature of the signal process is the presence of a  $\ell\omega$  pair having an invariant mass consistent with  $m_\tau = 1.777 \text{ GeV}/c^2$  [5] and a total energy equal to  $\sqrt{s}/2$  in the c.m. frame, along with other particles in the event whose properties are consistent with a  $\tau$  decay. Only the dominant decay mode of the  $\omega$  meson ( $\omega \rightarrow \pi^+\pi^-\pi^0$ ) is used in this analysis. The estimation of the background rate in the final sample comes from data only, while samples of Monte Carlo (MC) simulated events are used to obtain the signal reconstruction efficiency, the kinematic distributions of the signal and background events, to optimize the selection criteria and to study systematic uncertainties in the signal efficiency. Control samples with two identified electrons in the event are used to study background contamination from radiative Bhabha scattering, since the relevant cross-section is large, making it impractical to generate a sufficient number of simulated events.

The signal events are simulated with KK2F [6], where one  $\tau$  decays to  $\ell^\pm \omega$  according to two body phase space

and the other  $\tau$  decays according to measured branching fractions [7] simulated with TAUOLA [8]. The  $\mu^+\mu^-$  and  $\tau^+\tau^-$  background processes are generated using KK2F and TAUOLA, and  $q\bar{q}$  processes are generated using the EVTGEN [9] and JETSET [10] packages. The detector response for the MC events is simulated using the GEANT4 package [11]. Radiative corrections for signal and background processes are simulated using PHOTOS [12].

Events with four well-reconstructed tracks and zero total charge are selected, where no track pair is consistent with being a photon conversion in the detector material. Each event is divided into two hemispheres in the c.m. frame by a plane perpendicular to the thrust axis [13], calculated using all reconstructed charged and neutral particles. The events having 3-1 topology, where the signal-side hemisphere contains three tracks and the tag-side contains one track, are selected.

Photon candidates are required to have the measured energy in the EMC greater than 0.1 GeV to reduce the background originating from the  $e^+e^-$  colliding beams in the accelerator beam pipe. Pairs of these photons are combined to form  $\pi^0$  candidates, with the invariant mass in the range  $m(\gamma\gamma) \in [0.115, 0.150] \text{ GeV}/c^2$ . The  $\omega$  mesons are reconstructed from two oppositely charged pion candidates combined with a  $\pi^0$ , with the invariant mass in the range  $m(\pi^+\pi^-\pi^0) \in [0.760, 0.805] \text{ GeV}/c^2$ . In the  $\tau^\pm \rightarrow \ell^\pm \omega$  decay, two of the tracks in the signal-side hemisphere have the same charge. Each of these two same-sign tracks is combined independently with the opposite-sign track and the neutral pion to form two  $\omega$  candidates. The candidate with invariant mass nearest to the nominal  $\omega$  mass [7] is considered to be the signal  $\omega$ . The signal track that is not combined to form the  $\omega$  candidate is required to have a momentum in the laboratory frame greater than 0.5 GeV/c and to be identified as an electron or muon as appropriate, using *BABAR* particle identification techniques [14]. The three tracks in the signal-side hemisphere are fitted to a common vertex, and the photons from the  $\pi^0$  are assumed to originate from this vertex. The reconstructed  $\pi^0$  candidate from the signal  $\tau$  is constrained to the nominal  $\pi^0$  mass [7]. The  $\omega$  candidate is then combined with the lepton track to form the signal  $\tau$  candidate. The signal-side hemisphere may contain up to four photons so as to allow hadronic split-offs from the pion tracks in the EMC. Thus, there may be more than one  $\pi^0$  candidate, resulting in multiple  $\tau$  candidates. In this case, the  $\ell\omega$  combination with invariant mass closest to the nominal  $\tau$  mass is accepted as the signal  $\tau$  candidate. From a sample of  $1.6 \times 10^6$  generated signal MC events, all the reconstructed signal candidates are verified to have correct association with the truth-matched signal  $\tau$  decays.

Signal events are distinguished by two kinematic vari-

ables: the beam-energy constrained mass ( $m_{\text{EC}}$ ) and the energy difference  $\Delta E = E_\ell + E_\omega - \sqrt{s}/2$ , where  $E_\ell$  and  $E_\omega$  are energies of the lepton and the  $\omega$  in the c.m. frame. The  $m_{\text{EC}}$  is calculated from a fit to the reconstructed  $\tau$  candidate decay products with a constraint that the  $\tau$  energy is equal to  $\sqrt{s}/2$  in the c.m. frame. These two variables are weakly correlated and have non-Gaussian tails due to initial and final state radiation. For the signal MC events, the  $m_{\text{EC}}$  distribution peaks at  $m_\tau$ , while the  $\Delta E$  distribution peaks close to but below zero, primarily due to photon energy reconstruction effects producing a small negative offset in the reconstructed  $\tau$  energy. The peak positions ( $\hat{m}_{\text{EC}}$ ,  $\Delta\hat{E}$ ) and standard deviations ( $\sigma(m_{\text{EC}})$ ,  $\sigma(\Delta E)$ ) of the  $m_{\text{EC}}$  and  $\Delta E$  distributions for the reconstructed signal MC events are presented in Table I. To study signal-like events, a large box (LB) is defined in the  $m_{\text{EC}}$  vs.  $\Delta E$  plane as:  $m_{\text{EC}} \in [1.6, 2.0]$  GeV/ $c^2$  and  $\Delta E \in [-0.8, 0.4]$  GeV. To avoid experimenter bias, the number and the properties of data events falling within the  $\pm 3\sigma$  rectangular region in the  $m_{\text{EC}}-\Delta E$  plane, defined as the signal box (SB), are neither used to optimize the selection criteria nor to study systematic effects. The region inside the LB but outside the SB is called grand side band (GSB) and is used for estimation of the background contribution in the SB. The selection requirements are optimized to yield the lowest expected upper limit (UL) [15] derived from the events inside the SB under a background-only hypothesis.

To suppress non- $\tau$  backgrounds with radiation along the beam direction, the polar angle of the missing momentum with respect to beam axis ( $\theta_{\text{miss}}$ ) is required to lie within the detector acceptance:  $\cos\theta_{\text{miss}} \in [-0.76, 0.92]$ . The total c.m. momentum of all tracks and photon candidates in the tag-side must be less than 4.75 GeV/ $c$ .

The events are classified into four different categories depending on tag-side hemisphere properties: the particle identification for the track and the total neutral c.m. energy in the hemisphere ( $\Sigma E_{\text{neutral}}^{\text{CM}}$ ). If the tag-side track is identified as an electron or a muon it is categorized as an  $e$ -tag or a  $\mu$ -tag. Otherwise it is categorized as an  $h$ -tag or a  $\rho$ -tag, depending on whether  $\Sigma E_{\text{neutral}}^{\text{CM}}$  is less than or greater than 0.2 GeV. The  $e$ -tag events are not used in the final selection of  $\tau^\pm \rightarrow e^\pm\omega$  candidates, but are used as the control sample to estimate the Bhabha contribution to this decay mode.

The tag-side hemisphere is expected to contain a SM  $\tau$  decay characterized by the presence of one charged particle and one or two neutrinos. The missing mass due to the undetected neutrino(s) is reconstructed as  $m_\nu^2 = (P_\tau^{\text{tag}} - P_{\text{obs}}^{\text{tag}})^2$ , where  $P_\tau^{\text{tag}}$  and  $P_{\text{obs}}^{\text{tag}}$  are four-momenta in the c.m. frame. The energy and momentum components of  $P_\tau^{\text{tag}}$  are  $(\sqrt{s}/2, \sqrt{(s/4 - m_\tau^2)} \cdot \hat{n})$  where  $\hat{n}$  is the unit vector opposite in direction to the signal-side  $\tau$  momentum and  $P_{\text{obs}}^{\text{tag}}$  is the combined four-momentum in the c.m. frame of all the tracks and photon candidates observed in

the tag-side hemisphere. To reduce non- $\tau$  backgrounds, tag-dependent requirements on  $m_\nu^2$  are applied for the  $\tau^\pm \rightarrow \mu^\pm\omega$  candidates. For  $e$ -tags and  $\mu$ -tags,  $m_\nu^2$  must be in the range  $\in [-2.0, 2.5]$  GeV $^2/c^4$  whereas for  $h$ -tags and  $\rho$ -tags,  $m_\nu^2 \in [-1.2, 2.0]$  GeV $^2/c^4$  and  $m_\nu^2 \in [-2.0, 0.5]$  GeV $^2/c^4$ , respectively. For the  $\tau^\pm \rightarrow \mu^\pm\omega$  candidates, the ratio  $p_{\text{miss}}^T/\sqrt{s}$  in the c.m. frame is required to be greater than 0.061, where  $p_{\text{miss}}^T$  is the component of the missing momentum of the event transverse to the beam direction. For  $\tau^\pm \rightarrow e^\pm\omega$  candidates,  $p_{\text{miss}}^T/\sqrt{s}$  is required to be greater than 0.034.

After applying all the selection criteria for  $\tau^\pm \rightarrow e^\pm\omega$  and  $\tau^\pm \rightarrow \mu^\pm\omega$  decays, the number of data events surviving inside the GSB are 39 and 502, respectively, as shown in Fig. 1. The number of background events in the MC and control samples, in the same region and passing the same set of requirements as data, is  $35 \pm 6$  for  $\tau^\pm \rightarrow e^\pm\omega$  and  $564 \pm 26$  for  $\tau^\pm \rightarrow \mu^\pm\omega$  decay. Out of these MC background events in the  $\tau^\pm \rightarrow e^\pm\omega$  decay, the dominant contributions are from  $q\bar{q}$  (54%) and  $\tau^+\tau^-$  (34%); the rest arise from radiative Bhabha scattering. About 92% of the background in  $\tau^\pm \rightarrow \mu^\pm\omega$  decay is from  $\tau^+\tau^-$  events; within this category, 94% are due to the decay  $\tau^- \rightarrow 2\pi^-\pi^+\pi^0\nu$ , where one of the charged pions is misidentified as a muon. The number of background events in the  $\tau^\pm \rightarrow \mu^\pm\omega$  sample is more than  $\tau^\pm \rightarrow e^\pm\omega$  because of the larger mis-identification rate for a pion track to be identified as a muon than an electron.

The number of expected background events in the SB is extracted from an unbinned maximum likelihood fit to the  $m_{\text{EC}}$  and  $\Delta E$  distributions of data events inside the GSB, using a two-dimensional probability density function (PDF) made of a linear combination of PDFs representing each background component,  $e^+e^-$ ,  $\mu^+\mu^-$ ,  $\tau^+\tau^-$  and  $q\bar{q}$ . The MC event samples are used to determine each component PDF but the one describing radiative Bhabha events, for which a data control sample is used. Each PDF is obtained by interpolating the two-dimensional binned distribution of its respective sample using Gaussian weight terms that are fit with an adaptive kernel estimation procedure [16]. The expected background normalization is fixed to the amount of data events in the GSB, while the relative yields of the different background components are fitted to the background shape. The numbers of background events expected from this fit for various regions around the SB are compared with the numbers observed in Table II, and they confirm that the backgrounds in the data are adequately modeled.

Systematic uncertainties on the signal efficiency and the estimated background are considered in this measurement. The uncertainty due to knowledge of the efficiencies for trigger, for the tracking, and in the beam energy scale and spread is 1.4% for both decay modes. An uncertainty of 2.0% originates from uncertainties on the lepton track momentum and on the photon energy scale




FIG. 1: The selected candidates (dots) inside the large box region of the  $m_{\text{EC}}-\Delta E$  plane for  $\tau^\pm \rightarrow e^\pm\omega$  (left plot) and  $\tau^\pm \rightarrow \mu^\pm\omega$  (right plot) decays. The  $\pm 3\sigma$  signal box is shown by a dashed rectangle. The dark and light shading indicates contours containing 50% and 90% of the selected MC signal events, respectively. The signal box contains 67% of the selected MC signal events for  $\tau^\pm \rightarrow e^\pm\omega$  and 77% for  $\tau^\pm \rightarrow \mu^\pm\omega$  decay.

TABLE I: The peak positions and standard deviations of the  $m_{\text{EC}}$  and  $\Delta E$  distributions, obtained from the fit to the signal MC events. Also shown are the reconstruction efficiencies ( $\varepsilon$ ), the number of expected background (exp.) events and the observed (obs.) events inside the signal box, and the resulting upper limits at 90% confidence level (C.L.) including the systematic uncertainties.

Decay modes	$\hat{m}_{\text{EC}}$	$\sigma(m_{\text{EC}})$	$\Delta \hat{E}$	$\sigma(\Delta E)$	$\varepsilon$ (%)	SB events		UL ( $\times 10^{-7}$ )	
	MeV/ $c^2$	MeV/ $c^2$	MeV	MeV		exp.	obs.	exp.	obs.
$\tau^\pm \rightarrow e^\pm\omega$	$1777.4 \pm 0.1$	$6.8 \pm 0.1$	$-14.4 \pm 0.3$	$32.2 \pm 0.3$	$2.96 \pm 0.13$	$0.35 \pm 0.06$	0	1.4	1.1
$\tau^\pm \rightarrow \mu^\pm\omega$	$1777.7 \pm 0.1$	$6.4 \pm 0.1$	$-11.2 \pm 0.2$	$30.9 \pm 0.3$	$2.56 \pm 0.16$	$0.73 \pm 0.03$	0	1.7	1.0

TABLE II: The expected number of background events obtained from the fit to  $m_{\text{EC}} - \Delta E$  distributions within the  $\pm(3-5)\sigma$ ,  $\pm(5-7)\sigma$ ,  $\pm(7-9)\sigma$ ,  $\pm(9-11)\sigma$  and the combined  $\pm(3-11)\sigma$  nested rectangular regions centered around the signal box. Also shown are the number of observed events inside the corresponding regions.

Decay modes	# of events	$\pm(3-5)\sigma$	$\pm(5-7)\sigma$	$\pm(7-9)\sigma$	$\pm(9-11)\sigma$	$\pm(3-11)\sigma$
$\tau^\pm \rightarrow e^\pm\omega$	expected	$0.6 \pm 0.1$	$1.0 \pm 0.2$	$1.4 \pm 0.2$	$1.9 \pm 0.3$	$4.9 \pm 0.8$
	observed	0	0	1	2	3
$\tau^\pm \rightarrow \mu^\pm\omega$	expected	$1.9 \pm 0.1$	$3.9 \pm 0.2$	$6.7 \pm 0.3$	$12.1 \pm 0.5$	$24.6 \pm 1.1$
	observed	2	3	7	10	22

and resolution, which affect the position and spread of the  $\Delta E$  and  $m_{\text{EC}}$  distributions. There is a 3.3% uncertainty in the  $\pi^0$  reconstruction efficiency, the uncertainty in lepton identification is 1.1% for electrons and 4.5% for muons, and there is a 1% uncertainty on the number of  $\tau$  pairs produced. After combining these individual contributions in quadrature, the total systematic uncertainty on efficiency is 4.4% for  $\tau^\pm \rightarrow e^\pm\omega$  and 6.2% for  $\tau^\pm \rightarrow \mu^\pm\omega$ . The uncertainties on background estimation are determined by the background fit errors. The

uncertainty due to MC statistics is negligible.

The signal is simulated according to the two body phase space, i.e. with a uniform distribution of the cosine of the helicity angle with respect to the  $\tau$  spin. Since  $\tau$  pairs are produced with spin correlation, the event selection efficiency may be sensitive to the helicity angle distribution of the  $\tau^\pm \rightarrow \ell^\pm\omega$  decay, which depends on the model of the LFV interaction [17]. This effect is simulated by weighting the generated events to match the helicity angle distributions of both  $V - A$  and  $V + A$  in-

interactions and its consequences on the measured upper limit is found to be negligible.

The upper limits for  $\tau^\pm \rightarrow \ell^\pm \omega$  decays are calculated using  $\mathcal{B}_{UL}^{90} = N_{UL}^{90}/(2\mathcal{L}\sigma_{\tau\tau}\mathcal{B}\varepsilon)$ , where  $N_{UL}^{90}$  is the 90% C.L. upper limit on the number of signal events inside the SB,  $\mathcal{B}$  is the branching fraction [7] of the decay  $\omega \rightarrow \pi^+\pi^-\pi^0$  ( $\pi^0 \rightarrow \gamma\gamma$ ) and  $\varepsilon$  is the reconstruction efficiency of the signal decay mode under consideration. The expected and observed upper limits, including all contributions from systematic uncertainties, are calculated using the technique of Cousins and Highland [18] with the implementation of Barlow [19]. No signal is found, and the upper limits on the branching ratios are determined to be  $\mathcal{B}(\tau^\pm \rightarrow e^\pm \omega) < 1.1 \times 10^{-7}$  and  $\mathcal{B}(\tau^\pm \rightarrow \mu^\pm \omega) < 1.0 \times 10^{-7}$  at 90% confidence level, as shown in Table I.

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), MEC (Spain), and STFC (United Kingdom). Individuals have received support from the Marie Curie EIF (European Union) and the A. P. Sloan Foundation.

---

\* Deceased

† Now at Temple University, Philadelphia, Pennsylvania 19122, USA

‡ Now at Tel Aviv University, Tel Aviv, 69978, Israel

§ Also with Università di Perugia, Dipartimento di Fisica, Perugia, Italy

¶ Also with Università della Basilicata, Potenza, Italy

\*\* Also with Università di Sassari, Sassari, Italy

- [1] B. T. Cleveland *et al.*, *Astrophys. J.* **496**, 505 (1998); Y. Fukuda *et al.* [Super-Kamiokande Collab.], *Phys. Rev. Lett.* **81**, 1562 (1998); Q. R. Ahmad *et al.* [SNO Collab.], *Phys. Rev. Lett.* **89**, 011301 (2002); M. H. Ahn *et al.* [K2K Collab.], *Phys. Rev. Lett.* **90**, 041801 (2003).
- [2] A. Ilakovac, *Phys. Rev. D* **62**, 036010 (2000); T. Fukuyama, A. Ilakovac and T. Kikuchi, arXiv:hep-ph/0506295; A. Ilakovac, B. A. Kniehl and A. Pilaftsis, *Phys. Rev. D* **52**, 3993 (1995).
- [3] S. Banerjee, B. Pietrzyk, J. M. Roney and Z. Was, arXiv:0706.3235 [hep-ph], submitted to *Phys. Rev. D*.
- [4] B. Aubert *et al.* [*BABAR* Collab.], *Nucl. Instrum. Methods Phys. Res., Sect. A* **479**, 1 (2002).
- [5] J. Z. Bai *et al.* [BES Collab.], *Phys. Rev. D* **53**, 20 (1996).
- [6] B. F. Ward, S. Jadach, and Z. Was, *Nucl. Phys. Proc. Suppl.* **116**, 73 (2003).
- [7] W. M. Yao *et al.* [Particle Data Group], *J. Phys. G* **33**, 1 (2006).
- [8] S. Jadach, Z. Was, R. Decker, and J. H. Kuhn, *Comput. Phys. Commun.* **76**, 361 (1993).
- [9] D. J. Lange, *Nucl. Instrum. Methods Phys. Res., Sect. A* **462**, 152 (2001).
- [10] T. Sjöstrand, *Comput. Phys. Commun.* **82**, 74 (1994).
- [11] S. Agostinelli *et al.* [GEANT4 Collab.], *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
- [12] P. Golonka and Z. Was, *Eur. Phys. J. C* **45**, 97 (2006).
- [13] S. Brandt *et al.*, *Phys. Lett.* **12**, 57 (1964); E. Farhi, *Phys. Rev. Lett.* **39**, 1587 (1977).
- [14] B. Aubert *et al.* [*BABAR* Collab.], *Phys. Rev. D* **66**, 032003 (2002).
- [15] G. J. Feldman and R. D. Cousins, *Phys. Rev. D* **57**, 3873 (1998).
- [16] K. Cranmer, *Comput. Phys. Commun.* **136**, 198 (2001).
- [17] R. Kitano and Y. Okada, *Phys. Rev. D* **63**, 113003 (2001).
- [18] R. D. Cousins and V. L. Highland, *Nucl. Instrum. Methods Phys. Res., Sect. A* **320**, 331 (1992).
- [19] R. Barlow, *Comput. Phys. Commun.* **149**, 97 (2002).