

Universal entanglement entropy in 2D conformal quantum critical points

Benjamin Hsu,¹ Michael Mulligan,² Eduardo Fradkin,¹ and Eun-Ah Kim³

¹*Department of Physics, University of Illinois at Urbana-Champaign, Illinois 61801-3080, USA*

²*Department of Physics and SLAC, Stanford University, Stanford, California 94305, USA*

³*Department of Physics, Cornell University, Ithaca, New York 14853, USA*

(Dated: November 30, 2008)

We study the scaling behavior of the entanglement entropy of two dimensional conformal quantum critical systems, *i.e.* systems with scale invariant wave functions. They include two-dimensional generalized quantum dimer models on bipartite lattices and quantum loop models, as well as the quantum Lifshitz model and related gauge theories. We show that, under quite general conditions, the entanglement entropy of a large and simply connected sub-system of an infinite system with a smooth boundary has a universal finite contribution, as well as scale-invariant terms for special geometries. The universal finite contribution to the entanglement entropy is computable in terms of the properties of the conformal structure of the wave function of these quantum critical systems. The calculation of the universal term reduces to a problem in boundary conformal field theory

PACS numbers: 03.67.Mn, 11.25.Hf

I. INTRODUCTION

The non-local correlations of a quantum mechanical system are encoded in the behavior of the entanglement properties of its wave functions. A pure quantum state of a bipartite system $A \cup B$ defines a mixed state in the observed region A obtained from tracing out the degrees of freedom in the unobserved region B . The non-local correlations connecting regions A and B are encoded in the behavior of the von Neumann entanglement entropy, $S = -\text{Tr} \rho_A \ln \rho_A$, where ρ_A is the reduced density matrix of region A . The entanglement entropy of a local quantum field theory, relativistic or not is known to exhibit an “area law” scaling of the form $S \sim \mu \ell^{D-1}$ in spatial dimensions $D > 1$ where μ is a non-universal coefficient^{1,2}. There has been growing interest in the scaling behavior of the entanglement entropy at quantum critical points and in topological phases. The entanglement entropy of quantum critical systems in $D > 1$ should contain universal subleading terms, whose structure for a general quantum critical system is not yet known.

The scaling behavior of the entanglement entropy has only been studied in detail in quantum critical systems in $D = 1$ space dimension. Such systems are described by a $(1+1)$ -dimensional conformal field theory (CFT). In a $1+1$ -dimensional CFT, the entanglement entropy of a subsystem A of linear size ℓ of an otherwise infinite system (*i.e.* of linear size $L \rightarrow \infty$) obeys a logarithmic scaling law,³⁻⁷ $S \sim \frac{c}{3} \ln(\frac{\ell}{a}) + \dots$, where c is the *central charge* of the CFT, and a is the short distance cutoff. There has been a number of studies on topics related to this 1D logarithmic scaling form. For instance, a possible connection between this result and gravitational physics was suggested⁸. A similar logarithmic scaling behavior was found at infinite disorder fixed points of 1D random spin chains^{9,10}. The quantum entanglement of quantum impurity systems has also been studied.¹¹⁻¹⁵

In this paper, we consider the universal scaling form of the entanglement entropy at 2D conformal quantum

critical points (QCP) – two-dimensional quantum critical systems with scale-invariant many body wave functions. At a 2D conformal QCP, equal-time correlators of local operators coincide with the correlation functions of an appropriate 2D *classical* system at criticality (which is described by an Euclidean 2D CFT)¹⁶. The entanglement entropy of 2D conformal QCPs was first considered in Ref.[17], where a scaling form was found: $S = \mu \ell - \frac{c}{6}(\Delta\chi) \ln(\ell/a) + \dots$, where c is the central charge of the 2D Euclidean CFT associated with the norm squared of the wave function and $\Delta\chi$ is the change of the Euler characteristic χ , $\Delta\chi = \chi_{A \cup B} - \chi_A - \chi_B$. Notice for a region $A \subset B$ with a smooth boundary, $\Delta\chi = 0$ and hence the logarithmic term vanishes. Hence, if region A has a smooth boundary, there is no universal logarithmic term. In this case, we will show that instead there is a *finite*, $\mathcal{O}(1)$, *universal term* γ_{QCP} in the entanglement entropy at these quantum critical points, *i.e.*

$$S_{QCP} = \mu \ell + \gamma_{QCP} + \dots \quad (1.1)$$

Through explicit calculations and using general arguments based on CFT, we will show that γ_{QCP} has a topological meaning in the sense that it is determined by the contributions of the winding modes of the underlying CFT.

In a topological phase in 2D, the entanglement entropy scales as^{18,19}

$$S_{\text{topo}} = \alpha \ell - \gamma_{\text{topo}} + \mathcal{O}(\ell^{-1}), \quad (1.2)$$

where α is a non-universal coefficient and γ_{topo} , the *topological entanglement entropy*, is a topological invariant, the logarithm of the so-called total quantum dimension \mathcal{D} of the underlying topological field theory describing the topological phase.^{18,19} Topological phases have non-trivial ground state degeneracies on surfaces of non-trivial topology. The topological entanglement entropy γ_{topo} also depends on the global topology of the manifold, and on surfaces with non-trivial topology, on the degenerate ground state on that surface.²⁰

Although superficially similar, the finite universal contributions to the entanglement entropy in topological phases and conformal quantum critical points, γ_{topo} and γ_{QCP} , have a different origin and structure. In the case of a topological phase, γ_{topo} is in general determined by the modular S -matrix of the topological field theory of the topological phase.^{18–20} This modular S -matrix governs the transformation properties of the (degenerate) ground states of the topological phase on a torus under modular transformations, $\tau \rightarrow -1/\tau$, where τ is the modular parameter of the torus.²¹ However, we show below that for a general conformal quantum critical point, whose ground state wave function is given by the Gibbs weights of a Euclidean rational unitary CFT, the universal term γ_{QCP} is determined by the modular S -matrix associated with the norm squared of the wave function. Thus, the modular S -matrix of the topological phase and that of the wave functions of 2D conformal quantum critical points have a conceptually different origin. In particular, in all the cases we checked here, γ_{QCP} and γ_{topo} contribute with opposite signs to their respective entanglement entropies, as implied by the conventions we used in Eq.(1.1) and Eq.(1.2).

We will show that, when the logarithmic terms in the entanglement entropy cancel, the finite terms γ_{QCP} are universal and are determined not only by the central charge but also by the restrictions on the states imposed by the compactification conditions. Furthermore, the form of the result for the entanglement entropy of Eq.(2.12) implies a connection with boundary CFT, as developed by Cardy.^{22,23} Thus, in addition of it being determined by the central charge c , it must also depend on the operator content of the CFT. For the same reason, the structure of Eq.(2.12) also suggests a direct connection between this problem and the Affleck-Ludwig boundary entropy of 1D quantum CFTs.²⁴

The paper is organized as follows. In Section II we apply this approach first to the simpler case of the quantum Lifshitz model (and the related QDMs) on planar, cylindrical and toroidal geometries. These results apply to the QCPs of (generalized) quantum dimer model on bipartite lattices^{25–30} and in quantum eight-vertex models¹⁶. Through explicit calculations for various geometries, we show that that, when the logarithmic terms in the entanglement entropy cancel, and that the subleading finite terms γ_{QCP} are universal, determined not only by the central charge but also by the restrictions imposed by the compactification conditions. In Section III we generalize this result to all 2D conformal QCPs whose scale-invariant wave functions have norms that are the partition functions of 2D Euclidean Rational CFTs (RCFT), CFTs with a finite number of primary fields^{31,32}. More specifically, we show that the finite term in the entanglement entropy of the 2D wave function is determined by the change of the Affleck-Ludwig boundary entropy of the 1D CFT – a quantity determined by the modular S -matrix of the associated CFT and by the coefficients in the fusion rules. We also discuss specific examples of

this class including 2D quantum loop models³³ which, with the naive inner product, are known to be quantum critical.^{34,35} We also briefly discuss the quantum net models.^{33–36} In Section IV we conclude with a summary and a discussion on open questions. In particular, we comment on the implications of our results to the nature of related topological phases.

II. QUANTUM LIFSHITZ MODEL UNIVERSALITY CLASS

The quantum Lifshitz model¹⁶ (QLM) in two space dimensions is defined by the following Hamiltonian with an arbitrary parameter k :

$$H = \int d^2x \left[\frac{\Pi^2}{2} + \frac{1}{2} \left(\frac{k}{4\pi} \right)^2 (\nabla^2 \phi)^2 \right], \quad (2.1)$$

where ϕ is a scalar field $\Pi = \dot{\phi}$ is its canonical momentum conjugate to ϕ . The QLM Hamiltonian Eq.(2.1) defines a class of QCP's with dynamic critical exponent $z = 2$, and a continuous parameter k .

This remarkable property of the model is evident in the exactly known wave function for the ground state $|GS\rangle$ which is a superposition of all field configurations $\phi(x, y)$ with the configuration dependent weight¹⁶:

$$\Psi_{GS}[\phi] = \langle [\phi] | GS \rangle = \frac{1}{\sqrt{Z}} e^{-S[\phi]/2}, \quad (2.2)$$

with

$$S[\phi] = \int d^2x \frac{k}{4\pi} \left(\vec{\nabla} \phi(x) \right)^2 \quad (2.3)$$

and the norm squared of the state

$$Z = \|\Psi_{GS}\|^2 = \int D\phi e^{-S[\phi]}. \quad (2.4)$$

Notice Z is identical to the partition function for the Gaussian model, which defines free boson Euclidean CFT³⁷, albeit with the “stiffness” k . Hence Eq.(2.1) defines an infinite class of 2D conformal QCP's all associated with free boson CFTs.

The QLM can be viewed low energy effective field theory capturing universal aspects of various microscopic lattice models with ϕ playing the role of coarse grained height field^{16,38,39} with the “stiffness” k determined by the appropriate “microscopic” coupling constants^{16,29}. For such mapping to work, the constraints of the lattice models should be built in through compactification of the boson field ϕ by demanding all physical operators to be invariant under the shift of $\phi \rightarrow \phi + 2\pi r$ or equivalently all physical operators to take the form of vertex operators $e^{in\phi/r}$ for integer n . In subsection II C we will discuss specific examples of this mapping corresponding to particular values of k using the convention of fixing

$r = 1$. The examples will include so-called Rokhsar-Kivelson point (RK) of the quantum dimer model²⁵ and its generalizations^{30,40,41} and the *quantum* eight-vertex model¹⁶ special choices of the Baxter weight⁴². Since k can be varied in the QLM, this theory has an exactly *marginal* operator, resulting in continuously varying critical exponents (scaling dimensions) of the allowed (vertex) operators.⁴¹

A. Entanglement entropy and partition functions for 2D conformal QCPs

To investigate the universal finite terms in the entanglement entropy at 2D conformal QCPs, we will rely on the approach described in the work of Fradkin and Moore.¹⁷ They showed that $\text{tr} \rho_A^n$, where ρ_A is the (normalized) reduced density matrix of a region A , with $A \subset B$ separated by the boundary Γ , for the ground state Ψ_0 on $A \cup B$, is given by

$$\text{tr} \rho_A^n = \frac{Z_n}{Z^n} = \left(\frac{Z_A Z_B}{Z_{A \cup B}} \right)^{n-1}. \quad (2.5)$$

Here Z_n is the partition function of n copies of the equivalent 2D classical statistical mechanical system satisfying the constraint that their degrees of freedom are identified on the boundary Γ , and Z^n is the partition function for n decoupled systems. The partition functions on the r.h.s of Eq.(2.5) are $Z_A = \|\Psi_0^A\|^2$ with support on region A and $\|\Psi_0^B\|^2$ with support in region B , both satisfying generalized Dirichlet (*i.e.* fixed) boundary conditions on Γ of A and B , and $Z_{A \cup B} = \|\Psi_0\|^2$ is the norm squared for the full system. The entanglement entropy S is then obtained by an analytic continuation in n ,

$$\begin{aligned} S &= -\text{tr}(\rho_A \ln \rho_A) \\ &= -\lim_{n \rightarrow 1} \frac{\partial}{\partial n} \text{tr} \rho_A^n \\ &= -\log \left(\frac{Z_A Z_B}{Z_{A \cup B}} \right) \end{aligned} \quad (2.6)$$

Hence, the computation of the entanglement entropy is reduced to the computation of a ratio of partition functions in a 2D classical statistical mechanical problem, an Euclidean CFT in the case of a critical wave function, each satisfying specific boundary conditions.

In order to construct $\text{tr} \rho_A^n$, we need an expression for the matrix elements of the reduced density matrix $\langle \phi^A | \rho_A | \phi'^A \rangle$. Since the ground state wave function Eqs.(2.2) and (2.3) is a local function of the field $\phi(x)$, a general matrix element of the reduced density matrix is a trace of the density matrix of the pure state $\Psi_{GS}[\phi]$ over the degrees of freedom of the “unobserved” region B , denoted by $\phi^B(x)$. Hence the matrix elements of ρ_A

take the form

$$\begin{aligned} \langle \phi^A | \rho_A | \phi'^A \rangle &= \\ \frac{1}{Z} \int [D\phi^B] e^{-\left(\frac{1}{2} S^A(\phi^A) + \frac{1}{2} S^A(\phi'^A) + S^B(\phi^B) \right)}, \end{aligned} \quad (2.7)$$

where the degrees of freedom satisfy the *boundary condition* at the common boundary Γ :

$$BC_\Gamma : \quad \phi^B|_\Gamma = \phi^A|_\Gamma = \phi'^A|_\Gamma. \quad (2.8)$$

Proceeding with the computation of $\text{tr} \rho_A^n$, it is immediate to see that the matrix product requires the condition $\phi_i^A = \phi'_{i-1}^A$ for $i = 1, \dots, n$, and $\phi_n^A = \phi_1^A$ from the trace condition. Hence, $\text{tr} \rho_A^n$ takes the form

$$\begin{aligned} \text{tr} \rho_A^n &\equiv \frac{Z_n}{Z^n} \\ &= \frac{1}{Z^n} \int_{BC_\Gamma} \prod_i D\phi_i^A D\phi_i^B e^{-\sum_{i=1}^n (S(\phi_i^A) + S(\phi_i^B))} \end{aligned} \quad (2.9)$$

subject to the boundary condition BC_Γ of Eq.(2.8). Notice that the numerator, Z_n is the partition function on n systems whose degrees of freedom are identified in Γ but are otherwise independent. Also notice the absence of the factors of $1/2$ in the exponentials of Eq.(2.9).

The other important consideration is that the compactification condition requires that two fields that differ by $2\pi r$ be equivalent. Hence, the boundary condition of Eq.(2.8) is defined *modulo* $2\pi r$. (Equivalently, the proper form of the degrees of freedom is $e^{i\phi}$.) This means that one can alternatively define Z_n as a partition function for n systems which are decoupled *in the bulk* but have a boundary coupling of the form (in the limit $\lambda_\Gamma \rightarrow \infty$, which enforces the boundary condition)

$$S_\Gamma = -\oint_\Gamma \lambda_\Gamma \sum_{i=1}^n \cos(\phi_i - \phi_{i+1}). \quad (2.10)$$

Here the fields ϕ_i extend over the entire region $A \cup B$. Thus, this problem maps onto a boundary CFT for a system with n “replicas” coupled only through the boundary condition on the closed contour Γ , the boundary between the A and B regions.

For the special case of the free scalar field, one can simplify this further by taking linear combinations of the replica fields. Then the condition that the scalar fields ϕ_i agree with each other on Γ can be satisfied by forming $n-1$ relative coordinates $\varphi_i \equiv \phi_i - \phi_{i+1}$ ($i = 1, \dots, n-1$) that vanish (*mod* $2\pi r$) on Γ , and one “center of mass coordinate” field $\phi \equiv \frac{1}{\sqrt{n}} \sum_{i=1}^n \phi_i$ that is unaffected by the boundary Γ (reflecting the fact that nothing physical takes place at Γ). Hence, the computation of $\text{tr} \rho_A^n$ reduces to the product of two partition functions:

1. The partition function for the “center of mass” field ϕ ; since ϕ does not see the boundary Γ , this is just the partition function $Z_{A \cup B}$ for a single field in the entire system.
2. The partition function for the $n - 1$ fields φ_i which are independent from each other and vanish (*mod* $2\pi r$) on Γ . We denote this by $(Z_\Gamma^D)^{n-1}$. However, the fields φ_i on the A and B regions are effectively decoupled from each other. Hence, this partition function further factorizes to $Z_\Gamma^D = Z_A^D Z_B^D$, where Z_A^D and Z_B^D are the partition functions for a single field ϕ on A and B respectively, satisfying in each case Dirichlet (fixed) boundary conditions (*mod* $2\pi r$) at their common boundary Γ .

Thus, we can write the trace $\text{tr}\rho_A^n$ as

$$\text{tr}\rho_A^n = \frac{(Z_\Gamma^D)^{n-1} Z_{A \cup B}}{Z_{A \cup B}^n} = \left(\frac{Z_\Gamma^D}{Z_F}\right)^{n-1} = \left(\frac{Z_A^D Z_B^D}{Z_{A \cup B}}\right)^{n-1}. \quad (2.11)$$

Here the denominator factor, $Z_{A \cup B}^n$ comes from the normalization factors, and represents the partition function over the entire system. The entanglement entropy is then¹⁷

$$S = -\log Z_A^D - \log Z_B^D + \log Z_{A \cup B} \equiv F_A^D + F_B^D - F_{A \cup B}, \quad (2.12)$$

which, as indicated in the r.h.s of Eq. (??) reduces to the computation of the free energies F_A^D , F_B^D and $F_{A \cup B}$, for the equivalent 2D Euclidean CFT on regions A and B , each satisfying Dirichlet (fixed) boundary conditions on the common boundary Γ , and on the full system, $A \cup B$, respectively.

The behavior of the free energy of a CFT as a function of the system size ℓ has been studied in detail. The divergent terms, as $\ell \rightarrow \infty$, have the form^{43–45}

$$F(\ell) = f_0 \ell^2 + \sigma \ell - \frac{c}{6} \chi \ln\left(\frac{\ell}{a}\right) + \mathcal{O}(1) \quad (2.13)$$

provided the boundary Γ is smooth (and differentiable). Here, f_0 and σ are two non-universal quantities, and a is the short-distance cutoff; c and χ are, respectively, the central charge of the CFT and the Euler characteristic of the manifold. It follows from this result that the entanglement entropy for region A takes the form¹⁷

$$S = \alpha \ell - \frac{c}{6} (\Delta\chi) \ln\left(\frac{\ell}{a}\right) + \mathcal{O}(1). \quad (2.14)$$

provided the boundary Γ is smooth. In all the geometries we discuss, the change in the Euler characteristic vanishes, $\Delta\chi = 0$, and there is no logarithmic term. However we will show below that, if the logarithmic terms cancel, there exist a universal finite $\mathcal{O}(1)$ term, as well as other universal dependences on the geometry (such as aspect ratios). We will now extract these universal finite terms.

B. The Entanglement Entropy of the Quantum Lifshitz Universality Class

Here we calculate γ_{QCP} at QCPs of the QLM universality class defined by Eq.(2.1) for three different geometries: (i) a cylindrical geometry, (ii) a toroidal geometry, and (iii) a disk geometry. For the cylinder and disk we assume the Dirichlet boundary conditions at the open ends. We use the known results on the free boson partition function(2.4) for different topologies and boundary conditions^{31,32,46–49}, which are necessary for the calculation of entanglement entropy. It is useful to note that the action Eq.(2.3) for general value of the “stiffness” k turns into the standard form:

$$S[\varphi] = \frac{1}{8\pi} \int d^2x (\partial_\mu \varphi)^2, \quad (2.15)$$

upon a rescaling of the field $\sqrt{2k}\phi = \varphi$. If ϕ is compactified with radius $r = 1$, the rescaled field φ has an effective compactification radius $R = \sqrt{2kr^2}$. We find γ_{QCP} to depend linearly on $\ln R$ in all cases we consider.

1. The Cylinder




FIG. 1: Cylinder

Let us begin by considering first a system on a long cylinder of linear size L and circumference ℓ with $L \gg \ell$. Region A to be observed, is a cylinder of length L_A and circumference ℓ . The complement region, B , is a cylinder of length L_B (see Fig.1), also with circumference ℓ . We assume that the QLM wave function Eq.(2.2) and hence the associated 2D partition function Eq.(2.4) obey the Dirichlet boundary conditions at both ends of the cylinder, $A \cup B$.

From Eq.(??), the entanglement entropy $S_A = S_B \equiv S$ is given by

$$S = -\ln Z_{DD}^A(L_A, \ell) - \ln Z_{DD}^B(L_B, \ell) + \ln Z_{DD}^{A \cup B}(L_A + L_B, \ell) \quad (2.16)$$

Here $Z_{DD}(L, \ell)$ is the partition function of Eq.(2.4) for a boson with compactification radius R on cylinder of length L and circumference ℓ with Dirichlet boundary conditions on both ends, which is well known:⁴⁸

$$Z_{DD}(L, \ell) = \mathcal{N} \frac{1}{R} \frac{\vartheta_3\left(\frac{2\tau}{R^2}\right)}{\eta(q^2)} \quad (2.17)$$

where $R = \sqrt{2r^2k}$ is the effective compactification radius (as before), and \mathcal{N} is a non-universal regularization-dependent prefactor, responsible for the area and perimeter dependent terms in the free energy shown in Eq.(2.13). (There are no logarithmic terms for a cylinder or a torus as their Euler characteristic χ vanishes.) In Eq.(2.17) $\tau = i\frac{L}{\ell}$ is the modular parameter, encoding the geometry of the cylinder, and $q = e^{2\pi i\tau}$. The elliptic theta-function $\vartheta_3(\tau)$ and the Dedekind eta-function $\eta(q)$ are given by

$$\vartheta_3(\tau) = \sum_{n=-\infty}^{\infty} q^{\frac{n^2}{2}}, \quad \eta(q) = q^{\frac{1}{24}} \prod_{n=1}^{\infty} (1 - q^n). \quad (2.18)$$

The important feature of Eq.(2.17) is the factor $1/R$, the contribution of the winding modes of the compactified boson on the cylinder with Dirichlet boundary conditions.

Putting it all together, it is straightforward to find an expression for the entanglement entropy using Eq.(2.12). In general, the entanglement entropy depends on the geometry (*e.g.* the aspect ratios L/ℓ) of the cylinders, encoded in ratios of theta and eta functions. However, in the limit $L_A \gg \ell$, in which the length of the cylinders are long compared to their circumference, the entanglement entropy given by Eq.(2.16) and Eq.(2.17) takes a simple form

$$S = \mu\ell + \ln R, \quad (2.19)$$

where μ is a non-universal constant that depending on the regularization-dependent pre-factor \mathcal{N} of Eq.(2.17). Hence, there is a $\mathcal{O}(1)$ universal contribution to the entanglement entropy $\gamma_{QCP} = \ln R$ for the cylindrical geometry. The explicit dependence of γ_{QCP} on the effective effective compactification radius $R = \sqrt{2kr^2}$ shows that it is determined by the winding modes of the compactified boson and thus it is a universal quantity determined by the topology of the surface. In particular we find that the universal piece of the entanglement entropy, γ_{QCP} , for a compactified boson is a continuous function of the radius R , a consequence of the existence of an exactly marginal operator at this QCP. We find the similar relations for all topologies we considered. We will come back to this point in section II C, in the context of several microscopic models of interest.

2. The Torus

We now consider the case in which the full system $A \cup B$ is a torus for which the real part of the modulus $L/\ell \gg 1$, as shown in Fig.2. The two subsystems, A and B are now two cylinders, of length L_A and L_B respectively ($L = L_A + L_B$), both with the same circumference ℓ . We will thus need the partition function on a torus and on two cylinders (with both ends of the cylinders obeying Dirichlet boundary conditions.) The trace

$\text{tr} \rho_A^n$ now becomes

$$\text{tr} \rho_A^n = \left(\frac{Z_{DD}^A(L_A, \ell) Z_{DD}^B(L_B, \ell)}{Z_{\text{torus}}^{A \cup B}(L, \ell)} \right)^{n-1}. \quad (2.20)$$

The partition functions for the two cylinders, A and B has the form of Eq. (2.17). The partition function for the torus is^{31,32}

$$Z_{\text{torus}}(L, \ell) = \left(Z_{\text{cylinder}}^{NN} \left(\frac{L}{2}, \ell \right) \right)^2, \quad (2.21)$$

where $Z_{\text{cylinder}}^{NN}(\frac{L}{2}, \ell)$ is the partition function on a cylinder of length $\frac{L}{2}$ and circumference ℓ , with Neumann boundary conditions at both ends:

$$Z_{\text{cylinder}}^{NN} \left(\frac{L}{2}, \ell \right) = \mathcal{N} \sqrt{\frac{kr^2}{2}} \frac{\vartheta_3(\tau kr^2)}{\eta(q^2)}, \quad (2.22)$$

where $\tau = i\frac{L}{\ell}$ and $q = \exp(2\pi i\tau)$.

In the limit $L_A \gg \ell \gg a$ and $L_B \gg \ell \gg a$, the entanglement entropy for the toroidal geometry is

$$S = \mu\ell + 2 \ln \left(\frac{R^2}{2} \right). \quad (2.23)$$

Hence, for the toroidal geometry, the universal term is $\gamma_{QCP} = 2 \ln(kr^2) = 2 \ln(R^2/2)$. In Eq.(2.23) μ is, once again, a non-universal factor which depends on both the short distance regularization and boundary conditions (in fact, it is not equal to the constant we also called “ μ ” in the entanglement entropy for the case of the cylinder, Eq.(2.19).) As was the case for the cylindrical geometry, in the case of the torus γ_{QCP} is also determined by the contribution of the zero modes of the compactified boson to the partition functions. Thus, here too, γ_{QCP} depends on the effective boson radius $R = \sqrt{2kr^2}$. However, the different values of γ_{QCP} in Eq.(2.23) and Eq.(2.19) is due to the fact that on the torus all three partition functions have contributions from the zero modes.




FIG. 2: Torus

3. The Disk




FIG. 3: Disk

Finally, we compute the entanglement entropy for the disk geometry, shown in Fig.3. The line of argument used above applies here as well. This is the case discussed in Ref.[17], where it was found that the logarithmic term in the entanglement entropy cancels exactly if the boundary Γ is smooth. Here we compute the (subleading) finite universal piece.

To compute the entanglement entropy we need to compute three partition functions, on the two disks A and $A \cup B$, and on the annulus B , all with Dirichlet boundary conditions. These partition functions were computed in the literature long ago for an uncompactified boson.^{46,47} They can be obtained from the partition functions on cylinders, with Dirichlet-Dirichlet (for the annulus) and Dirichlet-Neumann (for the disks) boundary conditions by a conformal mapping $w = \frac{\ell}{2\pi} \ln z$, from the z complex plane to the cylinder (labeled by w). The partition function for the annulus (region B) of inner circumference ℓ and outer circumference L (with Dirichlet boundary conditions) is

$$Z_{DD}^B(L, \ell) = \mathcal{N} \sqrt{\frac{\pi}{\ln(L/\ell)}} \frac{1}{\sqrt{2kr^2}} \frac{\vartheta_3\left(\frac{\tau_B}{r^2 k}\right)}{\eta(q_B^2)}. \quad (2.24)$$

Except for the factor of $1/\sqrt{2kr^2}$, which is due to the zero modes of the compactified boson, this result agrees with those of Ref.[47]. In Eq.(2.24) we have used $q_B = e^{2\pi i \tau_B} = \frac{\ell}{L}$ (with the modular parameter $\tau_B = -\frac{i}{2\pi} \ln\left(\frac{L}{\ell}\right)$).

Similarly, the partition functions on the two disks, regions A and $A \cup B$, are conformally mapped to two infinitely long cylinders (as the UV cutoff $a \rightarrow 0$) with Neumann-Dirichlet boundary conditions. These parti-

tion functions are

$$Z_{\text{disk}} = 2^{-5/12} \pi^{1/4} \frac{\vartheta_4(\tau)}{\eta(q^2)}, \quad (2.25)$$

where $q = \left(\frac{a}{\ell}\right)^4, \left(\frac{a}{L}\right)^4$ for regions A and $A \cup B$, respectively, and τ is their corresponding modular parameter; $\vartheta_4(\tau)$ is the elliptic theta-function

$$\vartheta_4(\tau) = \sum_{n=-\infty}^{\infty} (-1)^n q^{\frac{n^2}{2}}. \quad (2.26)$$

The resulting entanglement entropy for the planar (disk) geometry is found to be

$$S = \frac{1}{2} \ln \left[\frac{1}{\pi} \ln \left(\frac{L}{\ell} \right) \right] + \ln R. \quad (2.27)$$

Hence, for the case of the disk there is also a universal finite piece in the entanglement entropy, $\gamma_{QCP} = \ln \sqrt{2kr^2} \equiv \ln R$. As in the cases discussed above (the cylinder and the torus), here too γ_{QCP} has a topological origin as it is due to the winding modes of the compactified boson. However, unlike the case of the cylinder and toroidal geometries, in the case of the disk there is also a dependence on the aspect ratio L/ℓ (the double logarithmic term), as already noted in Ref.[17]. (Note that we included the factor of $1/\pi$ in the double logarithm since it arises from the conformal mapping.)

C. Entanglement Entropy of Quantum Dimer Models and Related Systems

The results on the entanglement entropy of the preceding subsections apply to several “microscopic” systems of interest. The simplest of them is the quantum dimer model on bipartite lattices at the RK point (associated with the RK wave function of the QDM). As noted in Ref.[16], the RK point of the QDM maps onto the quantum Lifshitz model for a particular value of the radius $r = 1$ and stiffness $k = 2$ (in the notation used here.) This corresponds to a 2D Euclidean boson CFT at the free fermion radius. Of course, this is not an accident, since in this case the lattice partition functions can also be computed exactly by pfaffian methods,^{27,50,51} and hence it is a free Dirac fermion system.

Generalized quantum dimer models have been discussed recently.^{29,30,40} In these models the wave functions correspond to dimer models with weights that depend on the number of dimer pairs on the plaquettes. For a considerable range of values of these weights the system remains critical and can also be mapped onto a quantum Lifshitz model, albeit with a different stiffness connected with the presence of an exactly marginal operator. Thus, in these models the stiffness varies continuously as a function of the microscopic weights. This dependence, discussed in detail in Ref.[29], is of course

non-universal, as it depends on the microscopic structure of the system. Nevertheless, the critical exponents have a universal dependence on the stiffness. The same applies to the universal piece of the entanglement entropy γ_{QCP} , which can be read-off from the results presented in this section.

Similarly, the *quantum* eight-vertex model wave function¹⁶ also maps onto a free fermion problem for a special choice of weights.⁴² For general values of k the fermions are interacting (see the discussion below) but the effects only enter through an exactly marginal operator. The mapping of the quantum 2D eight-vertex model to the quantum Lifshitz model was shown in detail in Ref.[16] where the relation between the stiffness k of the compactified boson and the Baxter weights is given explicitly. k and the weight c in the Baxter wave function (along the six vertex line) are related by

$$\frac{\pi}{2k} = \cot^{-1} \sqrt{\frac{4}{c^4} - 1} \quad (2.28)$$

for a boson with compactification radius $r = 1$ or, equivalent, an effective radius $R = \sqrt{2kr^2}$.

The results of the preceding subsections on the entanglement entropy for the quantum Lifshitz model apply to the lattice models almost without change. Once the mapping of the stiffness to the microscopic parameters (as in the case of the quantum eight vertex model) is known, the universal piece, γ_{QCP} , can be read-off immediately. The only caveat here is that in lattice models it is impossible to have closed simply connected regions with smooth boundaries. The resulting paths of the effective coarse grained quantum Lifshitz model will always have singularities, such as corners, which contribute with a logarithmic dependence to the entanglement entropy (as discussed in Ref.[17]) rendering the finite terms generally non-universal. The cylinder and torus geometries are exceptional in this sense, and allow for a direct check of these ideas in microscopic models, either through an exact solution or by means of numerical computations.

We end this discussion by giving the results for the universal entanglement entropies γ_{QCP} for the Lifshitz universality class at the free fermion (or dimer) and Kosterlitz-Thouless transition of the dimer and Baxter (six vertex) wave functions for all three geometries. (See the summary of Table I.) At the “free dimer” point (the free fermion point of the dimer models) the stiffness $k = 2$ (corresponding to $c^2 = \sqrt{2}$ in the Baxter wave function), and the universal term of the entanglement entropy for a disk geometry is $\gamma_{QCP}^{\text{disk}} = \ln \sqrt{2kr^2} = \ln 2$. For the cylinder we found instead, $\gamma_{QCP}^{\text{cylinder}} = 2 \ln 2$, which, for this value of k , is also the same value for the torus, $\gamma_{QCP}^{\text{torus}}$. (Below we will discuss the relation of these results with the *topological* entanglement entropy of the nearby \mathbb{Z}_2 topological phase.) Away from the free dimer (or fermion) points, the stiffness k changes and so does the entanglement entropy. Thus, at the Kosterlitz-Thouless transition point of both the dimer and six vertex wave

functions (where the Baxter weight is $c = \sqrt{2}$), the stiffness is $k = 1$. (At this point the associated $c = 1$ CFT has an $SU(2)_1$ Kac-Moody current algebra, and the effective compactification radius here is $R = \sqrt{2}$.) The (finite) entanglement entropies now are $\gamma_{QCP}^{\text{torus}} = 2 \ln \sqrt{2}$, $\gamma_{QCP}^{\text{cylinder}} = 0$, and $\gamma_{QCP}^{\text{disk}} = \ln \sqrt{2}$.

R	cylinder	torus	disk
2 (RKpoint)	$2 \ln 2$	$2 \ln 2$	$\ln 2$
$\sqrt{2}$ (KTpoint)	$2 \ln \sqrt{2}$	0	$\ln \sqrt{2}$

TABLE I: Universal entanglement entropies γ_{QCP} of the lattice models in QLM universality class in the cylinder, torus, and disk geometries. γ_{QCP} based on calculations from QLM is quoted at the free fermion point (or RK point) $R = 2$, and at the Kosterlitz-Thouless ($SU(2)_1$) point, $R = \sqrt{2}$.

The only caveat in applying the calculation of γ_{QCP} in the QLM to microscopic models is that is impossible to have closed simply connected regions with smooth boundaries on a lattice. Hence the resulting paths of the effective coarse grained QLM will always have singularities (such as corners) which contribute a finite logarithmic dependence to the entanglement entropy.¹⁷ The cylinder and torus geometries are exceptional in this sense, and allow for a direct check of these ideas in microscopic models, either through an exact solution or by means of numerical computations.

III. GENERALIZED CONFORMAL QCPS ASSOCIATED WITH RCFT

We now generalize the application of Eq.(2.12) to the computation of the entanglement entropy to more general case of conformal QCPs, specifically those associated whose wave functions have an associated 2D Euclidean RCFT (a CFT with a finite number of primary fields.)

A. Entanglement entropy and Boundary Conformal Field theory

The ground state wave function for a conformal quantum critical point can be expressed as Gibbs weight associated with a 2D Euclidean CFT:

$$\Psi_{GS}[\phi] = \frac{1}{\sqrt{Z}} e^{-S[\phi]/2} \quad (3.1)$$

as in the case of the QLM discussed in the previous section. Hence there is a one-to-one mapping between the norm square of the wave function and the partition function of a local 2D Euclidean CFT, and also between the equal-time correlators of the operators of the 2D conformal QCP map onto the correlators of primary fields

of the 2D Euclidean CFT. Furthermore, we will also assume that the associated Euclidean CFT is *unitary* (the S -matrix to be defined below is unitary) and that it is a *RCFT*. The restriction to unitary RCFT allows us to exploit well developed technology for this large class of CFTs^{31,32}, especially that of operator product expansion (OPE) and of *modular S -matrix*, in calculation of γ_{QCP} .

The behavior of RCFTs with specified boundary conditions (especially their partition functions), is the subject of boundary conformal field theory, and was discussed extensively by Cardy^{22,52}. We will follow the approach and results of Cardy in this section. We also need to specify the boundary conditions at the ends of the cylinder, *i.e.* the *boundary states* of the boundary CFT.²² Let us denote these conformal boundary conditions by (α, β) . The associated (conformally invariant) boundary states $|a\rangle$ and $|b\rangle$ can be constructed for each CFT. On the other hand, at the common boundary Γ between the regions A and B , all $n - 1$ fields must obey fixed ('Dirichlet') boundary conditions. As shown by Cardy,²² this boundary condition is quite generally given by the boundary state $|0\rangle$ in the conformal block of the identity $\mathbf{1}$.

For simplicity, we will consider here only the geometries of a cylinder (with specific boundary conditions at each end) and a torus. As in Eq.(2.12) we will need to compute the free energies of region A , B and $A \cup B$ with fixed boundary conditions.

The partition function for a RCFT on a cylinder of length L and circumference ℓ , with boundary conditions a and b on the left and right ends respectively, $Z_{a/b}$, can be expressed in terms of the characters χ_i of the RCFT:

$$Z_{a/b} = \sum_j N_{ab}^j \chi_j \left(e^{-\pi\ell/L} \right), \quad (3.2)$$

where the integers N_{ab}^j are the fusion constants, the coefficients in the OPE of the RCFT,

$$\Phi_a \times \Phi_b = \sum_j N_{ab}^j \Phi_j. \quad (3.3)$$

The Virasoro characters χ_j are given by the trace over the descendants $|\Phi_j\rangle$ of the highest weight state, which are obtained by acting on it with the Virasoro generators \hat{L}_{-n} ($n > 0$):

$$\chi_j \left(e^{-\pi\ell/L} \right) = e^{\pi\ell c/24L} \text{tr}_a \left(e^{-\frac{\pi\ell}{L} \hat{L}_0} \right), \quad (3.4)$$

where c is the central charge of the CFT, \hat{L}_0 is the $n = 0$ Virasoro generator. Here the modular parameter is $\tau \equiv i\ell/2L$. Under a modular transformation $\tau \rightarrow -1/\tau$, which exchanges the Euclidean "space" and "time" dimensions of the cylinder (*i.e.* it flips the cylinder from the "horizontal" to the "vertical" position), the characters transform as

$$\chi_i \left(e^{-\pi\ell/L} \right) = S_i^j \chi_j \left(e^{-4\pi L/\ell} \right), \quad (3.5)$$

where S_i^j is the *modular S -matrix* of the RCFT. The modular S -matrix and the fusion coefficients are related by the Verlinde formula⁵³

$$N_{ab}^j = \sum_i \frac{S_j^i S_a^i S_b^i}{S_i^0}. \quad (3.6)$$

The limit of interest here is, once again, $L \gg \ell$. Under a modular transformation, the partition function of Eq.(3.2) becomes

$$Z_{a/b} = \sum_{i,j} N_{ab}^i S_i^j \chi_j \left(e^{-4\pi L/\ell} \right). \quad (3.7)$$

In the limit $\frac{\ell}{L} \rightarrow 0$, $Z_{a/b}$ is dominated by the the descendants of the identity $\mathbf{1}$ (up to exponentially small corrections). Hence, in this limit,

$$Z_{a/b} \rightarrow \sum_i N_{ab}^i S_i^0 \chi_0 \left(e^{-4\pi L/\ell} \right) \rightarrow e^{\frac{\pi Lc}{6\ell}} \sum_i N_{ab}^i S_i^0 \quad (3.8)$$

and $\ln Z_{a/b}$ becomes

$$\ln Z_{a/b} = \frac{\pi Lc}{6\ell} + \ln g_{ab}, \quad (3.9)$$

dropping UV singular (non-universal) terms. The quantity $\ln g_{ab}$ in Eq.(3.9) is the *boundary entropy* of a boundary RCFT introduced by Affleck and Ludwig²⁴, where the "ground state degeneracy" g_{ab} is given by

$$g_{ab} = \sum_i N_{ab}^i S_i^0. \quad (3.10)$$

Using Eq.(2.12), these standard results imply that the entanglement entropy of the 2D rational conformal QCP for a cylindrical geometry (see Fig.1). For boundary conditions a and b at the two ends associated with regions A and B , the entanglement entropy is

$$\begin{aligned} S &= -\ln \left(\frac{Z_A^{a0} Z_B^{0b}}{Z_{A \cup B}^{ab}} \right) \\ &= \mu\ell - \ln \left(\frac{\left(\sum_j N_{a0}^j S_j^0 \right) \left(\sum_k N_{0b}^k S_k^0 \right)}{\sum_l N_{ab}^l S_l^0} \right) \\ &= \mu\ell - \ln \left(\frac{g_{a0} g_{0b}}{g_{ab}} \right), \end{aligned} \quad (3.11)$$

where we explicitly used the fact that the state at the common boundary Γ should be fixed to be the *fixed* BC with boundary state $|0\rangle$.

The result Eq.(3.11) provides an explicit way to compute γ_{QCP} for the entire class of many-body wave functions at QCPs associated with RCFT in terms of the data of the RCFT:

$$\gamma_{QCP} = -\ln \left(\frac{\left(\sum_j N_{a0}^j S_j^0 \right) \left(\sum_k N_{0b}^k S_k^0 \right)}{\sum_l N_{ab}^l S_l^0} \right). \quad (3.12)$$

This is the main result of this section. It shows that γ_{QCP} is in general determined by the OPE coefficients N_{ba}^c (which encode the boundary conditions on the partition functions) and by the modular S -matrix, S_i^j , of the RCFT associated with the *norm squared of the many-body wave function* at the given QCP.

It is important to note that it is also possible to define a unitary S -matrix that governs the transformation properties of the *wave function* itself under a modular transformation. This modular S -matrix plays a central role in 2D topological phases and in topological field theories.^{18,21,54} However, only for topological theories these two S -matrices are the same and in general are different or even not be defined at all!) We will come back to this issue in the discussion section.

A particularly simple result is obtained for the case of a cylinder with fixed boundary conditions on both ends. In this case, Z_A , Z_B and $Z_{A \cup B}$ are cylinders with fixed boundary conditions, and hence the boundary states for all three cases are in the conformal block of the identity $\mathbf{1}$. Since in this case the only non-vanishing OPE coefficient is $N_{00}^0 = 1$, the universal term of the entanglement entropy, γ_{QCP} , depends only on the element S_0^0 of the modular S -matrix of the RCFT:

$$\gamma_{QCP} = -\ln S_0^0. \quad (3.13)$$

For the case in which the full region $A \cup B$ is a torus, we can use an analogue of Eq.(3.11) by writing the partition function $Z_{A \cup B}$ in the denominator of Eq.(3.11) as a modular invariant. In the limit of interest $L \gg \ell$, the denominator g_{ab} of Eq.(3.11) is replaced by a sum of terms with similar structure corresponding to a sum over boundary conditions (and twists) needed to represent the torus (see, for instance, Ref.[32]). Similarly, Eq.(3.11) can also be applied to the disk geometry upon a conformal mapping as it was done for the case of the compactified boson in section II B 3.

B. Applications

We will now discuss some examples of interest. In applying the results Eq.(3.12) to specific systems, one should keep in mind that that choice of the inner product of the 2D quantum theory can play a subtle role. As it was pointed out recently by Fendley³⁵, a scale invariant wave function does not necessarily imply scale invariance of the correlators. Their actual behavior depends also on the choice of inner product. Here we have assumed that the states labeled by the set of field configurations $\phi(x, y)$ form an orthogonal basis. Hence, the norm of the wave function is a sum over states with the local weights squared. However what matters is that the *matrix elements* (and in particular the norm of the states) be scale-invariant. A number of interesting counterexamples are known.⁵⁵ The QLM is a special case where such “naive” inner product maintains scale invariance. This is due to the existence of exactly marginal operators in the QLM.

Below we discuss four cases where the ground state wave function with the “naive” inner product describes QCPs: (i) a QCP associated with the 2D Ising CFT, (ii) the QCPs associated with compactified boson CFT, (iii) QCPs in quantum loop models^{33,34}, and (iv) quantum net models^{35,36,56,57}. (See footnote Ref.[58].)

1. The 2D Ising wave function

As an example of a system described by an RCFT we consider a 2D quantum spin system whose ground state wave function has for amplitudes the Gibbs weights of the 2D classical Ising model. This system is quantum critical if the square of the weights (which also have the form of a Gibbs weight for the 3D Ising model) are at the critical point of the 2D Ising model, the Onsager value.

The critical point of the 2D Ising model is the simplest RCFT. It has central charge $c = 1/2$, and three (bulk) primary fields: 1) the identity ($\mathbf{1}$, with conformal weight $h = 0$), 2) the energy density (ε , with conformal weight $h = 1/2$), and 3) the spin field (σ , with conformal weight $1/16$), which obey the operator algebra (OPE)

$$\begin{aligned} \varepsilon \times \varepsilon &= \mathbf{1} \\ \varepsilon \times \sigma &= \sigma \\ \sigma \times \sigma &= \mathbf{1} + \varepsilon. \end{aligned} \quad (3.14)$$

The critical Ising model has three possible boundary states:²² 1) the *spin up* state $|+\rangle$, 2) the *spin down* state $|-\rangle$, and 3) the *free* state $|f\rangle$. (Either the up or the down state can be regarded as the fixed boundary state.) These three boundary states, $|+\rangle$, $|-\rangle$, and $|f\rangle$ are in the conformal blocks of the identity $\mathbf{1}$ (denoted by $|\tilde{0}\rangle$), the energy density ε (denoted by $|\tilde{\frac{1}{2}}\rangle$), and the spin field σ (denoted by $|\tilde{\frac{1}{16}}\rangle$), respectively. The boundary states are given by²²

$$\begin{aligned} |+\rangle &\equiv |\tilde{0}\rangle = \frac{1}{\sqrt{2}}|0\rangle + \frac{1}{\sqrt{2}}|\varepsilon\rangle + \frac{1}{\sqrt[4]{2}}|\sigma\rangle \\ |-\rangle &\equiv |\tilde{\frac{1}{2}}\rangle = \frac{1}{\sqrt{2}}|0\rangle + \frac{1}{\sqrt{2}}|\varepsilon\rangle - \frac{1}{\sqrt[4]{2}}|\sigma\rangle \\ |f\rangle &\equiv |\tilde{\frac{1}{16}}\rangle = |0\rangle - |\varepsilon\rangle. \end{aligned} \quad (3.15)$$

The modular S -matrix is

$$S = \begin{pmatrix} \frac{1}{2} & \frac{1}{2} & \frac{1}{\sqrt{2}} \\ \frac{1}{2} & \frac{1}{2} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} & 0 \end{pmatrix}, \quad (3.16)$$

where the columns are labeled by the highest weights 0, $1/2$, and $1/16$, in that order.

The entanglement entropy for this wave function can now be computed, using the result of Eq.(3.11). We will take region $A \cup B$ to be a long cylinder of length L and

circumference ℓ , and regions A and B to be two cylinders of lengths L_A and L_B respectively, with the same circumference ℓ , and with $L = L_A + L_B$.

Let us take the boundary conditions at both ends of $A \cup B$ to be free. By a conformal mapping, this maps onto the disk. Back on the cylinder, the free boundary condition is described by the boundary state $|f\rangle$, which is in the conformal block of the primary field σ . On the other hand, at the boundary Γ between regions A and B , we have the fixed boundary condition, the up state $|+\rangle$. We readily find

$$\begin{aligned} g_{\sigma,1} &= N_{\sigma,0}^\sigma S_\sigma^0 = \frac{1}{\sqrt{2}} \\ g_{1,\sigma} &= N_{0,\sigma}^\sigma S_\sigma^0 = \frac{1}{\sqrt{2}} \\ g_{\sigma,\sigma} &= N_{\sigma,\sigma}^0 S_0^0 + N_{\sigma,\sigma}^\varepsilon S_\varepsilon^0 = 1. \end{aligned} \quad (3.17)$$

The universal term of the entanglement entropy, γ_{QCP} now is

$$\gamma_{QCP} = -\ln \frac{g_{ab}g_{bc}}{g_{ac}} = -\ln \frac{(S_\sigma^0)^2}{S_0^0 + S_\varepsilon^0} = \ln 2. \quad (3.18)$$

On the other hand, we could consider instead the case of fixed boundary conditions at both ends of the cylinder $A \cup B$. This corresponds to the boundary state $|\bar{0}\rangle$. Since the boundary condition on Γ is always fixed, γ_{QCP} is now

$$\gamma_{QCP} = -\ln S_0^0 = \ln 2. \quad (3.19)$$

In the case where $A \cup B$ is torus of large circumference L and small circumference ℓ (hence with modular parameter $\tau = i\ell/L$), the regions A and B are cylinders each of length L_A and L_B and circumference ℓ , with fixed boundary conditions at both ends. The partition function for the torus, Z_{AUB}^{torus} , is^{31,32}

$$Z_{AUB}^{\text{torus}} = \frac{1}{2} \left(\left| \frac{\vartheta_2(\tau)}{\eta(\tau)} \right| + \left| \frac{\vartheta_3(\tau)}{\eta(\tau)} \right| + \left| \frac{\vartheta_4(\tau)}{\eta(\tau)} \right| \right). \quad (3.20)$$

Using the modular invariance of Z on the torus ($\tau \rightarrow -1/\tau$), one finds that in the limit $L \gg \ell$, $Z_{AUB}^{\text{torus}} \rightarrow \frac{3}{2}$. Hence, in the case of the torus, γ_{QCP} is

$$\gamma_{QCP}^{\text{torus}} = -\ln \frac{(S_0^0)^2}{\frac{3}{2}} = \ln 6. \quad (3.21)$$

2. The compactified boson wave function

We can also use this approach to compute the entanglement entropy for the compactified boson wave function (the quantum Lifshitz state) discussed in the previous Section. However, unlike the explicit computation of the boson determinant presented in the previous section, a computation that can be done for any compactification radius R , the boundary CFT approach we are using in

this section only applies for a rational CFT. This restricts the compactification radius to be such that R^2 is a rational number. (The general case can be regarded as a limit.)

It is now straightforward to compute the entanglement entropy using Eq.(3.11). For this case we find $\gamma_{QCP} = -\ln S_0^0 = \ln R$, consistent with the results of the preceding section.

3. Quantum loop models

Quantum loop models are two-dimensional quantum systems whose Hilbert space is spanned by states labelled by loop configurations (or coverings) of a two-dimensional lattice. We will denote by $\{\mathcal{L}\}$ the set of these configurations. Conventionally, this set of states are taken to be a basis of the loop Hilbert space, and hence they are assumed to be linearly independent, complete and orthonormal, (with respect to the naively defined inner product.)

Quantum loop models were originally proposed as candidates for time-reversal invariant topological phases.^{33,59,60} Wave functions in the Hilbert space of (multi) loop configurations have the form

$$|\Psi_{(x,d)}\rangle = \sum_{\mathcal{L}} x^{L[\mathcal{L}]} d^{N[\mathcal{L}]} |\mathcal{L}\rangle. \quad (3.22)$$

Here $N[\mathcal{L}]$ is the number of loops in state (configuration) \mathcal{L} , $L[\mathcal{L}]$ is the length of loop in the configuration, d is the ‘‘loop fugacity’’, and x is the weight (fugacity) of a unit length of loop.

The candidate wave functions of a quantum loop model in a putative topological phase depends on the loop configuration but not on the length of the loops. The simplest such state is the ‘‘ d -isotopy’’ (multi) loop wave function^{59,60}

$$|\Psi_d\rangle = \sum_{\mathcal{L}} d^{N[\mathcal{L}]} |\mathcal{L}\rangle \quad (3.23)$$

obtained from $|\Psi_{(x,d)}\rangle$ by setting the fugacity of the unit length of loop $x = 1$. This is a generalization of Kitaev’s ‘‘Toric Code’’ wave function⁶¹ ($d = 1$), *i.e.* a \mathbb{Z}_2 gauge theory deep in its deconfined phase in $2 + 1$ dimensions. Another limit of interest is the ‘‘fully packed’’ state

$$|\Psi_{(\infty,d)}\rangle = \lim_{x \rightarrow \infty} \sum_{\mathcal{L}} x^{L[\mathcal{L}]} d^{N[\mathcal{L}]} |\mathcal{L}\rangle \quad (3.24)$$

obtained by setting $x \rightarrow \infty$, which forces the constraint that the loops cover the maximal allowable set of links on the lattice.

With the naively defined inner product, the norm squared of the d -isotopy state $|\Psi_d\rangle$, Eq.(3.23), is

$$Z(d^2) \equiv \|\Psi_d\|^2 = \sum_{\mathcal{L}} d^{2N[\mathcal{L}]}, \quad (3.25)$$

which is the same as the partition function of a 2D classical loop model on the same lattice, with a weight d^2 per loop. Likewise, the norm squared of the fully packed loop state $|\Psi_{(\infty,d)}\rangle$ is the partition function $Z(\infty, d^2)$ of the classical fully packed loop model, with fugacity d^2 , on the same lattice.

The partition functions of classical loop models on a 2D lattice have been studied extensively, particularly on the honeycomb lattice (for a detailed review see Refs.[37,62,63].) In the fully packed limit, the partition function $Z(\infty, d^2)$ is critical for $d \leq \sqrt{2}$. The universality classes of the fully packed loop models (on the honeycomb lattice) are rational *unitary* CFTs only for $d = 1$ (the $SU(2)_1$ RCFT) and $d = \sqrt{2}$ (the $SU(3)_1$ RCFT). For finite x , the partition function for the dense loop gas $Z(x, d^2)$ is also critical for $d \leq \sqrt{2}$. The universality classes are again rational unitary CFTs only for $d = 1$ and $d = \sqrt{2}$. The fixed point for the case $d = 1$ is equivalent to the statistics of the proliferated domain walls of the classical 2D Ising model at infinite temperature.³⁷ For $d = \sqrt{2}$ the dense and dilute loop gases have the same critical theory, the Kosterlitz-Thouless critical point, and hence also the $SU(2)_1$ RCFT.

We can now use the result in Eqs.(3.12) and (3.13) to compute the universal term of the entanglement entropy for the loop wave functions with $d = 1, \sqrt{2}$, on a cylinder with fixed boundary conditions (for the loops!). The modular S -matrices are known,^{20,31,32} and the needed S_0^0 matrix elements are $S_0^0 = \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{3}}$, for $SU(2)_1$ and $SU(3)_1$, respectively. The universal term γ_{QCP} of the entanglement entropy for each case is $\gamma_{QCP} = \ln \sqrt{2}, \ln \sqrt{3}, -\ln 2$ for the fully packed state at $d = 1$ (and also for the loop gas at $d = \sqrt{2}$), the fully packed loop state at $d = \sqrt{2}$, and the dense loop gas at $d = 1$ (corresponding to the Kitaev state), respectively. Here we have used a recent result on the behavior of of the dense loop model by Cardy⁶⁴ who showed (among many other things) that for $d = 1$ the partition function of the dense loop model on the cylinder $Z = 2$. We will see in the discussion section that this *negative* value, $\gamma = -\ln 2$, coincides with the direct computation of the *topological* entanglement entropy in the Kitaev wave function.^{18,19,65}

4. Quantum net models

Finally, we will briefly discuss the more interesting, but less understood problem of the wave functions for *quantum net models*^{35,36,56,57}. These states were proposed as candidates for a time-reversal invariant non-Abelian topological phase. The Hilbert space of quantum net models is spanned by the coverings of a lattice by configurations of nets, *i.e.* branching loops (with trivalent vertices). An interesting example is the chromatic polynomial state.⁵⁶ In this state, the nets are regarded as a configuration of domain walls of a Q -state Potts model. The weight of a given state $|\mathcal{L}\rangle$ is the chromatic poly-

nomial $\chi_Q[\mathcal{L}]$ of the configuration. The chromatic polynomial counts the number of ways of coloring regions of the lattice separated by domain walls of a Q -state 2D Potts model. They were first introduced in the computation of the low temperature expansion for the 2D Potts models (see, for instance, Ref.[42].) For non-integer Q , the chromatic polynomial can be computed by an iterative procedure.⁵⁶ The 2D Potts model is known to have a critical point for $Q \leq 4$.

Following Ref.[56], we consider the norm of the chromatic polynomial state with $Q \leq 4$. In order to compute the norm, we have to square the weight, resulting in a partition function involving the sum of the *square* of the chromatic polynomial. It is then natural to ask for a value of Q such that $\chi_Q^2[\mathcal{L}] \propto \chi_{Q_{\text{eff}}}[\mathcal{L}]$, for some Q_{eff} . Then the nets will be critical provided $Q_{\text{eff}} \leq 4$. It turns out⁵⁶ that, up to a suitably chosen fugacity for trivalent vertices⁵⁷, this property holds only for $\sqrt{Q} = \frac{1+\sqrt{5}}{2}$, the *Golden Ratio*, with $Q_{\text{eff}} = 2 + \frac{1+\sqrt{5}}{2} < 4$. Thus, for this state the nets are critical.

This case is interesting for several reasons. One is that strong arguments⁵⁶ suggest that it is possible to define for this wave function an excitation (a defect) which is denoted by τ , a Fibonacci anyon (not to be confused with the modular parameter!) with the fusion rule, $\tau \times \tau = \mathbf{1} + \tau$. Fibonacci anyons are of prime interest in the topological approach to quantum computation.⁶⁶ However, for this approach to work it is necessary that this state should describe a topological state, which requires that its local excitations (not the nets) be gapped. Fendley³⁵ has recently given strong arguments that imply that this state, with the naive inner product we use here, is not topological but a quantum critical state.

Another feature that makes this state interesting is that the correlations encoded in the norm of the state for $\sqrt{Q} = \frac{1+\sqrt{5}}{2}$ are described by a RCFT, the minimal model of the Friedan-Qiu-Shenker⁶⁷ series of unitary RCFTs at level $m = 9$, with central charge $c = \frac{14}{15}$. This minimal model has a large number of primaries (36) and has not been studied in detail. Nevertheless, its modular S -matrix is known (as it is for the entire series³¹). Although to the best of our knowledge the boundary CFT of this minimal model has not been investigated, we conjecture that the boundary state corresponding to the fixed boundary condition is the analog of the state $|\tilde{0}\rangle$ in the 2D critical Ising model (the $m = 3$ member of the same series.), *i.e.* the state in the conformal block of the identity.⁵² Thus, if we consider this state on a cylinder with fixed boundary conditions, the entanglement entropy for observing only half of the system, has a universal term γ_{QCP} of the form given in Eq.(3.13), and hence is given in terms of the S_0^0 element of the modular S -matrix of this RCFT:³¹

$$\gamma_{QCP} = -\ln S_0^0 = -\ln \left(\frac{\sin(\frac{\pi}{9})}{15 + 3\sqrt{5}} \right). \quad (3.26)$$

IV. CONCLUSIONS AND DISCUSSION

We have shown that at 2D conformal QCPs (with dynamical exponent $z = 2$), the entanglement entropy for a region with a smooth boundary quite generally has universal finite contributions which we denoted by γ_{QCP} :

$$S_{QCP} = \mu\ell + \gamma_{QCP}.$$

We studied the universal nature of γ_{QCP} with two complementary approaches for large classes of 2D conformal QCPs: First for the QLM universality class, we calculated γ_{QCP} explicitly in terms of the partition functions (that of compactified boson) associated with the norm squared of the wave function. Later we used known results from boundary CFT to show that γ_{QCP} is determined by the detailed structure of the associated RCFT encoded in the modular S -matrix and the OPE fusion coefficients for the primary fields. We also applied this general results to compute γ_{QCP} in several systems of interest: the quantum Lifshitz model, the generalized quantum dimer and quantum eight-vertex models, and quantum loop and net models.

However, we show below that for a general conformal quantum critical point, whose ground state wave function is given by the Gibbs weights of a Euclidean rational unitary CFT, the universal term γ_{QCP} is determined by the modular S -matrix associated with the norm squared of the wave function. Thus, the modular S -matrix of the topological phase and that of the wave functions of 2D conformal quantum critical points have a conceptually different origin. In particular, in all the cases we checked here, γ_{QCP} and γ_{topo} contribute with opposite signs to their respective entanglement entropies,

We note that while our result for the entanglement entropy has the *same form* as the entanglement entropy for a *topological phase*,^{18,19} the finite universal terms γ_{QCP} and γ_{topo} have a different origin and structure. In the case of a topological phase, γ_{topo} is in general determined by the modular S -matrix of the topological field theory of the topological phase.¹⁸⁻²⁰ This modular S -matrix governs the transformation properties of the ground state within the degenerate ground state Hilbert space of the topological phase under modular transformations on a torus: $\tau \rightarrow -1/\tau$, where τ is the modular parameter of the torus²¹. On the other hand, for 2D conformal QCPs whose ground state wave function is given by the Gibbs weights of a Euclidean rational unitary CFT, the universal term γ_{QCP} is determined by the modular S -matrix associated with the norm squared of the wave function

and the S -matrix connects between different boundary conditions. Hence the roles of the modular S -matrix in the computation of the universal $\mathcal{O}(1)$ terms to the entanglement entropy have conceptually different origin. Moreover, γ_{QCP} and γ_{topo} enter with opposite signs in their contributions to their respective entanglement entropies. In fact, in all the cases we looked at we found that $\gamma_{QCP} > 0$, except for the Kitaev state which is topological (and we recovered the known result.). It is unclear to us how general this difference actually is and, more importantly, if it has a deeper meaning.

It is also interesting to note the striking similarity of the structure of Eq.(3.12) (with its dependence on the S -matrix and the fusion rules) with the results of Fendley, Fisher and Nayak⁶⁸ for the change in the entanglement entropy of a 2D topological fluid, a fractional quantum Hall state, by the action of a point contact. Recently, Refs.[69,70] found finite universal terms in the entanglement entropy for 1 + 1 dimensional CFTs with a similar structure to what we found here in 2D conformal QCPs. Calculations of quantum fidelity in 1D also find a similar structure.^{71,72} Recent work by Li and Haldane⁷³ also raises the interesting possibility of computing the entanglement spectrum for a theory with a wave function described by a known CFT, but this is beyond the scope of this paper. Finally, it is interesting to consider how the entanglement entropy (and in particular the finite term γ_{QCP}) would evolve as one perturbed the system either by a finite non-zero temperature into the quantum critical regime, or by a relevant operator that drives the system into a nearby topologically ordered phase that can be accessed by local perturbations^{16,26,27,35,56} and to investigate possible connections with RCFT.⁷⁴⁻⁷⁶

Acknowledgments

We thank John Cardy, Paul Fendley, Greg Moore, and Joel Moore for their comments and suggestions. BH and MM thank the Les Houches Summer School for its hospitality. The work of EF and BF was supported by the National Science Foundation grant DMR 0758462 at the University of Illinois. MM was supported by the Stanford Institute for Theoretical Physics, the NSF under grant PHY-0244728, the DOE under contract DE-AC03-76SF00515, and an ARCS Fellowship. EAK was supported by the Stanford Institute for Theoretical Physics during a part of this work.

¹ M. Srednicki, Phys. Rev. Lett. **71**, 666 (1993).

² L. Bombelli, R. K. Koul, J. Lee, and R. D. Sorkin, Phys. Rev. D **34**, 373 (1986).

³ C. G. Callan and F. Wilczek, Phys. Lett. B **333**, 55 (1994).

⁴ C. Holzhey, F. Larsen, and F. Wilczek, Nucl. Phys. B **424**,

443 (1994).

⁵ P. Calabrese and J. Cardy, J. Stat. Mech. JSTAT **04**, P06002 (2004).

⁶ G. Vidal, J. L. Latorre, E. Rico, and A. Kitaev, Phys. Rev. Lett. **90**, 227902 (2003).

- ⁷ J. I. Latorre, E. Rico, and G. Vidal, *Quant. Inf. and Comp.* **4**, 048 (2004).
- ⁸ S. Ryu and T. Takayanagi, *Phys. Rev. Lett.* **96**, 181602 (2006).
- ⁹ G. Refael and J. E. Moore, *Phys. Rev. Lett.* **93**, 260602 (2004).
- ¹⁰ G. Refael and J. E. Moore, *Phys. Rev. B* **76**, 024419 (2007).
- ¹¹ A. Kopp, X. Jia, and S. Chakravarty, *Ann. Phys. (N. Y.)* **322**, 1466 (2007).
- ¹² A. Kopp and K. Le Hur, *Phys. Rev. Lett.* **98**, 220401 (2007).
- ¹³ N. Laflorencie, E. S. Sørensen, M.-S. Chang, and I. Affleck, *Phys. Rev. Lett.* **96**, 100603 (2006).
- ¹⁴ E. S. Sørensen, M.-S. Chang, N. Laflorencie, and I. Affleck, *J. Stat. Mech. JSTAT* **07**, P08003 (2007).
- ¹⁵ I. K. Affleck (2008), unpublished, Proceedings of the Les Houches Summer School 2008, arXiv:0809.3474.
- ¹⁶ E. Ardonne, P. Fendley, and E. Fradkin, *Ann. Phys. (N. Y.)* **310**, 493 (2004).
- ¹⁷ E. Fradkin and J. E. Moore, *Phys. Rev. Lett.* **97**, 050404 (2006).
- ¹⁸ A. Kitaev and J. Preskill, *Phys. Rev. Lett.* **96**, 110404 (2006).
- ¹⁹ M. Levin and X. G. Wen, *Phys. Rev. Lett.* **96**, 110405 (2006).
- ²⁰ S. Dong, E. Fradkin, R. G. Leigh, and S. Nowling, *J. High Energy Phys. JHEP* **05**, 016 (2008).
- ²¹ E. Witten, *Commun. Math. Phys.* **121**, 351 (1989).
- ²² J. L. Cardy, *Nucl. Phys. B* **275** [FS17], 200 (1986).
- ²³ J. L. Cardy, *Nucl. Phys. B* **270** [FS16], 186 (1986).
- ²⁴ I. Affleck and A. W. W. Ludwig, *Phys. Rev. Lett.* **67**, 161 (1991).
- ²⁵ D. S. Rokhsar and S. A. Kivelson, *Phys. Rev. Lett.* **61**, 2376 (1988).
- ²⁶ R. Moessner and S. L. Sondhi, *Phys. Rev. Lett.* **86**, 1881 (2001).
- ²⁷ P. Fendley, R. Moessner, and S. L. Sondhi, *Phys. Rev. B* **66**, 214513 (2002).
- ²⁸ E. Fradkin, D. A. Huse, R. Moessner, V. Oganesyan, and S. L. Sondhi, *Phys. Rev. B* **69**, 224415 (2004).
- ²⁹ S. Papanikolaou, E. Luijten, and E. Fradkin, *Phys. Rev. B* **76**, 134514 (2007).
- ³⁰ C. Castelnovo, C. Chamon, C. Mudry, and P. Pujol, *Ann. Phys. (N. Y.)* **318**, 316 (2005).
- ³¹ P. Ginsparg, in *Fields, Strings and Critical Phenomena*, edited by E. Brezin and J. Zinn-Justin (North-Holland, 1990), Les Houches Summer School in Theoretical Physics, Les Houches, France, 1988.
- ³² P. D. Francesco, P. Mathieu, and D. Sénéchal, *Conformal Field Theory* (Springer-Verlag, New York, 1997).
- ³³ M. Freedman, C. Nayak, and K. Shtengel, *Phys. Rev. Lett.* **94**, 066401 (2004).
- ³⁴ M. Troyer, S. Trebst, K. Shtengel, and C. Nayak (2008), unpublished, arXiv:0805.1933.
- ³⁵ P. Fendley, *Ann. Phys. (N.Y.)* **323**, 3113 (2008).
- ³⁶ M. Levin and X.-G. Wen, *Phys. Rev. B* **71**, 045110 (2005).
- ³⁷ B. Nienhuis, in *Phase Transitions and Critical Phenomena*, edited by C. Domb and J. Lebowitz (Academic Press, London, 1987), vol. 11, page 1.
- ³⁸ R. Moessner, S. L. Sondhi, and E. Fradkin, *Phys. Rev. B* **65**, 024504 (2002).
- ³⁹ C. L. Henley, *J. Stat. Phys.* **89**, 483 (1997).
- ⁴⁰ F. Alet, J. L. Jacobsen, G. Misguich, V. Pasquier, F. Mila, and M. Troyer, *Phys. Rev. Lett.* **94**, 235702 (2005).
- ⁴¹ S. Papanikolaou, K. S. Raman, and E. Fradkin, *Phys. Rev. B* **76**, 224421 (2007).
- ⁴² R. Baxter, *Exactly Solved Models in Statistical Mechanics* (Academic Press, 1982), and references therein.
- ⁴³ M. Kac, *Amer. Math. Monthly* **73**, 1 (1966).
- ⁴⁴ V. Privman, *Phys. Rev. B* **38**, 9261 (1988).
- ⁴⁵ J. L. Cardy and I. Peschel, *Nucl. Phys. B* **300** [FS22], 377 (1988).
- ⁴⁶ J. Polchinski, *Comm. Math. Phys.* **104**, 37 (1986).
- ⁴⁷ W. I. Weisberger, *Comm. Math. Phys.* **112**, 633 (1987).
- ⁴⁸ P. Fendley, H. Saleur, and N. P. Warner, *Nucl. Phys. B* **430** [FS], 577 (1994).
- ⁴⁹ S. Eggert and I. Affleck, *Phys. Rev. B* **46**, 19866 (1992).
- ⁵⁰ M. E. Fisher, *Phys. Rev.* **124**, 1664 (1961).
- ⁵¹ S. Samuel, *J. Math. Phys.* **21**, 2806 (1980).
- ⁵² J. L. Cardy, *Nucl. Phys. B* **324**, 581 (1989).
- ⁵³ E. P. Verlinde, *Nucl. Phys. B* **300**, 360 (1988).
- ⁵⁴ P. Bonderson, K. Shtengel, and J. K. Slingerland, *Phys. Rev. Lett.* **97**, 016401 (2006).
- ⁵⁵ For instance one can consider a 2D spin system whose ground state wave function is given by the Gibbs weights of a 2D classical Ising model at criticality. The norm of this state will correspond indeed to a 2D classical Ising model but it will be instead in the ordered phase as the effective “temperature” is half of the Onsager value. Similarly, if one takes a wave function whose amplitudes are the weights of an Euclidean Wess-Zumino-Witten (WZW) model (say for $SU(2)_k$), the norm squared of this state is actually the partition function of a non-linear sigma model rather than the WZW partition function and it is not conformally invariant. Alternatively, if one could take a wave function whose amplitudes are the weights of a gauged chiral WZW model, the norm of this state turns out to be indeed the partition function of a (non-chiral) WZW model¹⁶. However, since this wave function is that of a Chern-Simons gauge theory,²¹⁷ which is a topological field theory, and it does not describe a quantum critical system.
- ⁵⁶ P. Fendley and E. Fradkin, *Phys. Rev. B* **72**, 024412 (2005).
- ⁵⁷ L. Fidkowski, M. Freedman, C. Nayak, K. Walker, and Z. Wang (2006), unpublished, arXiv:cond-mat/0610583.
- ⁵⁸ The quantum loop and net models with scale invariant *wave functions* were originally proposed as candidates to exhibit topological phases non-Abelian fractional statistics^{16,36,56,57,607}. However, with “naive” inner products, they turned out to describe quantum critical states rather than gapped topological phases³⁴. These are recent entries to examples of conformal QCPs.
- ⁵⁹ M. H. Freedman, *Comm. Math. Phys.* **234**, 129 (2003).
- ⁶⁰ M. Freedman, C. Nayak, K. Shtengel, and K. Walker, *Ann. Phys.* **310**, 428 (2004).
- ⁶¹ A. Y. Kitaev, *Ann. Phys. (N. Y.)* **303**, 2 (2003).
- ⁶² J. Kondev and C. L. Henley, *Nucl. Phys. B* **464**, 540 (1996).
- ⁶³ J. Kondev, J. de Gier, and B. Nienhuis, *J. Phys. A* **29**, 6489 (1996).
- ⁶⁴ J. Cardy, *J. Stat. Phys.* **125**, 1 (2006).
- ⁶⁵ A. Hamma, R. Ionicioiu, and P. Zanardi, *Phys. Lett. A* **337**, 22 (2005).
- ⁶⁶ M. H. Freedman, A. Kitaev, M. J. Larsen, and Z. Wang, *Comm. Math. Phys.* **227**, 605 (2002).
- ⁶⁷ D. Friedan, Z. Qiu, and S. Shenker, *Phys. Rev. Lett.* **52**, 1575 (1984).
- ⁶⁸ P. Fendley, M. P. A. Fisher, and C. Nayak, *J. Stat. Phys.*

- 126**, 1111 (2007).
- ⁶⁹ M. Caraglio and F. Gliozzi (2008), unpublished, arXiv:0808.4094.
- ⁷⁰ S. Furukawa, V. Pasquier, and J. Shiraishi (2008), unpublished, arXiv:0809.5113v1.
- ⁷¹ D. F. Abasto, A. Hamma, and P. Zanardi, Phys. Rev. A **78**, 010301 (2008).
- ⁷² L. C. Venuti, H. Saleur, and P. Zanardi (2008), unpublished.
- ⁷³ H. Li and F. D. M. Haldane, Phys. Rev. Lett. **101**, 010504 (2008).
- ⁷⁴ P. Dorey, I. Runkel, R. Tateo, and G. Watts, Nucl. Phys. B **578**, 85 (2000).
- ⁷⁵ P. Dorey, D. Fioravanti, C. Rim, and R. Tateo, Nucl. Phys. B **696**, 445 (2004).
- ⁷⁶ D. R. Green, M. Mulligan, and D. Starr, Nucl. Phys. B **798**, 491 (2008), 0710.4348.