

SLAC-PUB-3059
LBL-15733
February 1983
(T/E)

Inclusive Electron Production in e^+e^- Annihilation at 29 GeV.*

M. E. Nelson, A. Blondel,⁽¹⁾ G. H. Trilling, G. S. Abrams, D. Amidei,
C. A. Blocker, A. M. Boyarski, M. Breidenbach, D. L. Burke,
W. Chinowsky, W. E. Dieterle,⁽²⁾ J. B. Dillon, J. M. Dorfan,
M. W. Eaton, G. J. Feldman, M. E. B. Franklin, G. Gidal, L. Gladney,
M. S. Gold, G. Goldhaber, L. J. Golding, G. Hanson, R. J. Hollebeek,
W. R. Innes, J. A. Jaros, A. D. Johnson, J. A. Kadyk, A. J. Lankford,
R. R. Larsen, B. LeClaire, M. Levi, N. Lockyer, V. Lüth, C. Matteuzzi,
J. F. Patrick,⁽³⁾ M. L. Perl, B. Richter, P. C. Rowson, T. Schaad,
H. M. Schellman, D. Schlatter, R. F. Schwitters, P. D. Sheldon,
J. Strait,⁽⁴⁾ C. de la Vaissière, J. M. Yelton, and C. Zaiser

Lawrence Berkeley Laboratory and Department of Physics
University of California, Berkeley, California 94720

Stanford Linear Accelerator Center
Stanford University, Stanford, California 94305

Department of Physics
Harvard University, Cambridge, Massachusetts 02138

(Submitted to Physical Review Letters)

* Work primarily supported by the Department of Energy under
Contracts No. DE-AC03-76SF00515, DE-AC03-76SF00098 and
DE-AC02-76ER03064.

- (1). Present address: LPNHE, Ecole Polytechnique, F-91128 Palaiseau,
France
- (2). Present address: University of Arizona, Tucson, AZ 85721
- (3). Present address: Fermilab, Batavia, IL 60510
- (4). Present address: University of Massachusetts, Amherst, MA 01002

ABSTRACT

We have measured the total momentum and transverse momentum spectra of electrons in e^+e^- annihilation at 29 GeV. The inclusive cross section is determined to be $14.4 \pm 1.6 \pm 5.2$ pb for momenta greater than 2 GeV/c. The average semi-electronic branching ratios of charm and bottom quarks are measured to be $(6.3 \pm 1.2 \pm 2.1)\%$ and $(11.6 \pm 2.1 \pm 1.7)\%$ respectively. The fragmentation function for bottom quarks is determined to be peaked at high z , with $\langle z \rangle_b = 0.75 \pm 0.05 \pm 0.04$.

Prompt lepton production in hadronic events from high energy e^+e^- annihilation is an excellent signal for the presence of hadrons containing charm (c) or bottom (b) quarks. The production rates and momentum spectra of such leptons depend on the weak decay semileptonic branching ratios and the momentum spectra of the parent hadrons. In turn the momentum spectra of these hadrons provide information on the fragmentation properties of c and b quarks. Simple kinematical considerations suggest that, as quark masses increase, hadrons containing the heavy quark carry an increasing fraction of its momentum.¹ This situation contrasts with the observation that light quarks fragment principally into low momentum hadrons. There is recent experimental evidence that charm quark fragmentation does lead to a harder momentum spectrum than is the case for light quark fragmentation.²⁻⁴ In this Letter we obtain the first experimental information on b quark fragmentation.

In this analysis we measure the total momentum and transverse momentum spectra of prompt electrons in hadronic events from e^+e^- annihilation at a center of mass (c.m.) energy of 29 GeV. The transverse momentum p_{\perp} is measured with respect to the thrust axis defined by all the charged particles in the event. The harder p_{\perp} distribution of electrons from bottom decays relative to charm decays allows us to separate the contributions of b and c quarks to the prompt electron signal.

The data, collected with the MARK II detector at the electron positron storage ring PEP, correspond to an integrated luminosity of 35 pb^{-1} . The MARK II detector has been described elsewhere,⁵ and we recall

here only the elements essential to our analysis. Charged particle momenta are measured over 76% of 4π in a cylindrical drift chamber in an axial magnetic field. The rms momentum resolution is given by $(\delta p/p)^2 \approx (.015)^2 + (.01p)^2$, where the momentum p is in units of GeV/c. Electron identification is accomplished over 64% of 4π by combining this momentum determination with measurements of energy deposition in a lead-liquid-argon calorimeter.⁶ In this calorimeter, spatial information is obtained by sampling energy depositions on strips which run in three different directions.

The electron-hadron separation algorithm is based on measurements of the ratio $r_i \equiv E_i/p$, where E_i is the energy deposition in one of three groupings, $i=1-3$, of layers in the calorimeter. Each of these groupings combines all layers in the first 8 radiation lengths which have the same strip orientation. To minimize the effects of neighboring particles, particularly photons, the energy deposition E_i is taken from a narrow lateral region, comparable in size to a strip width (≈ 4 cm), centered about the extrapolated particle trajectory. The algorithm demands that each value of r_i and that $\sum r_i$ be greater than an appropriate minimum value. The electron identification efficiency was determined with electrons from Bhabha events and photon conversions. This efficiency varies from 78% at 1 GeV/c to 93% at the highest momenta.

We determined the probability, as a function of p and p_{\perp} , that a hadron will be misidentified as an electron as follows. First, hadron interactions in our calorimeter were studied using data taken in a pion beam and using pions from the decay $\psi \rightarrow 2(\pi^+\pi^-)\pi^0$ in data taken at the SPEAR storage ring. Second, to measure the effect of accidental overlap

with nearby photons, we took advantage of the back-to-back jet topology of most hadronic events at 29 GeV. Charged tracks in these events were reversed and projected to the opposite jet. Energy depositions associated with the inverted track arise purely from accidental overlap. To determine overall misidentification probabilities we added hadron energy depositions, obtained from the beam test and ψ data, to the overlap contributions, and input the sums into the electron separation algorithm. The misidentification probabilities are typically 0.5%, but can be as large as 3% for a track of momentum 1 GeV/c in the core of a jet.

A hadronic event sample was selected by requiring a charged multiplicity of at least five and total detected energy (charged + neutral) greater than 25% of the c.m. energy. There were a total of 10691 such events. In this sample there are sources of real electrons which are not part of the prompt signal. A visual scan was performed to remove a small number of electron candidates arising from τ pair production, beam-gas interactions, and the process $e^+e^- \rightarrow e^+e^- + \text{hadrons}$. Background electrons from photon conversions and Dalitz decays were removed by a pair finding algorithm, which searched for low invariant mass combinations with oppositely charged tracks. This algorithm removes about 70% of the electrons coming from pairs while only removing 2% of the real prompt electrons. After removal of these backgrounds on an event by event basis, 930 electron candidates with $p > 1$ GeV/c remained. These were partitioned into 24 p, p_{\perp} bins.

In each p, p_{\perp} bin, we subtracted the remaining backgrounds and corrected for geometrical and identification efficiencies. The

background from misidentified hadrons was calculated from the number of observed charged hadron tracks in each bin and the appropriate misidentification probability. The background from remaining electron pairs was calculated using the number of electrons from identified pairs in each bin and the known efficiency of the pair finding algorithm. Fig. 1 shows the corrected signal in all 24 bins. Two sets of errors are plotted for each point. The smaller error bars represent the statistical errors. The larger ones show the quadratic sum of the statistical and systematic errors. The systematic errors are dominated by the uncertainty in the hadron misidentification probabilities. In three of the lowest momentum and transverse momentum bins [Fig. 1(a): $p < 3.0$ GeV/c and Fig. 1(b): $p < 2.0$ GeV/c] backgrounds account for almost 75% of the observed signal, thus these points have substantial systematic errors. In the remaining bins, however, the background levels are much less severe. Excluding these three bins, the background fractions for Fig. 1 (a)-(d) are 46%, 36%, 30%, and 23% respectively. Fig. 2 shows the total momentum and transverse momentum differential cross sections for prompt electrons with $p > 2$ GeV/c. The total inclusive cross section for this momentum range is determined to be $14.4 \pm 1.6 \pm 5.2$ pb.

We have performed a maximum likelihood fit to the observed populations in the various p, p_{\perp} bins, accounting for the signal above background in terms of the following contributions:

- (i) Bottom decays in $b\bar{b}$ events (b primary).
- (ii) Charm decays in $b\bar{b}$ events (c secondary)
- (iii) Charm decays in $c\bar{c}$ events (c primary)

We have excluded from the fit the three bins in which the background contributions strongly dominate. To represent contributions (i)-(iii) we have used a Monte Carlo simulation with a Feynman-Field hadronization model⁷ and gluon radiation as incorporated by Ali et al.⁸ We have verified that the semileptonic electron spectra produced by the heavy meson decay models in the Monte Carlo agree satisfactorily with measured spectra from the DELCO⁹ (charm) and CLEO¹⁰ (bottom) experiments.

We have parameterized the fragmentation function D_Q^H for quark Q into hadron H by an expression of the form:¹¹

$$D_Q^H(z) = \frac{A}{z[1 - 1/z - \epsilon_Q/(1-z)]^2} \quad (1)$$

where z = ratio of hadron energy to quark energy, A is a normalization factor, and ϵ_Q is a parameter. The average z of the distribution increases as ϵ_Q is reduced. It has been shown¹² that Eq. (1), with $\epsilon_Q \approx 0.25$, is a satisfactory representation of measured charmed meson momentum spectra.

We have fit the contributions (i)-(iii) in terms of $B_e(b)$ and $B_e(c)$, the average semi-electronic branching ratios in b quark and c quark decay, and ϵ_b and ϵ_c , the parameters in Eq. (1) for b and c fragmentation. Average quark semi-electronic branching ratios are equivalent to averages over all weakly decaying mesons and baryons, weighted by their relative populations. The ϵ_Q parameters characterize average fragmentation functions of quarks into these hadrons, weighted by the product of the populations and the semi-electronic branching ratios of the hadrons. These qualifications are particularly relevant

to the charmed mesons with different B_e for D^+ and D^0 .^{5,13} We have assumed that $B_e(c)$ is the same for contributions (ii) and (iii), and that all bottom decays lead to charm.^{10,14} Our results are not sensitive to reasonable deviations from these assumptions.

In the fit, we have fixed ϵ_c at 0.25 in accordance with previous measurements of c quark fragmentation.¹² For the remaining parameters we obtain $B_e(b) = (11.6 \pm 2.1 \pm 1.7)\%$, $B_e(c) = (6.3 \pm 1.2 \pm 2.1)\%$, and $\epsilon_b = 0.030^{+0.032}_{-0.018} \quad ^{+0.023}_{-0.014}$. The histograms in Fig. 1 show the results of the fit and the relative contributions of (i)-(iii) to the prompt electron signal in each p, p_\perp bin. The $\chi^2/D.F.$ of the fit is 14.0/18, and the model that we have used gives a very good representation of the data over all p and p_\perp studied. We have verified that the value chosen for ϵ_c is consistent with the data by performing an additional fit in which it was allowed to vary. The value obtained is compatible within errors with the input value of 0.25. The uncertainty in the charm fragmentation function is reflected in the systematic errors on ϵ_b . Since we chose the parameterization in Eq. (1) a priori, we have not established the detailed shape of the fragmentation function, but only its peaking at large z . We have also studied a parameterization different from Eq. (1), namely of the form $z^\alpha(1-z)$, and obtain qualitatively similar results. For either of these parameterizations, the average value of z_b is $\langle z \rangle_b = 0.75 \pm 0.05 \pm 0.04$.

In conclusion, we have measured the total momentum and transverse momentum spectra for prompt electrons in hadronic events in e^+e^- annihilation at 29 GeV. We have extracted information on c and b quark semi-electronic branching ratios and the b quark fragmentation function

based on a fit to these spectra. The values for the average semi-electronic branching ratios, $B_e(c) = (6.3 \pm 1.2 \pm 2.1)\%$ and $B_e(b) = (11.6 \pm 2.1 \pm 1.7)\%$, agree well with previous measurements^{5,9,10,13-15} even though they may represent averages over slightly different hadron populations. The value of ϵ_b and the corresponding average value of $\langle z \rangle_b = 0.75 \pm 0.05 \pm 0.04$ strongly support the theoretical expectations of a bottom quark fragmentation function which is peaked at large z .

This work was primarily supported by the Department of Energy under Contracts No. DE-AC03-76SF00515, DE-AC03-76SF00098 and DE-AC02-76ER03064.


REFERENCES

1. J. D. Bjorken, Phys. Rev. D17, 171 (1978); M. Suzuki, Phys. Lett. 71B, 139 (1977).
2. J. M. Yelton et al., Phys. Rev. Lett., 49, 430 (1982).
3. N. Abramowicz et al., Z. Phys. C15, 19 (1982).
4. W. B. Atwood, SLAC-PUB-2980, (1982).
5. R. Schindler et al., Phys. Rev. D24, 78 (1981).
6. G. S. Abrams, IEEE Trans. Nucl. Sci., NS27, 59 (1980).
7. R. D. Field and R. P. Feynman, Nucl. Phys. B136, 1 (1978).
8. A. Ali et al., Phys. Lett., 93B, 155 (1980).
9. W. Bacino et al., Phys. Rev. Lett. 43, 1073 (1979).
10. K. Chadwick et al., CLNS-82/546, (1982).
11. C. Peterson et al., Phys. Rev. D27, 105 (1983).
12. D. Schlatter, SLAC-PUB-2982, (1982).
13. W. Bacino et al., Phys. Rev. Lett. 45, 329 (1980).
14. L. J. Spencer et al., Phys. Rev. Lett. 47, 771 (1981).
15. R. Brandelik et al., Phys. Lett. 70B, 125 and 387 (1977); J. M. Feller et al., Phys. Rev. Lett. 40, 274 and 1677 (1978).


Figure Captions

Fig. 1 Prompt electron momentum spectra in 4 regions of transverse momentum p_{\perp} (GeV/c): (a) $p_{\perp} < 0.5$, (b) $0.5 < p_{\perp} < 1.0$, (c) $1.0 < p_{\perp} < 1.5$, and (d) $p_{\perp} > 1.5$. Two sets of error bars are shown for each data point. The smaller ones are statistical only. The larger ones are the statistical and systematic errors added in quadrature. The highest momentum bin includes all momenta ≥ 6 GeV/c. The histograms show the results of the fit. The three contributions shown are (i) b primary (solid), (ii) c secondary (diagonally hatched), and (iii) c primary (unshaded).

Fig. 2 Differential cross sections for prompt electrons with $p > 2$ GeV/c: (a) total momentum and (b) transverse momentum. Two sets of error bars are shown for each data point. The smaller ones are statistical only. The larger ones are the statistical and systematic errors added in quadrature.



2-83



4466A4

Fig. 2