

SLAC – PUB – 3148  
ITP – 800  
November 1985  
(T)

QUANTUM CHROMODYNAMIC  
EVOLUTION OF SIX-QUARK STATES\*

CHUENG-RYONG JI

*Stanford Linear Accelerator Center  
Stanford University, Stanford, California, 94305*

and

*Institute of Theoretical Physics, Department of Physics  
Stanford University, Stanford, California 94305*

and

STANLEY J. BRODSKY

*Stanford Linear Accelerator Center  
Stanford University, Stanford, California, 94305*

Submitted to *Physical Review D*

---

\* Work supported in part by the Department of Energy, contract DE-AC03-76SF00515 and by the National Science Foundation, grant PHY-81-07395.

## ABSTRACT

The evolution of six-quark color-singlet state distribution amplitudes is formulated as an application of perturbative quantum chromodynamics to nuclear wave functions. We derive and solve a set of coupled evolution equations for the deuteron S-wave amplitude. The solution of the evolution equations leads to a general matrix representation of anomalous dimensions which can be used to analyze the deuteron wavefunction at short distances.

## 1. Introduction

In the past few years a number of new applications of perturbative quantum chromodynamics to nuclear physics have been explored<sup>1</sup> including a qualitative description of the nuclear force in terms of quark exchange, and detailed predictions for the electromagnetic interactions of nuclei at large momentum transfer. Since the basic scale of QCD,  $\Lambda_{\overline{MS}}$ , is phenomenologically of order of a few hundred MeV or less, QCD predicts a transition from the traditional meson and nucleon degrees of freedom of nuclear physics to quark and gluon degrees of freedom at internucleon separations of a fermi or less. In addition, because of asymptotic freedom, perturbative QCD calculations should become relevant at momentum transfer scales of order of 1 GeV or even less.<sup>2</sup>

Recently, we have presented detailed QCD predictions for the asymptotic high- $Q^2$  behavior of the deuteron form factor which are, in principle, exact dynamical predictions of nuclear physics.<sup>3</sup> One of the most convenient and physical formalisms for analyzing exclusive processes with large transverse momenta is the QCD evolution formalism, based on a reformulation of the Bethe-Salpeter equation at equal light-cone time. In this paper, we present a detailed derivation of a set of six-quark evolution equations for the deuteron S-wave amplitude and a convenient way to solve the derived equations. We then construct a general matrix representation of the anomalous dimensions from explicit solutions of the evolution equations in order to analyze the deuteron wavefunction at short distances.

The evolution of the amplitude for simpler hadrons such as quark-antiquark meson<sup>4,5</sup> and three quark baryon<sup>4</sup> systems have already been formulated and solved. While these conventional hadrons have only one color singlet representa-

tion, the six-quark systems considered here have five independent color singlet representations. The formulation of the evolution equation for totally antisymmetric six-quark states is not trivial even though it is a natural extension of the three-quark case.<sup>4</sup> We have presented a general method for solving the QCD evolution equations which govern relativistic multi-quark wave functions.<sup>6</sup> We have also applied it to a four-quark toy system in  $SU(2)_C$  and derived some constraints on the effective force between two baryons.<sup>7</sup> However, since the antisymmetric representation of a multi-quark wave function must be constructed explicitly, it is hard in practice to solve the multi-quark evolution equation. In this paper we avoid this problem by exploiting the permutation symmetry of the evolution kernel.

In Sec. 2, a completely antisymmetric six-quark wave function is constructed and an example of an explicit representation is presented. In Sec. 3, we derive a set of evolution equations for the deuteron S-wave amplitude through a generalized kernel equation for a completely antisymmetric six-quark wave function. A convenient way to solve these coupled evolution equations is presented in Sec. 4. In Sec. 5, the general matrix representation for the anomalous dimension is obtained. Results for the leading anomalous dimension are given in detail. Discussions and conclusions are followed in Sec. 6. In Appendix A we describe a general method determining color singlet representations and explain the methods leading to the explicit representations given in Sec. 2. In Appendix B we present the color factor calculations and the detailed expressions for the orthogonal kernels which have specific permutation symmetries.

## 2. Six-Quark States

Six-quark states can be classified by their symmetries under  $SU(3)_C$  (color),  $SU(2)_T$  (isospin),  $SU(2)_S$  (spin), and spatial symmetry. Since the physical states are color singlets, the Young symmetry of the color singlet states of six-quark system is  $f_C = (222)$  or  $\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}$ . In the six-quark system, there are five independent color singlet states corresponding to five different Yamanouchi labels<sup>8</sup> of  $(222)$  symmetry. The explicit representations of the five independent color singlet states and their correspondence to Yamanouchi labels is given in Appendix A.

The completely antisymmetric six-quark representation  $|(1^6)[654321] \rangle_{f_T, f_{CSO}}$ <sup>9</sup> which has Young symmetries for isospin ( $SU(2)_T$ ) labelled by  $f_T$  is given by<sup>10</sup>

$$|(1^6)[654321] \rangle_{f_T, f_{CSO}} = \frac{1}{\sqrt{N_T}} \sum_{Y_T} \eta_T |f_T Y_T \rangle_{f_T} |\tilde{f}_T \tilde{Y}_T \rangle_{f_{CSO}} \quad , \quad (2.1)$$

where  $Y_T$  is the allowed Yamanouchi label of  $f_T$  ( $f_T$  symmetry has  $N_T$  different Yamanouchi labels) and the phase  $\eta_T = \pm 1 = (-1)^{Y_T}$  depends on whether  $Y_T$  is obtained from the Yamanouchi label with the indices in natural order by an even or odd number of transpositions. The dual symmetry states in the  $CSO$  space (represented by wavy lines) are used to construct the overall antisymmetric representations. It is convenient to construct a basis of completely antisymmetric six-quark representations from the combined color-spin symmetry. We thus introduce color-spin as an intermediate representation. For example, the specific projection of the the completely antisymmetric six-quark representation  $|(1^6)[654321] \rangle_{f_T, f_{CS}, f_O}$ <sup>9</sup> which has Young symmetries for color-spin ( $SU(6)_{CS}$ ),

and orbital symmetry labelled by  $f_{CS}$ , and  $f_O$  respectively, is represented by<sup>10</sup>

$$\begin{aligned}
| (1^6)[654321] \rangle_{f_T, f_{CS}, f_O} &= \frac{1}{\sqrt{N_T}} \sum_{Y_T} \sum_{Y_{CS}} \sum_{Y_O} \eta_T \langle f_{CS} Y_{CS}, f_O Y_O | \tilde{f}_T \tilde{Y}_T \rangle \\
&\times | f_T Y_T \rangle | f_{CS} Y_{CS} \rangle | f_O Y_O \rangle \quad , \quad (2.2)
\end{aligned}$$

where  $Y_{CS}$ , and  $Y_O$  are the allowed Yamanouchi labels of  $f_{CS}$ , and  $f_O$ , and  $\langle f_{CS} Y_{CS}, f_O Y_O | \tilde{f}_T \tilde{Y}_T \rangle$  are Clebsch - Gordon coefficients of the permutation group  $S_6$ . Now we decompose further  $f_{CS}$  into  $f_C$  and  $f_S$  using Clebsch - Gordon coefficients  $\langle f_C Y_C, f_S Y_S | f_{CS} Y_{CS} \rangle$  so that Eq. (2.1) becomes

$$\begin{aligned}
| (1^6)[654321] \rangle_{f_T, f_C, f_S, f_O} &= \frac{1}{\sqrt{N_T}} \sum_{Y_T} \sum_{Y_{CS}} \sum_{Y_O} \sum_{Y_C} \sum_{Y_S} (-1)^{Y_T} \\
&\times \langle f_{CS} Y_{CS}, f_O Y_O | \tilde{f}_T \tilde{Y}_T \rangle \langle f_C Y_C, f_S Y_S | f_{CS} Y_{CS} \rangle \\
&\times | f_T Y_T \rangle | f_C Y_C \rangle | f_S Y_S \rangle | f_O Y_O \rangle \quad . \quad (2.3)
\end{aligned}$$

Since our purpose is to formulate and solve a generalized evolution equation, it is useful to project Eq. (2.1) onto the light-cone momentum space of six-quarks, each carrying light-cone longitudinal momentum fraction  $x_i = (q_0^i + q_3^i)/(P_0 + P_3)$  of the deuteron's momentum  $P_\mu$  ( $\sum_{i=1}^6 x_i = 1$ ) and transverse momentum  $q_\perp^i$  ( $\sum_{i=1}^6 q_\perp^i = 0$ ). The corresponding light-cone wave function of six-quark system  $\Psi_d(x_i, q_\perp^i)$  is defined by

$$\begin{aligned}
\Psi_d(x_i, q_\perp^i) &= \langle x_i, q_\perp^i \rangle | (1^6)[654321] \rangle_{f_T, f_{CS}, f_O} \\
&= \frac{1}{\sqrt{N_T}} \sum_{Y_T=1}^{N_T} \sum_{Y_{CS}=1}^{N_{CS}} \sum_{Y_O=1}^{N_O} (-1)^{Y_T} \langle f_{CS} Y_{CS}, f_O Y_O | \tilde{f}_T \tilde{Y}_T \rangle \quad (2.4) \\
&\times \psi_o(x_i, q_\perp^i) | f_T Y_T \rangle | f_{CS} Y_{CS} \rangle \quad ,
\end{aligned}$$

where the orbital wave function is given by

$$\psi_o \left( x_i, q_{\perp}^{(i)} \right) = \langle x_i, q_{\perp}^{(i)} | f_O Y_O \rangle . \quad (2.5)$$

The remaining  $SU(2)_T$  and  $SU(6)_{CS}$  symmetries ( $|f_T Y_T \rangle$  and  $|f_{CS} Y_{CS} \rangle$ ) are given by specific tensor representations.

The probability amplitude for the constituents with light-cone momentum fraction  $\bar{x}_i$  to combine into the hadron with relative transverse momentum up to the scale  $Q^2$  is given by the distribution amplitude  $\Phi_d(x_i, Q)$  defined by

$$\Phi_d(x_i, Q) \equiv \int \prod_{i=1}^6 \left( \frac{d^2 q_{\perp}^{(i)}}{16\pi^3} \right) 16\pi^3 \delta^2 \left( \sum_i q_{\perp}^{(i)} \right) \Psi_d^{(Q)} \left( x_i, q_{\perp}^{(i)} \right) , \quad (2.6)$$

where the  $Q$  dependence of  $\Psi_d^{(Q)}$  comes from the renormalization of the quark fields.<sup>11</sup>

As an explicit example of a six-quark representation, we give a specific representation for  $f_T = (33)$ ,  $f_{CS} = (222)_C \times (6)_S$ , and  $f_O = (6)$  ( $T = 0$ ,  $S = S_Z = 3$ , and  $S$ -wave);

$$\begin{aligned}
\Phi_d(x_i, Q) = & \frac{a_0}{48\sqrt{5}} \left[ -\epsilon_{ijk}\epsilon_{lmn} (\epsilon_{ad}\epsilon_{be}\epsilon_{cf} + \epsilon_{ae}\epsilon_{bd}\epsilon_{cf} + \epsilon_{ad}\epsilon_{bf}\epsilon_{ce} + \epsilon_{af}\epsilon_{bd}\epsilon_{ce}) \right. \\
& + \epsilon_{ijl}\epsilon_{kmn} (\epsilon_{ac}\epsilon_{be}\epsilon_{df} + \epsilon_{ac}\epsilon_{bf}\epsilon_{de} + \epsilon_{ae}\epsilon_{bc}\epsilon_{df} + \epsilon_{af}\epsilon_{bc}\epsilon_{de}) \\
& - (\epsilon_{ijm}\epsilon_{kln} + \epsilon_{ijn}\epsilon_{klm}) (\epsilon_{ac}\epsilon_{bd}\epsilon_{ef} + \epsilon_{ad}\epsilon_{bc}\epsilon_{ef}) \\
& + (\epsilon_{ikm}\epsilon_{jln} + \epsilon_{ikn}\epsilon_{jlm} + \epsilon_{jkm}\epsilon_{iln} + \epsilon_{jkn}\epsilon_{ilm}) \epsilon_{ab}\epsilon_{cd}\epsilon_{ef} \\
& \left. - (\epsilon_{ikl}\epsilon_{jmn} + \epsilon_{jkl}\epsilon_{imn}) (\epsilon_{ab}\epsilon_{ce}\epsilon_{df} + \epsilon_{ab}\epsilon_{cf}\epsilon_{de}) \right] \\
& \times a_i^\dagger(1) b_j^\dagger(2) c_k^\dagger(3) d_l^\dagger(4) e_m^\dagger(5) f_n^\dagger(6) \\
& \times x_1 x_2 x_3 x_4 x_5 x_6 \left( \ln \frac{Q^2}{\Lambda^2} \right)^{-\gamma_0}, \tag{2.7}
\end{aligned}$$

where the indices  $i, j, \dots, n$  and  $a, b, \dots, f$  are the color ( $r, y, b$ ) and isospin ( $u, d$ ) indices, respectively. The  $\epsilon_{ijk}$ 's and  $\epsilon_{ab}$ 's are the completely antisymmetric Cartesian tensors of  $SU(3)_C$  and  $SU(2)_T$ . The coefficient  $a_0$  is the normalization of the orbital distribution amplitude (see Sec. 5). The leading anomalous dimension is  $\gamma_0$ . A detailed calculation of  $\gamma_0$  for the various six-quark states and the tensor representations in Eq. (2.7) will be given in Sec. 5 and the Appendix A, respectively.

In the following section, we will derive a set of evolution equations for the deuteron  $S$ -wave distribution amplitude. Since the deuteron is an isospin singlet and the  $S$ -wave is a symmetric orbital, the Young symmetry in each quantum space is given by  $f_T = (33)_T$ ,  $f_{CS} = (222)_{CS}$ , and  $f_O = (6)_O$ . Thus, an example of the ground-state  $S$ -wave of the deuteron distribution amplitude is represented by



$$|(1^6) \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array} \rangle_D = \frac{1}{\sqrt{5}} \sum_{Y_T} \sum_{Y_{CS}} \eta_T | \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \square & \square \\ \hline \square & \square \\ \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} Y_T \rangle | \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array} Y_{CS} \rangle | \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array} [111111]_O \rangle , \quad (2.8)$$

where  $(6)_O$  has only one Yamanouchi label  $[111111]_O$  [the Clebsch–Gordon coefficient in Eq. (2.2) is trivially given by 1]. We will concentrate on the the leading term in the high  $Q^2$  limit. It is an eigensolution of the evolution equation which will be derived in the next section. The leading term is the lowest power term of  $x_i$ -dependent polynomials i.e.  $x_1 x_2 x_3 x_4 x_5 x_6$ . For example, we can see that the basis element given by Eq. (2.7) is an eigensolution because only the symmetric tableau is allowed for the spin in  $S_Z = 3$  case. However, in general, and specifically for the deuteron, the eigensolutions will be given by mixing of basis elements in spin space.

### 3. Evolution Equations for the Deuteron

Each eigensolution of a six-quark state satisfies a kernel equation of the form

$$K|(1^6)[654321] \rangle_e = e|(1^6)[654321] \rangle_e , \quad (3.1)$$

where  $|(1^6)[654321] \rangle_e$  is an eigensolution with the eigenvalue  $e$  given by a linear combination of completely antisymmetric representations (basis elements). The kernel  $K$  is calculated to leading order in  $\alpha_s(Q^2)$  from one-gluon-exchange by using light-cone perturbation theory. It is given explicitly in Sec. 5. The pairwise one-gluon exchange diagrams are shown in Fig. 1. Since the isospin representation does not change through gluon exchange, the  $SU(2)_T$  symmetry has no mixing and is fixed. For example,  $f_T = (33)$  in the deuteron case. Also, the six-quark states are always color-singlet states i.e.  $f_C = (222)$ . Thus, in general, the

evolution equation has the following form in the basis of Eq. (2.3);

$$\begin{aligned}
& \sum_{f_{CS}} \sum_{f_S} \sum_{f_O} \sum_{Y_{CS}} \sum_{Y_C} \sum_{Y_S} \sum_{Y_O} C_{f_O f_S}^{f_{CS}} \langle f_C Y_C, f_S Y_S | f_{CS} Y_{CS} \rangle \\
& \quad \times \langle f_{CS} Y_{CS}, f_O Y_O | \tilde{f}_T \tilde{Y}_T \rangle K | f_C Y_C \rangle | f_S Y_S \rangle | f_O Y_O \rangle \\
& = e \sum_{f_{CS}} \sum_{f_S} \sum_{f_O} \sum_{Y_{CS}} \sum_{Y_C} \sum_{Y_S} \sum_{Y_O} C_{f_O f_S}^{f_{CS}} \langle f_C Y_C, f_S Y_S | f_{CS} Y_{CS} \rangle \\
& \quad \times \langle f_{CS} Y_{CS}, f_O Y_O | \tilde{f}_T \tilde{Y}_T \rangle | f_C Y_C \rangle | f_S Y_S \rangle | f_O Y_O \rangle .
\end{aligned} \tag{3.2}$$

The unknown coefficients  $C_{f_O f_S}^{f_{CS}}$  and the eigenvalue  $e$  are obtained by solving Eq. (3.2). In Eq. (3.2), the possible Young-tableaus  $f_{CS}$  and  $f_O$  in the sums are determined by the Clebsch–Gordan series of  $S_6$  to produce the  $CSO$  Young-tableau  $\tilde{f}_T$ . Likewise, the possible Young-tableaus  $f_S$  are determined to produce  $f_{CS}$  after combining with  $f_C$ . Each possible combination of  $f_S$ ,  $f_{CS}$  and  $f_O$  gives an equation (3.2). The combinations of  $f_S$ ,  $f_{CS}$  and  $f_O$  are given by coefficients of  $C_{f_O f_S}^{f_{CS}}$  and has a corresponding eigenvalue  $e$ .

In general there are many possible combinations of  $f_S$ ,  $f_{CS}$  and  $f_O$  contributing to Eq.(3.2), since only  $f_T$  and  $f_C$  are fixed and mixed Young symmetries are allowed. However, if we constrain ourselves to some special cases then only a few equations actually need to be solved. For example, the leading term in high  $Q^2$  limit of the  $S_Z = 3$  and  $T = 0$  amplitude has only one possible combination of  $f_S$ ,  $f_{CS}$  and  $f_O$  (i.e.  $f_T = (33)$ ,  $f_C = (222)$ ,  $f_S = (6)$ ,  $f_{CS} = (222)$ ,  $f_O = (6)$ ) and only one equation needs be constructed for this special case. Therefore, we can easily see that the explicit representation given by Eq. (2.7) is an eigensolution itself as we already mentioned in Sec. 2.

In any case, the unknown coefficients  $C_{f_O f_S}^{f_{CS}}$  and eigenvalue  $e$  in Eq. (3.2)

can be determined after a given set of equations (corresponding to the number of possible combinations of  $f_S$ ,  $f_{CS}$  and  $f_O$ ) are solved.

In this paper, we will concentrate on analyzing the asymptotic amplitude which dominates exclusive processes at large transverse momenta such as the asymptotic high- $Q^2$  behavior of the deuteron form factor. However, the general equation is given by Eq. (3.2), and the method which we present in the rest of this section can be applied to arbitrary cases. Since a deuteron is isospin singlet and the dominant degree of freedom in high- $Q^2$  limit is  $S$ -wave, the eigensolutions which we are considering have  $f_T = (33)$  and  $f_O = (6)$ . In the  $S_Z = 1$  case, one has mixing between  $f_S = (6)_S$  and  $f_S = (42)_S$ .

For the  $f_T = (33)_T, f_O = (6)_O$  case, Eq. (3.2) becomes

$$\begin{aligned}
& \sum_{f_S} \sum_{Y_C} \sum_{Y_S} C_{(6)f_S}^{(222)} < (222)_C Y_C, f_S Y_S | (222)_{CS} Y_{CS} > \\
& \quad \times K | (222)_C Y_C > | f_S Y_S > | (6)_O [111111]_O > \\
& = e \sum_{f_S} \sum_{Y_C} \sum_{Y_S} C_{(6)f_S}^{(222)} < (222)_C Y_C, f_S Y_S | (222)_{CS} Y_{CS} > \\
& \quad \times | (222)_C Y_C > | f_S Y_S > | (6)_O [111111]_O >,
\end{aligned} \tag{3.3}$$

where  $f_{CS} = \tilde{f}_T = (\tilde{3}\tilde{3}) = (222)$  and  $Y_{CS} = \tilde{Y}_T$  because  $< f_{CS} Y_{CS}, (6)_O [111111]_O | \tilde{f}_T \tilde{Y}_T > = \delta_{f_{CS} \tilde{f}_T} \times \delta_{Y_{CS} \tilde{Y}_T}$ .

Since we know from the Clebsch-Gordon series of  $S_6$  that the possible values of  $f_S$  are (6) and (42), we have two unknown coefficients  $C_{(6)(6)}^{(222)}$  and  $C_{(6)(42)}^{(222)}$  which must satisfy the normalization condition,

$$\left( C_{(6)(6)}^{(222)} \right)^2 + \left( C_{(6)(42)}^{(222)} \right)^2 = 1 \quad . \tag{3.4}$$

Because of the normalization condition (3.4), one can define two eigensolutions in terms of one mixing angle  $\theta$ . The corresponding eigenvalues are:

$$\begin{aligned}
e = e_1 \text{ when } C_{(6)(6)}^{(222)} &= \cos \theta, & C_{(6)(42)}^{(222)} &= \sin \theta \\
e = e_2 \text{ when } C_{(6)(6)}^{(222)} &= -\sin \theta, & C_{(6)(42)}^{(222)} &= \cos \theta.
\end{aligned}
\tag{3.5}$$

Furthermore, the kernel  $K$  has the factorized color factor corresponding to one gluon exchange

$$K = \sum_{i \neq j} \left( \frac{\vec{\lambda}_i}{2} \cdot \frac{\vec{\lambda}_j}{2} \right) V_{ij}, \tag{3.6}$$

where each term represent the kernel given by the interaction between the  $i^{th}$  and  $j^{th}$  quark and each component of the eight dimensional vector  $\vec{\lambda}$  is the Gell-Mann matrix of the  $SU(3)_C$  group. If we sandwich Eq. (3.5) between two color states which have  $Y_C = \alpha$  and  $Y_C = \beta$  respectively, we can define a  $5 \times 5$  matrix representation of  $K$  whose elements are given by

$$K_{\alpha\beta} \equiv \langle (222)_{C\alpha} | K | (222)_{C\beta} \rangle = \sum_{i \neq j}^6 C_{\alpha\beta}(i, j) V_{ij}, \tag{3.7}$$

where

$$C_{\alpha\beta}(i, j) = \langle (222)_{C\alpha} \left| \frac{\vec{\lambda}_i}{2} \cdot \frac{\vec{\lambda}_j}{2} \right| (222)_{C\beta} \rangle. \tag{3.8}$$

The most important observation in this formulation is that the kernel  $K$  is a linear combination of the operators  $\Theta_{fY}$  in color space each of which has a

definite Young symmetry  $f$  with Yamanouchi labels  $Y$ ;

$$K = \sum_{fY} K_{fY} \Theta_{fY}.$$

Therefore the kernel element  $K_{\alpha\beta}$  can be rewritten in terms of the kernel  $K_{fY}$

$$\begin{aligned} K_{\alpha\beta} &= \langle (222)_{C\alpha} | \sum_{fY} K_{fY} \Theta_{fY} | (222)_{C\beta} \rangle \\ &= \sum_f \sum_Y \langle (222)_{C\alpha}, fY | (222)_{C\beta} \rangle K_{fY} \end{aligned} \quad (3.9)$$

where the possible  $f$  which gives the non-zero Clebsch-Gordon coefficient  $\langle (222)_{C\alpha}, fY | (222)_{C\beta} \rangle$  is only (6) or (42). One can rewrite  $K_{fY}$  in terms of color factors  $C_{\alpha\beta}(i, j)$  and  $V_{ij}$ ;

$$K_{(6)[111111]} = \frac{1}{5} \sum_{\alpha} \sum_{i \neq j} C_{\alpha\alpha}(i, j) V_{ij} = -\frac{1}{5} C_F \sum_{i \neq j} V_{ij}, \quad \left( C_F = \frac{4}{3} \right) \quad (3.10)$$

$$\begin{aligned} K_{(42)Y} &= \frac{9}{5} \sum_{\alpha} \sum_{\beta} \langle (222)_{C\alpha}, (42)Y | (222)_{C\beta} \rangle C_{\alpha\beta}(i, j) V_{ij}, \\ &\quad (Y = 1, \dots, 9). \end{aligned} \quad (3.11)$$

In appendix B, we present details of the derivation of Eq. (3.10) and (3.11) and the color factors of Eq. (3.8). Note there are five possible labels  $Y_{CS}$  in correspondence with the labels  $\beta = 1, \dots, 5$  of  $Y_C$ . If we project Eq. (3.3) with  $Y_{CS} = \beta$  by a color singlet state  $Y_C = \alpha$  then we get a set of two equations using

Eqs. (3.5) through (3.11). One of them is given by

$$\begin{aligned}
& \cos \theta \left( \delta_{\alpha\beta} K_{(6)[111111]} + \sum_Y \langle (222)_C \alpha, (42)_Y | (222)_{CS} \beta \rangle K_{(42)Y} \right) \\
& \times |(6)_S [111111]_S \rangle | (6)_O [111111]_O \rangle \\
& + \sin \theta \left( \sum_{Y_S} \langle (222)_C \alpha, (42)_S Y_S | (222)_{CS} \beta \rangle K_{(6)[111111]} \right. \\
& \left. + \sum_{Y_S} \sum_Y \sum_{\gamma} \langle (222)_C \gamma, (42)_S Y_S | (222)_{CS} \beta \rangle \langle (222)_C \alpha, (42)_Y | (222)_C \gamma \rangle K_{(42)Y} \right) \\
& \times |(42)_S Y_S \rangle | (6)_O [111111]_O \rangle = e_1 \left( \cos \theta \delta_{\alpha\beta} |(6)_S [111111]_S \rangle | (6)_O [111111]_O \rangle \right. \\
& \left. + \sin \theta \sum_{Y_S} \langle (222)_C \alpha, (42)_S Y_S | (222)_{CS} \beta \rangle |(42)_S Y_S \rangle | (6)_O [111111]_O \rangle \right), \tag{3.12}
\end{aligned}$$

and another one is given by substitutions  $\theta \rightarrow \theta + \frac{\pi}{2}$  and  $e_1 \rightarrow e_2$  in Eq. (3.12).

Combining the two equations and using properties of Clebsch–Gordon coefficients we get the following set of evolution equations (we drop the trivial orbital factor  $| (6)_O [111111]_O \rangle$ ):

$$K_{(6)[111111]} |(6)_S [111111]_S \rangle = (e_1 \cos^2 \theta + e_2 \sin^2 \theta) |(6)_S [111111]_S \rangle \tag{3.13}$$

$$\frac{1}{9} \sum_Y K_{(42)Y} |K_{(42)S} Y \rangle = (e_1 - e_2) \cos \theta \sin \theta |(6)_S [111111]_S \rangle \tag{3.14}$$

$$K_{(42)Y} |(6)_S [111111]_S \rangle = (e_1 - e_2) \cos \theta \sin \theta |(42)_S Y \rangle \tag{3.15}$$

$$\begin{aligned}
K_{(6)[111111]}|(42)_s Y \rangle &+ \frac{9}{5} \sum_{\alpha} \sum_{\beta} \sum_{\gamma} \sum_{Y_S} \sum_{Y_K} \langle (222)_C \beta, (42)_s Y | (222)_C S \alpha \rangle \\
&\times \langle (222)_C \gamma, (42)_s Y_S | (222)_C S \beta \rangle \\
&\times \langle (222)_C \alpha, (42)_{Y_K} | (222)_C \gamma \rangle K_{(42)_{Y_K}} |(42)_s Y_S \rangle \\
&= (e_1 \sin^2 \theta + e_2 \cos^2 \theta) |(42)_s Y \rangle .
\end{aligned} \tag{3.16}$$

Since the operator order of the  $K_{fY}$  is irrelevant,  $[K_{(6)[111111]}, K_{(42)Y}] = 0$ , we can see from Eqs. (3.13) and (3.15)

$$K_{(6)[111111]}|(42)_s Y \rangle = (e_1 \cos^2 \theta + e_2 \sin^2 \theta) |(42)_s Y \rangle . \tag{3.17}$$

Furthermore, we can prove the following property of Clebsch Gordon coefficients;

$$\begin{aligned}
\sum_{\alpha} \sum_{\beta} \sum_{\gamma} \langle (222)_C \beta, (42)_s Y | (222)_C S \alpha \rangle \langle (222)_C \gamma, (42)_s Y_S | (222)_C S \beta \rangle \\
\times \langle (222)_C \alpha, (42)_{Y_K} | (222)_C \gamma \rangle = \frac{5\sqrt{53}}{108} \langle (42)_s Y_S, (42)_{Y_K} | (42)_s Y \rangle
\end{aligned} \tag{3.18}$$

Thus, if we combine Eqs. (3.16) through (3.18), then we get

$$\begin{aligned}
\sum_{Y_S} \sum_{Y_K} \frac{\sqrt{53}}{12} \langle (42)_s Y_S, (42)_{Y_K} | (42)_s Y \rangle K_{(42)_{Y_K}} |(42)_s Y_S \rangle \\
= (e_1 - e_2)(\sin^2 \theta - \cos^2 \theta) |(42)_s Y \rangle .
\end{aligned} \tag{3.19}$$

Eqs. (3.13), (3.14), (3.15), and (3.19) appear to be independent. However, we can see that only three combinations of  $e_1$ ,  $e_2$  and  $\theta$  can be determined in the above equations. So we need to solve three equations (for example, Eqs. (3.13), (3.15), and (3.19)) and the other equation Eq. (3.14) can be used to check the

results. We will describe a convenient way to solve these equations in the next section.

#### 4. Symmetry of Kernel Equations

The symmetry of the left and right hand sides of any of the equations derived in the last section should be conserved. This can be easily checked since each kernel and eigenstate has definite symmetry. If the kernel is symmetric, then the states of left and right hand side have the same symmetry. For the simple case where the  $x_i$ -dependence of the orbital distribution amplitude is given by the lowest power, such as the example we presented in the last section, all the equations become an eigenvalue equation where the eigenvalue is determined by  $S_Z$ . This explains why Eqs. (3.13) and (3.17) have the same eigenvalue  $e_1 \cos^2 \theta + e_2 \sin^2 \theta$ . For the symmetric kernel case, we can fix the color factor as  $-C_F/5$  (see Eq. (3.10)). We can generalize this procedure for any spin-orbit state and find the general matrix representation of the kernel in the basis of polynomials. This will be done in the next section.

However, the eigensolution of the symmetric kernel equation (3.13) is not a true solution of the whole system in general because there could be mixing i.e.  $\theta \neq 0$  in our example. We need to use other mixed symmetric kernel equations and find more constraints to determine  $e_1, e_2$  and  $\theta$ . In our example, we can find relations between three combinations of  $e_1, e_2$  and  $\theta$  by counting spin annihilation or surviving terms. [In the general case one must consider the  $x_i$ -dependent orbital distribution functions and their integration. One can find relations between combinations of eigenvalues and mixing angles case by case by counting the number of terms which are surviving or annihilated after the kernel operation.



The equation which we need to solve actually is the symmetric kernel equation presented in the next section.]

By counting spin terms, we find the following results for the leading anomalous dimension. If the symmetric kernel equation is given by

$$K_{(6)[111111]}|(6)_S[111111]_S \rangle = \gamma|(6)_S[111111]_S \rangle , \quad (4.1)$$

then

$$K_{(6)[111111]}|(42)_S Y \rangle = \gamma|(42)_S Y \rangle , \quad (4.2)$$

$$K_{(42)Y}|(6)_S[111111]_S \rangle = \frac{3\sqrt{6}}{16}\gamma|(42)_S[111111]_S \rangle , \quad (4.3)$$

$$\frac{1}{9}K_{(42)Y}|(42)_S Y \rangle = \frac{3\sqrt{6}}{16}\gamma|(6)_S[111111]_S \rangle , \quad (4.4)$$

and

$$\sum_{Y_S} \sum_{Y_K} \frac{\sqrt{53}}{12} \langle (42)_S Y_S, (42)_{Y_K} | (42)_S Y \rangle K_{(42)Y_K} | (42)_S Y_S \rangle = \frac{5}{48}\gamma | (42)_S Y \rangle . \quad (4.5)$$

Comparing with Eqs. (3.13), (3.14), (3.15), (3.17), and (3.19), we can solve  $e_1, e_2$ , and  $\theta$  in terms of  $\gamma$ . We find two solutions;

$$i) \quad e_1 = \frac{25}{16}\gamma, \quad e_2 = \frac{5}{8}\gamma, \quad \tan \theta = \frac{\sqrt{6}}{2} \quad (4.6)$$

$$ii) \quad e_1 = \frac{11}{8}\gamma, \quad e_2 = \frac{7}{16}\gamma, \quad \tan \theta = -\frac{\sqrt{6}}{3} .$$

However, if  $\gamma > 0$  then only *i*) solution is valid. The value of  $\gamma$  will be determined in the next section.

## 5. Symmetric Kernel Equation and Solution

The evolution equation with the symmetric kernel has the same spin-orbital symmetry  $f_{SO}$  and  $Y_{SO}$  in the left and right hand sides of the equation:

$$K_{(6)[111111]}|f_{SO}Y_{SO}\rangle = \Gamma|f_{SO}Y_{SO}\rangle, \quad (5.1)$$

where  $\Gamma$  is the eigenvalue of the symmetric kernel equation. In order to give a more explicit expression of Eq (5.1), we project both sides of Eq (5.1) to the light-cone momentum space of six-quarks as illustrated in Sec. 2. We then obtain a kernel equation for the spin-orbital wave function  $\psi_{SO}(x_i, q_\perp^i)$ ,

$$\psi_{SO}(x_i, q_\perp^i) = C_d \int [dy] \int [d^2 k_\perp^i] \langle x_i, q_\perp^i | \frac{1}{\Gamma} \sum_{k \neq \ell} V_{k\ell} | y_i, k_\perp^i \rangle \psi_{SO}(y_i, k_\perp^i) \quad (5.2)$$

where  $C_d = -C_F/5$ , and

$$[dy] = \delta \left( 1 - \sum_{i=1}^6 y_i \right) \prod_{i=1}^6 dy_i, \quad (5.3)$$

$$[d^2 k_\perp^i] = \prod_{i=1}^6 \left( \frac{d^2 k_\perp^i}{16\pi^3} \right) 16\pi^3 \delta^2 \left( \sum_i k_\perp^i \right).$$

Thus  $\psi_{SO}(x_i, q_\perp^i)$  is a linear combination of some orbital wave functions  $\phi_O(x_i, q_\perp^i)$  (see Eq. (2.5)) with some coefficients of spin tensor representations. By calculating the leading order diagrams shown in Fig. 1, one obtains an explicit expression of  $\langle x_i, q_\perp^i | \frac{1}{\Gamma} \sum_{k \neq \ell} V_{k\ell} | y_i, k_\perp^i \rangle$  for the spin-orbital wave function  $\psi(x_i, q_\perp^i)$  (the spin-orbit index  $SO$  is dropped here since each possible  $f_{SO}$  and  $Y_{SO}$  satisfies

the same equation):

$$\begin{aligned}
\psi(x_i, q_\perp^i) &= -2C_d \sum_{i \neq j} \frac{4\pi\alpha_s((q_\perp^i)^2)}{(q_\perp^i)^2} \prod_{k=1}^6 x_k \int [dy] \theta(y_i - x_i) \\
&\times \prod_{\ell \neq i, j}^6 \delta(y_\ell - x_\ell) \frac{y_j}{x_j} \left( \frac{\delta_{h_i \bar{h}_j}}{x_i + x_j} + \frac{1}{y_i - x_i} \right) \\
&\times \int [d^2 k_\perp^i] \frac{\psi(y_i, k_\perp^i)}{\prod_{k=1}^6 y_k} ,
\end{aligned} \tag{5.4}$$

where  $\alpha_s((q_\perp^i)^2) [\cong (4\pi/\beta)(\ell n(q_\perp^i)^2/\Lambda^2)^{-1}]$  is the QCD running coupling constant ( $\beta = 11 - 2/3n_f$ ,  $n_f$  is the effective number of flavors) and  $\delta_{h_i \bar{h}_j} = 1(0)$  when the constituents'  $\{i, j\}$  helicities are antiparallel (parallel). Eq. (5.4) has an infrared singularity at  $x_i = y_i$ . Thus, in order to obtain a well defined evolution equation of the six-quark system, we consider the quark distribution amplitude  $\phi(x_i, Q)$  which is the amplitude for finding constituents with longitudinal momenta  $x_i$  in the deuteron which are collinear up to the scale  $Q^2$ :

$$\phi(x_i, Q) \equiv \int^{Q^2} [d^2 q_\perp^{(i)}] \psi^{(Q)}(x_i, q_\perp^{(i)}) . \tag{5.5}$$

This definition is the same as Eq. (2.6) except that the only spin-orbital wave function  $\psi^{(Q)}(x_i, q_\perp^{(i)})$  is integrated instead of the total wave function  $\Psi_d^{(Q)}(x_i, q_\perp^{(i)})$ . By differentiating both sides of Eq. (5.5) with respect to  $Q^2$ , and combining with Eq. (5.4), we obtain the evolution equation of the six-quark system:

$$\prod_{k=1}^6 x_k \left[ \frac{\partial}{\partial \xi} + \frac{3C_F}{\beta} \right] \tilde{\phi}(x_i, Q) = -\frac{C_d}{\beta} \int_0^1 [dy] V(x_i, y_i) \tilde{\phi}(y_i, Q) , \tag{5.6}$$

where we use the definition  $\tilde{\phi}(x_i, Q)$  as

$$\phi(x_i, Q) = \prod_{k=1}^6 x_k \tilde{\phi}(x_i, Q) , \quad (5.7)$$

and the variable

$$\xi(Q^2) = \frac{\beta}{4\pi} \int_{Q_0^2}^{Q^2} \frac{dk^2}{k^2} \alpha_S(k^2) \sim \ln \left( \frac{\ln \frac{Q^2}{\Lambda^2}}{\ln \frac{Q_0^2}{\Lambda^2}} \right) , \quad (5.8)$$

and  $V(x_i, y_i)$  is given by

$$\begin{aligned} V(x_i, y_i) &= 2 \prod_{k=1}^6 x_k \sum_{i \neq j}^6 \theta(y_i - x_i) \prod_{\ell \neq i, j}^6 \delta(x_\ell - y_\ell) \frac{y_j}{x_j} \\ &\times \left( \frac{\delta_{h_i \bar{h}_j}}{x_i + x_j} + \frac{\Delta}{y_i - x_i} \right) = V(y_i, x_i) . \end{aligned} \quad (5.9)$$

By definition the factor  $\Delta$  in Eq. (5.9) means

$$\Delta \tilde{\phi}(y_i, Q) = \tilde{\phi}(y_i, Q) - \tilde{\phi}(x_i, Q) , \quad (5.10)$$

so we can see that the infrared singularity in Eq. (5.4) at  $x_i = y_i$  is completely cancelled by that in Eq. (5.5). We notice that this cancellation happens only when the correct value of deuteron color factor,  $C_d = -C_F/5$ , so that this is a good check of the correctness of  $C_d$ . We calculated  $C_d$  explicitly from the definition in the Appendix B.

Since the six-quark evolution equation (5.6) has a similar form to that of the three quark evolution equation,<sup>4</sup> we can solve this equation following the similar

methods to the three quark case. First, we separate the variables of  $\tilde{\phi}(x_i, Q)$  such as

$$\begin{aligned}\tilde{\phi}(x_i, Q) &= \tilde{\phi}(x_i) e^{-\gamma \xi} \\ &= \tilde{\phi}(x_i) \left( \ell n \frac{Q^2}{\Lambda^2} \right)^{-\gamma},\end{aligned}\tag{5.11}$$

and substitute into Eq. (5.6) so that Eq. (5.6) becomes [note that  $C_d = -C_F/5$ ]

$$\begin{aligned}\prod_{k=1}^6 x_k \left( -\gamma + \frac{3C_F}{\beta} \right) \tilde{\phi}(x_i) &= \frac{C_F}{5\beta} \int_0^1 [dy] V(x_i, y_i) \tilde{\phi}(y_i) \\ &\equiv \frac{C_F}{5\beta} V |\tilde{\phi} \rangle,\end{aligned}\tag{5.12}$$

where the equation is simply redefined by the quantum mechanical notation  $V |\tilde{\phi} \rangle$  and  $V(x_i, y_i)$  is given by Eq. (5.9). Next we expand  $\tilde{\phi}(x_i)$  in terms of eigenfunctions  $\tilde{\phi}_n(x_i)$ , so that the general solution of  $\tilde{\phi}(x_i, Q)$  is of the form

$$\tilde{\phi}(x_i, Q) = \sum_{n=0}^{\infty} a_n \tilde{\phi}_n(x_i) \left( \ell n \frac{Q^2}{\Lambda^2} \right)^{-\gamma_n},\tag{5.13}$$

where  $a_n$  are the coefficients and eigenvalues  $\gamma_n$  are corresponding to the anomalous dimensions of six-quark system. Since  $V(x_i, y_i)$  is both real and symmetric [ $V(x_i, y_i) = V(y_i, x_i)$ ], the  $\gamma_n$  are real. The  $\{\tilde{\phi}_n(x_i)\}_{n=0}^{\infty}$  are orthogonal with weight  $\omega(x_i) = x_1 x_2 x_3 x_4 x_5 x_6$ ,

$$\int_0^1 [dx] \omega(x_i) \tilde{\phi}_n^*(x_i) \tilde{\phi}_m(x_i) = K_n \delta_{nm},\tag{5.14}$$

where  $K_n$  are the normalizations. Since the  $\{\tilde{\phi}_n(x_i)\}_{n=0}^{\infty}$  form a completely orthonormal basis  $\left\{ \prod_{i=1}^5 x_i^{m_i} \right\}_{m_1, m_2, \dots, m_5=0}^{\infty}$ , where  $\sum_{i=1}^5 m_i = n$ , we expand  $V$  on

this polynomial basis. After some calculation, we obtain the following results:<sup>12</sup>

$$\begin{aligned}
\frac{1}{2} \frac{V \left| \prod_{i=1}^5 x_i^{m_i} \right\rangle}{\omega(x_i)} &= \left( \prod_{i=1}^5 x_i^{m_i} \right) \left[ \delta_{h_6 \bar{h}} \sum_{i=1}^3 \left( \frac{1}{m_i + 1} - \frac{1}{m_i + 2} \right) \right. \\
&+ \sum_{i=4}^5 \delta_{h_6 \bar{h}_i} \left( \frac{1}{m_i + 1} - \frac{1}{m_i + 2} \right) + \sum_{i=4}^5 \delta_{h_i \bar{h}} \sum_{j=1}^3 \frac{1}{m_i + m_j + 2} \\
&+ \delta_{h_4 \bar{h}_5} \frac{1}{m_4 + m_5 + 2} - 6 \sum_{i=1}^5 \sum_{k=2}^{m_i+1} \frac{1}{k} \left. \right] \\
&+ \sum_{i=1}^3 \sum_{\ell=1}^{m_i} \left( \frac{m_i - \ell + 1}{\ell(m_i + 1)} + \frac{\delta_{h_6 \bar{h}}}{(m_i + 1)(m_i + 2)} \right) x_i^{m_i - \ell} \\
&\times \sum_{\{j_k \neq i\}} \frac{\ell!}{(\ell - \sum_{k \neq i} j_k)! \prod_{k \neq i} (j_k!)} \prod_{k \neq i}^{5} (-1)^{\sum_{k \neq i} j_k} x_k^{m_k + j_k} \\
&+ \sum_{i=4}^5 \sum_{\ell=1}^{m_i} \left( \frac{m_i - \ell + 1}{\ell(m_i + 1)} + \frac{\delta_{h_6 \bar{h}_i}}{(m_i + 1)(m_i + 2)} \right) x_i^{m_i - \ell} \\
&\times \sum_{\{j_k\}} \frac{\ell!}{(\ell - \sum_{k \neq i} j_k)! \prod_{k \neq i} (j_k!)} \prod_{k \neq i}^{5} (-1)^{\sum_{k \neq i} j_k} x_k^{m_k + j_k} \\
&+ \sum_{i < j}^5 \left[ \sum_{\ell=1}^{m_j} x_i^{m_i + \ell} x_j^{m_j - \ell} \frac{m_j(m_j - 1) \cdots (m_j - \ell + 1)}{\ell(m_i + 2)(m_i + 3) \cdots (m_i + \ell + 1)} \right. \\
&\left. + \sum_{\ell=1}^{m_i} x_i^{m_i - \ell} x_j^{m_j + \ell} \frac{m_i(m_i - 1) \cdots (m_i - \ell + 1)}{\ell(m_j + 2)(m_j + 3) \cdots (m_j + \ell + 1)} \right] \prod_{k \neq i, j}^5 x_k^{m_k}
\end{aligned}$$

$$\begin{aligned}
& + \sum_{i=4}^5 \delta_{h_i \bar{h}} \sum_{j=1}^3 \frac{1}{m_i + m_j + 2} \times \left[ \sum_{\ell=1}^{m_j} x_i^{m_i + \ell} x_j^{m_j - \ell} \frac{m_j(m_j - 1) \cdots (m_j - \ell + 1)}{(m_i + 2)(m_i + 3) \cdots (m_i + \ell + 1)} \right. \\
& + \left. \sum_{\ell=1}^{m_i} x_i^{m_i - \ell} x_j^{m_j + \ell} \frac{m_i(m_i - 1) \cdots (m_i - \ell + 1)}{(m_j + 2)(m_j + 3) \cdots (m_j + \ell + 1)} \right] \prod_{k \neq i, j}^5 x_k^{m_k} \\
& + \delta_{h_4 h_5} \frac{1}{m_4 + m_5 + 2} \times \left[ \sum_{\ell=1}^{m_5} x_4^{m_4 + \ell} x_5^{m_5 - \ell} \frac{m_5(m_5 - 1) \cdots (m_5 - \ell + 1)}{(m_4 + 2)(m_4 + 3) \cdots (m_4 + \ell + 1)} \right. \\
& + \left. \sum_{\ell=1}^{m_4} x_4^{m_4 - \ell} x_5^{m_5 + \ell} \frac{m_4(m_4 - 1) \cdots (m_4 - \ell + 1)}{(m_5 + 2)(m_5 + 3) \cdots (m_5 + \ell + 1)} \right] \prod_{k=1}^3 x_k^{m_k} \\
& = 1/2 \sum_{\{n_i\}} \left| \prod_{k=1}^5 x_i^{n_i} > U_{\{n_i\}, \{m_i\}} \right. ,
\end{aligned} \tag{5.15}$$

where particles 1, 2, and 3 have helicity parallel to the hadron's helicity  $h$ , and the particles 4, 5, and 6 have helicities  $h_4, h_5$ , and  $h_6$ . The notation  $\sum_{\{j_k \neq i\}}$  means the summation over all possible integer  $j_k (k \neq i)$  as long as  $\sum_{k \neq i}^5 j_k \lesssim \ell$  is satisfied.  $U_{\{n_i\}, \{m_i\}}$  is a matrix representation of the linear operator  $\omega^{-1}V$  on the basis  $\{x_i^{m_i}\}$ . Since  $U_{\{n_i\}, \{m_i\}} = 0$  when  $\sum_{i=1}^5 n_i > \sum_{i=1}^5 m_i$ , the eigenfunctions are polynomials of degree  $n = \sum_{i=1}^5 m_i = 0, 1, 2, \dots$ . The corresponding eigenvalues are obtained by diagonalizing the matrix  $U_{\{n_i\}, \{m_i\}}$  with  $\sum_{i=1}^5 n_i = \sum_{i=1}^5 m_i = n$ . Several leading eigenvalues and the eigenfunctions are given in Table I.

From Table I, we see the leading anomalous dimension obtained from the symmetric kernel equation is given by

$$\gamma_0 = \frac{6}{5} \frac{C_F}{\beta} \text{ for } S_Z = 0, \quad \frac{7}{5} \frac{C_F}{\beta} \text{ for } S_Z = \pm 1 \quad . \tag{5.16}$$

Therefore,  $\gamma_0 > 0$  and the true leading anomalous dimension is given by

$$\gamma_0^{\text{real}} = \frac{3}{4} \frac{C_F}{\beta} \text{ for } S_Z = 0, \quad \frac{7}{8} \frac{C_F}{\beta} \text{ for } S_Z = \pm 1, \quad (5.17)$$

from the value of  $e_2$  in the *i*) case of Eq. (4.6).

## 6. Discussions and Conclusions

Harvey<sup>10</sup> has classified the color singlet six-quark states in terms of a physical cluster decomposition. Using his classification, the physical deuteron state [i.e. a bound state of two color singlet clusters] is represented as a linear combination of several different kinds of totally anti-symmetric color singlet six-quark states. For example, the two well separated nucleons  $|NN\rangle$  is given by<sup>10</sup>

$$|NN\rangle = \sqrt{\frac{1}{9}} |[6]\{33\rangle + \sqrt{\frac{4}{9}} |[42]\{33\rangle - \frac{\sqrt{4}}{9} |[42]\{51\rangle, \quad (6.1)$$

where the brackets [ ] and { } represent the orbital and spin-isospin symmetry [i.e.  $f_O$  and  $f_{TS}$  in our notations] and color symmetry (222) is abbreviated. However, this classification by itself does not include the dynamics of strong interactions between the constituents. In other words, the dynamics between the quarks inside the deuteron is not included.

Thus far, in this paper, we have formulated the dynamical evolution equation of six-quark systems and solved it to give the general form of the quark distribution amplitude  $\phi_d(x_i, Q)$  :

$$\phi_d(x_i, Q) = (\text{CTS})\phi(x_i, Q), \quad (6.2)$$

where (CTS) is tensor representation obtained from the Young symmetry of  $SU(3)_C$ ,  $SU(2)_T$  and  $SU(2)_S$  [one example is given by Eq. (2.7)], and the orbital



distribution amplitude  $\phi(x_i, Q)$  is given by

$$\phi(x_i, Q) = x_1 x_2 x_3 x_4 x_5 x_6 \sum_{n=0}^{\infty} a_n \tilde{\phi}_n(x_i) \left( \ln \frac{Q^2}{\Lambda^2} \right)^{-\gamma_n} . \quad (6.3)$$

We project Eq. (6.1) to momentum space:

$$\begin{aligned} \phi_{NN}(x_i, Q) &= \sqrt{\frac{1}{9}} \phi_{[6]\{33\}}(x_i, Q) + \sqrt{\frac{4}{9}} \phi_{[42]\{33\}}(x_i, Q) \\ &\quad - \sqrt{\frac{4}{9}} \phi_{[42]\{51\}}(x_i, Q) . \end{aligned} \quad (6.4)$$

In the limit  $Q \rightarrow \infty$ , the dependence of  $Q$  is determined by the leading anomalous dimension; all other terms which have non-leading anomalous dimensions are suppressed by logarithmic damping factors. However, as we can see from Table I, the orbital symmetry of the eigensolution which has the leading anomalous dimension cannot be [42] but is [6]. This means only the first term of Eq. (6.4) survives at the large  $Q$  limit. The  $NN$  amplitude itself is not sufficient. One can show that an 80 percent hidden-color state is necessary to saturate the normalization of six-quark amplitude when six quarks approach the same position in impact space  $b_{\perp} \rightarrow 0$ . We have called this new degree of freedom an anomalous state since it does not correspond to the usual nucleonic degrees of freedom of the nucleus. The physical implication of the anomalous state is discussed in our toy model analysis.<sup>7</sup>

The asymptotic behavior of the deuteron distribution amplitude is given by

$$\phi_D(x_i, Q) = a_0 x_1 x_2 x_3 x_4 x_5 x_6 \left( \frac{\ln Q^2 / \Lambda^2}{\ln Q_0^2 / \Lambda^2} \right)^{-\gamma_0} \quad (6.5)$$

where  $\gamma_0 = \frac{3}{4} \frac{C_F}{\beta}$  for  $S_Z = 0$  deuteron. The QCD predictions for high- $Q^2$  behavior of deuteron form factor and the form of the deuteron distribution amplitude at

short distances are given in Ref. 3. The fact that the six-quark state is 80 percent hidden color at small transverse separation implies that the deuteron form factors cannot be described at large  $Q^2$  by meson-nucleon degrees of freedom alone, and that the nucleon-nucleon potential is repulsive at short distances.<sup>3,7,13</sup>

## ACKNOWLEDGEMENTS

The calculation of the deuteron form factor using the QCD evolution formalism was done in collaboration with G. P. Lepage (see Ref. 3). We also wish to thank J. D. Walecka for helpful conversations.

## Appendix A

In this appendix we describe a general method for finding the color singlet representations for multihadron systems. This leads to the explicit representation of Eq. (2.7).

Group theoretically, color states of every quark and/or antiquark systems are represented by the multiple outer products:

$$(3)^p \times (3^*)^q \quad (A.1)$$

for the system of  $p$  quarks and  $q$  anti-quarks. The reducible tensor representations of (A.1) is decomposed into irreducible tensor representations and the resulting singlet representations provide the physical quark system.

Since the color singlet is invariant under  $SU(3)$  transformation, the only tensor representations<sup>14</sup> which are invariant under  $SU(3)$  transformation are the Kronecker delta and the completely antisymmetric Cartesian tensors:

$$\delta_j^i, \quad \epsilon_{ijk}, \quad \epsilon^{ijk}, \quad (A.2)$$

where the lower (upper) indices correspond to  $3(3^*)$  representations. This comes from the fact that only operations of contraction and anti-symmetrization commute with the  $SU(3)$  transformations on the mixed tensors. Thus, every color singlet representation can be represented by the products of the three tensors (A.2).

From this observation, we describe the rules to construct the color singlet tensor representations for arbitrary quark and/or antiquark systems such as (A.1):

1. Give each index to every 3 and 3\* representations. For example,  $p$  lower indices and  $q$  upper indices will be given to (A.1).
2. Assemble one possible product of Kronecker deltas and completely anti-symmetric Cartesian tensors to use up all indices considered in rule 1. For example, the  $p = 6$  and  $q = 0$  case needs the product of two antisymmetric Cartesian tensors.
3. Permute the upper and lower indices separately. For example,  $p = 6$  and  $q = 0$  case will give 10 possible different representations. However, note that they are not all independent.
4. To construct all the independent (orthogonal) representations follow the method of Schmidt's orthogonalization, where the inner product is defined as contraction. For example, in the  $p = 6$  and  $q = 0$  case, five independent representations are obtained:

$$\begin{aligned}
S_{ijklmn}^1 &= \frac{1}{12\sqrt{2}} (\epsilon_{ikm} \epsilon_{jln} + \epsilon_{ikn} \epsilon_{jlm} + \epsilon_{jkm} \epsilon_{iln} + \epsilon_{jkn} \epsilon_{ilm}) , \\
S_{ijklmn}^2 &= \frac{1}{4\sqrt{6}} (\epsilon_{ijm} \epsilon_{kln} + \epsilon_{ijn} \epsilon_{klm}) , \\
S_{ijklmn}^3 &= \frac{1}{4\sqrt{6}} (\epsilon_{ikl} \epsilon_{jmn} + \epsilon_{jkl} \epsilon_{imn}) , \\
S_{ijklmn}^4 &= \frac{1}{4\sqrt{2}} \left( \epsilon_{ijl} \epsilon_{kmn} - \frac{1}{3} \epsilon_{ijk} \epsilon_{lmn} \right) , \\
S_{ijklmn}^5 &= \frac{1}{6} \epsilon_{ijk} \epsilon_{lmn} ,
\end{aligned} \tag{A.3}$$

where  $S_{ijklmn}^\alpha$  ( $\alpha = 1, 2, \dots, 5$ ) are the five independent color singlet representations.

The correspondence between Eq. (A.3) and the five different Young tableau (or Yamanouchi labels) is as follows: If we denote  $i = 1, j = 2, \dots, n = 6$ , then the corresponding Young tableau with Eq. (A.3) are following:

$$\begin{array}{ccccc}
 \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 4 \\ \hline 5 & 6 \\ \hline \end{array} & 
 \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & 4 \\ \hline 5 & 6 \\ \hline \end{array} & 
 \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 5 \\ \hline 4 & 6 \\ \hline \end{array} & 
 \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & 5 \\ \hline 4 & 6 \\ \hline \end{array} & 
 \begin{array}{|c|c|} \hline 1 & 4 \\ \hline 2 & 5 \\ \hline 3 & 6 \\ \hline \end{array}
 \end{array}$$

$$\alpha = 1, \quad 2, \quad 3, \quad 4, \quad 5.$$

A similar method can be applied to construct the isospin singlet representations in  $SU(2)$ . The five independent isospin singlet tensors corresponding to the dual of the color singlet Young tableau shown in (A.3) are represented by

$$\begin{aligned}
 \begin{array}{|c|c|c|} \hline 1 & 3 & 5 \\ \hline 2 & 4 & 6 \\ \hline \end{array} &= \frac{1}{2\sqrt{2}} \epsilon_{ab}\epsilon_{cd}\epsilon_{ef}, \\
 \begin{array}{|c|c|c|} \hline 1 & 2 & 5 \\ \hline 3 & 4 & 6 \\ \hline \end{array} &= \frac{1}{2\sqrt{6}} (\epsilon_{ac}\epsilon_{bd}\epsilon_{ef} + \epsilon_{ad}\epsilon_{bc}\epsilon_{ef}), \\
 \begin{array}{|c|c|c|} \hline 1 & 3 & 4 \\ \hline 2 & 5 & 6 \\ \hline \end{array} &= \frac{1}{2\sqrt{6}} (\epsilon_{ab}\epsilon_{ce}\epsilon_{df} + \epsilon_{ab}\epsilon_{cf}\epsilon_{de}), \\
 \begin{array}{|c|c|c|} \hline 1 & 2 & 4 \\ \hline 3 & 5 & 6 \\ \hline \end{array} &= \frac{1}{6\sqrt{2}} (\epsilon_{ac}\epsilon_{be}\epsilon_{df} + \epsilon_{ac}\epsilon_{bf}\epsilon_{de} + \epsilon_{ae}\epsilon_{bc}\epsilon_{df} + \epsilon_{af}\epsilon_{bc}\epsilon_{de}), \\
 \begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 4 & 5 & 6 \\ \hline \end{array} &= \frac{1}{6} (\epsilon_{ad}\epsilon_{be}\epsilon_{cf} + \epsilon_{ae}\epsilon_{bf}\epsilon_{cd} + \epsilon_{af}\epsilon_{bd}\epsilon_{ce}),
 \end{aligned} \tag{A.4}$$

where the  $\epsilon_{ab}$ 's are the antisymmetric Cartesian tensors of  $SU(2)_T$  and the Young tableau are denoted by  $a = 1, b = 2, \dots, f = 6$ . If we multiply the color and isospin singlet tensors given by Eqs. (A.3) and (A.4), respectively, taking care of the phase factor<sup>8</sup>  $\eta_T$  for the Yamanouchi labels, we get the special case of the asymptotic deuteron representation, Eq. (2.7).

## Appendix B

The color factor defined in Eq. (3.8) can be explicitly expressed in terms of tensor notation, since we know the tensor representation for the multiplication of Gell-Mann matrices in Eq. (3.8) as <sup>14</sup>

$$\left(\frac{\vec{\lambda}}{2}\right)_{i'}^i \left(\frac{\vec{\lambda}}{2}\right)_{j'}^j = 1/2 \left( \delta_{j'}^i \delta_{i'}^j - 1/3 \delta_{i'}^i \delta_{j'}^j \right), \quad (B.1)$$

where the indices  $i$  and  $i'$  designate the color it  $i'$ th quark before and after a gluon exchange. Using the notation Eq. (A.3), the generalized color matrices ( $5 \times 5$ ) given by Eq. (3.8) can be obtained by

$$C_{\alpha\beta}(i, j) = S_{\dots i \dots j \dots}^{\alpha} \left(\frac{\vec{\lambda}}{2}\right)_{i'}^i \left(\frac{\vec{\lambda}}{2}\right)_{j'}^j S_{\dots i' \dots j' \dots}^{\beta}, \quad (B.2)$$

and Eq. (B.1). There are 15 such matrices and they are given by

$$C(1, 2) = \begin{bmatrix} \frac{1}{3} & 0 & 0 & 0 & 0 \\ 0 & -\frac{2}{3} & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{3} & 0 & 0 \\ 0 & 0 & 0 & -\frac{2}{3} & 0 \\ 0 & 0 & 0 & 0 & -\frac{2}{3} \end{bmatrix}$$

$$C(1, 3) = \begin{bmatrix} -\frac{5}{12} & -\frac{\sqrt{3}}{4} & 0 & 0 & 0 \\ -\frac{\sqrt{3}}{4} & \frac{1}{12} & 0 & 0 & 0 \\ 0 & 0 & -\frac{5}{12} & -\frac{\sqrt{3}}{4} & 0 \\ 0 & 0 & -\frac{\sqrt{3}}{4} & \frac{1}{12} & 0 \\ 0 & 0 & 0 & 0 & -\frac{2}{3} \end{bmatrix}$$

$$C(1,4) = \begin{bmatrix} -\frac{5}{12} & \frac{\sqrt{3}}{4} & 0 & 0 & 0 \\ \frac{\sqrt{3}}{4} & \frac{1}{12} & 0 & 0 & 0 \\ 0 & 0 & -\frac{5}{12} & -\frac{1}{4\sqrt{3}} & \frac{1}{\sqrt{6}} \\ 0 & 0 & -\frac{1}{4\sqrt{3}} & -\frac{7}{12} & -\frac{1}{3\sqrt{2}} \\ 0 & 0 & \frac{1}{\sqrt{6}} & -\frac{1}{3\sqrt{2}} & 0 \end{bmatrix}$$

$$C(1,5) = \begin{bmatrix} -\frac{5}{12} & 0 & 0 & -\frac{1}{4} & -\frac{1}{2\sqrt{2}} \\ 0 & -\frac{5}{12} & -\frac{1}{4} & -\frac{1}{2\sqrt{3}} & \frac{1}{2\sqrt{6}} \\ 0 & -\frac{1}{4} & -\frac{5}{12} & \frac{1}{2\sqrt{3}} & -\frac{1}{2\sqrt{6}} \\ -\frac{1}{4} & -\frac{1}{2\sqrt{3}} & \frac{1}{2\sqrt{3}} & -\frac{1}{12} & \frac{1}{6\sqrt{2}} \\ -\frac{1}{2\sqrt{2}} & \frac{1}{2\sqrt{6}} & -\frac{1}{2\sqrt{6}} & \frac{1}{6\sqrt{2}} & 0 \end{bmatrix}$$

$$C(1,6) = \begin{bmatrix} -\frac{5}{12} & 0 & 0 & \frac{1}{4} & \frac{1}{2\sqrt{2}} \\ 0 & -\frac{5}{12} & \frac{1}{4} & \frac{1}{2\sqrt{3}} & -\frac{1}{2\sqrt{6}} \\ 0 & \frac{1}{4} & -\frac{5}{12} & \frac{1}{2\sqrt{3}} & -\frac{1}{2\sqrt{6}} \\ \frac{1}{4} & \frac{1}{2\sqrt{3}} & \frac{1}{2\sqrt{3}} & -\frac{1}{12} & \frac{1}{6\sqrt{2}} \\ \frac{1}{2\sqrt{2}} & -\frac{1}{2\sqrt{6}} & -\frac{1}{2\sqrt{6}} & \frac{1}{6\sqrt{2}} & 0 \end{bmatrix}$$

$$C(2,3) = \begin{bmatrix} -\frac{5}{12} & \frac{\sqrt{3}}{4} & 0 & 0 & 0 \\ \frac{\sqrt{3}}{4} & \frac{1}{12} & 0 & 0 & 0 \\ 0 & 0 & -\frac{5}{12} & \frac{\sqrt{3}}{4} & 0 \\ 0 & 0 & \frac{\sqrt{3}}{4} & \frac{1}{12} & 0 \\ 0 & 0 & 0 & 0 & -\frac{2}{3} \end{bmatrix}$$

$$C(2,4) = \begin{bmatrix} -\frac{5}{12} & -\frac{\sqrt{3}}{4} & 0 & 0 & 0 \\ -\frac{\sqrt{3}}{4} & \frac{1}{12} & 0 & 0 & 0 \\ 0 & 0 & -\frac{5}{12} & \frac{1}{4\sqrt{3}} & -\frac{1}{\sqrt{6}} \\ 0 & 0 & \frac{1}{4\sqrt{3}} & -\frac{7}{12} & -\frac{1}{3\sqrt{2}} \\ 0 & 0 & -\frac{1}{\sqrt{6}} & -\frac{1}{3\sqrt{2}} & 0 \end{bmatrix}$$

$$C(2,5) = \begin{bmatrix} -\frac{5}{12} & 0 & 0 & \frac{1}{4} & \frac{1}{2\sqrt{2}} \\ 0 & -\frac{5}{12} & \frac{1}{4} & -\frac{1}{2\sqrt{3}} & \frac{1}{2\sqrt{6}} \\ 0 & \frac{1}{4} & -\frac{5}{12} & -\frac{1}{2\sqrt{3}} & \frac{1}{2\sqrt{6}} \\ \frac{1}{4} & -\frac{1}{2\sqrt{3}} & -\frac{1}{2\sqrt{3}} & -\frac{1}{12} & \frac{1}{6\sqrt{2}} \\ \frac{1}{2\sqrt{2}} & \frac{1}{2\sqrt{6}} & \frac{1}{2\sqrt{6}} & \frac{1}{6\sqrt{2}} & 0 \end{bmatrix}$$

$$C(2,6) = \begin{bmatrix} -\frac{5}{12} & 0 & 0 & -\frac{1}{4} & -\frac{1}{2\sqrt{2}} \\ 0 & -\frac{5}{12} & -\frac{1}{4} & \frac{1}{2\sqrt{3}} & -\frac{1}{2\sqrt{6}} \\ 0 & -\frac{1}{4} & -\frac{5}{12} & -\frac{1}{2\sqrt{3}} & \frac{1}{2\sqrt{6}} \\ -\frac{1}{4} & \frac{1}{2\sqrt{3}} & -\frac{1}{2\sqrt{3}} & -\frac{1}{12} & \frac{1}{6\sqrt{2}} \\ -\frac{1}{2\sqrt{2}} & -\frac{1}{2\sqrt{6}} & \frac{1}{2\sqrt{6}} & \frac{1}{6\sqrt{2}} & 0 \end{bmatrix}$$

$$C(3,4) = \begin{bmatrix} \frac{1}{3} & 0 & 0 & 0 & 0 \\ 0 & -\frac{2}{3} & 0 & 0 & 0 \\ 0 & 0 & -\frac{2}{3} & 0 & 0 \\ 0 & 0 & 0 & -\frac{1}{3} & \frac{\sqrt{2}}{3} \\ 0 & 0 & 0 & \frac{\sqrt{2}}{3} & 0 \end{bmatrix}$$

$$C(3,5) = \begin{bmatrix} -\frac{5}{12} & 0 & -\frac{\sqrt{3}}{4} & 0 & 0 \\ 0 & -\frac{5}{12} & 0 & \frac{1}{4\sqrt{3}} & -\frac{1}{\sqrt{6}} \\ -\frac{\sqrt{3}}{4} & 0 & \frac{1}{12} & 0 & 0 \\ 0 & \frac{1}{4\sqrt{3}} & 0 & -\frac{7}{12} & -\frac{1}{3\sqrt{2}} \\ 0 & -\frac{1}{\sqrt{6}} & 0 & -\frac{1}{3\sqrt{2}} & 0 \end{bmatrix}$$

$$C(3,6) = \begin{bmatrix} -\frac{5}{12} & 0 & \frac{\sqrt{3}}{4} & 0 & 0 \\ 0 & -\frac{5}{12} & 0 & -\frac{1}{4\sqrt{3}} & \frac{1}{\sqrt{6}} \\ \frac{\sqrt{3}}{4} & 0 & \frac{1}{12} & 0 & 0 \\ 0 & -\frac{1}{4\sqrt{3}} & 0 & -\frac{7}{12} & -\frac{1}{3\sqrt{2}} \\ 0 & \frac{1}{\sqrt{6}} & 0 & -\frac{1}{3\sqrt{2}} & 0 \end{bmatrix}$$



$$\begin{aligned}
C(4,5) &= \begin{bmatrix} -\frac{5}{12} & 0 & \frac{\sqrt{3}}{4} & 0 & 0 \\ 0 & -\frac{5}{12} & 0 & \frac{\sqrt{3}}{4} & 0 \\ \frac{\sqrt{3}}{4} & 0 & \frac{1}{12} & 0 & 0 \\ 0 & \frac{\sqrt{3}}{4} & 0 & \frac{1}{12} & 0 \\ 0 & 0 & 0 & 0 & -\frac{2}{3} \end{bmatrix} \\
C(4,6) &= \begin{bmatrix} -\frac{5}{12} & 0 & -\frac{\sqrt{3}}{4} & 0 & 0 \\ 0 & -\frac{5}{12} & 0 & -\frac{\sqrt{3}}{4} & 0 \\ -\frac{\sqrt{3}}{4} & 0 & \frac{1}{12} & 0 & 0 \\ 0 & -\frac{\sqrt{3}}{4} & 0 & \frac{1}{12} & 0 \\ 0 & 0 & 0 & 0 & -\frac{2}{3} \end{bmatrix} \\
C(5,6) &= \begin{bmatrix} \frac{1}{3} & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{3} & 0 & 0 & 0 \\ 0 & 0 & -\frac{2}{3} & 0 & 0 \\ 0 & 0 & 0 & -\frac{2}{3} & 0 \\ 0 & 0 & 0 & 0 & -\frac{2}{3} \end{bmatrix}
\end{aligned} \tag{B.3}$$

Using the results (B.3) and the formula (3.10) and (3.11), we can find expressions for the kernels in terms of  $V_{ij}$ . Since  $C_{\alpha\alpha}(i,j)$  is independent of  $i$  and  $j$  and given by

$$C_{\alpha\alpha}(i,j) = -\frac{2}{3} \times 3 + \frac{1}{3} \times 2 = -C_F, \tag{B.4}$$

the symmetric kernel is given by

$$K_{(6)[111111]} = C_d \sum_{i \neq j} V_{i,j}, \tag{B.5}$$

where  $C_d = -C_F/5$ .

The other kernels can also be obtained by using the Clebsch–Gordon coeffi-

icients of  $S_6$  and (B.3). The results are given by

$$K_{fY} = N_{fY} \sum_{i \neq j} A_{ij} V_{ij} , \quad (B.6)$$

where  $N_{fY}$  and  $A_{ij}$  are summarized in the Table II.

## References

1. S. J. Brodsky and C-R. Ji SLAC-PUB-3747, and references therein.
2. S. J. Brodsky and G. P. Lepage, Phys. Rev. D24, 1808(1981).
3. S. J. Brodsky, C-R. Ji and G. P. Lepage, Phys. Rev. Lett. 51, 83 (1983).
4. G. P. Lepage and S. J. Brodsky, Phys. Rev. D22, 2157 (1980).
5. V. N. Baier and A. G. Grozin, Nucl. Phys. B192, 476 (1981).
6. S. J. Brodsky and C-R. Ji, SLAC-PUB-3706.
7. S. J. Brodsky and C-R. Ji, SLAC-PUB-3707.
8. M. Hammermesh, Group Theory, (Addison-Wesley, Reading, MA, 1962).
9. [654321] is the Yamanouchi label of  $(1^6)$  symmetry; See Ref. 8.
10. M. Harvey, Nucl. Phys. A352, 301 (1981); 326 (1981); Nucl. Phys. A424, 428 (1984).
11. Detailed calculations and the origin of the logarithmic Q dependence are given in Ref. 4.
12. In Eq. (5.15), we use a more compact form for the coefficient of  $x_i^{m_i+\ell} x_j^{m_j-\ell}$  than that given in Ref. 4, since we can prove that

$$-\sum_{j=0}^i \binom{m}{j} \binom{m-j}{m-i} (-1)^j \sum_{k=2}^{n+j+1} \frac{1}{k} = \frac{m(m-1)\cdots(m-i+1)}{i(n+2)(n+3)\cdots(n+i+1)} .$$

Also we correct typographical error in Appendix D of Ref. 4:  $\frac{m-i+1+\delta_{h_2\bar{h}}}{i(m+2)}$

should read  $\frac{m-i+1}{i(m+1)} + \frac{\delta_{h_2\bar{h}}}{(m+1)(m+2)}$  .

13. Qualitative QCD-based arguments for a repulsive N-N potential at short distances are given in C. Detar, HU-TFT-82-6 (1982); M. Harvey (Ref. 10);

K. Maltman and N. Isgur, Phys. Rev. Lett. 50, 1827 (1983); R. L. Jaffe, Phys. Rev. Lett. 24, 228 (1983); G. E. Brown, in Erice1981, Proceedings, Quarks and the Nucleus. The possibility that the deuteron form factor is dominated at large momentum transfer by hidden color components is discussed in V. A. Matveev and P. Sorba, Nuovo Cim. 45A, 257 (1978); Nuovo Cim. Lett. 20, 435 (1977).

14. P. Carruthers, Introduction to Unitary Symmetry (Interscience Publishers Inc., New York, 1966).

**TABLE I**

Solutions of the evolution equation Eq. (5.12) for total helicity  $|3h+h_4+h_5+h_6| = 0$  ( $\phi^{\uparrow\uparrow\uparrow\downarrow\downarrow}$ ), 1 ( $\phi^{\uparrow\uparrow\uparrow\uparrow\downarrow}$ ,  $\phi^{\uparrow\uparrow\uparrow\downarrow\downarrow}$ ,  $\phi^{\uparrow\uparrow\uparrow\downarrow\uparrow}$ ), 2 ( $\phi^{\uparrow\uparrow\uparrow\downarrow\uparrow}$ ,  $\phi^{\uparrow\uparrow\uparrow\uparrow\uparrow}$ ,  $\phi^{\uparrow\uparrow\uparrow\downarrow\uparrow}$ ), and 3 ( $\phi^{\uparrow\uparrow\uparrow\uparrow\uparrow}$ ) cases. The procedure for the systematic derivation of the  $\tilde{\phi}_n$  is given in Sec. 5.

	$\gamma_n / \left( \frac{C_F}{\beta} \right)$	$N$	$a_{00000}^{(0)}$	$a_{10000}^{(1)}$	$a_{01000}^{(1)}$	$a_{00100}^{(1)}$	$a_{00010}^{(1)}$	$a_{00001}^{(1)}$
$\phi^{\uparrow\uparrow\uparrow\downarrow\downarrow}$	$\frac{6}{5}$	11!						
	$\frac{13}{5}$	$\frac{13!}{4}$		1	-1			
	$\frac{13}{5}$	$\frac{13!}{12}$		1	1	-2		
	$\frac{13}{5}$	$\frac{13!}{4}$					1	-1
	$\frac{13}{5}$	$\frac{13!}{12}$	2	-2	-2	-2	-3	-3
	$\frac{14}{5}$	$\frac{13!}{12}$	1	-2	-2	-2		
$\phi^{\uparrow\uparrow\uparrow\uparrow\downarrow}$								
$\phi^{\uparrow\uparrow\uparrow\downarrow\uparrow}$	$\frac{7}{5}$	11!						
$\phi^{\uparrow\uparrow\uparrow\downarrow\uparrow}$								
$\phi^{\uparrow\uparrow\uparrow\uparrow\downarrow}$								
$\phi^{\uparrow\uparrow\uparrow\downarrow\uparrow}$	2	11!						
$\phi^{\uparrow\uparrow\uparrow\downarrow\uparrow}$								
$\phi^{\uparrow\uparrow\uparrow\uparrow\uparrow}$	3	11!						

$$\tilde{\phi}_n(x_i) = \sqrt{N} \sum_{\{m_i\}} a_{m_1 \dots m_5}^{(n)} \prod_{i=1}^5 x_i^{m_i} \left( n = \sum_{i=1}^5 m_i \right)$$


TABLE II.

The coefficients  $A_{ij}$  for all  $(i, j)$  pairs and the normalization factor  $N_{fY}$  defined in Eq. (B.6). Since color symmetry is fixed by (222), the possible symmetry of the kernel is  $f = (6)$  or  $(42)$ .

$K_{fY}$	$N_{fY}$	(56)	(46)	(45)	(36)	(35)	(26)	(25)	(16)	(15)	(23)	(24)	(34)	(14)	(13)	(12)
$K_{(6)[111111]}$	$-\frac{4}{15}$	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
$K_{(42)[221111]}$	$\frac{\sqrt{6}}{20}$	12	-3	-3	-3	-3	-3	-3	-3	-3	2	2	2	2	2	2
$K_{(43)[212111]}$	$\frac{\sqrt{10}}{20}$		9	-3	-3	1	-3	1	-3	1	1	-2	-2	-2	2	2
$K_{(42)[211211]}$	$\frac{\sqrt{5}}{10}$				6	-2	-3	1	-3	1	-1	1	-2	1	-1	-2
$K_{(43)[211121]}$	$\frac{\sqrt{15}}{10}$						3	-1	-3	1	-1	-1	0	1	1	
$K_{(42)[122111]}$	$\frac{1}{\sqrt{5}}$			3		-1		-1		-1	1	-1	-1	-1	1	1
$K_{(42)[121211]}$	$\frac{1}{\sqrt{10}}$					4		-2		-2	-1	1	-2	1	-1	2
$K_{(42)[121121]}$	$\sqrt{\frac{3}{10}}$							2		-2	-1	-1		1	1	
$K_{(42)[112211]}$	$\sqrt{\frac{2}{10}}$										-1	-1	2	-1	-1	2
$K_{(42)[112121]}$	$\sqrt{\frac{3}{10}}$										-1	1		-1	1	

## Figure Captions

Fig. 1. The leading order contributions to the kernel of the six quark wave function in light cone perturbation theory. The longitudinal momentum of particle  $i$  ( $i = 1, 2, \dots, 6$ ) before and after the interactions is  $y_i$  and  $x_i$  respectively; the  $i$  and  $j$  particles interact with transverse momentum transfer of order  $Q$ . Fifteen diagrams are included by summation over  $i$  and  $j$  with  $i \neq j$ . The Feynman rules of light cone perturbation theory in light-cone gauge are summarized in Ref. 4.



6-82  
4586A1

Fig. 1