

SLAC-PUB-3231

October 1983

(T/E)

**Search for Singly Produced Supersymmetric Electrons  
in  $e^+e^-$  Interactions\***

E. Fernandez, W. T. Ford, A. L. Read, Jr., and J. G. Smith  
*Department of Physics*  
*University of Colorado, Boulder, Colorado 80309*

R. De Sangro, A. Marini, I. Peruzzi, M. Piccolo, and F. Ronga  
*Laboratori Nazionali Frascati dell' I.N.F.N.*  
*Frascati, Italy*

H. T. Blume, R. B. Hurst, J. P. Venutti,  
H. B. Wald, and Roy Weinstein  
*Department of Physics*  
*University of Houston, Houston, Texas 77004*

H. R. Band, M. W. Gettner, G. P. Goderre  
O. A. Meyer, J. H. Moromisato, W. D. Shambroom, and E. von Goeler  
*Department of Physics*  
*Northeastern University, Boston, Massachusetts 02115*

W. W. Ash, G. B. Chadwick, S. H. Clearwater, R. W. Coombes,  
H. S. Kaye, K. H. Lau, R. E. Leedy, H. L. Lynch, R. L. Messner,  
S. J. Michalowski<sup>(a)</sup>, K. Rich, D. M. Ritson, L. J. Rosenberg,  
D. E. Wiser, and R. W. Zdarko  
*Department of Physics and Stanford Linear Accelerator Center*  
*Stanford University, Stanford, California 94305*

D. E. Groom, H. Y. Lee, and E. C. Loh  
*Department of Physics*  
*University of Utah, Salt Lake City, Utah 84112*

M. C. Delfino, B. K. Heltsley, J. R. Johnson, T. L. Lavine,  
T. Maruyama, and R. Prepost  
*Department of Physics*  
*University of Wisconsin, Madison, Wisconsin 53706*

Submitted to Physical Review Letters

---

\* Work supported in part by the Department of Energy, contracts DE-AC02-81ER40025 (CU), DE-AC03-76SF00515 (SLAC) and DE-AC02-76ER00881 (UW); by the National Science Foundation, contracts NSF-PHY82-15133 (UH), NSF-PHY82-15413 and NSF-PHY82-15414 (NU), and NSF-PHY80-06504 (UU); and by I.N.F.N.

## ABSTRACT

A search for supersymmetric electron production via the reaction  $e^+e^- \rightarrow e^\pm \tilde{\gamma} \tilde{e}^\mp$  followed by the decay  $\tilde{e}^\mp \rightarrow e^\mp \tilde{\gamma}$  has been performed with the MAC detector at PEP. The absence of candidates in a sample corresponding to an integrated luminosity of  $36.4 \text{ pb}^{-1}$  is used to set a new lower limit on the  $\tilde{e}$  mass of  $22.4 \text{ GeV}/c^2$  at the 95% confidence level.

One of the most striking predictions of supersymmetric theories<sup>1</sup> is that for every familiar particle of the standard model there should be supersymmetric partners with spin differing by 1/2 unit and, in the case of exact supersymmetry, identical mass. Since none of these particles has yet been found, the symmetry, if valid, is apparently broken at low energies. There is no universal agreement, however, about the mechanism responsible for supersymmetry breaking and it is therefore important to search for supersymmetric particles in current experiments. The spin-0 partners of the electron, muon and tau ( $\tilde{e}$ ,  $\tilde{\mu}$ ,  $\tilde{\tau}$ ), could be pair-produced in  $e^+e^-$  annihilations if the beam energy exceeded their mass.<sup>2</sup> Their decay would lead to distinctive noncoplanar  $e^+e^-$  or  $\mu^+\mu^-$  final states. Experiments conducted at PETRA<sup>3</sup> and PEP<sup>4</sup> have established upper bounds on these processes leading to lower limits on the mass of the  $\tilde{e}$ ,  $\tilde{\mu}$  and  $\tilde{\tau}$ . The most stringent limit on the  $\tilde{e}$  mass using this technique<sup>5</sup> is  $m_{\tilde{e}} > 17.8 \text{ GeV}/c^2$ . As suggested by several authors,<sup>2,6</sup>  $\tilde{e}$ 's with masses larger than the beam energy but smaller than the center of mass energy, could be produced singly in  $e^+e^-$  interactions in the process  $e^+e^- \rightarrow e^\pm \tilde{\gamma} \tilde{e}^\mp$  where  $\tilde{\gamma}$  is the supersymmetric partner of the photon.

A search for this process has been completed using the MAC detector at PEP. The MAC detector<sup>7</sup> consists of a cylindrical drift chamber surrounded by a hexagonal barrel of electromagnetic calorimeters, scintillation counters, and hadronic calorimeters. Planar endcap scintillators and calorimeters extend the coverage to small angles from the beam. Outside the central and endcap calorimeters, 3 to 6 layers of drift chambers provide muon identification and momentum measurement. The central drift chamber has ten layers of drift cells inside a solenoid which provides a 5.7 kG magnetic field. Momentum resolution is  $\delta p/p \simeq 0.065 p \sin\theta$  for  $23^\circ < \theta < 157^\circ$ , where  $\theta$  is the angle with respect to the beam axis. The electromagnetic calorimeter consists of alternating planes of lead and proportional chambers totaling 14 radiation lengths of material. Each calorimeter sextant is segmented into 32 azimuthal sectors and 3 layers in depth. Charge division in each of the segments is used to measure the axial position of the showers. The energy resolution for electromagnetic showers has been measured with Bhabha events as  $\delta E/E = 20\%/\sqrt{E}$ . The hadronic calorimeters are of similar construction, with steel absorbers instead of lead. The endcap calorimeters are made of alternating layers of steel and proportional chambers with electromagnetic energy resolution given by  $\delta E/E = 45\%/\sqrt{E}$ . A single plane of scintillators is located inside each endcap at a depth of 8 radiation lengths.

The MAC trigger for single electron or photon showers uses energy sums from the central electromagnetic calorimeter. Sums corresponding to the three layers of radial readout and  $30^\circ$  azimuthal sectors are made and combined again to form sextant energy sums. A timing discriminator<sup>8</sup> on each sextant sum produces a

pre-trigger if that sum is greater than 2 GeV and within a 150 nanosecond timing gate. The full trigger further requires that at least two adjacent layers in any sector each have more than 0.3 GeV energy deposited. Only triggers with energy greater than 2.5 GeV in the central electromagnetic calorimeter were logged. The efficiency of this trigger was measured from the real data sample, using  $e^+e^- \rightarrow e^+e^-\gamma$  events that have only one particle in the central section and satisfy other MAC triggers. The trigger efficiency increases with energy, rising from 92% at 3 GeV to greater than 99% above 6 GeV.

The cross section for the process  $e^+e^- \rightarrow e^\pm \tilde{\gamma} \tilde{e}^\mp$  has been calculated by Gaillard et al.<sup>6</sup> Their calculation assumes that the dominant contribution to the cross section comes from the interaction of one of the beam electrons with a quasi-real photon radiated by the other beam electron, producing a  $\tilde{\gamma}$  and a  $\tilde{e}$ . The beam electron that radiates the photon, is scattered by a small angle and is therefore not observed in the detector.<sup>9</sup> The  $\tilde{e}$  is assumed to decay promptly into an electron and a  $\tilde{\gamma}$  with a 100% branching ratio. The decay electron has a nearly isotropic angular distribution and is fairly energetic. These characteristics reflect the fact that the  $\tilde{e}$  is a heavy particle which decays isotropically in its rest frame and is moving slowly in the laboratory frame. The energy distribution of the electron, shown in the inset of Fig. 1 for our center of mass energy of 29 GeV and for several  $\tilde{e}$  masses, was calculated for  $|\cos\theta| < 0.75$ , where  $\theta$  is the angle of the electron with respect to the beam axis. Approximately 75% of the total cross section satisfies this cut. Assuming the  $\tilde{\gamma}$ 's are undetected,<sup>10</sup> this electron is the only observed final state particle.

The single electron sample was obtained by selecting events with one and only one track in the central drift chamber, with  $|\cos\theta| < 0.75$ , and with associated calorimeter hits consistent with an electromagnetic shower of energy greater than 3 GeV. For this search region typically 95% of the electron's energy is deposited in the central shower chamber. Events with showers in the central or endcap calorimeters greater than 1 GeV and not correlated with the electron track were removed from the sample. Events with endcap scintillator hits were also removed. The energy and angular distribution of the final sample are shown in figures 1 and 2 respectively. The total number of events is  $1565 \pm 40$  where the error is only statistical. No event is observed with an electron energy of more than 6 GeV.

Background single electron events come primarily from  $e^+e^-\gamma$  final states where one electron and the photon are not seen in the detector. Two-photon processes and tau events can also contribute to the background. The MAC detector is very efficient at detecting energetic ( $> 2$  GeV) electrons or photons over its entire angular range (98% of  $4\pi$ ). Although the detector has small dead regions between adjacent calorimeter segments, in the central part these regions are smaller than the shower size, thus ensuring very efficient detection of electromagnetic showers. In the endcap regions, shower detection is complemented by the scintillation counter layer located near shower maximum. Due to the complicated detector geometry at small angles, the minimum polar angle at which showers can be detected varies from  $9^\circ$  to  $12^\circ$  depending on the azimuthal angle. Except for this low polar angle region the efficiency for detecting energetic electromagnetic showers is essentially 100%. All background processes then have

at least two particles at small polar angles, severely restricting the energy of the electron observed in the search region.

The energy and angular distribution of single electrons from the  $e^+e^- \rightarrow e^+e^-\gamma$  reaction was estimated using the Monte Carlo program of Berends and Kleiss<sup>11</sup> to generate  $ee\gamma$  events which were then run through a detector simulation program.<sup>12</sup> These Monte Carlo events were analyzed with the same cuts used in the data sample. The resulting energy and angular distributions were corrected by folding in the trigger inefficiency as a function of energy. In addition a 5% loss of events due to the effect of dead channels in the energy measurement and a 1% loss due to accidental endcap calorimeter and scintillator hits were included in obtaining the curves shown in figures 1 and 2. From the Monte Carlo analysis, the total number of single electron events expected for the luminosity of  $36.4 \pm 0.6 \text{ pb}^{-1}$  is  $1640 \pm 230$ . The error includes the statistical errors due to the finite number of Monte Carlo generated events, the uncertainties in the correction factors mentioned above and an additional 10% error due to uncertainty in the absolute calorimeter energy calibration. Also included is an 8% error due to the uncertainty of modeling the detector geometry at small angles. Contributions due to  $\tau^+\tau^-\gamma$ , other tau events, and two photon processes are negligible compared to the  $e^+e^-\gamma$  contribution. The  $e^+e^-\gamma$  final state entirely accounts for the observed single electron events within errors. In particular, no events are predicted with electron energy above 6 GeV in agreement with our experimental observation.

The observation of no events in the region  $E_e > 6 \text{ GeV}$  and  $|\cos\theta| < 0.75$  corresponds to an upper limit on the cross section for  $e^+e^- \rightarrow e^\pm \tilde{\gamma} \tilde{e}^\mp$  of 0.08

pb. within our acceptance. Extrapolating to full acceptance using the angular dependence of reference 6, this limit is 0.11 pb. An overall detection and analysis efficiency of 95% in this region was used to correct for the losses previously mentioned. The number of single electron events expected is shown in Fig. 3 as a function of the  $\tilde{e}$  mass. At the 95% confidence level  $\tilde{e}$  masses less than 22.4 GeV are excluded.<sup>13</sup> Comparable limits have recently been reported by the Mark II collaboration<sup>14</sup> based upon an analysis similar to the one presented here, and by this group<sup>5</sup> based upon the preliminary results of a search for the reaction  $e^+e^- \rightarrow \tilde{\gamma}\tilde{\gamma}\gamma$ <sup>15</sup> where the experimental signature is a single photon in the final state.

In conclusion, we have measured the energy and angular distribution of single electron events in the MAC detector at PEP. These distributions are consistent with the expected QED process  $e^+e^- \rightarrow e^+e^-\gamma$  where only one electron is observed. No signal from the reaction  $e^+e^- \rightarrow e^\pm\tilde{\gamma}\tilde{e}^\mp$  was found, which allows us to set a new lower limit on the  $\tilde{e}$  mass of 22.4 GeV/ $c^2$  at the 95% confidence level.

One of us (E. F.) thanks B. LeClaire for discussions about the paper by Gaillard et al.<sup>6</sup> We are also grateful to the PEP staff for providing the record luminosities that made this experiment possible.

This work was supported in part by the Department of Energy under contract numbers DE-AC02-81ER40025 (CU), DE-AC03-76SF00515 (SLAC), and DE-AC02-76ER00881 (UW); by the National Science Foundation under contract

numbers NSF-PHY82-15133 (UH), NSF-PHY82-15413 and NSF-PHY82-15414 (NU), and NSF-PHY80-06504 (UU); and by I.N.F.N.



## REFERENCES


- (a) Present address: Mechanical Engineering Department, Stanford University, Stanford, CA 94305
1. J. Wess and B. Zumino, Nucl. Phys. B70, 39 (1974); P. Fayet and S. Ferrara, Phys. Rep. 32C, 249 (1977); P. Fayet, talk at the XVIth Rencontre de Moriond (1981).
  2. G. R. Farrar and P. Fayet, Phys. Lett., 89B, 191 (1980).
  3. JADE Collab., W. Bartel *et al.*, Phys. Lett. 114B, 211 (1982); CELLO Collab., W. Bartel *et al.*, Phys. Lett. 114B, 287 (1982); MARK J Collab., B. Adeva *et al.*, Phys. Lett. 115B, 345 (1982); TASSO Collab., R. Brandelik *et al.*, Phys. Lett. 117B, 365 (1982).
  4. MAC Collab., E. Fernandez *et al.*, SLAC-PUB-3144 (1983), to be published in Phys. Rev. D.
  5. S. Yamada, talk at 1983 Int. Symp. on Lepton and Photon Interactions at High Energies, Cornell Univ., Aug. 1983.
  6. M. K. Gaillard, L. Hall and I. Hinchliffe, Phys. Lett. 116B, 279 (1982)
  7. W. T. Ford in *Proceedings of the International Conference on Instrumentation for Colliding Beams*, edited by W. Ash, SLAC report No. SLAC-250, 1982. Roy Weinstein, in *American Institute of Physics proceeding No. 98, Particles and Fields subseries, No. 29, p. 126* (1982).
  8. B. Gottschalk, Nucl. Instruments and Methods 190, 67 (1981).
  9. In a recent publication, M. Kuroda *et al.*, Phys. Lett. 127B, 467 (1983), have calculated the fraction of the cross section for which the scattered electron has  $|\cos\theta| < 0.8$  for a beam energy of 20 GeV. From their figures we have estimated that this fraction is negligible for our

beam energy and the  $\tilde{e}$  masses of interest.

10. The  $\tilde{\gamma}$  mass and lifetime are very model dependent. The calculation of reference 6 is applicable only if the  $\tilde{\gamma}$  mass is much smaller than the  $\tilde{e}$  mass. If this is the case, most models would give a  $\tilde{\gamma}$  decay length too long to be observed in the detector.
11. F. A. Berends and R. Kleiss, Nucl. Phys. B177, 237 (1981) and Nucl. Phys. B178, 141 (1981).
12. Electromagnetic showers are simulated by EGS, described in R. L. Ford and W. R. Nelson, SLAC Report No. SLAC-210, 1978. Hadronic cascades are simulated by HETC, described in T. W. Armstrong, in *Computer Techniques in Radiation Transport and Dosimetry*, edited by W. R. Nelson and T. M. Jenkins. (Plenum Press, New York, 1980).
13. The calculation of reference 6 assumes that the mass of the supersymmetric partners of the left-handed and right-handed components of the electron are identical. If only one of them has mass less than the center of mass energy the expected number of events will be reduced by a factor of two and the limit would be  $m_{\tilde{e}} > 20.7 \text{ GeV}/c^2$ .
14. L. Gladney *et al.* SLAC PUB-3178, 1983
15. J. Ellis and J. S. Hagelin, Phys. Lett. 122B, 303 (1983).

## FIGURE CAPTIONS


1. Energy distribution of the single electron sample selected as described in the text. The curve is the Monte Carlo prediction for single electron events coming from the  $ee\gamma$  final state. The inset shows the energy distribution of single electrons that would result from the  $e^+e^- \rightarrow e^\pm \tilde{\gamma} \tilde{e}^\mp$  reaction.
2. Angular distribution of the single electron events selected as described in the text. The curve is the Monte Carlo prediction for single electron events coming from the  $ee\gamma$  final state.
3. Number of single electron events with  $|\cos\theta| < 0.75$  expected from the reaction  $e^+e^- \rightarrow e^\pm \tilde{\gamma} \tilde{e}^\mp$  as a function of the selectron mass. An overall detection efficiency of 95% was used.



9-83

4671A1

Fig. 1



9-83

4671A2

Fig. 2




Fig. 3