

EFFECTS OF CHARGED HIGGS IN τ DECAY***Yung Su Tsai**Stanford Linear Accelerator Center,
Stanford University, Stanford, CA 94309**ABSTRACT**

An experiment to test the effect of charged Higgs exchange in τ decay is proposed. It is pointed out that in the decay $\tau^- \rightarrow \nu_\tau + \pi^- + \pi^0$ the effect due to Higgs exchange can be obtained by observing the angular distribution of the π^- in the rest frame of the $\pi^- + \pi^0$. For each value of invariant mass of the $\pi^- + \pi^0$ system, the π^- must have a unique p-wave angular distribution independent of the strong interaction if it is due to W^- exchange alone. Any s-wave interference with this p-wave angular distribution can be attributed to the scalar exchange.

Submitted to *Phys. Rev. D*; also contributed to the
Tau-Charm Factory Workshop, Stanford, CA, May 23-27, 1989.

*Work supported by Department of Energy contract DE-AC03-76SF00515.

1. INTRODUCTION

In the Standard Model there is one neutral Higgs particle and no charged ones. In most other models²⁻⁴ charged Higgs-particles are required. Thus the discovery of effects due to a charged Higgs will demonstrate that the Standard Model-needs to be revised. In this paper we propose an experiment to test the effects of charged Higgs exchange through two π -decay modes of τ . In the Standard Model the decay $\tau^- \rightarrow \nu_\tau + \pi^- + \pi^0$ proceeds via W^- exchange as shown in Fig. 1(a). If a charged Higgs exists, it can also proceed via H^- exchange as shown in Fig. 1(b).

This channel was chosen for the following reasons:

1. It has a relatively simple structure to be analyzed experimentally; both π^- and π^0 ($\rightarrow 2\gamma$) can be identified and momentum analyzed.
2. $\pi^- + \pi^0$ state via W^- exchange can be either $J^P = 1^-$ or 0^+ . But the state $J^P = 0^-$ is forbidden because the vector current is conserved.⁵ Thus the existence of a 0^+ state for $\pi^- + \pi^0$ from τ^- decay must go through something other than W^- . The $\pi^- \pi^0 W^-$ and $\pi^- \pi^0 H^-$ -vertex form factors are functions of $q^2 = (q_1 + q_2)^2$ only. Thus in the rest frame of $q_1 + q_2$ for a given value of q^2 , the p -wave due to W^- exchange has a unique angular distribution:

$$\cos^2 \theta + \frac{q^2}{m_\tau^2 - q^2} \quad , \quad (1.1)$$

where θ is the angle between π^- and τ^- in the rest frame of $q_1 + q_2$. This distribution is independent of form factors coming from the strong interaction. The effects of adding the Higgs exchange is to add an s-wave term $a(q^2)$ and the interference term $b(q^2) \cos \theta$ to Eq. (1.1); thus Eq. (1.1) is changed into:

$$\cos^2 \theta + \frac{q^2}{m_\tau^2 - q^2} + a(q^2) + b(q^2) \cos \theta \quad . \quad (1.2)$$

The functions $a(q^2)$ and $b(q^2)$ depend upon the details of the strong interaction, as well as the Higgs mass m_H and the coupling constants of Higgs to $\tau\nu_\tau$ and to

the quarks. The $a(q^2)$ is probably too small to be observable because it comes from the square of the Higgs exchange term [see Fig. 1(b)]. With the proposed Tau-Charm Factory,⁹ one is supposed to get more than 10^7 τ events, and thus one should be able to observe $b(q^2)$ even if it is as small as 10^{-2} .

CALCULATIONS

Let us denote the matrix element for the W exchange diagram, Fig. 1(a), by:

$$M_W = \bar{u}(p_2) \not{p}(1 - \gamma_5) u(p_1) F_W(q^2) , \quad (2.1)$$

where $p = q_1 - q_2$, $q = q_1 + q_2$ and $F_W(q^2)$ is a function of q^2 representing the effect due to the W propagator and the p-wave $\pi\pi$ form factor. The matrix element for the Higgs exchange, Fig. 1(b), can be written as:

$$M_H = \bar{u}(p_2) u(p_1) F_H(q^2) , \quad (2.2)$$

where $F_H(q^2)$ is a function of q^2 representing the effect due to the Higgs propagator and the s-wave $\pi\pi$ form factor. The decay rate can then be written as:

$$\begin{aligned} \Gamma (\tau^- \rightarrow \nu_\tau + \pi^- + \pi^0) &= \frac{1}{2m_\tau} \frac{1}{(2\pi)^5} \int \frac{d^3 p_2}{2E_2} \int \frac{d^3 q_1}{2\omega_1} \int \frac{d^3 q_2}{2\omega_2} \\ &\times \delta^4(p_1 - p_2 - q_1 - q_2) |M_W + M_H|^2 , \end{aligned} \quad (2.3)$$

where

$$|M_W + M_H|^2 = |F_W|^2 F_{11} + |F_H|^2 F_{00} + 2Re(F_W^* F_H) F_{10} , \quad (2.4)$$

$$\begin{aligned} F_{11} &= \frac{1}{2} Tr(\not{p}_1 + m_\tau) (1 + \gamma_5) \not{p} \not{p}_2 \not{p} (1 - \gamma_5) \\ &= 2 (q^2 - 4m_\pi^2) \frac{(m_\tau^2 - q^2)^2}{q^2} \left[\cos^2 \theta + \frac{q^2}{m_\tau^2 - q^2} \right] , \end{aligned} \quad (2.5)$$

$$F_{00} = \frac{1}{2} Tr(\not{p}_1 + m_\tau) \not{p}_2$$

$$= m_\tau^2 - q^2 \quad , \quad (2.6)$$

$$\begin{aligned} F_{10} &= \frac{1}{2} \text{Tr} (\not{\mathbf{p}}_1 + m_\tau) (1 + \gamma_5) \not{\mathbf{p}} \not{\mathbf{p}}_2 \\ &= -m_\tau \sqrt{q^2 - 4m_\pi^2} \frac{(m_\tau^2 - q^2)}{\sqrt{q^2}} \cos \theta \quad , \end{aligned} \quad (2.7)$$

where θ is the angle between \mathbf{p}_1 and \mathbf{q}_1 in the rest frame of $q = q_1 + q_2$. F_{11} , F_{00} and F_{10} are independent of the detail of strong interaction; they come only from properties of angular momenta of the $\pi^- \pi^0$ system.

From Eqs. (2.4) through (2.7) we see that $a(q^2)$ and $b(q^2)$ of Eq. (1.2) are given by:

$$a(q^2) = \frac{q^2 |F_H|^2}{2 (q^2 - 4m_\pi^2) (m_\tau^2 - q^2) |F_W|^2} \quad , \quad (2.8)$$

and

$$b(q^2) = \frac{-2 \text{Re}(F_W^* F_H) m_\tau \sqrt{q^2}}{\sqrt{q^2 - 4m_\pi^2} (m_\tau^2 - q^2) |F_W|^2} \quad . \quad (2.9)$$

Now the functions F_H and F_W are dependent on the strong interactions and the detail of the Higgs mechanism. The $a(q^2)$ is probably too small to be observable because it is proportional to $|F_H|^2 / |F_W|^2$. The $b(q^2)$ is proportional to F_H / F_W , so let us concentrate on this term. Equation (2.9) shows that $b(q^2)$ is large when $q^2 \rightarrow m_\tau^2$. Since the $\pi^+ \pi^0$ system is charged, it cannot be in $I = 0$ state. Also, the s-wave $\pi^+ \pi^0$ system cannot be in $I = 1$ state because it must obey Bose statistics; thus it must have $I = 2$. We do not see any resonance having these quantum numbers, so we may assume that F_H is mostly real. F_W is dominated by ρ . At the peak of the resonance, F_W is purely imaginary; hence, $b(q^2)$ is small at the ρ peak.

In order to see how $b(q^2)$ may behave, let us assume that F_W is dominated⁶ by ρ , and F_H is a simple H exchange with a reasonable form factor assumed for the $\pi\pi$ system:

$$F_W = g_{\tau\nu\rho} g_{\rho\pi\pi} \frac{1}{(q_1 + q_2)^2 - m_\rho^2 + i\Gamma_\rho m_\rho} \quad , \quad (2.10)$$

where $g_{\rho\pi\pi}$ is related⁶ to the ρ width by:

$$\Gamma_{\rho}(q^2) = \frac{g_{\rho\pi\pi}^2}{48\pi} \frac{q^3}{m_{\rho}^2} \left(1 - \frac{4m_{\pi}^2}{q^2}\right)^{3/2} . \quad (2.11)$$

From

$$\Gamma_{\rho} [m_{\rho}^2 = (0.77 \text{ GeV})^2] = 0.153 \text{ GeV} ,$$

we have

$$g_{\rho\pi\pi} = 6.088 . \quad (2.12)$$

The $g_{\tau\nu\rho}$ can be obtained either. from $e^+e^- \rightarrow \rho$ using *CVC*, or from the partial width of $\tau \rightarrow \nu_{\tau} + \rho$.

The partial decay width of $\tau \rightarrow \nu_{\tau} + \rho$ is given by⁷:

$$\begin{aligned} \Gamma(\tau \rightarrow \rho^- + \nu) &= \frac{g_{\tau\nu\rho}^2}{8\pi} \frac{m_{\tau}^3}{m_{\rho}^2} \left(1 - \frac{m_{\rho}^2}{m_{\tau}^2}\right)^2 \left(1 + \frac{2m_{\rho}^2}{m_{\tau}^2}\right)^2 , \quad (2.13) \\ &= \Gamma(\tau \rightarrow \text{all}) B(\tau \rightarrow \rho^- + \nu) \\ &= \frac{6.582 \times 10^{-25} \text{ GeV}}{3.04 \times 10^{-13}} \times 0.223 \\ &= 0.4828 \times 10^{-12} \text{ GeV} . \end{aligned}$$

From which we obtain:

$$g_{\tau\nu\rho} = 1.13 \times 10^{-6} . \quad (2.14)$$

Combining Eqs. (2.12) and (2.14) we obtain:

$$g_{\tau\nu\rho} g_{\rho\pi\pi} = 6.88 \times 10^{-6} , \quad (2.15)$$

which is very close to the value given by *CVC* and the vector dominance,*

$$g_{\tau\nu\rho} g_{\rho\pi\pi} = m_{\rho}^2 G_F \cos\theta_c = 6.73 \times 10^{-6} . \quad (2.16)$$

The form factor for the Higgs, F_H , is highly model-dependent and almost nothing about it is known. In the Standard Model, the Higgs-particle is coupled to the lepton or quark with coupling constant proportional to the mass of the lepton or quark:

$$m_i = 2^{3/4} \sqrt{G_f} \ , \quad (2.17)$$

where m_i is the mass of the lepton or quark to which the Higgs is coupled, and $G_F = 1.166 \times 10^{-5} \text{ GeV}^{-2}$ is the Fermi coupling constant. In some non-standard theory,^{2,3,4} the Higgs coupling can be proportional to the mass of the heaviest lepton, m_τ , or the heaviest quark, m_t , depending upon whether it is coupled to leptons or quarks. In this case, the Higgs-particle does not necessarily couple more favorably to heavier particles, and $F_H(q^2)$ can be written as:

$$F_H(q^2) = \mathcal{O} \left(\frac{G_F m_\tau m_t m_s}{m_H^2} \right) \ , \quad (2.18)$$

where the Higgs mass comes from the propagator of the Higgs-particle, m_s represents some strong interaction mass in order to make $a(q^2)$ and $b(q^2)$ in Eqs. (2.8) and (2.9) dimensionless. From Eqs. (2.10), (2.16) and (2.18), we see that $b(q^2) = \mathcal{O}(m_\tau m_t / m_H^2)$ and $a(q^2) = \mathcal{O}(m_\tau^2 m_t^2 / m_H^4)$. Since $\cos \theta$ in Eq. (1.2) integrates to zero, $b(q^2)$ does not contribute to the total rate of $\tau^- \rightarrow \nu_\tau + \pi^- + \pi^0$. A limit on the magnitude of $a(q^2)$ can be obtained by comparing the experimental rate of $\tau^- \rightarrow \nu_\tau + \pi^- + \pi^0$ and the prediction of CVC. The difference could be as large as 4%. However, the rate of $\tau^- \rightarrow \nu_\tau + \pi^- + \pi^0$ is dominated by the ρ resonance peak where $a(q^2)$ is small. Thus, in the nonresonance region, $a(q^2)$ could be as large as 10% and $b(q^2)$ could be as large as 30%.

In conclusion, the proposed Tau-Charm Factory⁹ can be used to investigate the detailed structure of the charged Higgs effect using the decay mode $\tau^- \rightarrow \nu_\tau + \pi^- + \pi^0$ described above.

REFERENCES

1. S. Weinberg, Phys. Rev. Lett. 19, 1264 (1967);
A. Salam, "Elementary Particle Theory: Relativistic Groups and Analyticity," (Nobel Symposium No. S), N. Svartholm, ed. (Amquist and Wiksale, Stockholm, 1968), p. 367.
2. E. Derman, Phys. Lett. **78B**, **497** (1978).
3. J. Kim and G. Segre, Phys. Lett. **78B**, 75 (1978).
4. J. F. Donoghue and L. F. Li, Phys. Rev. **19D**, **945** (1979).
5. Y. S. Tsai, Phys. Rev. **4D**, 2821 (1971).
6. Eq. (2.14) of Ref. 5.
7. Eq. (2.8) of Ref. 5.
8. Eqs. (2.9) and (2.10) of Ref. 5.
9. Proc. of Tau-Charm Factory Workshop (1989), Perl et al., eds. (Stanford Linear Accelerator Center, Stanford, CA, 1989).





Fig. 1. Feynman diagrams for $\tau^- \rightarrow \nu_\tau + \pi^- + \pi^0$ via W^- and H^- exchanges.