SLAC-PUB-5403 **DESY 91–001** HEN - 334 December 1990 (T/E)

Measurement of the branching ratios for the decays $\tau \rightarrow \text{hadron } \pi^{\circ}\nu \text{ and } \tau \rightarrow \text{hadron } \pi^{\circ}\pi^{\circ}\nu.$

D. Antreasyan⁹, H.W. Bartels⁵, D. Besset¹¹, Ch. Bieler⁸, J.K. Bienlein⁵, A. Bizzeti⁷, E.D. Bloom¹², I. Brock³, K. Brockmüller⁵, R. Cabenda¹¹, A. Cartacci⁷, M. Cavalli-Sforza², R. Clare¹², A. Compagnucci⁷, G. Conforto⁷, S. Cooper^{a,12}, R. Cowan¹¹, D. Coyne², A. Engler³, K. Fairfield¹², G. Folger⁶. A. Fridman^{b,12}, J. Gaiser¹², D. Gelphman¹², G. Glaser⁶, G. Godfrey¹², K. Graaf⁸, F.H. Heimlich^{7,8}, F.H. Heinsius⁸, R. Hofstadter^{†,12}, J. Irion⁹, Z. Jakubowski⁴, H. Janssen^{c,10}, K. Karch⁵, S. Keh¹³, T. Kiel⁸, H. Kilian¹³, I. Kirkbride¹², T. Kloiber⁵, M. Kobel⁶, W. Koch⁵, A.C. König¹⁰, K. Königsmann^{c,13}, R.W. Krämer³, S. Krüger⁸, G. Landi⁷, R. Lee¹², S. Leffler¹², R. Lekebusch⁸ A.M. Litke¹², W. Lockman¹², S. Lowe¹², B. Lurz⁶, D. Marlow³, H. Marsiske^{5,12}, W. Maschmann⁸, P. McBride⁹, F. Messing³, W.J. Metzger¹⁰, H. Meyer⁵, B. Monteleoni⁷, B. Muryn⁴, R. Nernst⁸, B. Niczyporuk¹², G. Nowak⁴, C. Peck¹, P.G. Pelfer⁷, B. Pollock¹², F.C. Porter¹, D. Prindle³, P. Ratoff¹, M. Reidenbach¹⁰ B. Renger³, C. Rippich³, M. Scheer¹³, P. Schmitt¹³, J. Schotanus¹⁰, J. Schütte⁶, A. Schwarz¹², D. Sievers⁸, T. Skwarnicki⁵, V. Stock⁸, K. Strauch⁹, U. Strohbusch⁸, J. Tompkins¹², H.J. Trost⁵, B. van Uitert¹², R.T. Van de Walle¹⁰, H. Vogel³, A. Voigt⁵, U. Volland⁶, K. Wachs⁵, K. Wacker¹², W. Walk¹⁰, H. Wegener⁶, D.A. Williams^{2,9}, P. Zschorsch⁵

THE CRYSTAL BALL COLLABORATION

¹ California Institute of Technology^d, Pasadena, CA 91125, USA

² University of California at Santa Cruz^e, Santa Cruz, CA 95064, USA

³ Carnegie-Mellon University^f, Pittsburgh, PA 15213, USA

⁴ Cracow Institute of Nuclear Physics, PL-30055 Cracow, Poland

⁵ Deutsches Elektronen Synchrotron DESY, D-2000 Hamburg, Germany

⁶ Universität Erlangen-Nürnberg⁹, D-8520 Erlangen, Germany

⁷ INFN and University of Firenze, I-50100 Firenze, Italy

⁸ Universität Hamburg, I. Institut für Experimentalphysik^h, D-2000 Hamburg, Germany ⁹ Harvard Universityⁱ, Cambridge, MA 02138, USA

¹⁰ University of Nijmegen and NIKHEF^k, NL-6525 ED Nijmegen, The Netherlands ¹¹ Princeton University¹, Princeton, NJ 08544, USA

¹² Department of Physics^m, HEPL, and Stanford Linear Accelerator Centerⁿ,

Stanford University, Stanford, CA 94305, USA

¹³ Universität Würzburg^o, D-8700 Würzburg, Germany

Abstract

The Crystal Ball detector at the DORIS II storage ring at DESY has been used to measure the branching ratios for the decay modes $\tau^{\pm} \rightarrow h^{\pm} \pi^{\circ} \nu$ and $\tau^{\pm} \rightarrow h^{\pm} \pi^{\circ} \pi^{\circ} \nu$, where h^{\pm} is any charged hadron. The results are

$$BR(\tau^{\pm} \to h^{\pm} \pi^{\circ} \nu) = (22.0 \pm 0.8 \pm 1.9)\%$$

$$BR(\tau^{\pm} \to h^{\pm} \pi^{\circ} \pi^{\circ} \nu) = (5.7 \pm 0.5 \pm \frac{1.7}{1.0})\%$$

The first result is in good agreement with the present world average. The decay mode $h^{\pm}\pi^{\circ}\pi^{\circ}\nu$ is reconstructed in τ decays for the first time. Its branching ratio, however, is somewhat lower than the corresponding world average, and therefore tends to increase the 1-prong problem.

(Submitted to Physics Letter B)

t) Deceased

^{a)} Present Address: Max-Planck-Institut für Physik, D-8000 München 40, Germany

b) Permanent Address: DPHPE, Centre d'Etudes Nucléaires de Saclay, F-91191 Gif sur Yvette, France

c) Present Adress: CERN, CH-1211 Geneva 23, Switzerland

d) Supported by the U.S. Department of Energy, contract No. DE-AC03-81ER40050

and by the National Science Foundation, grant No. PHY75-22980

e) Supported by the National Science Foundation, grant No. PHY85-12145

f) Supported by the U.S. Department of Energy, contract No. DE-AC02-76ER03066

g Supported by the German Bundesministerium für Forschung und Technologie, contract No. 054 ER 11P(5)

h) Supported by the German Bundesministerium für Forschung und Technologie, contract No. 054 HH 11P(7)

and by the Deutsche Forschungsgemeinschaft

ⁱ) Supported by the U.S. Department of Energy, contract No. DE-AC02-76ER03064

k) Supported by FOM-NWO

¹) Supported by the U.S. Department of Energy, contract No. DE-AC02-76ER03072

and by the National Science Foundation, grant No. PHY82-08761

^m) Supported by the U.S. Department of Energy, contract No. DE-AC03-76SF00326

and by the National Science Foundation, grant No. PHY81-07396

ⁿ⁾ Supported by the U.S. Department of Energy, contract No. DE-AC03-76SF00515

o) Supported by the German Bundesministerium für Forschung und Technologie, contract No. 054 WU 11P(1)

1. Introduction

Since the discovery of the τ lepton¹ in $e^+e^- \rightarrow \tau^+\tau^-$, a large number of groups at various laboratories have investigated its properties.² All evidence available to date supports the validity of the standard model and the treatment of the τ lepton as the sequential partner of the muon and electron.

The 1-prong decays (the decays to a single charged particle) of the τ , however, still present puzzling aspects. The measured exclusive branching ratios agree well with their predictions. However, their sum is smaller than the measured inclusive 1-prong branching ratio.³⁻⁷ It is not clear whether this discrepancy indicates as yet undetected decays or is simply due to poor measurements, in particular of the modes involving several neutral particles.

The Crystal Ball detector is an excellent apparatus to detect photons (and e^{\pm}) with high efficiency and to determine their energy and direction with high precision. Therefore it is well suited to investigate τ decay modes involving π° 's by explicit reconstruction of the latter. In this paper we present the results of a high statistics analysis of the τ decay modes $\tau^{\pm} \rightarrow h^{\pm}\pi^{\circ}\nu$ and $\tau^{\pm} \rightarrow h^{\pm}\pi^{\circ}\pi^{\circ}\nu$, where h^{\pm} is a charged hadron.

2. Detector

The Crystal Ball detector has been described in detail elsewhere^{8,9} and its properties are only briefly summarized here. It is a nonmagnetic calorimeter designed to measure precisely the energies and directions of electromagnetically showering particles. The main part of the detector consists of a spherical shell of 672 NaI(Tl) crystals covering 93% of 4π sr. The length of each crystal corresponds to about 16 radiation lengths and to about 1 nuclear interaction length. An additional 5% of the solid angle is covered by endcaps, consisting of 40 NaI(Tl) crystals; these endcaps, however, do not allow as accurate a measurement of the energy and direction of a particle and are only used to veto events having particles outside the main detector.

For electromagnetically showering particles the measurement of energy and direction is made using thirteen contiguous crystals in the main detector. This procedure yields an energy resolution given by $\sigma_E/E = (2.7 \pm 0.2)\%/\sqrt[4]{E/GeV}$. For such particles the angular resolution in the polar angle^{*} with respect to the beam axis is $\sigma_{\theta} = 2^{\circ}$ to 3°, the resolution improving as the energy increases.

^{*}The z axis is in the direction of the e^+ beam, the origin is at the center of the overlapping beam bunches, and θ is the angle with respect to the z axis.

Proportional tube chambers surrounding the beam pipe detect charged particles. Depending on the run period, the chambers consisted of three or four double layers of tubes. The inner (outer) layer covers 98% (78%) of 4π sr. Charge division readout allows a determination of the z-position to $\sigma = 1.5\%$ of the length of the tube, *i.e.*, from 9.8 mm for the inner layer to 5.5 mm for the outer layer.

Photons, electrons and positrons yield a rather symmetric lateral energy deposition pattern in the NaI(Tl), with typically 70% of the energy in one crystal and about 98% within a group of thirteen contiguous crystals. A charged particle is identified by a track in the tube chambers that is correlated with an energy deposition in the calorimeter. Muons and charged hadronic particles that do not undergo a strong interaction deposit energy by ionization only. Minimum ionizing particles typically deposit about 200 MeV in one or two crystals. If an energetic charged hadron interacts strongly while traversing the ball, the deposited energy is in general much larger than 200 MeV and the pattern of the hadronic shower is very irregular compared to that of an electromagnetic shower.

3. Monte Carlo Simulation

We made a Monte Carlo simulation of $\tau\tau$ events to determine the efficiencies of our selection criteria. The KORALB program¹⁰ was used to generate $e^-e^+ \rightarrow \tau^-\tau^+$ events. This program includes radiative corrections up to $O(\alpha^3)$ and effects due to spin correlations between the τ 's. The τ 's are decayed by the package TAUOLA,¹¹ which includes all the standard τ decay modes. It has been found that the decay of the τ into two (or three) pions proceeds dominantly via ρ (or a_1) resonance production.^{4,12} Resonance production is therefore assumed in the Monte Carlo simulation. Our selection criteria, however, are chosen such that the efficiency does not strongly depend on this assumption.

The generated events are passed through a complete detector simulation including the following features: (a) Electromagnetically showering particles are simulated by the program EGS 3.¹³ (b) Hadronic interactions in the detector are simulated by an improved version of the GHEISHA 6 program.^{14,15} (c) Extra energy deposited in the crystals by beam-related background is taken into account by adding special background events to Monte Carlo events. These background events are obtained by triggering on one in every 10⁷ beam crossings, with no other condition imposed. (d) Events are reconstructed using our standard software and subjected to the same cuts as the data.

4. Data Sample

The data used in this analysis were collected at the DORIS II storage ring at DESY in the center-of-mass energy $(E_{\rm CM})$ range from 9.4 to 10.6 GeV. The data sample corresponds to an integrated luminosity of 256 pb⁻¹ and a corresponding number of produced $e^+e^- \rightarrow \tau^+\tau^-$ events of $N_{\tau} = (267\pm11)\times10^3$. The luminosity is measured⁹ using large angle Bhabha scattering events with an accuracy better than 3%. The number of produced $\tau\tau$ pairs is calculated using the measured luminosity and the radiatively corrected QED cross section¹⁰ for $e^+e^- \rightarrow \tau^+\tau^-$ in the above energy range. For the data taken on the $\Upsilon(1S)$ and $\Upsilon(2S)$ resonances, the additional contribution from $\Upsilon \rightarrow \tau\tau$ is calculated from the number of Υ 's produced and their leptonic branching ratios.

5. Analysis Method

To identify $e^+e^- \rightarrow \tau^+\tau^-$ events we require that one of the τ 's decays to one charged particle and nothing else visible: $\tau \rightarrow e\nu\overline{\nu}$, $\tau \rightarrow \mu\nu\overline{\nu}$ or $\tau \rightarrow h\nu$, where his a charged π or K. We refer to the charged particle as the 'tag'. We are looking for events where the other τ decays into $h^{\pm}\pi^{\circ}\nu$ or $h^{\pm}\pi^{\circ}\pi^{\circ}\nu$.[†] Therefore, we select low multiplicity events with a jet-jet topology: a single charged particle on one side, and a charged particle plus some associated photons on the other side of the event.

In our experiment we do not measure the π^{\pm} energy; therefore, it is impossible to reconstruct the ρ and a_1 resonances. The determination of the branching ratios for $\tau^{\pm} \rightarrow h^{\pm} \pi^{\circ} \nu$ and $\tau^{\pm} \rightarrow h^{\pm} \pi^{\circ} \pi^{\circ} \nu$ is performed by counting the τ decays with one or two π° 's. We account for feeddown from other $\tau\tau$ channels into the channels of interest, e.g., feeddown of $\tau\tau \rightarrow \rho\nu$, $\rho\overline{\nu}$ into $\tau\tau \rightarrow tag$, $a_1\nu$; and of $\tau\tau \rightarrow tag$, $a_1\nu$ into $\tau\tau \rightarrow tag$, $\rho\nu$. The latter process occurs if one of the π° 's from the a_1 decay misses the main detector or when a π^{\pm} and π° from a highly boosted ρ (from $a_1^{\pm} \rightarrow \rho^{\pm} \pi^{\circ}$) are not resolved by the detector.

A π° is reconstructed from its two decay photons. For low-momentum π° 's the two photons have a large enough opening angle that their energy depositions in the calorimeter are well separated. Such π° 's are reconstructed from the invariant mass of two well-measured photons; we refer to this as method (1).

For π° energies above 500 MeV, it becomes increasingly likely that the photon

[†]We use ' h^{\pm} ' to denote a charged hadron. Charged pions and kaons cannot be distinguished in our experiment. Most of the charged hadrons in τ decay are pions, a small contribution from kaons, however, is expected.⁷

showers overlap, giving a single elongated energy deposition. Such 'merged' π° 's are identified by method (2), which is an analysis of the shape of the energy deposition, yielding the invariant mass and the direction cosines of the parent π° . This method¹⁶ makes use of the second moment, S, of the energy distribution, which is calculated using $G = \frac{1}{2} \sum F(G = f(x))^2$ (1)

$$S = \frac{1}{E} \sum_{i} E_{i} \left(\hat{n}_{i} - \langle \hat{n} \rangle \right)^{2} \tag{1}$$

where E_i is the energy deposited in crystal *i* and \hat{n}_i is the unit vector formed by the direction cosines of crystal *i*. The sum runs over all the crystals in the energy cluster under consideration; $E = \sum_i E_i$ is the total energy of the cluster; and $\langle \hat{n} \rangle = \sum_i E_i \hat{n}_i / E$ is the direction of its center-of-gravity. The invariant mass M of the shower is given by $M^2 = E^2 - E^2 \langle \hat{n} \rangle^2 = E^2 S$. Using Monte Carlo simulations it was found that an estimate for the real invariant mass (also called shower mass) of particle X, decaying into two photons, can be calculated using the expression

$$M_X^2 = E^2(S - S_\gamma) \tag{2}$$

where $S_{\gamma} = 0.004$ is the average second moment of a single photon shower.

In the search for $\tau^{\pm} \to h^{\pm} \pi^{\circ} \pi^{\circ} \nu$, the highest detection efficiency is obtained by selecting one π° via method (1) and the other via method (2). This is simply a consequence of the energy distribution of the π° 's in this process. In the search for $\tau^{\pm} \to \pi^{\pm} \pi^{\circ} \nu$, the highest efficiency is obtained with method (2). However, use of this method would result in a heavy contamination by events containing a highly energetic photon, e.g., $e^+e^- \to e^+e^-\gamma$, $\mu^+\mu^-\gamma$. Estimation and subtraction of such background is expected to result in a large systematic error. Therefore, we have chosen to use method (1) for this process.

6. General cuts

We start with a set of general cuts to select the desired $\tau\tau$ sample from a large data sample including radiative Bhabha and $\mu^+\mu^-$ events, continuum hadron events, two-photon events, Υ decays and events originating from beam-gas and beam-wall collisions. This first set of cuts is the same for both the $\tau^{\pm} \rightarrow h^{\pm}\pi^{\circ}\nu$ and the $\tau^{\pm} \rightarrow$ $h^{\pm}\pi^{\circ}\pi^{\circ}\nu$ analyses. The cuts below are designed to reduce the above background as much as possible while retaining a maximum acceptance for the channels of interest.

• We require exactly two charged particles in the main detector ($|\cos \theta| < 0.85$). These particles must have a deposited energy of at least 75 MeV, and at most $0.85 \times E_{\text{beam}}$ each. The angle between their directions must be larger than 90°. In addition, at least one of the charged particles must be isolated from neutral particles; that is, there must not be a neutral particle (with an energy of more than 50 MeV) within 60° of the charged particle direction.

- The total energy $(E_{\rm vis})$ measured in the main detector is required to satisfy $2 \,{\rm GeV} < E_{\rm vis} < 0.75 \times E_{\rm CM}$. The lower limit, apart from reducing the machine related background and two photon events, also ensures that the trigger efficiency is 100%. Further, we require the energy deposited in the endcaps to be less than 100 MeV.
- To obtain events having a large energy flow perpendicular to the beam directions, we require that the transverse energy E_T be greater than 1500 MeV. The transverse energy is defined as E_T = ∑_i E_i sin θ_i, where E_i and θ_i are the energy deposited in, and the polar angle of, crystal i, respectively.
- The events must have a jet-jet topology; this is achieved by requiring the second Fox-Wolfram moment¹⁷ to be greater than 0.4. This moment is defined as $H_2 = \sum_{i,j} E_i E_j (3\cos^2 \alpha_{ij} 1)/2(\sum_k E_k)^2$, where E_i is the energy deposition in crystal *i* and α_{ij} is the angle between crystals *i* and *j*.

Further selection criteria depend on the specific τ decay mode under study.

7. τ decays with one π°

We first discuss the selection of $\tau\tau$ events where one of the τ 's decays into $h^{\pm}\pi^{\circ}\nu$. Recall (section 5) that we reconstruct such a π° using two well-separated photons. Therefore, we require two neutral particles that must fulfill the following conditions:

- For their energy (E_{γ_i}) and polar angle (θ_{γ_i}) , we require $E_{\gamma_i} > 50 \text{ MeV}$ and $|\cos \theta_{\gamma_i}| < 0.85$, respectively. Their lateral energy deposition patterns must be consistent with that of electromagnetically showering particles. Furthermore, the sum of their energies, $E_{\gamma_1} + E_{\gamma_2}$, is required to be smaller than 1200 MeV, because it is more likely that a π° above 1200 MeV produces a 'merged' shower in the calorimeter. In addition, we require the angle between the directions of γ_1 and γ_2 to be smaller than 120°.

In addition to these two photons, we allow at most one neutral particle with an energy less than $50 \,\text{MeV}$ in the main detector. We thus allow at most one soft radiative photon. Further, we require:

- The pseudo-invariant mass of the $h^{\pm}\gamma\gamma$ system to be smaller than 1400 MeV (*i.e.*, well below the τ mass). The term 'pseudo' refers to the fact that we do not measure the real h^{\pm} energy. In the calculation of the invariant mass, the energy deposited by the charged hadron is used instead.
- The tagging particle to be a highly energetic electron or positron (as expected from τ → evv); that is, (a) the lateral energy deposition must be consistent with that of an electromagnetically showering particle, and (b) its energy must be greater than 1000 MeV.

Tagging on an e^{\pm} results in events with a typical $\tau\tau$ signature: a lepton on one side, hadrons on the other. We do not lose much efficiency by this cut because most of the events where the tag is a μ^{\pm}, π^{\pm} or K^{\pm} were already (unavoidably) rejected by the lower limit on the visible energy E_{vis} . By requiring an energy deposition of at least 1000 MeV for the tagging particle, we avoid the problem that most of the energy needed to pass the lower limit on E_{vis} comes from the $h^{\pm}\pi^{\circ}$ system. Since the selected photons are relatively soft, the absence of this cut would have biased the events such that the charged pion deposits a lot of energy in the calorimeter. This would have lead to a large systematic error because the uncertainty in the Monte Carlo simulation for the h^{\pm} interaction in the NaI increases with the deposited h^{\pm} energy.

Figure 1 shows the distribution of the invariant mass of the two photons for the events that passed the above cuts. To obtain the number of $\tau\tau$ events with one π° , the signal plus background is fitted with a Gaussian on top of a third order polynomial for the background. The π° mass thus obtained was consistent with the known value of the π° mass, and the width of the distribution with the resolution of our detector. From the fit we find 800 ± 28 events. From a Monte Carlo simulation of continuum hadron $(q\bar{q})$ events,¹⁸ we expect the continuum to contribute 21 ± 14 events to the peak. No evidence for other background not originating from $\tau\tau$ events was found. The number of $\tau\tau$ events with one π° thus becomes 779 ± 31 .

The selected data sample does, however, contain events originating from τ decays involving an a_1 as described in section 5; their contribution is also determined by a Monte Carlo simulation. Significant feeddown comes from the following decay channels: $\tau \tau \rightarrow e \nu \overline{\nu} a_1 \overline{\nu}$, $\pi(K) \nu a_1 \overline{\nu}$, and $\rho \nu a_1 \overline{\nu}$, where the a_1 subsequently decays



Figure 1. The two photon invariant mass distribution from $\tau\tau$ events where one τ decayed into $h\pi^{\circ}\nu$ and the other τ into $e\nu\overline{\nu}$.

into $\rho\pi^{\circ}$. The fractions of such events that pass our cuts for $\tau^{\pm} \rightarrow \rho^{\pm}\nu \rightarrow \pi^{\pm}\pi^{\circ}\nu$ are: $f_{ea_1} = (0.34 \pm 0.04)\%$, $f_{\pi(K)a_1} = (0.07 \pm 0.03)\%$ and $f_{\rho a_1} = (0.04 \pm 0.02)\%$, respectively. Because the actual feeddown is dependent on the branching ratio for $\tau^{\pm} \rightarrow a_1^{\pm}\nu \rightarrow \pi^{\pm}\pi^{\circ}\pi^{\circ}\nu$ the former can only be determined after (or simultaneously with) the determination of the latter.

8. τ decays with two π° 's

We now proceed with the selection of $\tau\tau$ events where one of the τ 's decays to two neutral pions. Recall (section 5) that we reconstruct one of the π° 's using the shower-mass technique and the other π° using two well-separated photons. In addition to the set of general cuts described in sect. 6, we make the following cuts:

- We require exactly three neutral energy clusters, at least one with E > 500 MeV and the other(s) with E > 50 MeV, in the main ball. The cluster with the highest shower mass M and with E > 500 MeV is the 'merged'- π° candidate. The other two clusters are the single-photon candidates.
- In addition to these energy clusters, we allow at most one neutral particle with an energy less than 50 MeV in the main detector (to allow for a radiative

photon).

• The single-photon candidates must have lateral energy deposition patterns consistent with being a single photon. Just as in the selection of events containing only one π° , we require that $E_{\gamma_1} + E_{\gamma_2} < 1200 \text{ MeV}$ and that the angle between the directions of γ_1 and γ_2 be smaller than 120°.

We now introduce some cuts to ensure to a high degree that the π° and both photons originate from the decay of the same τ . These cuts mainly select against feeddown of events of the type $\tau \tau \rightarrow \rho \nu \rho \overline{\nu}$.

- The direction of the total momentum of the neutral particles, $\vec{p}_{neut} = \vec{p}_{\pi^\circ} + \vec{p}_{\gamma_1} + \vec{p}_{\gamma_2}$, must be close to one of the charged particles, *i.e.*, if \vec{n}_{ch} is the direction of a charged particle, then for at least one of the charged particles one must have $(\vec{p}_{neut} \cdot \vec{n}_{ch})/|\vec{p}_{neut}| > 0.2$. In addition, the other charged particle should be at least 120° away from the direction of \vec{p}_{neut} .
- The pseudo-invariant mass of the $h^{\pm}\pi^{\circ}\gamma\gamma$ system must be smaller than 1400 MeV.



Figure 2. The two photon invariant mass $M(\gamma, \gamma)$, vs. the shower mass, M(shower), in $\tau\tau$ events after all the cuts described in the text. Events clustering around the π° mass on both axes come from $\tau^{\pm} \rightarrow h^{\pm}\pi^{\circ}\pi^{\circ}\nu$.



Figure 3. The invariant mass of the two photons in $\tau\tau$ events after one π° has already been identified by the shower mass technique.

We do not make any additional cuts on the tagging particle such as those used in the $\tau^{\pm} \rightarrow h^{\pm} \pi^{\circ} \nu$ analysis. Such cuts are unnecessary here. The bias of the selected events towards high energy depositions from charged pions (as described in the last item of the previous section) does not occur here, because we now have two neutral pions that deposit their total energy in the calorimeter.

In Figure 2 we plot the the invariant mass of the two photons vs. the shower mass of the 'merged'- π° candidate. Clear clustering of $\pi^{\circ}\pi^{\circ}$ events is observed. After a cut on the shower mass (90 MeV < M < 180 MeV), we make a projection on the $M(\gamma, \gamma)$ axis (Figure 3). This $\gamma\gamma$ mass distribution is fitted with a Gaussian π° signal plus a linear polynomial for the background. The π° mass from the fit was consistent with the known value and the width with the resolution of our detector. The number of $\tau\tau$ events with two π° 's thus found is 133 ± 12 events. This number contains a small contribution from the process $\tau\tau \rightarrow \rho\nu\rho\overline{\nu}$: From a Monte Carlo sample of 11000 events only two events survived the selection criteria for $\tau^{\pm} \rightarrow h^{\pm}\pi^{\circ}\pi^{\circ}\nu$. Again, the actual feeddown can only be given after (or simultaneously with) the determination of $BR(\tau^{\pm} \rightarrow h^{\pm}\pi^{\circ}\nu)$. No evidence of other background to the peak was found.

$\tau^{\pm} \rightarrow \pi^{\pm} \pi^{\circ} \nu$	efficiencies	$\tau^{\pm} \rightarrow \pi^{\pm} \pi^{\circ} \pi^{\circ} \nu$	efficiencies
$\epsilon_{e ho}$	$3.36 \pm 0.09~\%$	ϵ_{ea_1}	$1.05 \pm 0.07~\%$
$\epsilon_{\mu ho}$	< 0.01%	$\epsilon_{\mu a_1}$	$0.72 \pm 0.09~\%$
$\epsilon_{\pi(K) ho}$	$0.19 \pm 0.03~\%$	$\epsilon_{\pi(K)a_1}$	$0.79 \pm 0.08~\%$
$\epsilon_{ ho ho}$	$0.25 \pm 0.05 \ \%$	$\epsilon_{ ho a_1}$	$0.10 \pm 0.03 ~\%$

Table 1. Efficiencies of the selection criteria for $\tau^{\pm} \rightarrow h^{\pm}\pi^{\circ}\nu$ and $\tau^{\pm} \rightarrow h^{\pm}\pi^{\circ}\pi^{\circ}\nu$ of the contributing $\tau\tau$ channels. The error given is due to limited Monte Carlo statistics.

9. Results

If $N(\pi^{\circ})$ is the number of observed $\tau\tau$ events with one π° , the branching ratio for $\tau^{\pm} \rightarrow h^{\pm}\pi^{\circ}\nu$ is given by

$$BR(\tau^{\pm} \to h^{\pm} \pi^{\circ} \nu) = \frac{N(\pi^{\circ}) - N^{\log}(\pi^{\circ})}{2 \left[\epsilon_{\iotaag,\rho} + \frac{1}{2} \epsilon_{\rho\rho} \cdot BR(\tau^{\pm} \to h^{\pm} \pi^{\circ} \nu) \right] N_{\tau}}$$
(3)

where $N^{\text{bg}}(\pi^{\circ})$ is the feeddown from $\tau \tau \to \text{tag}, a_1 \nu$ events (section 7), *i.e.*,

$$N^{\mathrm{bg}}(\pi^{\mathrm{o}}) = 2 \left[BR(\tau \to e\nu\overline{\nu}) \cdot f_{ea_{1}} + BR(\tau \to \pi(K)\nu) \cdot f_{\pi(K)a_{1}} + BR(\tau^{\pm} \to h^{\pm}\pi^{\mathrm{o}}\nu) \cdot f_{\rho a_{1}} \right] BR(\tau^{\pm} \to h^{\pm}\pi^{\mathrm{o}}\pi^{\mathrm{o}}\nu) N_{\tau} , \qquad (4)$$

and

$$\epsilon_{\text{tag},\rho} = BR(\tau \to e\nu\overline{\nu}) \cdot \epsilon_{e\rho} + BR(\tau \to \mu\nu\overline{\nu}) \cdot \epsilon_{\mu\rho} + BR(\tau \to \pi(K)\nu) \cdot \epsilon_{\pi(K)\rho}$$

= (0.62 ± 0.02)%

 $N_{\tau} = (267 \pm 11) \times 10^3$ is the total number of $\tau \tau$ events in our full data sample. For the branching ratios of the tagging particles we use the Particle Data Group values,⁷ $BR(\tau \rightarrow e\nu\overline{\nu}) = (17.7 \pm 0.4)\%$, $BR(\tau \rightarrow \mu\nu\overline{\nu}) = (17.8 \pm 0.4)\%$ and $BR(\tau \rightarrow \pi(K)\nu) = (11.7 \pm 0.5)\%$. The selection efficiencies $\epsilon_{\text{tag}_i,\rho}$ and $\epsilon_{\text{tag}_i,a_1}$ for the $\tau\tau$ channels contributing to our selected data samples are shown in Table 1. The subscripts ' ρ ' and ' a_1 ' of the efficiencies ϵ above are used to remind the reader that ρ and a_1 dominance is assumed in the simulated τ decays. The efficiencies are defined in the following way: e.g., $\epsilon_{e\rho}$ is the ratio of $\tau\tau$ Monte Carlo events (one τ decaying into $e\nu\overline{\nu}$ and the other into $\rho\nu$) accepted by our selection criteria to the total number of such events.

The branching ratio for $\tau^{\pm} \to h^{\pm} \pi^{\circ} \pi^{\circ} \nu$ is given by

$$BR(\tau^{\pm} \to h^{\pm} \pi^{\circ} \pi^{\circ} \nu) = \frac{N(\pi^{\circ} \pi^{\circ}) - N^{\mathrm{bg}}(\pi^{\circ} \pi^{\circ})}{2 \epsilon_{\mathrm{tag},a_{1}} N_{\tau}}$$
(5)

where $N(\pi^{\circ}\pi^{\circ})$ is the number of $\tau\tau$ events that passed our selection criteria, $N^{\text{bg}}(\pi^{\circ}\pi^{\circ})$ is the feeddown from $\tau\tau \to \rho\nu\rho\overline{\nu}$ (section 8),

$$N^{\rm bg}(\pi^{\circ}\pi^{\circ}) = \frac{2}{11000} \left[BR(\tau^{\pm} \to h^{\pm}\pi^{\circ}\nu) \right]^2 N_{\tau} , \qquad (6)$$

and

$$\begin{aligned} \epsilon_{\text{tag},a_1} &= BR(\tau \to e\nu\overline{\nu}) \cdot \epsilon_{ea_1} + BR(\tau \to \mu\nu\overline{\nu}) \cdot \epsilon_{\mu a_1} \\ &+ BR(\tau \to \pi(K)\nu) \cdot \epsilon_{\pi(K)a_1} + BR(\tau^{\pm} \to h^{\pm}\pi^{\circ}\nu) \cdot \epsilon_{\rho a_1} \;. \end{aligned}$$

Eqs (3) – (6) are solved. The result for the feeddown is $N^{\text{bg}}(\pi^{\circ}) = 24 \pm 5$ and $N^{\text{bg}}(\pi^{\circ}\pi^{\circ}) = 2.3 \pm 1.5$. For the branching ratios we obtain $BR(\tau^{\pm} \rightarrow h^{\pm}\pi^{\circ}\nu) = (22.0 \pm 0.8 \pm 1.2)\%$ and $BR(\tau^{\pm} \rightarrow h^{\pm}\pi^{\circ}\pi^{\circ}\nu) = (5.7 \pm 0.5 \pm 0.5)\%$. The systematic error includes the error due to limited Monte Carlo statistics, the uncertainty on the total number $\tau\tau$ events (N_{τ}) , and the errors on the branching ratios for the tagging particles e, μ, π or K.

A nice check on the $\tau^{\pm} \to h^{\pm} \pi^{\circ} \pi^{\circ} \nu$ analysis can be performed by making subsamples according to the nature of the tagging particle.[‡] Restricting the tagging particle to a highly energetic e^{\pm} (originating from $\tau \to e\nu\overline{\nu}$), we obtain $BR(\tau^{\pm} \to h^{\pm}\pi^{\circ}\pi^{\circ}\nu) = (5.0 \pm 0.8 \pm 0.5)\%$. By instead requiring the tag to be a mininum ionizing particle (*i.e.*, originating from $\tau \to \mu\nu\overline{\nu}$ or $\tau \to \pi(K)\nu$), we find a consistent value, $BR(\tau^{\pm} \to h^{\pm}\pi^{\circ}\pi^{\circ}\nu) = (5.5 \pm 0.8 \pm 0.7)\%$. The systematic error only includes the sources mentioned in the previous paragraph. These consistent results give us confidence that there is indeed no significant background in our data sample.

The most important additional systematic errors for the $\tau^{\pm} \rightarrow h^{\pm}\pi^{\circ}\nu$ analysis are: (i) An uncertainty in the charged tagging efficiency (*i.e.*, the efficiency that a charged particle is identified as charged by the tube chamber system): 5%; (ii) An uncertainty in the estimation of the background from continuum hadron events $(q\bar{q})$: 2.5%; (iii) An uncertainty arising from variation of the fit interval and the background function (see fig. 1): 2%; (iv) An uncertainty due to variation (within reasonable limits) of the selection criteria: 3%. Adding these systematic errors quadratically, the total error for $BR(\tau^{\pm} \rightarrow h^{\pm}\pi^{\circ}\nu)$, including the ones mentioned above, becomes 8.4%.

The most important additional systematic errors for the $\tau^{\pm} \rightarrow h^{\pm} \pi^{\circ} \pi^{\circ} \nu$ analysis are: Uncertainties of 5%, another 5%, and 3% as described by the items (i) - (iii) above, and 12.6% due to variation of the selection criteria. The latter error is

¹Such a test is not possible for the $\tau^{\pm} \rightarrow h^{\pm} \pi^{\circ} \nu$ analysis because there the tagging particle is required to be an e^{\pm} (cf. sect. 7).

dominated by a small discrepancy between the experimental data and the Monte Carlo simulation of the distribution $\vec{p}_{neut} \cdot \vec{n}_{ch} / |\vec{p}_{neut}|$ (defined in section 8), *i.e.*, an angular distribution of the $\pi^{\pm}\pi^{\circ}\pi^{\circ}$ system. The cuts on this quantity were made rather loose in order not to become too dependent on the specific Monte Carlo model (which assumes a_1 dominance in the $\pi^{\pm}\pi^{\circ}\pi^{\circ}$ system). A maximum difference of 10% was found by using a Monte Carlo sample in which the τ decays directly into $h^{\pm}\pi^{\circ}\pi^{\circ}\nu$. Adding all the systematic errors in quadrature gives a total systematic error for $BR(\tau^{\pm} \to h^{\pm}\pi^{\circ}\pi^{\circ}\nu)$ of 17.1% or an absolute systematic error of 1.0%.

With an efficiency for the $\tau^{\pm} \to \pi^{\pm}\pi^{\circ}\pi^{\circ}\nu$ analysis at the level of 1%, a very good understanding of the acceptance of the detector is needed to achieve an accurate final result. Indeed, uncertainties in the Monte Carlo determination of the acceptance dominate the systematic error.

Earlier preliminary Crystal Ball analyses,^{19,20} using essentially the same data as in the present analysis, found $BR(\tau^{\pm} \rightarrow h^{\pm}\pi^{\circ}\pi^{\circ}\nu) = (7.0 \pm 0.7 \pm 1.4)\%$ and $(7.4 \pm 0.6 \pm 1.3)\%$. The sensitivity of the result to the Monte Carlo details is the main reason that these previous preliminary results have not been published. In the interim, the analysis presented here was performed in conjunction with an extensive reexamination of the Monte Carlo used to calculate the acceptance. These studies resulted in a number of improvements¹⁵ to the Monte Carlo, accounting in part for the noticeable difference in the value of the branching ratio between the previous preliminary analyses and the present one. It must be noted that all three analyses are in good agreement with each other for $BR(\tau^{\pm} \rightarrow h^{\pm}\pi^{\circ}\nu)$.

As described above, an estimate of the absolute systematic error yields $\pm 1\%$. This error does not cover the central values of the previous preliminary analyses. We believe that the best current estimate of the branching ratio from our data is the new value presented in this paper. However, we choose to increase the size of the systematic error to cover the central values of the previous preliminary analyses. This leads to a final absolute systematic error of $\pm \frac{1.7}{1.0}\%$.

10. Conclusions

We have measured the branching ratios for the τ decay modes $\tau^{\pm} \to h^{\pm} \pi^{\circ} \nu$ and $\tau^{\pm} \to h^{\pm} \pi^{\circ} \pi^{\circ} \nu$, where h^{\pm} is a charged hadron. The results are

$$BR(\tau^{\pm} \to h^{\pm} \pi^{\circ} \nu) = (22.0 \pm 0.8 \pm 1.9)\%$$
$$BR(\tau^{\pm} \to h^{\pm} \pi^{\circ} \pi^{\circ} \nu) = (5.7 \pm 0.5 \pm \frac{1.7}{1.0})\%$$

As discussed in section 5, the charged hadron may be a π or K. Subtracting a contribution⁷ of $(0.7 \pm 0.1)\%$ for the branching ratio of the decay $\tau^{\pm} \to K^{*\pm}\nu \to K^{\pm}\pi^{\circ}\nu$, and assuming ρ dominance in the $\pi\pi^{\circ}$ system, we obtain $BR(\tau^{\pm} \to \rho^{\pm}\nu \to \pi^{\pm}\pi^{\circ}\nu) = (21.3 \pm 0.8 \pm 1.9)\%$. For the two- π° mode there is a kaon contribution⁷ of 0.1% due to the decay chain $\tau^{\pm} \to K^{*\pm}\nu \to K_S^{\circ}\pi^{\pm}\nu \to \pi^{\pm}\pi^{\circ}\pi^{\circ}\nu$. Thus, assuming a_1 dominance, $BR(\tau^{\pm} \to a_1^{\pm}\nu \to \pi^{\pm}\pi^{\circ}\pi^{\circ}\nu) = (5.6 \pm 0.5 \pm \frac{1.7}{1.0})\%$. It should be noted that, although we assumed ρ and a_1 dominance in the efficiency determination, our selection criteria are chosen such that the result does not critically depend on this hypothesis.

Our experiment is the first to measure the branching ratio of $\tau^{\pm} \rightarrow \pi^{\pm} \pi^{\circ} \pi^{\circ} \nu$ by reconstruction of both neutral pions from their decay photons.

Our value for $BR(\tau^{\pm} \rightarrow h^{\pm}\pi^{\circ}\nu)$ (and $BR(\tau^{\pm} \rightarrow \rho^{\pm}\nu)$) is in good agreement with other measurements. The result for $BR(\tau^{\pm} \rightarrow \pi^{\pm}\pi^{\circ}\pi^{\circ}\nu)$ is somewhat lower than the present world average⁷ of $(7.5 \pm 0.9)\%$. Furthermore, it is about 1.7 standard deviations below the recent CELLO result²¹ of $(10.0 \pm 1.5 \pm 1.1)\%$ which was among the results essential to the observed absence of the 1-prong problem in their data. The discrepancy between their and our result for this mode suggests that the problem(s) with τ decay modes are not yet solved. In particular, our result tends to increase the 1-prong problem.³⁻⁷

Acknowledgments

We would like to thank the DESY and SLAC directorates for their support. This experiment would not have been possible without the dedication of the DORIS machine group as well as the experimental support groups at DESY. The visiting groups thank the DESY laboratory for its hospitality. Z.J., B.N., and G.N. thank DESY for financial support. E.D.B., R.H., and K.S. have benefitted from financial support from the Humboldt Foundation. K. Königsmann acknowledges support from the Heisenberg Foundation.

References

¹SLAC, LBL collab.: M.L.Perl et al., Phys. Rev. Lett. **35** (1975) 1489.

²For a recent review see, for example,

B.C.Barish and R.Stroynowski, Phys. Rep. 157 (1988) 1.

K.K.Gan and M.L.Perl, Int. J. Mod. Phys. A3 (1988) 531.

³T.N.Truong, Phys. Rev. D 30 (1984) 1509.

⁴F.G.Gilman and S.H. Rhie, Phys. Rev. D 31 (1985) 1066.

⁵H.J.Gerber, Proc. International Europhysics Conference on High Energy Physics, Uppsala, 1987, p.937.

⁶K.G.Hayes and M.L.Perl, Phys. Rev. D 38 (1988) 3351.

⁷Particle Data Group: Review of Particle Properties, Phys. Lett. B 239 (1990) 1.

⁸E.D.Bloom and C.W.Peck, Ann. Rev. Nucl. Part. Sci. 33 (1983) 143; Crystal Ball collab.: D.Antreasyan et al., Phys. Rev. D 36 (1987) 2633.

⁹K.Wachs, Ph.D. thesis, University of Hamburg, 1988;
Crystal Ball collab.: K.Wachs et al., Zeit. Phys.C 42 (1989) 33.

¹⁰S.Jadach and Z.Was, Comp. Phys. Comm. **36** (1985)191.

¹¹Part of the Monte Carlo data was generated with version 1.4 of TAUOLA, another part with version 1.5. The main difference between these versions is the inclusion of radiative (QED) corrections to the decay products of the τ . No significant differences however, were observed in the efficiency determination when using one or the other subsample. For TAUOLA 1.5 see,

S.Jadach and Z.Was, preprint CERN-TH-5855;

S.Jadach, J.H. Kühn and Z. Was, preprint CERN-TH-5856

¹²For the ρ resonance, see *e.g.*,

CELLO collab.: H.J.Behrend et al., Zeit. Phys.C 23 (1984) 103.

Mark II collab.: J.M.Yelton et al., Phys. Rev. Lett. 56 (1986) 812;

For the a_1 resonance, see *e.g.*,

ARGUS collab.: H.Albrecht et al., Zeit. Phys.C 33 (1986) 7 and

Phys. Lett. B 250 (1990) 164. See also,

J.H.Kühn and A. Santa Maria, Zeit. Phys.C 48 (1990) 445.

¹³R.Ford and W.Nelson, SLAC-210 (1978).

¹⁴H.Fesefeldt, Aachen preprint, PITHA-85/02.

¹⁵M.Kobel and Z.Jakubowski, 'A verified upgrade of the GHEISHA 6/7 simulation of particle interactions', DESY Report 89-165 (1989) and Nucl.Inst.Meth.A (in press).

¹⁶See also:

Crystal Ball collab.: P.Schmitt et al., Zeit. Phys.C 40 (1988) 199;

R.A.Lee, Ph.D. thesis Stanford University, SLAC-282 (1985);

R.L.Walker, Proceedings of the Summer Institute on Particle Physics, July 29 - August 10, 1974, SLAC-179 (vol. II), 1 (1974).

¹⁷G.C.Fox and S.Wolfram, Phys. Rev. Lett. 41 (1978) 1581.

¹⁸T.Sjöstrand, Comp. Phys. Comm. **39** (1986) 347;
 T.Sjöstrand and M.Bengtsson, Comp. Phys. Comm. **43** (1987) 367.

¹⁹T.Skwarnicki, Search for τ decays to the η meson, DESY 87-166 (1987), SLAC-PUB-4506 (1987).

²⁰S.T.Lowe, Proceedings of the International Symposium on Production and Decay of Heavy Flavors, eds. E.D.Bloom and A.Fridman, Annals of the New York Academy of Sciences Volume 535, 527(1988).

²¹CELLO collab.: H.J.Behrend et al., Zeit. Phys.C 46 (1990) 537.